FULL TITLE ASP Conference Series, Vol. **VOLUME**, c **YEAR OF PUBLICATION** **NAMES OF EDITORS**

A Bayesian Approach Accounting for Stochastic Fluctuations in Stellar Cluster Properties

M.Fouesneau & A.Lancon

O bærvatoire A stronom ique (UM R 7550), U niversite de Strasbourg & CNRS, 11 rue de l'Universite, 67000 Strasbourg, France

A bstract. The integrated spectro-photom etric properties of star clusters are subject to large cluster-to-cluster variations. They are distributed in non trivial ways around the average properties predicted by standard population synthesism odels. This results from the stochastic mass distribution of the nite (sm all) number of lum incus stars in each cluster, stars which may be either particularly blue or particularly red. The color distributions are broad and usually far from G aussian, especially for young and intermediate age clusters, as found in interacting galaxies. When photom etric measurements of clusters are used to estimate ages and masses in conjunction with standard models, biases are to be expected. We present a Bayesian approach that explicitly accounts for stochasticity when estimating ages and masses of star clusters that cannot be resolved into stars. Based on M onte-C arb simulations, we are starting to explore the probability distributions of star cluster properties obtained given a set ofmulti-wavelength photom etric data.

1 Introduction

O bærving star clusters o ers a powerful tool for studies of the form ation history of galaxies. C lusters can be measured out to much further distances than individual eld stars, and they provide complementary information to the integrated light emitted by galaxy elds. Star formation and cluster formation are closely related, although the interplays between clusters and the eld remain to be properly understood. The age distribution of star clusters may be a record of the star formation history of galaxies. Their age and mass distributions are especially important when we wish to understand what happens during collisions or mergers of gas-rich galaxies, where the interaction is well known to trigger the formation of stars.

For clusters that are not resolved into stars, estim ates of ages, m asses and other intrinsic properties are based on integrated photom etry or spectra. The standard approach is to compare these observed properties with synthetic spectra produced by population synthesis codes that assume a continuously populated stellar initial m ass function (IM F).

However the energy distribution of clusters is dom inated by the light of a nite (often small) number of massive stars, whereas the mass is mostly produced by early evolved stars with relatively faint lum incus contributions. As an example, red supergiants are the predom inant source of near-IR light in young clusters (6-60 M yr). These stars are intrinsically rare. As a combined result of the Initial M ass Function (IM F) and the short duration of the relevant evolu-



Figure 1. Stochastic properties of star clusters at solar metallicity. Left: The dots represent photom etric properties of individual clusters containing 10^3 and 10^5 stars each. The solid lines show the corresponding age sequences from \standard" predictions, i.e. rescaled versions of population synthesis predictions strictly valid only for a very large number of stars (gure inspired by B ruzual 2002). R ight: D ensity distribution of models in our catalog, constructed assum ing a power law with a index -2 for the cluster mass function.

tionary phases, we expect on average only one red supergiant for a total cluster mass of 10^4 M for a standard IM F with a lower cut-o of 0.1 M (Lancon et al. 2008). Hence a 10^4 M star cluster may contain zero, one or a few red supergiants, which will result in multi-modal probability distributions for integrated colors and uxes (Chiosi, Bertelli, & Bressan 1988; Lancon & Mouhcine 2000; Cervino & Luridiana 2004; Lancon & Fouesneau 2009). Large spreads result from the above-mentioned small numbers of lum inous stars, the so-called stochastic uctuations (Figure 1). It requires clusters with several 10^6 M to narrow down the uctuations to 5% in the K-band. In this case, colors and uxes follow uni-modal distributions and G aussian approxim ations become to the total cluster of the several to the colors and uxes follow uni-modal distributions and G aussian approximations become to the colors and to the colors and the co

M assive clusters for which the G aussian approximation holds exist in very small numbers, because cluster mass functions tend to fall rapidly with increasing mass (W him one et al. 1999; Bastian et al. 2009). On the other hand, many low and intermediate mass clusters have already been observed with space-based and large ground-based telescopes. It is thus very important to develop methods that explicitly deal with the stochastic uctuations.

2 E stim ates of Star C luster P roperties

The m ethod we are developing follows a Bayesian approach. It is a close analog to the one introduced by K au m ann et al. (2003) for the study of star form ation histories in the Sloan D igital Sky Survey. However the variety of observable properties has completely di erent origins in both contexts: stochasticity at a given age, m ass and m etallicity, plays a predom inant role here while it is negligible in m ost galaxies taken as a whole.

We have started a cam paign of Monte-Carlo (MC) simulations (Fig.1), that is used as a catalog for establishing the joint probability distributions of the intrinsic parameters of a cluster (age, total num ber of stars or total mass, metallic-



Figure 2. Example of a posteriori probability distributions of age, m ass and extinction, for a synthetic cluster <code>\observed"</code> in U, B, V, I, J and K with 1- photom etric errors of 0.05 m agnitudes. The prior is that the cluster m ass distributions in galaxies vary as M², and that the cluster age distributions in galaxies are constant in logarithm ic age units (only clusters that survived dynam ical disruption m atter here). The true properties of the <code>\observed"</code> cluster are indicated as triangles, the m ost probable values as derived from the <code>\observations"</code> are shown as sm all circles.

ity, extinction), given a set of photom etric m easurem ents and their uncertainties (Fouesneau et al., in prep). The total mass is not a simple scaling factor. It enters the problem even when only colors are available because it a ects the color probability distributions of clusters of a given age and m etallicity.

A m ong the assumptions in the m ethod are the stellar IM F, and the adopted stellar evolutionary tracks and stellar spectral libraries. The m ain a priori of the B ayesian inversion are the underlying cluster form ation history (number of clusters per age bin in the catalog), and the cluster m ass function. In applications to real cluster populations, several a priori distributions should be tested and com pared. The e ects of dynam ical evolution on the IM F and on the cluster m ass function also need to be discussed.

W e generate integrated photom etric properties of individual clusters with a modi ed version of PEGASE (Fioc & Rocca-Volm erange 1999) that im plements a discrete IM F. The models take into account the nebular continuum and line em ission of clusters that contain ionizing stars.

The assumption that errors in the observational data are G aussian leads to a simple expression for the a posteriori probability distribution of an intrinsic property, such as the age or mass of an individual cluster. The probability for an intrinsic property X to be located in an interval $[x_1;x_2]$, given the photom etric m easurem ents $Y = fY_k g_{k2} [_{l:n}]$, is:

$$P(X \ 2 \ [x_1;x_2] \ jY) = \frac{X \ Y}{\frac{1}{2 \ k}} \exp \left(\frac{(Y_k \ Y_{k,M_i})^2}{2 \ k}\right)^2 P(M_i); \quad (1)$$

where P (M $_{\rm i}$) is the probability of the individual model M $_{\rm i}$ to be chosen and is the norm alization constant. Through the factors P (M $_{\rm i}$), this expression adapts to any prior m ass or age distribution.

Fouesneau & Lancon

As an example, Figure 2 shows recovered age, mass and extinction probability distributions based on pseudo-observational data, i.e. on the synthetic magnitudes of one of the clusters in the MC-model catalog (modied by the addition of reasonable noise). Although six absolute uxes across the spectrum are used, the possible age and mass ranges remain signi cantly broad, and asymmetric. We are in the process of establishing the systematic dierences between the actual properties of a cluster and various estimates thereof (best² model, most probable values, and values recovered using the classical approach with a continuously populated IMF). We will also reinvestigate which minimal combination of observations constrains the intrinsic properties most e ciently.

Varying the m etallicity, the allow ed extinction range or the a priori distributions of m ass and age will signi cantly change the results. We are still exploring the dierent e ects, especially those of m etallicity and extinction. We are also extending the MC-m odel catalog by adding new massive clusters, which will be necessary to test the e ect of adopting atter a priori cluster mass distributions.

3 Conclusion

A though massive clusters contain in portant information on active episodes of star formation in galaxies, the need to study smaller clusters is increasing as the amount of available observational data grows. Intrinsically rare but lum inous stars are responsible for large stochastic uctuations in the integrated energy distribution of these objects. Hence, new tools that deal with stochastic uctuations explicitly need to be developed. In this context we have in plemented a B ayesian approach that, given a set of photom etric data, provides most probable cluster parameters as well as con dence intervals. The ongoing study of system – atic di erences between true and estimated values should allow us to correct any signi cant biases in previous studies.

A cknow ledgm ents. We thank M.Fioc formaking his updated version of Pegase available formodication in advance of publication.

References

Bastian, N., Trancho, G., Konstantopoulos, I.S., & Miller, B.W. 2009, ApJ, 701, 607
Bruzual, G.A. 2002, Extragalactic Star C lusters, 207, 616
Cervino, M., & Luridiana, V. 2004, A & A, 413, 145
Chiosi, C., Bertelli, G., & Bressan, A. 1988, A & A, 196, 84
Fioc, M., & Rocca-Volmerange, B. 1999, arX is astro-ph/9912179
Kau mann, G., et al. 2003, M NRAS, 341, 33
Lancon, A., & Mouhcine, M. 2000, Massive Stellar C lusters, 211, 34
Lancon, A., Gallagher, J.S., III, Mouhcine, M., Sm ith, L.J., Ladjal, D., & de G rijs, R. 2008, A & A, 486, 165
Lancon, A., & Fouesneau, M. 2009, arX is 0903.4557
W hitm ore, B.C., Zhang, Q., Leitherer, C., Fall, S.M., Schweizer, F., & Miller, B.W. 1999, AJ, 118, 1551