```
**FULL TITLE **
ASP Conference Series, Vol. **VOLUME **, c **YEAR OF PUBLICATION **
**NAMES OF EDITORS **
```

M id-Infrared Evidence for Accelerated Evolution in C om pact G roup G alaxies

L isa M ay W alker¹, K elsey E . Johnson¹, Sarah C . G allagher², John E . H ibbard³, A nn E . H ornschem eier⁴, Jane C . C harlton⁵, T hom as H . Jarrett⁶

¹D ept of A stronom y, University of Virginia, Charlottesville, VA 22904; ²D ept of Physics and A stronom y, University of W estern Ontario, London, ON N 6A 3K 7 Canada; ³NRAO, Charlottesville, VA 22903; ⁴Laboratory for X - Ray A strophysics, NA SA G oddard Space F light Center, G reenbelt, M D 20771; ⁵D ept of A stronom y and A strophysics, Pennsylvania State University, University Park, PA 16802; ⁶Spitzer Science Center, California Institute of Technology, Pasadena, CA 91125

A bstract. We nd evidence for accelerated evolution in compact group galaxies from the distribution in mid-infrared colorspace of 42 galaxies from 12 Hickson C om pact G roups (H C G s) compared to the the distributions of several other samples including the LV L+ SINGS galaxies, interacting galaxies, and galaxies from the C om a Cluster. We nd that the HCG galaxies are not uniform ly distributed in colorspace, as well as quantitative evidence for a gap. Galaxies in the infall region of the C om a cluster also exhibit a non-uniform distribution and a less well de ned gap, which m ay re ect a similarity with the compact group environm ent. N either the C om a C enter or interacting samples show evidence of a gap, leading us to speculate that the gap is unique to the environm ent of high galaxy density where gas has not been fully processed or stripped.

1 Sam ples

The HCG dataset, taken from Johnson et al. (2007), comprises 42 galaxies from 12 groups. The groups contain varying amounts of HI, and span the three classi cation categories discussed in Johnson et al. (2007). The most HI gas-rich groups with log (M_{HI})=log (M_{dyn}) 0:9 are classi ed as type I, the HI gas-poor groups with log (M_{HI})=log (M_{dyn}) < 0:8 as type III, while type II contains the interm ediate groups.

The LVL data, discussed in D ale et al. (2009) consists of 211 galaxies within 11 M pc. The SINGS data, as described in D ale et al. (2007), consist of 71 galaxies. We combined the LVL and SINGS galaxies to create a control sample, referred to as LVL+SINGS. The 35 galaxies from the Spitzer Spirals, Bridges, and Tails Interacting G alaxy Survey (hereafter referred to as the interacting sample) are comprised of otherwise relatively isolated binary galaxy systems, whose members are tidally disturbed (Sm ith et al. 2007). The C om a sample, discussed in Jenkins et al. (2007), is comprised of galaxies from two elds. The

rst eld is located in the center of the cluster, where the galaxy density is very high. The second eld is the infall region, located near 0.4 virial radii at the



Figure 1. Left: Color-colorplot of HCG galaxies. The plus signs are galaxies from type I (HIgas-rich) groups, the triangles correspond to galaxies from type II groups, and the squares represent galaxies from type III (HIgas-poor) groups. The low er-left region of the plot contains galaxies whose light is dom inated by norm al stellar populations, while active galaxies (i.e. star-form ing) reside in the upper right. Right: Color-colorplot of com parison sam ples. The grey plus signs are the HCG galaxies, and the error bars in the bottom right indicate typical errors for each sam ple. The dashed lines indicate the cropping values, so that the analysis was done only over the range of colorspace occupied by both the HCG galaxies and the com parison sam ple.

X-ray secondary peak, where the galaxy density is still higher than eld density (Jenkins et al. 2007). Our HCG sample has very few galaxies with lum inosities below log (L_{4:5} [erg=s]) = 40:9. Therefore, to compare the HCG galaxies to similar galaxy populations from the other samples we only consider galaxies with lum inosities greater than this.

2 Colorspace

G alaxies with blue M IR colors have IR SED sconsistent with being dom inated by stellar light. G alaxies with ongoing SF tend to have red M IR colors indicative of PAH emission and warm /hot dust. The gap between these colors in HCG colorspacem ay be due to rapid evolution through the stage during which galaxies have interm ediate M IR colors. The dotted lines indicate the range of com parison used for the statistical tests.

3 Statistical Tests

The KS test comparing the HCG galaxies with a uniform distribution clearly illustrates the gap, manifested as the nearly-horizontal portion of the CDF. The two-distribution KS tests yield values for the infallC om a sam ple high enough that we cannot reject the hypothesis that they are drawn from the sam e parent distribution as the HCG s over the range of com parison.



Figure 2. Left: K S test for the rotated H CG distribution against a m odel of uniform distribution. The m axim um deviation of the H CG CDF (solid line) from a uniform distribution (dashed line) is D, indicated by the vertical line. The nearly at portion highlighted in grey reveals the gap. R ight: K S test for the rotated com parison samples against the H CG galaxies.

4 G ap R egion

The gap region is de ned by HCG galaxies bounding gap at $_{C_{M IR}} = 0.31$ and 0.38, indicated by the gray box in Figure 2. Since the gap represents a de cit of galaxies over this color range, its signature is a attening in the CDFs. We excluded the interacting sample due to insu cient data blueward of gap The HCG sample has the most pronounced gap, followed by Com a Infall.

Table 1. Slopes of the CDFs of all sam ples over the region of the HCG gap.

Sample	Slope
НСG	0.104
LVL + SINGS	808.0
Coma (center)	0.544
Coma (infall)	0.215

5 CMDs

The morphology of the CMDs recets the criteria used to de neeach sample. For example, both Coma samples have a signi cant concentration of galaxies with blue MIR colors, characteristic of galaxies with little to no SF, while the samples biased toward actively star forming systems show a scattering of galaxies with red MIR colors. The CMD of the HCGs is a composite of these two types, but has a decit of galaxies with intermediate MIR colors! a transitional phase between no PAHs and strong PAH features.



Figure 3. Left: CMD for the HCG galaxies. The dotted line indicates the minimum luminosity required for inclusion. The gap is indicated by the grey stripe. It is noteworthy that there does not seem to be a color-lum inosity correlation. Right: CMD for the comparison samples. The criteria used to de ne each sample is apparent in the morphology of the CMD s.

6 D iscussion

W e suggest that the distribution in M IR colorspace, and particular the occurance of a gap in the color distribution, likely re ect di erent star form ation states, with galaxies ranging from active/e cient (red M IR colors) to passive/ine cient (blue M IR colors). The distribution in colorspace suggests that galaxies in the HCG environm ent m ove quickly from active to passive states, in a m anner sim ilar to, but m ore dram atic than, the environm ent on the outskirts of clusters. G alaxies from both m ore dense (C om a center) and less dense (e.g. LVL+SINGS) environm ents have a continuous distribution of system s between active and passive states.

The presence of the gap in the M IR colorspace distribution of the HCGs combined with the fact that the gap is not present in less dense environments indicates that this type of local environment signicantly in uncessed galaxy properties. In order to understand the processes that a lect galaxy evolution, we need to understand how gas is processed in the interstellar medium and intragroup medium in these environments. Compact groups are clearly an important part of understanding galaxy assembly and hierarchical structure formation.

References

- Dale, D.A. et al. 2007, ApJ, 655, 863
- Dale, D.A. et al. 2009, subm itted
- Jenkins, L.P., Hornschem eier, A.E., Mobasher, B., Alexander, D.M., & Bauer, F.E., 2007 ApJ, 666, 846
- Johnson, K.E., Hibbard, J.E., Gallagher, S.C., Charlton, J.C., Hornshem eier, A.E., Jarrett, T.H., & Reines, A.E. 2007, AJ, 134, 1522
- Sm ith, B.J., Struck, C., Hancock, M., Appleton, P.N., Charmandaris, V., & Reach, W.T.2007, AJ, 133, 791