

\*\*FULL TITLE\*\*  
 ASP Conference Series, Vol. \*\*VOLUME\*\*, c \*\*YEAR OF PUBLICATION\*\*  
 \*\*NAMES OF EDITORS\*\*

## Models of the Intergalactic Gas in Stephan's Quintet

Jeong-Sun Hwang<sup>1</sup>, Curtis Struck<sup>1</sup>, Florent Renaud<sup>2,3</sup>, and Philip Appleton<sup>4</sup>

<sup>1</sup>Department of Physics and Astronomy, Iowa State University, Ames, IA 50011, USA

<sup>2</sup>Observatoire Astronomique and CNRS UMR 7550, Université de Strasbourg, 11 rue de l'Université, F-67000 Strasbourg, France

<sup>3</sup>Institut für Astronomie der Univ. Wien, Turkenschanzstr. 17, A-1180 Vienna, Austria

<sup>4</sup>NASA Herschel Science Center (NHSC), California Institute of Technology, Mailcode 100-22, Pasadena, CA 91125, USA

**Abstract.** We use smoothed particle hydrodynamics (SPH) models to study the large-scale morphology and dynamical evolution of the intergalactic gas in Stephan's Quintet, and compare to multiwavelength observations. Specifically, we model the formation of the hot X-ray gas, the large-scale shock, and emission line gas as the result of NGC 7318b colliding with the group. We also reproduce the N-body model of Renaud and Appleton for the tidal structures in the group.

### 1 Introduction

Stephan's Quintet (hereafter SQ) is a compact group of galaxies discovered by Edouard Stephan in 1877. Since it is relatively close and shows many interesting features of galaxy-galaxy and galaxy-intergalactic medium (IGM) interactions SQ has been a popular object of study. Its multiple interactions, however, make difficult to know its dynamical history and interpret some observed aspects. In order to study SQ's dynamical evolution and the cause of the large-scale shock structure in its IGM seen in multiwavelength observations we have performed numerical simulations of SQ. In this proceedings we first review SQ's morphological and kinematical structures that we tried to simulate and then present our preliminary model results. Finally we discuss some adjustment of our current model to obtain better results.

### 2 Past Study of SQ

The optical morphology of SQ is shown in Figure 1. NGC 7320 is a foreground galaxy which has a considerably smaller redshift than others (Burbridge & Burbridge 1961). Assuming a group distance of 94 Mpc and a Hubble constant of  $70 \text{ km s}^{-1} \text{ Mpc}^{-2}$ , the group except NGC 7318b has a redshift about  $6600 \text{ km s}^{-1}$  and NGC 7318b is coming toward us with the relative velocity of about  $900 \text{ km s}^{-1}$ .

Two long tidal tails extending from NGC 7319 toward NGC 7320c are visible. The southern tail, which is fainter, passes behind the foreground galaxy NGC 7320 and runs almost parallel with the northern tail. NGC 7319 is known to be a Seyfert 2 galaxy.

The VLA HI observations reveal that most of the HI gas resides outside of the galaxies (Williams, Yun, & Verdes-Montenegro 2002). The observed HI clouds along the two optical parallel tails are thought to be the result of one or more past encounters of NGC 7320c with the group (Moles, Sulentic, & Marquez 1997; Sulentic et al. 2001). The past encounters might have stripped and removed most of the gas from NGC 7319 to the intergalactic space. NGC 7318b is colliding with the IGM which has formed during the past encounters (Sulentic et al. 2001; Xu et al. 2003). This ongoing collision may produce the large-scale shock in the IGM observed in radio continuum (Allen & Hartsuiker 1972; van der Hulst & Rots 1981), X-rays (Pietsch et al. 1997; Trinchieri et al. 2003; O’Sullivan et al. 2009) and mid-IR H<sub>2</sub> emission (Appleton et al. 2006).



Figure 1. The optical morphology of SQ. North is up and east to the left. Credit: NOAO/AURA/NSF

### 3 Numerical Simulations of SQ

Renaud and Appleton had performed N-body simulations of SQ using mostly NEMO, a stellar dynamics toolbox (Teuben 1995). Considering observational constraints they deduced probable ranges of initial conditions for their model galaxies and reproduced SQ’s general morphology (Renaud & Appleton 2009). In order to add them hydrodynamical effects to the N-body model and study dynamical evolution of SQ we have produced an SPH code by modifying the SPH code of Struck (1997) which was designed for studying collisions between galaxy pairs (see Struck 1997 for details). We performed the simulation with four strongly interacting members. Each of our model galaxies has disks with gas and collisionless star particles and a rigid dark matter halo. We took the initial parameters from the N-body model and adjusted them (mainly initial velocities of model galaxies) to use in the SPH model which has more extended dark matter halo potentials. We used total 81120 (40320 gas + 40800 star) particles and the mass ratio of 1 (NGC 7319) : 0.7 (NGC 7318a) : 0.6 (NGC 7318b) : 0.2 (NGC 7320c).

Our preliminary results of the SPH simulation are as follows. Figure 2a and b show the production of two long tidal tails at an earlier time step. Since

the effect of NGC 7318b to the group is small at early times we have put NGC 7318b into the simulation at this time step. Figure 2.c and d represent current morphology of SQ and the collision between NGC 7318b and the IGM from the model.




Figure 2. Snapshots of the model galaxies at an early time step (top panels) and time near the present (bottom panels). The left and right panels show the distributions of gas and star particles respectively. Particles originating from NGC 7319, NGC 7318a, NGC 7318b, and NGC 7320c are shown with red, green, yellow, and cyan dots respectively. The center positions of four galaxies are marked with crosses. (a) and (b): Particles originating from NGC 7319 and NGC 7320c are shown for simplicity. The two long parallel tidal tails have developed. The  $(x, y, z)$  values of the centers of NGC 7319, NGC 7318a, NGC 7318b and NGC 7320c are  $(0, 0, 0)$ ,  $(-26, 34, 24)$ ,  $(20, 11, -98)$ , and  $(-47, 25, -24)$  respectively. (c) and (d): Particles originating from all four galaxies are shown. NGC 7318b is colliding with the group. The  $(x, y, z)$  values of the centers of NGC 7319, NGC 7318a, NGC 7318b and NGC 7320c are  $(0, 0, 0)$ ,  $(44, -7, -22)$ ,  $(26, -3, 12)$ , and  $(-25, 21, 4)$  respectively.

## 4 Discussion and Future Work

The general morphology of SQ was reproduced except the position of the southern tail and inner structures of spiral members. With our extended halo potentials the southern tail was placed too far out, so NGC 7318b did not hit many particles in the tidal tails. We will adjust it to make a more direct collision between the galaxy and the IGM. Also, to obtain well-matched results in radial velocities and temperatures of gas particles (not shown in this article) with those of HI, radio continuum, and X-ray observations, we will be working on fine tuning the initial parameters and cooling in the models.

Acknowledgments. We thank the conference organizers for providing an opportunity to discuss this topic and financial support. This work has been supported by NASA Chandra grant AR 90010B.

## References

- Allen, R. J., & Hartsuiker, J. W. 1972, *Nature*, 239, 324  
 Appleton, P. N., Xu, K. C., Reach, W., Dopita, M. A., Gao, Y., Lu, N., Popescu, C. C., Sulentic, J. W., Tu, S. R. J., & Yun, M. S. 2006, *ApJ*, 639, L51  
 Burbidge, E. M., & Burbidge, G. R. 1961, *ApJ*, 134, 244  
 Moles, M., Sulentic, J. W., & Marquez, I. 1997, *ApJ*, 485, L69  
 O'Sullivan, E., Giacintucci, S., Vrtilak, J. M., Raychaudhury, S., & David, L. P. 2009, *ApJ*, 701, 1560  
 Pietsch, W., Trinchieri, G., Arp, H., & Sulentic, J. W. 1997, *A & A*, 322, 89  
 Renaud, F., & Appleton, P. N. 2009, in preparation  
 Struck, C. 1997, *ApJS*, 113, 269  
 Sulentic, J. W., Rosado, M., Dultzin-Hacyan, D., Verdes-Montenegro, L., Trinchieri, G., Xu, C., & Pietsch, W. 2001, *AJ*, 122, 2993  
 Trinchieri, G., Sulentic, J., Breitschwerdt, D., & Pietsch, W. 2003, *A & A*, 401, 173  
 Teuben, P. 1995, in *ASP Conf. Ser. 77, Astronomical Data Analysis Software and Systems IV*, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes (San Francisco: ASP), 398  
 van der Hulst, J. M., & Rots, A. H. 1981, *AJ*, 86, 1775  
 Williams, B. A., Yun, M. S., & Verdes-Montenegro, L. 2002, *AJ*, 123, 2417  
 Xu, C. K., Lu, N., Condon, J. J., Dopita, M., & Tu, S. R. J. 2003, *ApJ*, 595, 665