

Diagonalization of boundary transfer matrix for the $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model

February 13, 2022

Takeo KOJIMA

*Department of Mathematics, College of Science and Technology, Nihon University,
Surugadai, Chiyoda-ku, Tokyo 101-0062, JAPAN*

Abstract

We construct a free field realization of the ground state of the boundary transfer matrix for the $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model. Using this ground state and type-II vertex operator, we have a diagonalization of the boundary transfer matrix.

1 Introduction

The vertex operator approach [1, 2] provides a powerful method to study solvable models. In [1] authors diagonalized the XXZ-transfer matrix on infinite spin chain by vertex operators. Sklyanin [3] begin systematic approach to boundary condition generalization in the framework of the algebraic Bethe ansatz. In periodic boundary condition, we construct a family of commuting transfer matrix from a solution of the Yang-Baxter equation. Sklyanin showed that similar construction is possible with the aid of a solution of the boundary Yang-Baxter equation. In open boundary condition, we construct a family of commuting transfer matrix from both a solution of the Yang-Baxter equation and a solution of the boundary Yang-Baxter equation. In

[6] the authors diagonalized the boundary XXZ-transfer matrix on semi-infinite spin chain by vertex operators. The vertex operator approach was extended to higher-rank boundary XXZ-model in [9]

The vertex operator approach to solvable model was originally formulated for vertex type model such as the XXZ-model [1], and then extended to the face type model such as the Andrews-Baxter-Forrester (ABF) model [4, 10]. The ABF model is described by the elliptic algebra $U_{q,p}(\widehat{sl}(2, \mathbb{C}))$. The $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model we are going to study in this paper were introduced in [5], as the higher-rank generalization of the ABF model introduced in [4]. In this paper we diagonalize the boundary transfer matrix for the $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model by vertex operator approach. We construct a free field realization of the ground-state of the commuting boundary transfer matrix. Using this groundstate and type-II vertex operators for the elliptic algebra, we get a diagonalization of the boundary transfer matrix of the $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model. The result of this paper gives a higher-rank generalization of the ABF model [7].

The text is organized as follows. In section 2 we recall the boundary $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model, and introduce the boundary transfer matrix. In section 3 we recall the free field realization of the vertex operators. In section 4 we construct a free field realization of the ground-state of the boundary transfer matrix, and give a diagonalization of the boundary transfer matrix by using the type-II vertex operators.

2 Boundary $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model

We recall the boundary $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model.

2.1 Bulk Boltzmann weights

The $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model has two parameters x and r . We assume $0 < x < 1$ and $r \geq 5$ ($r \in \mathbb{Z}$). We set the elliptic theta function $[u]$ by

$$[u] = x^{\frac{u^2}{r} - u} \Theta_{x^{2r}}(x^{2u}). \quad (2.1)$$

Here we have used

$$\Theta_q(z) = (q, q)_\infty (z; q)_\infty (q/z; q)_\infty, \quad (2.2)$$

$$(z; q_1, q_2, \dots, q_m)_\infty = \prod_{j_1, j_2, \dots, j_m=0}^{\infty} (1 - q_1^{j_1} q_2^{j_2} \cdots q_m^{j_m} z). \quad (2.3)$$

Let $\epsilon_\mu (1 \leq \mu \leq 3)$ be the orthonormal basis of \mathbb{R}^3 with the inner product $(\epsilon_\mu | \epsilon_\nu) = \delta_{\mu,\nu}$. Let us set $\bar{\epsilon}_\mu = \epsilon_\mu - \epsilon$ where $\epsilon = \frac{1}{3} \sum_{\nu=1}^3 \epsilon_\nu$. The type $\widehat{sl}(3, \mathbb{C})$ weight lattice is the linear span of $\bar{\epsilon}_\mu$.

$$P = \sum_{\mu=1}^3 \mathbb{Z} \bar{\epsilon}_\mu. \quad (2.4)$$

Note $\sum_{\mu=1}^3 \bar{\epsilon}_\mu = 0$. Let the simple root $\alpha_\mu = \bar{\epsilon}_\mu - \bar{\epsilon}_{\mu+1}$ ($\mu = 1, 2$). For $a \in P$ we set

$$a_{\mu,\nu} = a_\mu - a_\nu, \quad a_\mu = (a + \rho | \bar{\epsilon}_\mu), \quad (2.5)$$

where $\rho = 2\bar{\epsilon}_1 + \bar{\epsilon}_2$. The Boltzmann weights $W \left(\begin{array}{cc} a & b \\ c & d \end{array} \middle| u \right)$ are given by

$$W \left(\begin{array}{cc} a + 2\bar{\epsilon}_\mu & a + \bar{\epsilon}_\mu \\ a + \bar{\epsilon}_\mu & a \end{array} \middle| u \right) = r_1(u), \quad (2.6)$$

$$W \left(\begin{array}{cc} a + \bar{\epsilon}_\mu + \bar{\epsilon}_\nu & a + \bar{\epsilon}_\mu \\ a + \bar{\epsilon}_\nu & a \end{array} \middle| u \right) = r_1(u) \frac{[u][a_{\mu,\nu} - 1]}{[u - 1][a_{\mu,\nu}]}, \quad (2.7)$$

$$W \left(\begin{array}{cc} a + \bar{\epsilon}_\mu + \bar{\epsilon}_\nu & a + \bar{\epsilon}_\nu \\ a + \bar{\epsilon}_\nu & a \end{array} \middle| u \right) = r_1(u) \frac{[u - a_{\mu,\nu}][1]}{[u - 1][a_{\mu,\nu}]}. \quad (2.8)$$

Otherwise are zero. The function $r_1(u)$ is given by

$$r_1(u) = z^{\frac{r-1}{r} \frac{2}{3}} \frac{h_1(z^{-1})}{h(z)}, \quad h_1(z) = \frac{(x^2 z; x^{2r}, x^6)_\infty (x^{2r+4} z; x^{2r}, x^6)_\infty}{(x^{2r} z; x^{2r}, x^6)_\infty (x^6 z; x^{2r}, x^6)_\infty}. \quad (2.9)$$

Here we have used $z = x^{2u}$. The Boltzmann weights satisfy the following relations.

(1) Yang-Baxter equation :

$$\begin{aligned} & \sum_g W \left(\begin{array}{cc} d & e \\ c & g \end{array} \middle| u_1 \right) W \left(\begin{array}{cc} c & g \\ b & a \end{array} \middle| u_2 \right) W \left(\begin{array}{cc} e & f \\ g & a \end{array} \middle| u_1 - u_2 \right) \\ &= \sum_g W \left(\begin{array}{cc} g & f \\ b & a \end{array} \middle| u_1 \right) W \left(\begin{array}{cc} d & e \\ g & f \end{array} \middle| u_2 \right) W \left(\begin{array}{cc} d & g \\ c & b \end{array} \middle| u_1 - u_2 \right). \end{aligned} \quad (2.10)$$

(2) The first inversion relation :

$$\sum_g W \left(\begin{array}{cc} c & g \\ b & a \end{array} \middle| -u \right) W \left(\begin{array}{cc} c & d \\ g & a \end{array} \middle| u \right) = \delta_{b,d}. \quad (2.11)$$

(3) The second inversion relation :

$$\sum_g G_g W \left(\begin{array}{cc} g & b \\ d & c \end{array} \middle| 3 - u \right) W \left(\begin{array}{cc} g & d \\ b & a \end{array} \middle| u \right) = \delta_{a,c} \frac{G_b G_d}{G_a}. \quad (2.12)$$

where $G_a = [a_{1,2}][a_{1,3}][a_{2,3}]$. The Boltzmann weights are related to the elliptic algebra $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$.

2.2 Boundary Boltzmann weights

In [8] the boundary Boltzmann weights $K \left(\begin{array}{c|c} b & u \\ a & \\ c & \end{array} \right)$ are given by

$$K \left(\begin{array}{c|c} k & u \\ k + \bar{\epsilon}_\mu & \\ k & \end{array} \right) = z^{\frac{2}{r}(-\frac{r-1}{3} + (\bar{\epsilon}_1|k))} \frac{h(z)}{h(z^{-1})} \frac{[c-u][k_{1,\mu} + c + u]}{[c+u][k_{1,\mu} + c - u]}, \quad (c \in \mathbb{R}; \mu = 1, 2, 3). \quad (2.13)$$

Otherwise are zero. The function $h(z)$ is given in (4.18). They satisfy the boundary Yang-Baxter equation,

$$\begin{aligned} & \sum_{f,g} W \left(\begin{array}{c|c} c & f \\ b & a \end{array} \middle| u-v \right) W \left(\begin{array}{c|c} c & d \\ f & g \end{array} \middle| u+v \right) K \left(\begin{array}{c|c} g & u \\ f & a \end{array} \right) K \left(\begin{array}{c|c} e & v \\ d & g \end{array} \right) \\ &= \sum_{f,g} W \left(\begin{array}{c|c} c & d \\ f & e \end{array} \middle| u-v \right) W \left(\begin{array}{c|c} c & f \\ b & g \end{array} \middle| u+v \right) K \left(\begin{array}{c|c} e & u \\ f & g \end{array} \right) K \left(\begin{array}{c|c} g & v \\ b & a \end{array} \right). \end{aligned} \quad (2.14)$$

2.3 Vertex operator

Following the general scheme of algebraic approach in solvable lattice models, we give the type-I vertex operators. Let us consider the corner transfer matrices $A(z), B(z), C(z), D(z)$ which represent NW, SW, SE, NE quadrants, respectively. The space $\mathcal{H}_{l,k}$ of the eigenvectors of $A(z)$ is parametrized by $l, k \in P$. Let us introduce the type-I vertex operator $\Phi^{(a,b)}(z)$. We denote by $\Phi^{(k+\bar{\epsilon}_j,k)}(z^{-1})$ the half-infinite transfer matrix extending to infinity in the north. This is an operator

$$\Phi^{(k+\bar{\epsilon}_j,k)}(z^{-1}) : \mathcal{H}_{l,k} \rightarrow \mathcal{H}_{l,k+\bar{\epsilon}_j}.$$

The operator $\Phi^{(a,b)}(z) = 0$ for $a - b \neq \bar{\epsilon}_j$. Similarly, we introduce dual type-I vertex operator $\Phi^{*(a,b)}(z)$. We denote by $\Phi^{*(k,k+\bar{\epsilon}_j)}(z)$ the half-infinite transfer matrix extending to infinity in the west. This is an operator

$$\Phi^{*(k,k+\bar{\epsilon}_j)}(z) : \mathcal{H}_{l,k+\bar{\epsilon}_j} \rightarrow \mathcal{H}_{l,k}.$$

The operator $\Phi^{(a,b)}(z) = 0$ for $b - a \neq \bar{\epsilon}_j$. They satisfy the following relations.

(1) Commutation relation

$$\Phi^{(c,b)}(z_1) \Phi^{(b,a)}(z_2) = \sum_d W \left(\begin{array}{c|c} c & d \\ b & a \end{array} \middle| u_1 - u_2 \right) \Phi^{(c,d)}(z_2) \Phi^{(d,a)}(z_1). \quad (2.15)$$

(2) Inversion relation

$$\sum_g \Phi^{*(a,g)}(z) \Phi^{(g,a)}(z) = 1, \quad \Phi^{(a,b)}(z) \Phi^{*(b,c)}(z) = \delta_{a,c}. \quad (2.16)$$

Later we give a free field realization of the vertex operators acting on the bosonic Fock space.

2.4 Boundary transfer matrix

We define the boundary transfer matrix

$$T_B^{(k)}(z) = \sum_{j=1,2,3} \Phi^{*(k,k+\bar{\epsilon}_j)}(z) K \left(\begin{array}{c|c} k & \\ \hline k + \bar{\epsilon}_j & u \\ \hline k & \end{array} \right) \Phi^{(k+\bar{\epsilon}_j,k)}(z^{-1}). \quad (2.17)$$

The boundary Yang-Baxter equation implies the commutativity.

$$[T_B^{(k)}(z_1), T_B^{(k)}(z_2)] = 0. \quad (2.18)$$

Our problem of this paper is to diagonalize the boundary transfer matrix $T_B^{(k)}(z)$.

3 Free field realization

We give a free field realization of the vertex operators [11, 12].

3.1 Boson

We set the bosonic oscillators β_m^i , ($i = 1, 2; m \in \mathbb{Z}$) by

$$[\beta_m^j, \beta_n^k] = \begin{cases} m \frac{[(r-1)m]_x [2m]_x}{[rm]_x [3m]_x} \delta_{m+n,0} & (j = k) \\ -mx^{3m} \operatorname{sgn}(j-k) \frac{[(r-1)m]_x [m]_x}{[rm]_x [3m]_x} \delta_{m+n,0} & (j \neq k). \end{cases} \quad (3.1)$$

Here the symbol $[a]_x = \frac{x^a - x^{-a}}{x - x^{-1}}$. Let us set β_m^3 by $\sum_{j=1}^3 x^{-2jm} \beta_m^j = 0$. The above commutation relations are valid for all $1 \leq j, k \leq 3$. We also introduce the zero-mode operators P_α, Q_α , ($\alpha \in P$) by

$$[iP_\alpha, Q_\beta] = (\alpha|\beta), \quad (\alpha, \beta \in P). \quad (3.2)$$

In what follows we deal with the bosonic Fock space $\mathcal{F}_{l,k}(l, k \in P)$ generated by $\beta_{-m}^j (m > 0)$ over the vacuum vector $|l, k\rangle$:

$$\mathcal{F}_{l,k} = \mathbb{C}[\{\beta_{-1}^j, \beta_{-2}^j, \dots\}_{j=1,2,3} | l, k\rangle, \quad (3.3)$$

where

$$\begin{aligned}\beta_m^j |k, l\rangle &= 0, \quad (m > 0), \\ P_\alpha |l, k\rangle &= \left(\alpha \left| \sqrt{\frac{r}{r-1}} l - \sqrt{\frac{r-1}{r}} k \right. \right) |l, k\rangle, \\ |l, k\rangle &= e^{i\sqrt{\frac{r}{r-1}} Q_l - \sqrt{\frac{r-1}{r}} Q_k} |0, 0\rangle.\end{aligned}$$

3.2 Vertex operator

We give a free field realization of the type-I vertex operators [11], associated with the elliptic algebra $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ [13, 14]. Let us set $P(z), Q(z), R_-^j(z), S_-^j(z) (j = 1, 2)$ by

$$P(z) = \sum_{m>0} \frac{1}{m} \beta_{-m}^1 z^m, \quad Q(z) = - \sum_{m>0} \frac{1}{m} \beta_m^1 z^{-m}, \quad (3.4)$$

$$R_-^j(z) = - \sum_{m>0} \frac{1}{m} (\beta_{-m}^j - \beta_{-m}^{j+1}) x^{jm} z^m, \quad S_-^j(z) = \sum_{m>0} \frac{1}{m} (\beta_m^j - \beta_m^{j+1}) x^{-jm} z^{-m}. \quad (3.5)$$

Let us set the basic operators $U(z), F_{\alpha_1}(z), F_{\alpha_2}(z)$ on the Fock space $\mathcal{F}_{l,k}$.

$$U(z) = z^{\frac{r-1}{3r}} e^{-i\sqrt{\frac{r-1}{r}} Q_{\bar{\epsilon}_1}} z^{-\sqrt{\frac{r-1}{r}} P_{\bar{\epsilon}_1}} e^{P(z)} e^{Q(z)}, \quad (3.6)$$

$$F_{\alpha_j}(z) = z^{\frac{r-1}{r}} e^{i\sqrt{\frac{r-1}{r}} Q_{\alpha_j}} z^{\sqrt{\frac{r-1}{r}} P_{\alpha_j}} e^{R_-^j(z)} e^{S_-^j(z)}. \quad (3.7)$$

In what follows we set

$$\pi_\mu = \sqrt{r(r-1)} P_{\bar{\epsilon}_\mu}, \quad \pi_{\mu\nu} = \pi_\mu - \pi_\nu. \quad (3.8)$$

Then $\pi_{\mu\nu}$ acts on $\mathcal{F}_{l,k}$ as an integer $(\epsilon_\mu - \epsilon_\nu |rl - (r-1)k)$. We give the free field realization of the type-I vertex operators.

$$\Phi_1(z) = U(z), \quad (3.9)$$

$$\Phi_2(z) = \oint_{C_1} \frac{dw_1}{w_1} U(z) F_{\alpha_1}(w_1) \frac{[v_1 - u + \frac{1}{2} - \pi_{1,2}]}{[v_1 - u - \frac{1}{2}]}, \quad (3.10)$$

$$\Phi_3(z) = \oint_{C_2} \oint_{C_2} \frac{dw_1}{w_1} \frac{dw_2}{w_2} U(z) F_{\alpha_1}(w_1) F_{\alpha_2}(w_2) \frac{[v_1 - u + \frac{1}{2} - \pi_{1,3}]}{[v_1 - u - \frac{1}{2}]} \frac{[v_2 - v_1 + \frac{1}{2} - \pi_{2,3}]}{[v_2 - v_1 - \frac{1}{2}]}. \quad (3.11)$$

Here we set $z = x^{2u}, w_j = x^{2v_j} (j = 1, 2)$. We take the integration contours to be simple closed curves around the origin satisfying

$$C_1 : x|z| < |w_1| < x^{-1}|z|,$$

$$C_2 : x|z| < |w_1| < x^{-1}|z|, \quad x|w_1| < |w_2| < x^{-1}|w_1|.$$

We identify $\Phi^{(k+\bar{\epsilon}_j, k)}(z) = \Phi_j(z)$.

Let us introduce the type-II vertex operators [12], associated with the elliptic algebra $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$. The type-II vertex operators represents the excitations. Let us set $r^* = r - 1$. Let us set the elliptic theta function $[u]^*$ by

$$[u]^* = x^{\frac{u^2}{r^*} - u} \Theta_{x, 2r^*}(x^{2u}). \quad (3.12)$$

Let us set $P^*(z), Q^*(z), R_+^j(z), S_+^j(z) (j = 1, 2)$ by

$$P^*(z) = - \sum_{m>0} \frac{[rm]_x}{m[r^*m]_x} \beta_{-m}^1 z^m, \quad Q^*(z) = \sum_{m>0} \frac{[rm]_x}{m[r^*m]_x} \beta_m^1 z^{-m}, \quad (3.13)$$

$$R_+^j(z) = \sum_{m>0} \frac{[rm]_x}{m[r^*m]_x} (\beta_{-m}^j - \beta_{-m}^{j+1}) x^{jm} z^m, \quad S_+^j(z) = - \sum_{m>0} \frac{[rm]_x}{m[r^*m]_x} (\beta_m^j - \beta_m^{j+1}) x^{-jm} z^{-m}. \quad (3.14)$$

Let us set the basic operators $V(z), E_{\alpha_1}(z), E_{\alpha_2}(z)$ on the Fock space $\mathcal{F}_{l,k}$.

$$V(z) = z^{\frac{r}{3r^*}} e^{i\sqrt{\frac{r}{r^*}} Q_{\bar{\epsilon}_1}} z^{\sqrt{\frac{r}{r^*}} P_{\bar{\epsilon}_1}} e^{P^*(z)} e^{Q^*(z)}, \quad (3.15)$$

$$F_{\alpha_j}(z) = z^{\frac{r}{r^*}} e^{-i\sqrt{\frac{r}{r^*}} Q_{\alpha_j}} z^{-\sqrt{\frac{r}{r^*}} P_{\alpha_j}} e^{R_+^j(z)} e^{S_+^j(z)}. \quad (3.16)$$

We give a free field realization of the type-II vertex operators.

$$\Psi_1^*(z) = V(z), \quad (3.17)$$

$$\Psi_2^*(z) = \oint_{\tilde{C}_1} \frac{dw_1}{w_1} V(z) E_{\alpha_1}(w_1) \frac{[v_1 - u - \frac{1}{2} + \pi_{1,2}]^*}{[v_1 - u + \frac{1}{2}]^*}, \quad (3.18)$$

$$\Psi_3^*(z) = \oint \oint_{\tilde{C}_2} \frac{dw_1}{w_1} \frac{dw_2}{w_2} V(z) E_{\alpha_1}(w_1) E_{\alpha_2}(w_2) \frac{[v_1 - u - \frac{1}{2} + \pi_{1,3}]^*}{[v_1 - u + \frac{1}{2}]^*} \frac{[v_2 - v_1 - \frac{1}{2} + \pi_{2,3}]^*}{[v_2 - v_1 + \frac{1}{2}]^*}. \quad (3.19)$$

The integration contours \tilde{C}_1 encloses the poles at $w_1 = x^{-1+2sr^*} z, (s = 0, 1, 2, \dots)$, but not the poles at $w_1 = x^{1-2sr^*} z, (s = 0, 1, 2, \dots)$. The integration of $\Psi_3^*(z)$ is carried out in the order w_2, w_1 along the contours \tilde{C}_2 which encloses the poles the poles at $w_{j+1} = x^{-1+2sr^*} w_j, (s = 0, 1, 2, \dots; j = 1, 2)$, but not the poles at $w_{j+1} = x^{1-2sr^*} w_j, (s = 0, 1, 2, \dots; j = 1, 2)$. Here we set $w_0 = z$. Let us set the type-II Boltzmann weights $W^* \left(\begin{array}{cc|c} a & b & u \\ c & d & \end{array} \right)$ are given by

$$W^* \left(\begin{array}{cc|c} a + 2\bar{\epsilon}_\mu & a + \bar{\epsilon}_\mu & u \\ a + \bar{\epsilon}_\mu & a & \end{array} \right) = r_1^*(u), \quad (3.20)$$

$$W^* \left(\begin{array}{cc|c} a + \bar{\epsilon}_\mu + \bar{\epsilon}_\nu & a + \bar{\epsilon}_\mu & u \\ a + \bar{\epsilon}_\nu & a & \end{array} \right) = r_1^*(u) \frac{[u]^* [a_{\mu,\nu} - 1]^*}{[u - 1]^* [a_{\mu,\nu}]^*}, \quad (3.21)$$

$$W^* \left(\begin{array}{cc|c} a + \bar{\epsilon}_\mu + \bar{\epsilon}_\nu & a + \bar{\epsilon}_\nu & u \\ & a + \bar{\epsilon}_\nu & a \end{array} \right) = r_1^*(u) \frac{[u - a_{\mu,\nu}]^*[1]^*}{[u - 1]^*[a_{\mu,\nu}]^*}. \quad (3.22)$$

Otherwise are zero. The function $r_1^*(u)$ is given by

$$r_1^*(u) = z^{\frac{r}{r^*} \frac{2}{3}} \frac{h_1^*(z^{-1})}{h^*(z)}, \quad h_1^*(z) = \frac{(z; x^{2r^*}, x^6)_\infty (x^4 z; x^{2r^*}, x^6)_\infty}{(x^{2r^*} z; x^{2r^*}, x^6)_\infty (x^6 z; x^{2r^*}, x^6)_\infty}. \quad (3.23)$$

The Boltzmann weights satisfy the following relations.

(1) Yang-Baxter equation :

$$\begin{aligned} & \sum_g W^* \left(\begin{array}{cc|c} d & e & u_1 \\ & c & g \end{array} \right) W^* \left(\begin{array}{cc|c} c & g & u_2 \\ & b & a \end{array} \right) W^* \left(\begin{array}{cc|c} e & f & u_1 - u_2 \\ & g & a \end{array} \right) \\ &= \sum_g W^* \left(\begin{array}{cc|c} g & f & u_1 \\ & b & a \end{array} \right) W^* \left(\begin{array}{cc|c} d & e & u_2 \\ & g & f \end{array} \right) W^* \left(\begin{array}{cc|c} d & g & u_1 - u_2 \\ & c & b \end{array} \right). \end{aligned} \quad (3.24)$$

(2) The first inversion relation :

$$\sum_g W^* \left(\begin{array}{cc|c} c & g & -u \\ & b & a \end{array} \right) W^* \left(\begin{array}{cc|c} c & d & u \\ & g & a \end{array} \right) = \delta_{b,d}. \quad (3.25)$$

(3) The second inversion relation :

$$\sum_g G_g^* W^* \left(\begin{array}{cc|c} g & b & 3 - u \\ & d & c \end{array} \right) W^* \left(\begin{array}{cc|c} g & d & u \\ & b & a \end{array} \right) = \delta_{a,c} \frac{G_b^* G_d^*}{G_a^*}. \quad (3.26)$$

where $G_a^* = [a_{1,2}]^*[a_{1,3}]^*[a_{2,3}]^*$. The type-II vertex operators satisfy the following relations.

$$\Phi_{\mu_1}^*(z_1) \Phi_{\mu_2}^*(z_2) = \sum_{\epsilon_{\mu'_1} + \epsilon_{\mu'_2} = \epsilon_{\mu_1} + \epsilon_{\mu_2}} W^* \left(\begin{array}{cc|c} k + \bar{\epsilon}_{\mu_1} + \bar{\epsilon}_{\mu_2} & k + \bar{\epsilon}_{\mu'_1} & u_2 - u_1 \\ & k + \bar{\epsilon}_{\mu_2} & k \end{array} \right) \Phi_{\mu'_2}^*(z_2) \Phi_{\mu'_1}^*(z_1). \quad (3.27)$$

The type-I and type-II vertex operators commute modulo the "energy function" $\chi(z)$.

$$\Phi_{\mu_1}(z_1) \Psi_{\mu_2}^*(z_2) = \chi(z_1/z_2) \Psi_{\mu_2}^*(z_2) \Phi_{\mu_1}(z_1). \quad (3.28)$$

Here we have set

$$\chi(z) = z^{\frac{2}{3}} \frac{(x^5 z^{-1}; x^6)_\infty (xz; x^6)_\infty}{(xz^{-1}; x^6)_\infty (x^5 z; x^6)_\infty}. \quad (3.29)$$

4 Boundary state

In this section we construct the free field realization of the ground state of the boundary transfer matrix of the $U_{q,p}(\widehat{sl}(3, \mathbb{C}))$ ABF model, which give higher-rank generalization of [7]. We construct the free field realization of the ground state on the Fock space $\mathcal{F}_{l,k}$. Precisely the Fock space $\mathcal{F}_{l,k}$ is different from the space of the state $\mathcal{H}_{l,k}$. The space of state $\mathcal{H}_{l,k}$ is obtained by the cohomological argument from the Fock space $\mathcal{F}_{l,k}$ by using the so-called screening operators. In this paper we omit the detailed of the screening operators and consider the operators acting on the Fock space $\mathcal{F}_{l,k}$.

4.1 Bogoliubov transformation

The commutation relations of bosons β_m^j are not symmetric. Hence it is convenient to introduce new generators of bosons α_m^1, α_m^2 whose commutation relations are symmetric.

$$\alpha_m^1 = x^{-m}(\beta_m^1 - \beta_m^2), \quad \alpha_m^2 = x^{-2m}(\beta_m^2 - \beta_m^3). \quad (4.1)$$

They satisfy the following commutation relations.

$$[\alpha_m^j, \alpha_n^k] = \begin{cases} m \frac{[(r-1)m]_x}{[rm]_x} \frac{[2m]_x}{[m]_x} \delta_{m+n,0} & (j=k) \\ -m \frac{[(r-1)m]_x}{[rm]_x} \delta_{m+n,0} & (j \neq k). \end{cases} \quad (4.2)$$

Let us set

$$F_0 = - \sum_{m>0} \frac{1}{m} \frac{[rm]_x}{[(r-1)m]_x [3m]_x} ([2m]_x \alpha_{-m}^1 \alpha_{-m}^1 + 2[m]_x \alpha_{-m}^1 \alpha_{-m}^2 + [2m]_x \alpha_{-m}^2 \alpha_{-m}^2). \quad (4.3)$$

The adjoint action of e^{F_0} has the effect of a Bogoliubov transformation,

$$e^{-F_0} \alpha_m^j e^{F_0} = \alpha_m^j - \alpha_{-m}^j \quad (m > 0, j = 1, 2), \quad (4.4)$$

$$e^{-F_0} \alpha_{-m}^j e^{F_0} = \alpha_{-m}^j \quad (m > 0, j = 1, 2), \quad (4.5)$$

and

$$e^{-F_0} \beta_m^1 e^{F_0} = -\beta_{-m}^1 \quad (m > 0), \quad (4.6)$$

$$e^{-F_0} \beta_m^2 e^{F_0} = (x^{2m} - 1) \beta_{-m}^1 - x^{2m} \beta_{-m}^2 \quad (m > 0), \quad (4.7)$$

$$e^{-F_0} \beta_{-m}^j e^{F_0} = \beta_{-m}^j \quad (m > 0, j = 1, 2). \quad (4.8)$$

Proposition 4.1

$$e^{-F_0} e^{Q(z)} e^{F_0} = \sqrt{\frac{(x^{2r+4} z^{-2}; x^{2r}, x^6)_\infty (x^2 z^{-2}; x^{2r}, x^6)_\infty}{(x^{2r} z^{-2}; x^{2r}, x^6)_\infty (x^6 z^{-2}; x^{2r}, x^6)_\infty}} e^{P(1/z)} e^{Q(z)}, \quad (4.9)$$

$$e^{-F_0} e^{S_-^j(z)} e^{F_0} = \sqrt{\frac{(1-z^{-2})(x^{2r-2} z^{-2}; x^{2r})_\infty}{(x^2 z^{-2}; x^{2r})_\infty}} e^{R_-^j(1/z)} e^{S_-^j(z)} \quad (j = 1, 2). \quad (4.10)$$

4.2 Boundary state

Let us set the bosonic operator F by

$$F = F_0 + F_1, \quad (4.11)$$

where F_0 is given in (4.3) and F_1 is given by

$$F_1 = \sum_{m>0} (D_1(m)\beta_{-m}^1 + D_2(m)\beta_{-m}^2). \quad (4.12)$$

Here we set

$$\begin{aligned} D_1(m) &= \frac{x^{-m}[(r-2c+2-2\pi_{1,3})m]_x - [(r-2c-1)m]_x + x^{(r-2c-2\pi_{1,2})m}[m]_x}{m[(r-1)m]_x} \\ &\quad - \theta_m \left(\frac{x^{-\frac{m}{2}}[m]_x [rm/2]_x^+}{m[(r-1)m/2]_x} \right), \end{aligned} \quad (4.13)$$

$$D_2(m) = \frac{x^m[(r-2c+2-2\pi_{1,3})m]_x - [(r-2c-2\pi_{1,2})m]_x}{m[(r-1)m]_x} - \theta_m \left(\frac{x^m[rm/2]_x^+ [m/2]_x}{m[(r-1)m/2]_x} \right), \quad (4.14)$$

where

$$[a]_x^+ = a^x + a^{-x}, \quad \theta_m(x) = \begin{cases} x & (m : \text{even}) \\ 0 & (m : \text{odd}) \end{cases}.$$

Let us set the vector

$$|B\rangle_{l,k} = e^F |l, k\rangle. \quad (4.15)$$

We call this vector $|B\rangle_{l,k}$ the boundary state.

Proposition 4.2 *The boundary state $|B\rangle_{l,k}$ has the following properties.*

$$e^{Q(z)} |B\rangle_{l,k} = h(z) e^{P(1/z)} |B\rangle_{l,k}, \quad (4.16)$$

$$e^{S_-^j(w)} |B\rangle_{l,k} = g_j(w) e^{R_-^j(1/w)} |B\rangle_{l,k}, \quad (j = 1, 2). \quad (4.17)$$

Here the fuctions $h(z), g_j(w) (j = 1, 2)$ are given by

$$\begin{aligned} h(z) &= \frac{(x^{2r+4}z^{-2}; x^{2r}, x^{12})_\infty (x^8z^{-2}; x^{2r}, x^{12})_\infty}{(x^{12}z^{-2}; x^{2r}, x^{12})_\infty (x^{2r}z^{-2}; x^{2r}, x^{12})_\infty} \\ &\times \frac{(x^{2r+2c+4}z^{-1}; x^{2r}, x^{12})_\infty (x^{2c+2}z^{-1}; x^{2r}, x^{12})_\infty}{(x^{2r-2c}z^{-1}; x^{2r}, x^{12})_\infty (x^{2c+6}z^{-1}; x^{2r}, x^{12})_\infty} \\ &\times \frac{(x^{2r+6-2c+2\pi_{1,2}}z^{-1}; x^{2r}, x^{12})_\infty (x^{2c+2\pi_{1,2}}z^{-1}; x^{2r}, x^{12})_\infty}{(x^{2r+4-2c-2\pi_{1,2}}z^{-1}; x^{2r}, x^{12})_\infty (x^{2c+2+2\pi_{1,2}}z^{-1}; x^{2r}, x^{12})_\infty} \end{aligned}$$

$$\times \frac{(x^{2r+6-2c+2\pi_{1,3}}z^{-1}; x^{2r}, x^{12})_{\infty} (x^{2c+2\pi_{1,3}}z^{-1}; x^{2r}, x^{12})_{\infty}}{(x^{2r+4-2c-2\pi_{1,3}}z^{-1}; x^{2r}, x^{12})_{\infty} (x^{2c+2+2\pi_{1,3}}z^{-1}; x^{2r}, x^{12})_{\infty}}, \quad (4.18)$$

$$g_1(w) = (1 - 1/w^2) \frac{(x^{2c+1}/w; x^{2r})_{\infty} (x^{2r-2c-2\pi_{1,2}+1}/w; x^{2r})_{\infty}}{(x^{2c+2\pi_{1,2}-1}/w; x^{2r})_{\infty} (x^{2r-2c-1}/w; x^{2r})_{\infty}}, \quad (4.19)$$

$$g_2(w) = (1 - 1/w^2) \frac{(x^{2c+2\pi_{1,2}}/w; x^{2r})_{\infty} (x^{2r-2c+2-2\pi_{1,3}}/w; x^{2r})_{\infty}}{(x^{2c+2\pi_{1,3}-2}/w; x^{2r})_{\infty} (x^{2r-2c-2\pi_{1,2}}/w; x^{2r})_{\infty}}. \quad (4.20)$$

The parameters $l, k \in P$ are determined by the boundary conditions. The parameter k represents the central height. The parameter l represents the asymptotic boundary height. In what follows we consider the case $k = l \in P$ for simplicity. For more general $l, k \in P$, there exist similar boundary state. We construct the eigenvector of the commuting boundary transfer matrix, $[T_B^{(k)}(z_1), T_B^{(k)}(z_2)] = 0$. The following is the main result of this paper.

Theorem 4.3 *The boundary state $|B\rangle_{k,k}$ is the eigenvector of boundary transfer matrix $T_B^{(k)}(z)$.*

$$T_B^{(k)}(z)|B\rangle_{k,k} = |B\rangle_{k,k}. \quad (4.21)$$

Corollary 4.4 *Using the type-II vertex operators, we get general eigenvectors of the boundary transfer matrix $T_B^{(k)}(z)$.*

$$\begin{aligned} & T_B^{(k)}(z) \cdot \Psi_{\mu_1}^*(\xi_1) \cdots \Psi_{\mu_M}^*(\xi_M) |B\rangle_{k,k} \\ &= \prod_{j=1}^M \chi(1/z\xi_j) \chi(\xi_j/z) \cdot \Psi_{\mu_1}^*(\xi_1) \cdots \Psi_{\mu_M}^*(\xi_M) |B\rangle_{k,k}, \end{aligned} \quad (4.22)$$

where $\chi(z)$ is given by (3.29).

Here we sketch proof of main theorem 4.3. Acting the vertex operator $\Phi^{(k+\bar{\epsilon}_j, k)}(z)$ to the condition $T_B^{(k)}|B\rangle_{k,k} = |B\rangle_{k,k}$ from the left, we get the following necessary and sufficient condition.

$$z^{\frac{1}{r}(-\frac{r-1}{3}+(\bar{\epsilon}_1|k))} h(z)[c-u][\pi_{1,j}+c+u] \Phi_j(z^{-1}) |B\rangle_{k,k} = (z \leftrightarrow z^{-1}) \quad (j=1, 2, 3). \quad (4.23)$$

Here we used the inversion relation of the type-I vertex operators. In the case of $j=1$, after some calculation, LHS becomes the following

$$h(z)h(z^{-1})[c-u][c+u]e^{P(z)+P(1/z)}|B\rangle_{k,k},$$

which is invariant $z \leftrightarrow z^{-1}$. Hence the relation (4.23) for $j = 1$ holds. In the case of $j = 2, 3$, after some calculation as similar as [9], the relation (4.23) are reduced to the following theta identity.

$$\begin{aligned} & \frac{[v+k+c-\frac{1}{2}][v-c-\frac{1}{2}]}{[-v+k+c-\frac{1}{2}][-v-c-\frac{1}{2}]} \\ = & \frac{[c-u][k+c+u][u-v+\frac{1}{2}-k][-v-u+\frac{1}{2}]-[c+u][k+c-u][-u-v+\frac{1}{2}-k][-v+u+\frac{1}{2}]}{[c-u][k+c+u][u+v+\frac{1}{2}-k][v-u+\frac{1}{2}]-[c+u][k+c-u][-u+v+\frac{1}{2}-k][v+u+\frac{1}{2}]}. \end{aligned} \quad (4.24)$$

Later the author will write complete proof of this theorem, $U_{q,p}(\widehat{sl}(N, \mathbb{C}))$ version and generalization of asymptotic boundary condition $l \in P$ in the another place [15]. Our proof is different from those given in [7] even for $U_{q,p}(\widehat{sl}(2, \mathbb{C}))$ case.

Acknowledgement

The author would like to thank the organizing committee of VIII-th International Workshop Lie theory and its applications in Physics, held in Varna, Bulgaria 2009. This work is partly supported by the Grant-in Aid for Scientific Research **C**(21540228) from Japan Society for the Promotion of Science.

A Some formule

$$\begin{aligned} h(z) &= \exp \left(- \sum_{m>0} \frac{1}{2m} \frac{[(r-1)m]_x [2m]_x}{[rm]_x [3m]_x} z^{-2m} - \sum_{m>0} \frac{[(r-1)m]_x}{[rm]_x} \left(\frac{[2m]_x}{[3m]_x} D_1(m) - \frac{x^{-3m} [m]_x}{[3m]_x} D_2(m) \right) z^{-m} \right), \\ g_1(z) &= \exp \left(- \sum_{m>0} \frac{1}{2m} \frac{[(r-1)m]_x (x^m + x^{-m})}{[rm]_x} z^{-2m} + \sum_{m>0} \frac{[(r-1)m]_x}{[rm]_x} (D_1(m) - x^{-2m} D_2(m)) z^{-m} \right), \\ g_2(z) &= \exp \left(- \sum_{m>0} \frac{1}{2m} \frac{[(r-1)m]_x (x^m + x^{-m})}{[rm]_x} z^{-2m} + \sum_{m>0} \frac{[(r-1)m]_x}{[rm]_x} D_2(m) x^{-m} z^{-m} \right). \end{aligned}$$

References

- [1] B.Davis, O.Foda, M.Jimbo, T.Miwa and A.Nakayashiki, Diagonalization of the XXZ Hamiltonian by vertex operators, *Commun.Math.Phys.* **151**, 89- (1993).

- [2] M.Jimbo and T.Miwa, Algebraic Analysis of Solvable Lattice Models, CBMS Regional Conference Series in Mathematics, Vol **85**, Providence, RI:AMS, (1994).
- [3] E.K.Sklyanin, Boundary condition for integrable quantum system, *J.Phys.***A21**, 2375- (1988).
- [4] G.E.Andrews, R.J.Baxter and P.J.Forrester, Eight-vertex SOS model and generalized Rogers-Ramanujan-type identities, *J.Stat.Phys.***35**, 193- (1984).
- [5] M.Jimbo, T.Miwa and M.Okado, Local state probabilities of solvable lattice models : an $A_{n-1}^{(1)}$ family, *Nucl.Phys.***B300** [FS22], 74- (1988).
- [6] M.Jimbo,R.Kedem,T.Kojima,H.Konno and T.Miwa, XXZ chain with a boundary, *Nucl.Phys.***B441**[FS], 437- (1995).
- [7] T.Miwa and R.Weston, Boundary ABF models, *Nucl.Phys.***B486**[PM], 517- (1997).
- [8] M.T.Batchelor, V.Fridkin, A.Kuniba, Y.K.Zhou, Solutions of the reflection equation for face and vertex models associated with $A_n^{(1)}, B_n^{(1)}, C_n^{(1)}$ and $A_n^{(2)}$, *Phys.Lett.***B376** no.4, 266- (1996).
- [9] H.Furutsu and T.Kojima, The $U_q(\widehat{sl_n})$ analogue of the XXZ chain with a boundary, *J.Math.Phys.* **41**, no.7, 4413- (2000).
- [10] S.Lukyanov and Ya.Pugai, Multi-point local height probabilities in the integrable RSOS model,*Nucl.Phys.***B473**[FS], 631- (1996).
- [11] Y.Asai, M.Jimbo, T.Miwa, Ya.Pugai, Bosonization of vertex operators for the $A_{n-1}^{(1)}$ face model, *J.Phys.***A29**, 6595- (1996).
- [12] H.Furutsu, T.Kojima and Y.Quano, Type-II vertex operators for the $A_{n-1}^{(1)}$ face model, *Int.J.Mod.Phys.***A15**,no.10, 1533- (2000).
- [13] M.Jimbo, H.Konno, S.Odake and J.Shiraishi : Elliptic Algebra $U_{q,p}(\widehat{sl_2})$: Drinfeld currents and vertex operators, *Commun.Math.Phys.***199**, 605- (1999).
- [14] T.Kojima and H.Konno : The Elliptic Algebra $U_{q,p}(\widehat{sl_N})$ and the Drinfeld Realization of the Elliptic Quantum Group $\mathcal{B}_{q,\lambda}(\widehat{sl_N})$, *Commun.Math.Phys.***237**, 405- (2003).
- [15] T.Kojima , in preparation.