

SCATTERING ABOVE ENERGY NORM OF SOLUTIONS OF A LOGLOG ENERGY-SUPERCRITICAL SCHRÖDINGER EQUATION WITH RADIAL DATA

TRISTAN ROY

ABSTRACT. We prove scattering of $\tilde{H}^k := \dot{H}^k(\mathbb{R}^n) \cap \dot{H}^1(\mathbb{R}^n)$ - solutions of the loglog energy-supercritical Schrödinger equation $i\partial_t u + \Delta u = |u|^{\frac{4}{n-2}} u \log^c(\log(10 + |u|^2))$, $0 < c < c_n$, $n \in \{3, 4\}$, with radial data $u(0) = u_0 \in \tilde{H}^k := \dot{H}^k(\mathbb{R}^n) \cap \dot{H}^1(\mathbb{R}^n)$, $k > \frac{n}{2}$. This is achieved, roughly speaking, by extending Bourgain's argument [1] (see also Grillakis [5]) and Tao's argument [10] in high dimensions.

1. INTRODUCTION

We shall study the solutions of the following Schrödinger equation in dimension n , $n \in \{3, 4\}$:

$$(1) \quad i\partial_t u + \Delta u = |u|^{\frac{4}{n-2}} u g(|u|)$$

with $g(|u|) := \log^c(\log(10 + |u|^2))$, $0 < c < c_n$ and ¹

$$(2) \quad c_n := \begin{cases} \frac{1}{5772}, & n = 3 \\ \frac{1}{8024}, & n = 4 \end{cases}$$

This equation has many connections with the following power-type Schrödinger equation, $p > 1$

$$(3) \quad i\partial_t v + \Delta v = |v|^{p-1} v$$

(3) has a natural scaling: if v is a solution of (3) with data $v(0) := v_0$ and if $\lambda \in \mathbb{R}$ is a parameter then $v_\lambda(t, x) := \frac{1}{\lambda^{\frac{2}{p-1}}} v\left(\frac{t}{\lambda^2}, \frac{x}{\lambda}\right)$ is also a solution of (3) but with data $v_\lambda(0, x) := \frac{1}{\lambda^{\frac{2}{p-1}}} v_0\left(\frac{x}{\lambda}\right)$. If $s_p := \frac{n}{2} - \frac{2}{p-1}$ then the \dot{H}^{s_p} norm of the initial data is invariant under the scaling: this is why (3) is said to be \dot{H}^{s_p} -critical. If $p = 1 + \frac{4}{n-2}$ then (3) is \dot{H}^1 (or energy) critical. The energy-critical Schrödinger equation

$$(4) \quad i\partial_t u + \Delta u = |u|^{\frac{4}{n-2}} u$$

has received a great deal of attention. Cazenave and Weissler [2] proved the local well-posedness of (4): given any $u(0)$ such that $\|u(0)\|_{\dot{H}^1} < \infty$ there exists, for

¹we shall prove global well-posedness and scattering of radial solutions to (1). The computations show that these properties hold for functions g that do not grow faster than $x \rightarrow \log^c \log(10 + |x|^2)$ with $c < c_n$ but not for functions g that grow faster (i.e $c \geq c_n$). The values of c_n are determined by technical computations but do not have a particular physical meaning.

some t_0 close to zero, a unique $u \in \mathcal{C}([0, t_0], \dot{H}^1) \cap L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, t_0])$ satisfying (4) in the sense of distributions

$$(5) \quad u(t) = e^{it\Delta} u(0) - i \int_0^t e^{i(t-t')\Delta} \left[|u(t')|^{\frac{4}{n-2}} u(t') \right] dt'$$

Bourgain [1] proved global existence and scattering of radial solutions in the class $\mathcal{C}(\mathbb{R}, \dot{H}^1) \cap L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\mathbb{R})$ in dimension $n = 3, 4$. He also proved this fact that for smoother solutions. Another proof was given by Grillakis [5] in dimension $n = 3$. The radial assumption for $n = 3$ was removed by Colliander-Keel-Staffilani-Takaoka-Tao [4]. This result was extended to $n = 4$ by Rickman-Visan [7] and to $n \geq 5$ by Visan [11]. If $p > 1 + \frac{4}{n-2}$ then $s_p > 1$ and we are in the energy supercritical regime. The global existence of \tilde{H}^k -solutions in this regime is an open problem. Since for all $\epsilon > 0$ there exists $c_\epsilon > 0$ such that $\left| |u|^{\frac{4}{n-2}} u \right| \lesssim \left| |u|^{\frac{4}{n-2}} u g(|u|) \right| \leq c_\epsilon \max(1, \| |u|^{\frac{4}{n-2} + \epsilon} u \|)$ then the nonlinearity of (1) is said to be barely supercritical.

In this paper we are interested in establishing global well-posedness and scattering of $\tilde{H}^k := \dot{H}^k(\mathbb{R}^n) \cap \dot{H}^1(\mathbb{R}^n)$ - solutions of (1) for $n \in \{3, 4\}$. First we prove a local-wellposed result. The local well-posedness theory for (1) and for \tilde{H}^k -solutions can be formulated as follows

Proposition 1. “Local well-posedness ” *Let $n \in \{3, 4\}$ and $k > \frac{n}{2}$. Let M be such that $\|u_0\|_{\tilde{H}^k} \leq M$. Then there exists $\delta := \delta(M) > 0$ small such that if $T_l > 0$ (T_l =time of local existence) satisfies*

$$(6) \quad \|e^{it\Delta} u_0\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \leq \delta$$

then there exists a unique

$$(7) \quad u \in \mathcal{C}([0, T_l], \tilde{H}^k) \cap L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l]) \cap L_t^{\frac{2(n+2)}{n}} D^{-1} L_x^{\frac{2(n+2)}{n}}([0, T_l]) \\ \cap L_t^{\frac{2(n+2)}{n}} D^{-k} L_x^{\frac{2(n+2)}{n}}([0, T_l])$$

such that

$$(8) \quad u(t) = e^{it\Delta} u_0 - i \int_0^t e^{i(t-t')\Delta} \left(|u(t')|^{\frac{4}{n-2}} u(t') g(|u(t')|) \right) dt'$$

is satisfied in the sense of distributions. Here $D^{-\alpha} L^r := \dot{H}^{\alpha, r}$ endowed with the norm $\|f\|_{D^{-\alpha} L^r} := \|D^\alpha f\|_{L^r}$.

This allows to define the notion of maximal time interval of existence I_{max} , that is the union of all the open intervals I containing 0 such that (8) holds in the class $\mathcal{C}(I, \tilde{H}^k) \cap L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I) \cap L_t^{\frac{2(n+2)}{n}} D^{-1} L_x^{\frac{2(n+2)}{n}}(I) \cap L_t^{\frac{2(n+2)}{n}} D^{-k} L_x^{\frac{2(n+2)}{n}}(I)$. Next we prove a criterion for global well-posedness:

Proposition 2. “Global well-posedness: criterion” *If $|I_{max}| < \infty$ then*

$$(9) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I_{max})} = \infty$$

These propositions are proved in Section 2. With this in mind, global well-posedness follows from an *a priori* bound of the form

$$(10) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([-T, T])} \leq f(T, \|u_0\|_{\tilde{H}^k})$$

for arbitrarily large time $T > 0$. In fact we shall prove that the bound does not depend on time T : this is the preliminary step to prove scattering.

The main result of this paper is:

Theorem 3. *The solution of (1) with radial data $u(0) := u_0 \in \tilde{H}^k$, $n \in \{3, 4\}$, $k > \frac{n}{2}$ and $0 < c < c_n$ exists for all time T . Moreover there exists a scattering state $u_{0,+} \in \tilde{H}^k$ such that*

$$(11) \quad \lim_{t \rightarrow \infty} \|u(t) - e^{it\Delta} u_{0,+}\|_{\tilde{H}^k} = 0$$

and there exists C depending only on $\|u_0\|_{\tilde{H}^k}$ such that

$$(12) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\mathbb{R})} \leq C(\|u_0\|_{\tilde{H}^k})$$

Remark 1. *This implies global regularity² since by the Sobolev embedding $\|u\|_{L_t^\infty L_x^\infty(\mathbb{R})} \lesssim \|u\|_{L_t^\infty \tilde{H}^k(\mathbb{R})}$ for $k > \frac{n}{2}$.*

We recall some estimates. The pointwise dispersive estimate is $\|e^{it\Delta} f\|_{L^\infty(\mathbb{R}^n)} \lesssim \frac{1}{|t|^{\frac{n}{2}}} \|f\|_{L^1(\mathbb{R}^n)}$. Interpolating with $\|e^{it\Delta} f\|_{L^2(\mathbb{R}^n)} = \|f\|_{L^2(\mathbb{R}^n)}$ we have the well-known generalized pointwise dispersive estimate:

$$(13) \quad \|e^{it\Delta} f\|_{L^p(\mathbb{R}^n)} \lesssim \frac{1}{|t|^{\frac{n}{2}(\frac{1}{2} - \frac{1}{p})}} \|f\|_{L^{p'}(\mathbb{R}^n)}$$

Here $2 \leq p \leq \infty$ and p' is the conjugate of p . We recall some useful Sobolev inequalities:

$$(14) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} \lesssim \|Du\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2n(n+2)}{n^2+4}}(J)}$$

and

$$(15) \quad \|u\|_{L_t^\infty L_x^\infty(J)} \lesssim \|u\|_{L_t^\infty \tilde{H}^k(J)}.$$

If u is a solution of $i\partial_t u + \Delta u = G$, $u(t=0) := u_0$ on J with $0 \in J$ and with data $u_0 \in H^k$ then the Strichartz estimates (see for example [6]) yield

$$(16) \quad \begin{aligned} & \|u\|_{L_t^\infty \dot{H}^j(J)} + \|D^j u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)} + \|D^j u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2n(n+2)}{n^2+4}}(J)} \\ & \lesssim \|D^j G\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}(J)} + \|u_0\|_{\dot{H}^j} \end{aligned}$$

²By global regularity we mean “if the data is radial and Schwartz, then the solution is infinitely differentiable for all time.” It is well-known that if for all time we have a finite bound of the L^∞ norm of the solution, then we have global regularity.

if $j \in \{1, k\}$; if $t_0 \in J$ then we write

$$(17) \quad u(t) = u_{l,t_0}(t) + u_{nl,t_0}(t)$$

with u_{l,t_0} denoting the linear part starting from t_0 , i.e

$$(18) \quad u_{l,t_0}(t) := e^{i(t-t_0)\Delta} u(t_0)$$

and u_{nl,t_0} denoting the nonlinear part starting from t_0 , i.e

$$(19) \quad u_{nl,t_0}(t) := -i \int_{t_0}^t e^{i(t-s)\Delta} G(s) ds.$$

If u is a \tilde{H}^k -solution of (1) on J with $k > \frac{n}{2}$ ³ and if $t \in J$, then it has a finite energy⁴

$$(20) \quad E(u(t)) := \frac{1}{2} \int_{\mathbb{R}^n} |\nabla u(t, x)|^2 + \int_{\mathbb{R}^n} F(u, \bar{u})(t, x) dx$$

with

$$(21) \quad F(z, \bar{z}) := \int_0^{|z|} t^{\frac{n+2}{n-2}} g(t) dt$$

Indeed

$$(22) \quad \begin{aligned} \left| \int_{\mathbb{R}^n} F(u, \bar{u})(t, x) dx \right| &\lesssim \|u(t)\|_{L^{\frac{2n}{n-2}}}^{\frac{2n}{n-2}} g(\|u(t)\|_{L^\infty}) \\ &\lesssim \|u(t)\|_{\dot{H}^1}^{\frac{2n}{n-2}} g(\|u(t)\|_{\tilde{H}^k}) : \end{aligned}$$

this follows from a simple integration by part

$$(23) \quad F(z, \bar{z}) \sim |z|^{\frac{2n}{n-2}} g(|z|)$$

combined with (15). A simple computation shows that the energy is conserved, or, in other words, that $E(u(t)) = E(u_0) = E$ ⁵. Let χ be a smooth, radial function supported on $|x| \leq 2$ such that $\chi(x) = 1$ if $|x| \leq 1$. If $x_0 \in \mathbb{R}^n$, $R > 0$ and u is an \tilde{H}^k solution of (1) then we define the mass within the ball $B(x_0, R)$

$$(24) \quad \text{Mass}(B(x_0, R), u(t)) := \left(\int_{B(x_0, R)} |u(t, x)|^2 dx \right)^{\frac{1}{2}}$$

Recall (see [10]) that^{6*}

$$(25) \quad \text{Mass}(B(x_0, R), u(t)) \lesssim R \sup_{t' \in [0, t]} \|\nabla u(t')\|_{L^2}$$

and that its derivative satisfies

³Let I be an interval. In the sequel we say that u is an \tilde{H}^k -solution on I if for all $(t_0, t) \in I^2$ such that $t_0 \leq t$ (7) and (8) hold with $[0, T_1]$ (resp. the interval of integration) replaced with I (resp. $[t_0, t]$).

⁴Hence the norm \tilde{H}^k controls the energy. In other words, it is above the energy norm.

⁵More precisely, the computation holds for smooth solution (i.e solutions in \tilde{H}^p with exponents p large enough). Then $E(u(t)) = E(u_0)$ holds for an \tilde{H}^k -solution by a standard approximation argument with smooth solutions.

⁶(26) also holds if u is a solution of the linear Schrödinger equation with data in \tilde{H}^k .

$$(26) \quad |\partial_t \text{Mass}(u(t), B(x_0, R))| \lesssim \frac{\sup_{t' \in [0, t]} \|\nabla u(t')\|_{L^2}}{R}$$

Now we set up some notation. We write $a \ll b$ if $a \leq \frac{1}{100}b$, $a \gg b$ if $a \geq 100b$ and $a \sim b$ if $\frac{1}{100}b \leq a \leq 100b$, $a \ll_E b$ if $a \leq \frac{1}{100 \max(1, E)^{100n}}b$ (Here n is the dimension of the space), $a \gg_E b$ if $a \geq 100 \max(1, E)^{100n}b$, $a \lesssim_E b$ if $a \leq 100(\max(1, E))^{100n}b$ and $a \sim_E b$ if $\frac{1}{100} \max(1, E)^{100n}b \leq a \leq 100 \max(1, E)^{100n}b$. We say that \tilde{C} is the constant determined by $a \lesssim b$ (or $a \lesssim_E b$) if it is the smallest constant C (or $C = C(E)$) that satisfies $a \leq Cb$. If u is a function then u_h is the function defined by $x \rightarrow u_h(x) := u(x - h)$. If $x \in \mathbb{R}$ then $x+ = x + \epsilon$ for $0 < \epsilon \ll 1$. Let $j \in \mathbb{N}$. If J is an interval then we define

$$(27) \quad Q_j(J, u) := \|u\|_{L_t^\infty \tilde{H}^j(J)} + \|Du\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} + \|D^j u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} + \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)}$$

Remark 2. If $j = k$ then we write $Q(J, u)$ instead of $Q_j(J, u)$.

If X is a normed vector space endowed with the norm $\|\cdot\|_X$ and $R > 0$, then $\mathcal{B}(X, R) := \{y \in X, \|y\|_X \leq R\}$.

Now we explain how this paper is organized. In Section 3 we prove the main result of this paper, i.e Theorem 3. The proof relies upon the following bound of $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}}$ on an arbitrarily long time interval

Proposition 4. “Bound of $L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}$ norm” Let u be a radial \tilde{H}^k -solution of (1) on a compact interval J . There exist three constants $C_1 \gg_E 1$, $C_2 \gg_E 1$, and $a_n > 0$ such that if $\|u\|_{L_t^\infty \tilde{H}^k(J)} \leq M$ for some $M \gg 1$, then

$$(28) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} \leq (C_1 g^{a_n}(M))^{C_2 g^{b_n+}(M)}$$

with b_n such that

$$(29) \quad b_n := \begin{cases} 5772, & n = 3 \\ \frac{8024}{3}, & n = 4 \end{cases}$$

By combining this bound with the Strichartz estimates, we can prove, by induction, that in fact this norm and other norms (such as $\|u\|_{L_t^\infty \tilde{H}^k(J)}$, $\|Du\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)}$, etc.) can be bounded only by a constant only depending on the norm of the initial data for $\frac{n}{2} < k < \frac{2+n}{n-2}$. This already shows (by Proposition 2) global well-posedness of the \tilde{H}^k -solutions of (1) for this range of ks . In fact we show that that these bounds imply a linear asymptotic behaviour of the solutions, or, in other words, scattering. Proposition 9 allows to prove global well-posedness and scattering of solutions of (1) for the full range, i.e $k > \frac{n}{2}$. The rest of the paper is devoted to prove Proposition 4. First we prove a weighted Morawetz-type estimate: it shows, roughly speaking, that the $L_t^{\frac{2n}{n-2}} L_x^{\frac{2n}{n-2}}$ norm of the solution cannot concentrate around the

origin on long time intervals. Then we modify arguments from Bourgain [1], Grilakis [5] and mostly Tao [10]. We divide J into subintervals $(J_l)_{1 \leq l \leq L}$ such that the $L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}$ norm of u is small but also substantial. We prove that, on most of these intervals, the mass on at least one ball concentrates. By using the radial assumption, we prove that in fact the mass on a ball centered at the origin concentrates. This implies, by using the Morawetz-type estimate that there exists a significant number of intervals (in comparison with L) that concentrate around a point \bar{t} and such that the mass concentrates around the origin. But, by Hölder, this implies that L is finite: if not it would violate the fact that the $L_t^\infty L_x^{\frac{2n}{n-2}}$ norm of the solution is bounded by some power of the energy. The process involves several tuning parameters. The fact that these parameters depend on the energy is not important; however, it is crucial to understand how they depend on $g(M)$ since this will play a prominent role in the choice of c_n for which we have global well-posedness and scattering of \tilde{H}^k -solutions of (1) (with $g(|u|) := \log^c(\log(10 + |u|^2))$ and $c < c_n$): see the proof of Theorem 3, Section 3.

2. LOCAL WELL-POSEDNESS AND CRITERION FOR GLOBAL WELL-POSEDNESS

In this section we prove Proposition 1 and Proposition 2.

2.1. Proof of Proposition 1. This is done by a modification of standard arguments to establish a local well-posedness theory for (4).

We define

$$(30) \quad X := \mathcal{C}([0, T_l], \tilde{H}^k) \cap L_t^{\frac{2(n+2)}{n}} D^{-1} L_x^{\frac{2(n+2)}{n}}([0, T_l]) \cap L_t^{\frac{2(n+2)}{n}} D^{-k} L_x^{\frac{2(n+2)}{n}}([0, T_l]) \cap L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])$$

and, for some $C > 0$ to be chosen later,

$$(31) \quad X_1 := \mathcal{B} \left(\mathcal{C}([0, T_l], \tilde{H}^k) \cap L_t^{\frac{2(n+2)}{n}} D^{-1} L_x^{\frac{2(n+2)}{n}}([0, T_l]) \cap L_t^{\frac{2(n+2)}{n}} D^{-k} L_x^{\frac{2(n+2)}{n}}([0, T_l]), 2CM \right)$$

and

$$(32) \quad X_2 := \mathcal{B} \left(L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l]), 2\delta \right)$$

$X_1 \cap X_2$ is a closed space of the Banach space X : therefore it is also a Banach space.

$$(33) \quad \begin{aligned} \Psi &:= X_1 \cap X_2 \rightarrow X_1 \cap X_2 \\ u \rightarrow \Psi(u) : t &\rightarrow \Psi(u)(t) := e^{it\Delta} u_0 - i \int_0^t e^{i(t-t')\Delta} \left(|u|^{\frac{4}{n-2}}(t') u(t') g(|u(t')|) \right) dt' \end{aligned}$$

- Ψ maps $X_1 \cap X_2$ to $X_1 \cap X_2$

By the fractional Leibnitz rule (see Appendix with $F(x) := \log^c \log(10 + x)$, $G(x, \bar{x}) := |x|^{\frac{4}{n-2}} x$ and $\beta := \frac{4}{n-2}$) and (15) we have

$$(34) \quad \left\| D^j(|u|^{\frac{4}{n-2}}ug(|u|)) \right\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \lesssim \|D^j u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}([0, T_l])} \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} g(\|u\|_{L_t^\infty \tilde{H}^k([0, T_l])})$$

if $j = 1$. Now assume that $j = k$. By Proposition 8 we get

$$\left\| D^j(|u|^{\frac{4}{n-2}}ug(|u|)) \right\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \lesssim \delta^{\frac{4}{n-2}} \langle M \rangle^{\bar{C}}.$$

Therefore ⁷ by the Strichartz estimates (16) and the Sobolev embedding (15) we have

$$(35) \quad \|u\|_{L_t^\infty \tilde{H}^k([0, T_l])} + \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}([0, T_l])} + \|D^k u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}([0, T_l])} \lesssim M + \delta^{\frac{4}{n-2}} \langle M \rangle^{\bar{C}}$$

Moreover

$$(36) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} - \|e^{it\Delta} u_0\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \lesssim \left\| D(|u|^{\frac{4}{n-2}}ug(|u|)) \right\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \lesssim \delta^{\frac{4}{n-2}} M g(M)$$

so that

$$(37) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} - \delta \lesssim \delta^{\frac{4}{n-2}} M g(M)$$

Therefore if let C be equal to the maximum of the constants determined by (35) and (37), then we see that $\Psi(X_1 \cap X_2) \subset X_1 \cap X_2$, provided that $\delta = \delta(M) > 0$ is small enough.

- Ψ is a contraction. Indeed, by the fundamental theorem of calculus and Proposition 8

$$(38) \quad \begin{aligned} & \|\Psi(u) - \Psi(v)\|_X \\ & \lesssim \sum_{j=1} \left\| D^j(u - v) \right\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \left(g\left(\|u\|_{L_t^\infty \tilde{H}^k([0, T_l])}\right) + g\left(\|v\|_{L_t^\infty \tilde{H}^k([0, T_l])}\right) \right) \\ & \quad \left(\|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} + \|v\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \right) \\ & \quad + \left(\sum_{\substack{j \in \{1, k\} \\ \delta_1 + \delta_2 = \frac{4}{n-2}}} \left\| D^j(w_\tau^{\delta_1} \overline{w_\tau}^{\delta_2} g(|w_\tau|)) \right\|_{L_t^{\frac{n+2}{3}} L_x^{\frac{n+2}{3}}([0, T_l])} \right. \\ & \quad \left. + \sum_{\substack{j \in \{1, k\} \\ \delta_1 + \delta_2 = \frac{4}{n-2} \\ \delta_3 + \delta_4 = 2}} \left\| D^j(w_\tau^{\delta_1} \overline{w_\tau}^{\delta_2} w_\tau^{\delta_3} \overline{w_\tau}^{\delta_4} \tilde{g}'(|w_\tau|^2)) \right\|_{L_t^{\frac{n+2}{3}} L_x^{\frac{n+2}{3}}([0, T_l])} \right) \|u - v\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T_l])} \\ & \lesssim \delta \langle M \rangle^{\bar{C}} \|u - v\|_X \end{aligned}$$

⁷In the sequel we allow \bar{C} to change from one line to the other one.

and if $\delta = \delta(M) > 0$ is small enough then Ψ is a contraction.

2.2. Proof of Proposition 2. Again, this is done by a modification of standard arguments used to prove a criterion of global well-posedness of (3) (See [9] for similar arguments). Let $\frac{n}{2} < k' < \min\left(k, \frac{n+2}{n-2}\right)$. Assume that $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I_{max})} < \infty$. Then

- First step: $Q_{k'}(I_{max}, u) < \infty$. Indeed, let $0 < \epsilon \ll 1$. Let C be the constant determined by \lesssim in (16). We may assume without loss of generality that $C \gg \max\left(\|u_0\|_{\tilde{H}^k}^{100}, \frac{1}{\|u_0\|_{\tilde{H}^k}^{100}}\right)$. We divide $I_{max} \cap [0, \infty)$ into subintervals $(I_j)_{1 \leq j \leq J}$ such that $0 \in I_1$,

$$(39) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I_j)} = \frac{\epsilon}{g^{\frac{n-2}{4}} ((2C)^j \|u_0\|_{\tilde{H}^k})}$$

if $1 \leq j < J$ and

$$(40) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I_J)} \leq \frac{\epsilon}{g^{\frac{n-2}{4}} ((2C)^J \|u_0\|_{\tilde{H}^k})}$$

Notice that such a partition always exists since, for J large enough,

$$(41) \quad \sum_{j=1}^{J-1} \frac{\epsilon^{\frac{2(n+2)}{n-2}}}{g^{\frac{n-2}{2}} ((2C)^j \|u_0\|_{\tilde{H}^k})} \gtrsim \sum_{j=1}^{J-1} \frac{1}{\log((2C)^j \|u_0\|_{\tilde{H}^k})} = \sum_{j=1}^{J-1} \frac{1}{j \log(2C) + \log(\|u_0\|_{\tilde{H}^k})} \geq \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I_{max})}$$

By the fractional Leibnitz rule (see Appendix A) and (16) we have for some positive constant C'

$$(42) \quad \begin{aligned} Q_{k'}(I_1, u) &\leq C\|u_0\|_{\tilde{H}^{k'}} + C\|D(|u|^{\frac{4}{n-2}} ug(|u|))\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}(I_1)} + C\|D^{k'}(|u|^{\frac{4}{n-2}} ug(|u|))\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}(I_1)} \\ &\leq C\|u_0\|_{\tilde{H}^{k'}} + C' C(\|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(I_1)} + \|D^{k'} u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(I_1)}) \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I_1)} \\ &\quad g(\|u\|_{L_t^\infty \tilde{H}^{k'}(I_1)}) \\ &\leq C\|u_0\|_{\tilde{H}^{k'}} + 2C' C Q_{k'}(I_1, u) \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(I_1)} g(Q_{k'}(I_1, u)) \end{aligned}$$

and by a continuity argument⁸, $Q_{k'}(I_1, u) \leq 2C\|u_0\|_{\tilde{H}^{k'}}$. By iteration $Q_{k'}(I_j, u) \leq (2C)^j \|u_0\|_{\tilde{H}^{k'}}$. Therefore $Q_{k'}(I_{max}, u) < \infty$, proceeding similarly on $I_{max} \cap (-\infty, 0]$.

- Second step. We write $I_{max} = (a_{max}, b_{max})$. Choose $\bar{t} < b_{max}$ close enough to b_{max} so that $\|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}([\bar{t}, b_{max}])} \ll \delta$ and $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([\bar{t}, b_{max}])} \ll \delta$, with δ defined in Proposition 1. We have

⁸Let $K \subset I_1$. Then (42) also holds if I_1 is replaced with K .

$$\begin{aligned}
(43) \quad & \|e^{i(t-\bar{t})\Delta} u(\bar{t})\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([\bar{t}, b_{max}])} \leq \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([\bar{t}, b_{max}])} + \\
& C' C \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}([\bar{t}, b_{max}])} \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([\bar{t}, b_{max}])} \\
& g(\|u\|_{L_t^\infty \tilde{H}^{k'}(I_{max})}) \\
& \leq \frac{3\delta}{4}.
\end{aligned}$$

Also observe that $\|e^{i(t-\bar{t})\Delta} u(\bar{t})\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([\bar{t}, b_{max}])} \lesssim \|u(\bar{t})\|_{\dot{H}^1} < \infty$.

Hence by the monotone convergence theorem, there exists $\epsilon > 0$ such that $\|e^{i(t-\bar{t})\Delta} u(\bar{t})\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}[\bar{t}, b_{max}+\epsilon]} \leq \delta$. Hence contradiction with Proposition 1.

3. PROOF OF THEOREM 3

The proof makes use of Proposition 4 and is made of two steps:

- finite bound of $\|u\|_{L_t^\infty \tilde{H}^k(\mathbb{R})}$, $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\mathbb{R})}$, $\|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(\mathbb{R})}$ and $\|D^k u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(\mathbb{R})}$ for $\frac{n}{2} < k < \frac{n+2}{n-2}$. By time reversal symmetry⁹ and by monotone convergence it is enough to find, for all $T \geq 0$, a finite bound of all these norms restricted to $[0, T]$ and the bound should not depend on T . We define

$$(44) \quad \mathcal{F} := \left\{ T \in [0, \infty) : \sup_{t \in [0, T]} Q([0, t], u) \leq M_0 \right\}$$

We claim that $\mathcal{F} = [0, \infty)$ for M_0 , a large constant (to be chosen later) depending only on $\|u_0\|_{\tilde{H}^k}$. Indeed

- $0 \in \mathcal{F}$.
- \mathcal{F} is closed by continuity
- \mathcal{F} is open. Indeed let $T \in \mathcal{F}$. Then, by continuity there exists $\delta > 0$ such that for $T' \in [0, T+\delta]$ we have $Q([0, T']) \leq 2M_0$. In view of (28), this implies, in particular, that

$$(45) \quad \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([0, T'])} \leq C_1 (g^{a_n}(2M_0))^{C_2 g^{b_n} + (2M_0)}$$

Let $J := [0, a]$ be an interval. We get from (16), Proposition 7, and the Sobolev inequality $\|u\|_{L_t^\infty L_x^\infty(J)} \lesssim \|u\|_{L_t^\infty \tilde{H}^k(J)}$

⁹i.e if $t \rightarrow u(t, x)$ is a solution of (1) then $t \rightarrow \bar{u}(-t, x)$ is also a solution of (1).

$$\begin{aligned}
(46) \quad Q(J, u) &\lesssim \|u_0\|_{\tilde{H}^k} + \left(\|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)} + \|D^k u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)} \right) \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} \\
&g \left(\|u\|_{L_t^\infty \tilde{H}^k(J)} \right) \\
&\lesssim \|u_0\|_{\tilde{H}^k} + Q(J, u) \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} g(Q(J, u))
\end{aligned}$$

Let C be the constant determined by \lesssim in (46). We may assume without loss of generality that $C \gg \max \left(\|u_0\|_{\tilde{H}^k}^{100}, \frac{1}{\|u_0\|_{\tilde{H}^k}^{100}} \right)$. Let $0 < \epsilon \ll 1$. Notice that if J satisfies $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} = \frac{\epsilon}{g^{\frac{n-2}{4}} (2C \|u_0\|_{\tilde{H}^k})}$ then a simple continuity argument shows that

$$(47) \quad Q(J, u) \leq 2C \|u_0\|_{\tilde{H}^k}$$

We divide $[0, T']$ into subintervals $(J_i)_{1 \leq i \leq I}$ such that $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J_i)} = \frac{\epsilon}{g^{\frac{n-2}{4}} ((2C)^i \|u_0\|_{\tilde{H}^k})}$, $1 \leq i < I$ and $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J_I)} \leq \frac{\epsilon}{g^{\frac{n-2}{4}} ((2C)^I \|u_0\|_{\tilde{H}^k})}$. Notice that such a partition exists by (45), the definition of g^{10} and the following inequality

$$\begin{aligned}
(48) \quad (C_1 g^{a_n} (2M_0))^{C_2 g^{b_n} + (2M_0)} &\gtrsim \sum_{i=1}^{I-1} \frac{1}{g^{\frac{n+2}{2}} ((2C)^i \|u_0\|_{\tilde{H}^k})} \\
&\geq \sum_{i=1}^{I-1} \frac{1}{\log^{\frac{(n+2)c}{2}} (\log(10 + (2C)^{2i} \|u_0\|_{\tilde{H}^k}^2))} \\
&\gtrsim \sum_{i=1}^{I-1} \frac{1}{\log^{\frac{(n+2)c}{2}} (2i \log(2C) + 2 \log(\|u_0\|_{\tilde{H}^k}))} \\
&\gtrsim \|u_0\|_{\tilde{H}^k} \sum_{i=1}^{I-1} \frac{1}{i^{\frac{1}{2}}} \\
&\gtrsim \|u_0\|_{\tilde{H}^k} I^{\frac{1}{2}}
\end{aligned}$$

Moreover, by iterating the procedure in (46) and (47) we get

$$(49) \quad Q([0, T'], u) \lesssim (2C)^I \|u_0\|_{\tilde{H}^k}$$

Therefore by (48) there exists $C' = C'(\|u_0\|_{\tilde{H}^k})$

$$(50) \quad \log I \lesssim \log(C') + C_2 \log^{(b_n+c)}(\log(10 + 4M_0^2)) \log(C_1 \log^{a_n c}(\log(10 + 4M_0^2)))$$

and for $M_0 = M_0(\|u_0\|_{\tilde{H}^k})$ large enough

¹⁰Recall that $g(x) := \log^c \log(10 + x^2)$

$$(51) \quad \begin{aligned} & \log(C') + C_2 \log^{(b_n+)^c}(\log(10+4M_0^2)) \log(C_1 \log^{a_n c}(\log(10+4M_0^2))) \\ & \ll \log\left(\frac{\log\left(\frac{M_0}{\|u_0\|_{\tilde{H}^k}}\right)}{\log(2C)}\right) \end{aligned}$$

since (recall that $c < \frac{1}{b_n}$)

$$(52) \quad \frac{\log(C') + C_2 \log^{(b_n+)^c}(\log(10+4M_0^2)) \log(C_1 \log^{a_n c}(\log(10+4M_0^2)))}{\log\left(\frac{\log\left(\frac{M_0}{\|u_0\|_{\tilde{H}^k}}\right)}{\log(2C)}\right)} \rightarrow_{M_0 \rightarrow \infty} 0.$$

- Finite bound of $Q(\mathbb{R}, u)$ for all $k > \frac{n}{2}$: this follows from Proposition 9.
- Scattering: it is enough to prove that $e^{-it\Delta}u(t)$ has a limit as $t \rightarrow \infty$ in \tilde{H}^k . If $t_1 < t_2$ then by dualizing (16) with $G = 0$ (more precisely the estimate $\|D^j u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}([t_1, t_2])} \lesssim \|u_0\|_{\dot{H}^j}$) we get from Propositions 7 and 8

$$(53) \quad \begin{aligned} & \|e^{-it_1\Delta}u(t_1) - e^{-it_2\Delta}u(t_2)\|_{\tilde{H}^k} \\ & \lesssim \|D^k\left(|u|^{\frac{4}{n-2}}ug(|u|)\right)\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}([t_1, t_2])} + \|D\left(|u|^{\frac{4}{n-2}}ug(|u|)\right)\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}([t_1, t_2])} \\ & \lesssim \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_1, t_2])} \end{aligned}$$

and we conclude from the previous step that given $\epsilon > 0$ there exists $A(\epsilon)$ such that if $t_2 \geq t_1 \geq A(\epsilon)$ then $\|e^{-it_1\Delta}u(t_1) - e^{-it_2\Delta}u(t_2)\|_{\tilde{H}^k} \leq \epsilon$. The Cauchy criterion is satisfied. Hence scattering.

4. PROOF OF PROPOSITION 4

The proof relies upon a Morawetz type estimate that we prove in the next subsection:

Lemma 5. “*Morawetz type estimate*” Let u be an \tilde{H}^k -solution of (1) on a compact interval I . Let $A > 1$. Then

$$(54) \quad \int_I \int_{|x| \leq A|I|^{\frac{1}{2}}} \frac{\tilde{F}(u, \bar{u})(t, x)}{|x|} dx dt \lesssim EA|I|^{\frac{1}{2}}$$

with

$$(55) \quad \tilde{F}(u, \bar{u})(t, x) := \int_0^{|u|(t, x)} s^{\frac{n+2}{n-2}} \left(\frac{4}{n-2} g(s) + s g'(s) \right) ds$$

We prove now Proposition 4, following closely an argument in [10].

Step 1

We divide the interval $J = [t_1, t_2]$ into subintervals $(J_l := [\bar{t}_l, \bar{t}_{l+1}])_{1 \leq l \leq L}$ such that

$$(56) \quad \left\| u \right\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J_l)} = \eta_1$$

$$(57) \quad \left\| u \right\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J_L)} \leq \eta_1$$

with $0 < c_1 := c_1(E) \ll_E 1$ and $\eta_1 := \frac{c_1(E)}{g^{\frac{2(n+2)}{6-n}}(M)}$. It is enough to find an upper

bound of L that would depend on the energy E and M . In view of (28), we may replace WLOG the “ \leq ” sign with the “ $=$ ” sign in (57).

Notice that the value of this parameter, along with the values of the other parameters η_2 , η_3 and η are not chosen randomly: they are the largest ones (modulo the energy) such that all the constraints appearing throughout the proof are satisfied. Indeed, if we consider for example η_1 , we basically want to minimize $L\eta_1$. If we go throughout the proof without assigning any value to η_1 we realize that basically $L \lesssim \left(\frac{1}{\eta_1}\right)^{\frac{1}{\eta_1}}$ and therefore $L\eta_1$ is bounded by a smaller expression as η_1 grows.

Step 2

We first prove that some norms on these intervals J_l are bounded by a constant that depends on the energy.

Result 1. *We have*

$$(58) \quad \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J_l)} \lesssim_E 1$$

Proof.

$$(59) \quad \begin{aligned} \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J_l)} &\lesssim \|Du(\bar{t}_l)\|_{L^2} + \|D(|u|^{\frac{4}{n-2}}ug(|u|))\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}(J_l)} \\ &\lesssim E^{\frac{1}{2}} + \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J_l)}^{\frac{4}{n-2}} \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J_l)} g(M) \end{aligned}$$

Therefore, by a continuity argument¹¹, we conclude that $\|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J_l)} \lesssim_E 1$.

□

Result 2. *Let $\tilde{J} := [\tilde{t}_1, \tilde{t}_2] \subset J$ be such that*

$$(60) \quad \frac{\eta_1}{2} \leq \left\| u \right\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\tilde{J})} \leq \eta_1$$

Then

$$(61) \quad \|u_{l, \tilde{t}_j}\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\tilde{J})} \gtrsim \eta_1$$

for $j \in \{1, 2\}$.

¹¹Let $K \subset J_l$. Then (59) also holds if J_l is replaced with K .

Proof. By Result 1 we have

$$\begin{aligned}
 (62) \quad \|u - u_{l, \bar{t}_j}\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\bar{J})} &\lesssim \|D(|u|^{\frac{4}{n-2}} u g(|u|))\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}(\bar{J})} \\
 &\lesssim \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(\bar{J})} \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\bar{J})} g(M) \\
 &\lesssim_E \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\bar{J})} g(M) \\
 &\ll \eta_1^{\frac{n-2}{2(n+2)}}
 \end{aligned}$$

Therefore (61) holds. \square

Step 3

We define the notion of exceptional intervals and the notion of unexceptional intervals. Let

$$(63) \quad \eta_2 := \begin{cases} c_2 (\eta_1 g^{-1}(M))^{22}, & n = 3 \\ c_2 (\eta_1^{35} g^{-28}(M))^{\frac{1}{3}}, & n = 4 \end{cases}$$

with $0 < c_2 \ll_E c_1$. An interval $J_{l_0} = [\bar{t}_{l_0}, \bar{t}_{l_0+1}]$ of the partition $(J_l)_{1 \leq l \leq L}$ is exceptional if

$$(64) \quad \|u_{l, t_1}\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J_{l_0})} + \|u_{l, t_2}\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J_{l_0})} \geq \eta_2$$

Notice that, in view of the Strichartz estimates (16), it is easy to find an upper bound of the cardinal of the exceptional intervals:

$$(65) \quad \text{card} \{J_l : J_l \text{ exceptional}\} \lesssim_E \eta_2^{-1}$$

Step 4

Now we prove that on each unexceptional subintervals J_l there is a ball for which we have a mass concentration.

Result 3. “Mass Concentration” *There exists an $x_l \in \mathbb{R}^n$, two constants $0 < c \ll_E 1$ and $C \gg_E 1$ such that for each unexceptional interval J_l and for $t \in J_l$*

- if $n = 3$

$$(66) \quad \text{Mass} \left(u(t), B(x_l, C g^{\frac{13}{3}}(M) |J_l|^{\frac{1}{2}}) \right) \geq c g^{-\frac{13}{3}}(M) |J_l|^{\frac{1}{2}}$$

- if $n = 4$

$$(67) \quad \text{Mass} \left(u(t), B(x_l, C g^{\frac{17}{3}}(M) |J_l|^{\frac{1}{2}}) \right) \geq c g^{-\frac{17}{3}}(M) |J_l|^{\frac{1}{2}}$$

Proof. By time translation invariance¹² we may assume that $\bar{t}_l = 0$. By using the pigeonhole principle and the reflection symmetry (if necessary)¹³ we may assume that

$$(68) \quad \int_{\frac{|J_l|}{2}}^{|J_l|} \int_{\mathbb{R}^n} |u(t, x)|^{\frac{2(n+2)}{n-2}} dx dt \geq \frac{\eta_1}{4}$$

By the pigeonhole principle there exists t_* such that $[(t_* - \eta_3)|J_l|, t_*|J_l|] \subset [0, \frac{|J_l|}{2}]$ (with $\eta_3 \ll 1$) and

$$(69) \quad \int_{(t_* - \eta_3)|J_l|}^{t_*|J_l|} \int_{\mathbb{R}^n} |u(t, x)|^{\frac{2(n+2)}{n-2}} dx dt \lesssim \eta_1 \eta_3$$

$$(70) \quad \int_{\mathbb{R}^n} |u_{l, t_1}((t_* - \eta_3)|J_l|, x)|^{\frac{2(n+2)}{n-2}} dx \lesssim \frac{\eta_2}{|J_l|}$$

Applying Result 2 to (68) we have

$$(71) \quad \int_{t_*|J_l|}^{|J_l|} \int_{\mathbb{R}^n} |e^{i(t-t_*|J_l|)\Delta} u(t_*|J_l|, x)|^{\frac{2(n+2)}{n-2}} dx dt \gtrsim_E \eta_1$$

By Duhamel formula we have

$$(72) \quad \begin{aligned} u(t_*|J_l|) &= e^{i(t_*|J_l|-t_1)\Delta} u(t_1) - i \int_{t_1}^{(t_*-\eta_3)|J_l|} e^{i(t_*|J_l|-s)\Delta} (|u(s)|^{\frac{4}{n-2}} u(s) g(|u(s)|)) ds \\ &\quad - i \int_{(t_*-\eta_3)|J_l|}^{t_*|J_l|} e^{i(t_*|J_l|-s)\Delta} (|u(s)|^{\frac{4}{n-2}} u(s) g(|u(s)|)) ds \end{aligned}$$

and, composing this equality with $e^{i(t-t_*|J_l|)\Delta}$ we get

$$(73) \quad \begin{aligned} e^{i(t-t_*|J_l|)\Delta} u(t_*|J_l|) &= u_{l, t_1}(t) - i \int_{t_1}^{(t_*-\eta_3)|J_l|} e^{i(t-s)\Delta} (|u(s)|^{\frac{4}{n-2}} u(s) g(|u(s)|)) ds \\ &\quad - i \int_{(t_*-\eta_3)|J_l|}^{t_*|J_l|} e^{i(t-s)\Delta} (|u(s)|^{\frac{4}{n-2}} u(s) g(|u(s)|)) ds \\ &= u_{l, t_1}(t) + v_1(t) + v_2(t) \end{aligned}$$

We get from a variant of the Strichartz estimates (16) and the Sobolev inequality (14)

$$(74) \quad \begin{aligned} &\|v_2\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}} \cap L_t^\infty D^{-1} L_x^2([t_*|J_l|, |J_l|])} \\ &\lesssim \|D(|u|^{\frac{4}{n-2}} u g(|u|))\|_{L_t^{\frac{2(n+2)}{n+4}} L_x^{\frac{2(n+2)}{n+4}}([[(t_*-\eta_3)|J_l|, t_*|J_l|])} \\ &\lesssim \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}([[(t_*-\eta_3)|J_l|, t_*|J_l|])} \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([[(t_*-\eta_3)|J_l|, t_*|J_l|])} \\ &\quad g(\|u\|_{L_t^\infty \dot{H}^k([[(t_*-\eta_3)|J_l|, |J_l|])}) \\ &\lesssim_E (\eta_1 \eta_3)^{\frac{2}{n+2}} g(M) \\ &\ll \eta_1^{\frac{n-2}{2(n+2)}} \end{aligned}$$

¹²i.e if u is a solution of (1) and $t_0 \in \mathbb{R}$ then $(t, x) \rightarrow u(t - t_0, x)$ is also a solution of (1).

¹³if u is a solution of (1) then $(t, x) \rightarrow \bar{u}(-t, x)$ is also a solution of (1).

Notice also that $\eta_2 \ll \eta_1$ and that J_l is non-exceptional. Therefore $\|u_{l,t_1}\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])} \ll \eta_1$ and combining this inequality with (74) and (71) we conclude that the $L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])$ norm of v_1 on $[t_*|J_l|, |J_l|]$ is bounded from below:

$$(75) \quad \|v_1\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])} \gtrsim \eta_1$$

By (16), (73) and (74) we also have an upper bound of the $L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}$ norm of v_1 on $[t_*|J_l|, |J_l|]$

$$(76) \quad \|v_1\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|]) \cap L_t^\infty D^{-1} L_x^2([t_*|J_l|, |J_l|])} \lesssim_E 1$$

Now we use a lemma that is proved in Subsection 4.1.

Lemma 6. “*Regularity of v_1* ” We have

$$(77) \quad \|v_{1,h} - v_1\|_{L_t^\infty L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])} \lesssim_E |h|^\alpha |J_l|^\beta \gamma$$

with

- $\alpha = \frac{1}{5}$ if $n = 3$; $\alpha = 1$ if $n = 4$
- $\beta = -\frac{1}{5}$ if $n = 3$; $\beta = -\frac{2}{3}$ if $n = 4$
- $\gamma = g^{\frac{2}{15}}(M)$ if $n = 3$; $\gamma = g^{\frac{1}{3}}(M)$ if $n = 4$;

Denote by $v_{1,h}^{av}(x) := \int \chi(y) v_1(x + |h|y) dy$ with χ a bump function with total mass equal to one and such that $\text{supp}(\chi) \subset B(0, 1)$. Then

$$(78) \quad \begin{aligned} \|v_{1,h}^{av} - v_1\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])} &\lesssim |J_l|^{\frac{n-2}{2(n+2)}} \|v_{1,h}^{av} - v_1\|_{L_t^\infty L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])} \\ &\lesssim_E |h|^\alpha |J_l|^{\beta + \frac{n-2}{2(n+2)}} \gamma \end{aligned}$$

Therefore if h satisfies $|h| := c_3 |J_l|^{-\frac{(\beta + \frac{n-2}{2(n+2)})}{\alpha}} \gamma^{-\frac{1}{\alpha}} \eta_1^{\frac{n-2}{2(n+2)\alpha}}$ with $0 < c_3 \ll_E 1$ then

$$(79) \quad \|v_{1,h}^{av}\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])} \gtrsim \eta_1^{\frac{n-2}{2(n+2)}}$$

Now notice that by the Duhamel formula $v_1(t) = u_{l,(t_* - \eta_3)|J_l|}(t) - u_{l,t_1}(t)$ and therefore, by the Strichartz estimates (16) and the conservation of energy, $\|v_1\|_{L_t^\infty L_x^{\frac{2n}{n-2}}([t_*|J_l|, |J_l|])} \lesssim_E$

1. From that we get $\|v_{1,h}^{av}\|_{L_t^{\frac{2n}{n-2}} L_x^{\frac{2n}{n-2}}([t_*|J_l|, |J_l|])} \lesssim_E |J_l|^{\frac{n-2}{2n}}$ and, by interpolation,

$$(80) \quad \|v_{1,h}^{av}\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}([t_*|J_l|, |J_l|])} \lesssim \|v_{1,h}^{av}\|_{L_t^\infty L_x^\infty([t_*|J_l|, |J_l|])}^{\frac{2}{n+2}} \|v_{1,h}^{av}\|_{L_t^{\frac{2n}{n-2}} L_x^{\frac{2n}{n-2}}([t_*|J_l|, |J_l|])}^{\frac{n}{n+2}}$$

and, in view of (79)

$$(81) \quad \|v_{1,h}^{av}\|_{L_t^\infty L_x^\infty([t_*, |J_l|, |J_l|])} \gtrsim |J_l|^{-\frac{n-2}{4}} \eta_1^{\frac{n-2}{4}}$$

Writing $Mass(v(t), B(x, r)) = r^{\frac{n}{2}} \left(\int_{|y| \leq 1} |v(t, x + ry)|^2 dy \right)^{\frac{1}{2}}$ we deduce from Cauchy Schwartz and (81) that there exists $\check{t}_l \in [t_*, |J_l|, |J_l|]$ and $x_l \in \mathbb{R}^n$ such that

$$(82) \quad Mass(v_1(\check{t}_l), B(x_l, |h|)) \gtrsim |J_l|^{-\frac{n-2}{4}} \eta_1^{\frac{n-2}{4}} |h|^{\frac{n}{2}}$$

Therefore, by (26) we see that if $R = C_3(E) \eta_1^{\frac{2-n}{4}} |J_l|^{\frac{2+n}{4}} |h|^{-\frac{n}{2}}$ with $C_3 \gg_E 1$ then

$$(83) \quad Mass(v_1((t_* - \eta_3)|J_l|), B(x_l, R)) \gtrsim |J_l|^{-\frac{n-2}{4}} \eta_1^{\frac{n-2}{4}} |h|^{\frac{n}{2}}$$

Notice that $u((t_* - \eta_3)|J_l|) = u_{l,t_1}((t_* - \eta_3)|J_l|) - i v_1((t_* - \eta_3)|J_l|)$. By Hölder inequality, (63), and (70)

$$(84) \quad \begin{aligned} Mass(u_{l,t_1}((t_* - \eta_3)|J_l|), B(x_l, R)) &\lesssim R^{\frac{2n}{n+2}} \frac{\eta_2^{\frac{n-2}{2(n+2)}}}{|J_l|^{\frac{n-2}{2(n+2)}}} \\ &\ll |J_l|^{-\frac{n-2}{4}} \eta_1^{\frac{n-2}{4}} |h|^{\frac{n}{2}} \end{aligned}$$

Therefore $Mass(u((t_* - \eta_3)|J_l|), B(x_l, R)) \sim Mass(v_1((t_* - \eta_3)|J_l|), B(x_l, R))$. Applying again (26) we get

$$(85) \quad Mass(u(t), B(x_l, R)) \gtrsim |J_l|^{-\frac{n-2}{4}} \eta_1^{\frac{n-2}{4}} |h|^{\frac{n}{2}}$$

for $t \in J_l$. Putting everything together we get (66) and (67). \square

Next we use the radial symmetry to prove that, in fact, there is a mass concentration around the origin.

Step 5

Result 4. “Mass concentration around the origin” *There exists a positive constant $\ll_E 1$ (that we still denote by c to avoid too much notation) and a constant $\tilde{C} \gg_E 1$ such that on each unexceptional interval J_l we have*

- if $n = 3$

$$(86) \quad Mass\left(u(t), B(0, \tilde{C} g^{\frac{169}{3}}(M) |J_l|^{\frac{1}{2}})\right) \geq c g^{-\frac{13}{3}}(M) |J_l|^{\frac{1}{2}}$$

- if $n = 4$

$$(87) \quad Mass\left(u(t), B(0, \tilde{C} g^{51}(M) |J_l|^{\frac{1}{2}})\right) \geq c g^{-\frac{17}{3}}(M) |J_l|^{\frac{1}{2}}$$

Proof. We deal with the case $n = 4$. The case $n = 3$ is treated similarly and the proof is left to the reader.

Let $A := \tilde{C} g^{51}(M)$ for some $\tilde{C} \gg_E C$ (Recall that C is defined in (67)). There are (a priori) two options:

- $|x_l| \geq \frac{A}{2}|J_l|^{\frac{1}{2}}$. Then there are at least $\frac{A}{100Cg^{\frac{17}{3}}(M)}$ rotations of the ball $B(x_l, Cg^{\frac{17}{3}}(M)|J_l|^{\frac{1}{2}})$ that are disjoint. Now, since the solution is radial, the mass on each of these balls B_j is equal to that of the ball $B(x_l, Cg^{\frac{17}{3}}(M)|J_l|^{\frac{1}{2}})$. But then by Hölder inequality we have

$$(88) \quad \|u(t)\|_{L^2(B_j)}^{\frac{2n}{n-2}} \leq \|u(t)\|_{L^{\frac{2n}{n-2}}(B_j)}^{\frac{2n}{n-2}} \left(Cg^{\frac{17}{3}}(M)|J_l|^{\frac{1}{2}} \right)^{\frac{2n}{n-2}}$$

and summing over j we see from the equality $\|u(t)\|_{L^{\frac{2n}{n-2}}}^{\frac{2n}{n-2}} \lesssim E$ that

$$(89) \quad \frac{A}{100Cg^{\frac{17}{3}}(M)} \left(cg^{-\frac{17}{3}}(M)|J_l|^{\frac{1}{2}} \right)^{\frac{2n}{n-2}} \leq E \left(Cg^{\frac{17}{3}}(M)|J_l|^{\frac{1}{2}} \right)^{\frac{2n}{n-2}}$$

must be true. But with the value of A chosen above we see that this inequality cannot be satisfied if \tilde{C} is large enough. Therefore this scenario is impossible.

- $|x_l| \leq \frac{A}{2}|J_l|^{\frac{1}{2}}$. Then by (67) and the triangle inequality, we see that (87) holds.

□

Remark 3. In order to avoid too much notation we will still write in the sequel C for \tilde{C} in (87).

Step 6

Combining the inequality (87) to the Morawetz type inequality found in Lemma 5 we can prove that at least one of the intervals J_l is large. More precisely

Result 5. “One of the intervals J_l is large ” There exists a positive constant $\ll_E 1$ (that we still denote by c to avoid too much notation) and $\tilde{l} \in [1, \dots, L]$ such that

- if $n = 3$

$$(90) \quad |J_{\tilde{l}}| \geq cg^{-\frac{2860}{3}}(M)|J|$$

- if $n = 4$

$$(91) \quad |J_{\tilde{l}}| \geq cg^{-\frac{1972}{3}}(M)|J|$$

Proof. Again we shall treat the case $n = 4$. The case $n = 3$ is left to the reader.

There are two options:

- J_l is unexceptional. Let $R := Cg^{51}(M)|J_l|^{\frac{1}{2}}$. By Hölder inequality (in space), by integration in time we have

$$(92) \quad \int_{J_l} \int_{B(0,R)} \frac{|u(t,x)|^{\frac{2n}{n-2}}}{|x|} dx dt \geq |J_l| Mass^{\frac{2n}{n-2}}(u(t), B(0, R)) R^{\frac{2-3n}{n-2}}$$

After summation over l we see, by (87) and (54) that

$$(93) \quad \sum_{l=1}^L |J_l| \left(g^{-\frac{17}{3}}(M) |J_l|^{\frac{1}{2}} \right)^{\frac{2n}{n-2}} \left(C g^{51}(M) |J_l|^{\frac{1}{2}} \right)^{\frac{2-3n}{n-2}} \\ \lesssim E |J|^{\frac{1}{2}} g^{51}(M)$$

and after rearranging, we see that

$$(94) \quad \sum_{l=1}^L |J_l|^{\frac{1}{2}} g^{-\frac{986}{3}}(M) \lesssim E |J|^{\frac{1}{2}}$$

- J_l is exceptional. In this case by (65) and

$$(95) \quad \sum_{l=1}^L |J_l|^{\frac{1}{2}} \lesssim_E \eta_2^{-1} \sup_{1 \leq l \leq L} |J_l|^{\frac{1}{2}} \\ \lesssim_E \eta_2^{-1} |J|^{\frac{1}{2}}$$

Therefore, writing $\sum_{l=1}^L |J_l|^{\frac{1}{2}} \geq \frac{|J|}{\sup_{1 \leq l \leq L} |J_l|^{\frac{1}{2}}}$, we conclude that there exists a constant $\ll_E 1$ (still denoted by c) and $\tilde{l} \in [1, \dots, L]$ such that (91) holds. \square

Step 7

We use a crucial algorithm due to Bourgain [1] to prove that there are many of those intervals that concentrate.

Result 6. “Concentration of intervals” Let

$$(96) \quad \eta := \begin{cases} c g^{-\frac{2860}{3}}(M), & n = 3 \\ c g^{-\frac{1972}{3}}(M), & n = 4 \end{cases}$$

There exist a time \bar{t} , $K > 0$ and intervals J_{l_1}, \dots, J_{l_K} such that

$$(97) \quad |J_{l_1}| \geq 2|J_{l_2}| \dots \geq 2^{K-1}|J_{l_K}| \dots \geq 2^{K-1}|J_{l_K}|,$$

$$(98) \quad \text{dist}(\bar{t}, J_{l_k}) \leq \eta^{-1}|J_{l_k}|,$$

and

$$(99) \quad K \geq -\frac{\log(L)}{2 \log(\frac{\eta}{8})}.$$

Proof. There are several steps

- (1) By Result 5 there exists an interval J_{l_1} such that $|J_{l_1}| \geq \eta|J|$. We have $\text{dist}(t, J_{l_1}) \leq |J| \leq \eta^{-1}|J_{l_1}|$, $t \in J$.
- (2) Remove all the intervals J_l such that $|J_l| \geq \frac{|J_{l_1}|}{2}$. By the property of J_{l_1} , there are at most $2\eta^{-1}$ intervals satisfying this property and consequently there are at most $4\eta^{-1}$ remaining connected components resulting from this removal.

- (3) If $L \leq 100\eta^{-1}$ then we let $K = 1$ and we can check that (99) is satisfied. If not: one of these connected components (denoted by K_1) contains at least $\frac{\eta}{8}L$ intervals. Let L_1 be the number of intervals making K_1 .
- (4) Apply (1) again: there exists an interval J_{l_2} such that $|J_{l_2}| \geq \eta|K_1|$ and $\text{dist}(t, J_{l_2}) \leq |K_1| \leq \eta^{-1}|J_{l_2}|$, $t \in K_1$. Apply (2) again: remove all the intervals J_l such that $|J_l| \geq \frac{|J_{l_2}|}{2}$. By the property of J_{l_2} , there are at most $2\eta^{-1}$ intervals to be removed and there are at most $4\eta^{-1}$ remaining connected components. Apply (3) again: if $L_1 \leq 100\eta^{-1}$ then we let $K = 2$ and we can check that (99) is satisfied, since K_1 contains at least $\frac{\eta}{8}L$ intervals; if $L_1 \geq 100\eta^{-1}$ then one of the connected components (denoted by K_2) contains at least $\frac{\eta}{8}L_1$ intervals. Let L_2 be the number of intervals making K_2 . Then $L_2 \geq (\frac{\eta}{8})^2 L$.
- (5) We can iterate this procedure K times until $L_{K-1} \leq 100\eta^{-1}$. It is not difficult to see that K satisfies (99), since $L_{K-1} \geq (\frac{\eta}{8})^{K-1} L$.

□

Step 8

We prove that $L < \infty$, by using Step 7 and the conservation the energy. More precisely

Result 7. “finite bound of L ” *There exist two constants $C_1 \gg_E 1$ and $C_2 \gg_E 1$ such that*

- if $n = 3$

$$(100) \quad L \leq \left(C_1 g^{\frac{2860}{3}}(M) \right)^{C_2 g^{5772+}(M)}$$

- if $n = 4$

$$(101) \quad L \leq \left(C_1 g^{\frac{1972}{3}}(M) \right)^{C_2 g^{\frac{8024}{3}+}(M)}$$

Proof. Again we shall prove this result for $n = 4$. The case $n = 3$ is left to the reader. Let $R_{l_k} := Cg^{663}(M)|J_{l_k}|^{\frac{1}{2}}$. By Result 3 we have

$$(102) \quad \text{Mass}(u(t), B(x_{l_k}, R_{l_k})) \geq cg^{-\frac{17}{3}}(M)|J_{l_k}|^{\frac{1}{2}}$$

for all $t \in J_{l_k}$. Even if it means redefining C ¹⁴ then we see, by (26) and (98) that (102) holds for $t = \bar{t}$ with c substituted for $\frac{c}{2}$. On the other hand we see that by (25) that¹⁵

$$(103) \quad \sum_{k'=k+N}^K \int_{B(x_{l_{k'}}, R_{l_{k'}})} |u(\bar{t}, x)|^2 dx \leq \left(\frac{1}{2^N} + \frac{1}{2^{N+1}} \dots + \frac{1}{2^{K-k}} \right) ER_{l_k}^2 \\ \leq \frac{1}{2^{N-1}} ER_{l_k}^2$$

¹⁴i.e making it larger than its original value modulo a multiplication by some power of $\max(1, E)$

¹⁵Notation: $\sum_{k'=k+N}^K a_{k'} = 0$, if $k + N > K$

Now we let $N = C' \log(g(M))$ with $C' \gg_E 1$ so that $\frac{ER_k^2}{2^{N-1}} \leq \frac{1}{8}c^2g^{-\frac{34}{3}}(M)|J_{l_k}|$. By (102) we have

$$(104) \quad \sum_{k'=k+N}^K \int_{B(x_{l_{k'}}, R_{l_{k'}})} |u(\bar{t}, x)|^2 dx \leq \frac{1}{2} \int_{B(x_{l_k}, R_{l_k})} |u(\bar{t}, x)|^2 dx$$

Therefore

$$(105) \quad \begin{aligned} \int_{B(x_{l_k}, R_{l_k}) \setminus \bigcup_{k'=k+N}^K B(x_{l_{k'}}, R_{l_{k'}})} |u(\bar{t}, x)|^2 dx &\geq \frac{1}{2} \int_{B(x_{l_k}, R_{l_k})} |u(\bar{t}, x)|^2 dx \\ &\geq \frac{c^2g^{-\frac{34}{3}}(M)}{4} |J_{l_k}| \end{aligned}$$

and by Hölder inequality, there exists a positive constant $\ll_E 1$ (that we still denote by c) such that

$$(106) \quad \int_{B(x_{l_k}, R_{l_k}) \setminus \bigcup_{k'=k+N}^K B(x_{l_{k'}}, R_{l_{k'}})} |u(\bar{t}, x)|^{\frac{2n}{n-2}} dx \geq cg^{-\frac{8024}{3}}(M)$$

and after summation over k , we

$$(107) \quad \frac{K}{N} cg^{-\frac{8024}{3}}(M) \lesssim E$$

since $\sum_{k=1}^K \chi_{B(x_{l_k}, R_{l_k}) \setminus \bigcup_{k'=k+N}^K B(x_{l_{k'}}, R_{l_{k'}})} \leq N$ and $\|u(t)\|_{L^{\frac{2n}{n-2}}}^{\frac{2n}{n-2}} \lesssim E$. Rearranging we see from (99) that there exist two constants $C_1 \gg_E 1$ and $C_2 \gg_E 1$ such that

$$(108) \quad L \leq \left(C_1 g^{\frac{1972}{3}}(M) \right)^{C_2 \log(g(M))g^{\frac{8024}{3}}(M)}$$

We see that (101) holds.

Step 9

This is the final step. Recall that there are L intervals J_l and that on each of these intervals we have $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)} = \eta_1$. Therefore, there are two constants $\gg_E 1$ (that we denote by C_1 and C_2) such that (28) holds. \square

4.1. Proof of Lemma 6. In this subsection we prove Lemma 6. There are two cases

- $n = 3$

By the fundamental theorem of calculus (and the inequality $\|Du\|_{L_t^\infty L_x^2([t_*, |J_l|, |J_l|])} \lesssim E^{\frac{1}{2}}$) we have

$$(109) \quad \|u_h - u\|_{L_t^\infty L_x^2([t_*, |J_l|, |J_l|])} \leq E^{\frac{1}{2}} |h|$$

Moreover, by Sobolev (and the inequality $\|u\|_{L_t^\infty L_x^6([t_*, |J_l|, |J_l|])} \lesssim E^{\frac{1}{6}}$) we have

$$(110) \quad \|u_h - u\|_{L_t^\infty L_x^6([t_*, |J_l|, |J_l|])} \leq E^{\frac{1}{6}}$$

Therefore, by interpolation of (109) and (110), we get

$$(111) \quad \|u_h - u\|_{L_t^\infty L_x^3([t_*, |J_l|, |J_l|])} \leq E^{\frac{1}{3}} |h|^{\frac{1}{2}}$$

Now, by the fundamental theorem of calculus, the inequality $|x|g'(|x|) \lesssim g(|x|)$, (23) and (20) we have

$$(112) \quad \begin{aligned} \| |u(s)|^{\frac{4}{n-2}} u(s) g(|u(s)|) - |u_h(s)|^{\frac{4}{n-2}} u_h(s) g(|u_h(s)|) \|_{L^1} &\lesssim \|u_h(s) - u(s)\|_{L^3} \|u(s) g^{\frac{n-2}{2n}}(|u(s)|)\|_{L^6}^4 \\ &\|g^{\frac{n-2}{n}}(|u(s)|)\|_{L^\infty} \\ &\lesssim_E g^{\frac{n-2}{n}}(M) |h|^{\frac{1}{2}} \end{aligned}$$

and, by the dispersive inequality (13) we conclude that

$$(113) \quad \|v_{1,h} - v_1\|_{L_t^\infty L_x^\infty([t_*, |J_l|, |J_l|])} \lesssim_E \eta_3^{-\frac{1}{2}} |J_l|^{-\frac{1}{2}} g^{\frac{n-2}{n}}(M) |h|^{\frac{1}{2}}$$

Interpolating this inequality with

$$(114) \quad \begin{aligned} \|v_{1,h} - v_1\|_{L_t^\infty L_x^6([t_*, |J_l|, |J_l|])} &= \|u_{l,(t_*-\eta_3)|J_l|,h} - u_{l,t_1,h} - (u_{l,(t_*-\eta_3)|J_l|} - u_{l,t_1})\|_{L_t^\infty L_x^6([t_*, |J_l|, |J_l|])} \\ &\lesssim E^{\frac{1}{2}} \end{aligned}$$

we get (77).

- $n = 4$ By the fundamental theorem of calculus we have

$$(115) \quad \|v_{1,h} - v_1\|_{L_t^\infty L_x^{\frac{2(n+2)}{n-2}}([t_*, |J_l|, |J_l|])} \lesssim \|Dv_1\|_{L_t^\infty L_x^{\frac{2(n+2)}{n-2}}([t_*, |J_l|, |J_l|])} |h|$$

But, by interpolation

$$(116) \quad \begin{aligned} \|Dv_1\|_{L_t^\infty L_x^{\frac{2(n+2)}{n-2}}([t_*, |J_l|, |J_l|])} &\lesssim \|Dv_1\|_{L_t^\infty L_x^2([t_*, |J_l|, |J_l|])}^{\frac{2}{n+2}} \|Dv_1\|_{L_t^\infty L_x^{\frac{2n}{n-4}}([t_*, |J_l|, |J_l|])}^{\frac{n}{n+2}} \\ &\lesssim_E \|Dv_1\|_{L_t^\infty L_x^{\frac{2n}{n-4}}([t_*, |J_l|, |J_l|])}^{\frac{n}{n+2}} \end{aligned}$$

So it suffices to estimate $\|Dv_1\|_{L_t^\infty L_x^{\frac{2n}{n-4}}([t_*, |J_l|, |J_l|])}$. By (20), (23) and Result

1 we have

$$(117) \quad \begin{aligned} \|D(|u|^{\frac{4}{n-2}} u g(|u|))\|_{L_s^\infty L_x^{\frac{2n}{n+4}}([t_1, (t_*-\eta_3)|J_l|])} &\lesssim \|Du\|_{L_s^\infty L_x^2([t_1, (t_*-\eta_3)|J_l|])} \|u g^{\frac{n-2}{2n}}(|u|)\|_{L_s^\infty L_x^{\frac{2n}{n-2}}([t_1, (t_*-\eta_3)|J_l|])}^{\frac{4}{n-2}} \\ &g^{\frac{n-2}{n}}(\|u\|_{L_t^\infty \tilde{H}^k([t_1, (t_*-\eta_3)|J_l|])}) \\ &\lesssim_E g^{\frac{n-2}{n}}(M) \end{aligned}$$

and by combining (117) with the dispersive inequality (13) we have

$$\begin{aligned}
(118) \quad \|Dv_1\|_{L_t^\infty L_x^{\frac{2n}{n-4}}([t_*|J_l|, |J_l|])} &\lesssim \left\| \int_{t_1}^{(t_*-\eta_3)|J_l|} \|De^{i(t-s)\Delta}(|u(s)|^{\frac{4}{n-2}}u(s)g(|u(s)|))\|_{L_x^{\frac{2n}{n-4}}} ds \right\|_{L_t^\infty([t_*|J_l|, |J_l|])} \\
&\lesssim \left\| \int_{t_1}^{(t_*-\eta_3)|J_l|} \frac{1}{|t-s|^2} \|D(|u(s)|^{\frac{4}{n-2}}u(s)g(|u(s)|))\|_{L_x^{\frac{2n}{n+4}}} ds \right\|_{L_t^\infty([t_*|J_l|, |J_l|])} \\
&\lesssim g^{\frac{n-2}{n}}(M)\eta_3^{-1}|J_l|^{-1}
\end{aligned}$$

We conclude from (116) and (118) that (77) holds.

4.2. Proof of Lemma 5. By (1) we have ¹⁶

$$(119) \quad \partial_t \Im(\partial_k u \bar{u}) = \Re \left[|u|^{\frac{4}{n-2}} \bar{u} g(|u|) \partial_k u - \partial_k (|u|^{\frac{4}{n-2}} u g(|u|)) \right] + \Re \left(\Delta (\partial_k u) \bar{u} - \overline{\Delta u} \partial_k u \right)$$

Moreover

$$(120) \quad \frac{1}{2} \partial_k \Delta (|u|^2) = 2 \partial_j \Re(\partial_k u \overline{\partial_j u}) - \Re(\partial_k u \Delta \bar{u}) + \Re(u \Delta \overline{\partial_k u})$$

Therefore, adding (119) and (120) leads to

$$(121) \quad \partial_t \Im(\partial_k u \bar{u}) = -2 \partial_j \Re(\partial_k u \overline{\partial_j u}) + \frac{1}{2} \partial_k \Delta (|u|^2) + \Re \left[|u|^{\frac{4}{n-2}} \bar{u} g(|u|) \partial_k u - \partial_k (|u|^{\frac{4}{n-2}} u g(|u|)) \bar{u} \right]$$

It remains to understand $\Re \left[|u|^{\frac{4}{n-2}} \bar{u} g(|u|) \partial_k u - \partial_k (|u|^{\frac{4}{n-2}} u g(|u|)) \bar{u} \right]$. We write

$$(122) \quad \Re \left[|u|^{\frac{4}{n-2}} \bar{u} g(|u|) \partial_k u - \partial_k (|u|^{\frac{4}{n-2}} u g(|u|)) \bar{u} \right] = A_1 + A_2$$

with

$$(123) \quad A_1 := \Re \left[|u|^{\frac{4}{n-2}} \bar{u} g(|u|) \partial_k u \right]$$

and

$$(124) \quad A_2 := -\Re \left(\partial_k (|u|^{\frac{4}{n-2}} u g(|u|)) \bar{u} \right)$$

We are interested in finding a function $F_1 : \mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}$, continuously differentiable such that $F_1(z, \bar{z}) = \overline{F_1(z, \bar{z})}$, $F_1(0, 0) = 0$ and $A_1 = \partial_k F_1(u, \bar{u})$. Notice that the first condition implies in particular that $\partial_{\bar{z}} F_1(z, \bar{z}) = \overline{\partial_z F_1(z, \bar{z})}$. Therefore we get, after computation

$$\begin{aligned}
(125) \quad \partial_z F_1(z, \bar{z}) &= \frac{|z|^{\frac{4}{n-2}} \bar{z} g(|z|)}{2} \\
\partial_{\bar{z}} F_1(z, \bar{z}) &= \frac{|z|^{\frac{4}{n-2}} z g(|z|)}{2}
\end{aligned}$$

and by the fundamental theorem of calculus, if such a function exists, then

¹⁶Throughout this subsection, all the computations are done for smooth solutions. Then (54) holds for an \tilde{H}^k -solution by a standard approximation argument with smooth solutions.

$$\begin{aligned}
(126) \quad F_1(z, \bar{z}) &= \int_0^1 F_1'(tz, t\bar{z}) \cdot (z, \bar{z}) dt \\
&= 2\Re \int_0^1 \partial_z F_1(tz, t\bar{z}) z dt \\
&= \int_0^1 |tz|^{\frac{4}{n-2}} t |z|^2 g(t|z|) dt
\end{aligned}$$

and, after a change of variable, we get

$$(127) \quad F_1(z, \bar{z}) = \int_0^{|z|} t^{\frac{n+2}{n-2}} g(t) dt$$

Conversely it is not difficult to see that F_1 satisfies all the required conditions.

We turn now to A_2 . We can write

$$(128) \quad A_2 = A_{2,1} + A_{2,2}$$

with

$$(129) \quad A_{2,1} := -\Re \left(\partial_u (|u|^{\frac{4}{n-2}} u g(|u|)) \bar{u} \partial_k u \right)$$

and

$$(130) \quad A_{2,2} := -\Re \left(\partial_{\bar{u}} (|u|^{\frac{4}{n-2}} u g(|u|)) \bar{u} \partial_k u \right)$$

Again we search for a function $F_{2,1} : \mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}$ and continuously differentiable such that $F_{2,1}(z, \bar{z}) = \overline{F_{2,1}(z, \bar{z})}$ and $A_{2,1} = \partial_k F_{2,1}(u, \bar{u})$. By identification we have

$$\begin{aligned}
(131) \quad \partial_z F_{2,1}(z, \bar{z}) &= -\frac{|z|^{\frac{4}{n-2}} \bar{z} \left(\left(\frac{2}{n-2} + 1 \right) g(|z|) + \frac{g'(|z|)|z|}{2} \right)}{2} \\
\partial_{\bar{z}} F_{2,1}(z, \bar{z}) &= -\frac{|z|^{\frac{4}{n-2}} z \left(\left(\frac{2}{n-2} + 1 \right) g(|z|) + \frac{g'(|z|)|z|}{2} \right)}{2}
\end{aligned}$$

and by the fundamental theorem of calculus

$$\begin{aligned}
(132) \quad F_{2,1}(z, \bar{z}) &= \int_0^1 F_{2,1}'(tz, t\bar{z}) \cdot (z, \bar{z}) dt \\
&= \int_0^1 2\Re (\partial_z F_{2,1}(tz, t\bar{z}) z) dt \\
&= -\int_0^1 |tz|^{\frac{4}{n-2}} \left(\left(\frac{2}{n-2} + 1 \right) g(|tz|) + \frac{g'(|tz|)|tz|}{2} \right) t |z|^2 dt
\end{aligned}$$

and, after a change of variable, we get

$$(133) \quad F_{2,1}(z, \bar{z}) = -\int_0^{|z|} t^{\frac{n+2}{n-2}} \left(\left(\frac{2}{n-2} + 1 \right) g(t) + \frac{tg'(t)}{2} \right) dt$$

Again, we can easily check that $F_{2,1}$ satisfies all the required conditions. By using a similar process we can prove that

$$(134) \quad A_{2,2} = \partial_k F_{2,2}(u, \bar{u})$$

with

$$(135) \quad F_{2,2}(z, \bar{z}) = -\int_0^{|z|} t^{\frac{n+2}{n-2}} \left(\frac{2}{n-2} g(t) + \frac{tg'(t)}{2} \right) dt$$

Therefore we get the local momentum conservation identity

$$(136) \quad \partial_t \Im(\partial_k u \bar{u}) = -2\partial_j \Re(\partial_k u \overline{\partial_j u}) + \frac{1}{2}\partial_k \Delta(|u|^2) - \partial_k \left(\tilde{F}(u, \bar{u}) \right)$$

with $\tilde{F}(u, \bar{u})$ defined in (55). This identity has a similar structure to the local momentum conservation that for a solution v of the energy-critical Schrödinger equation

$$(137) \quad \partial_t \Im(\partial_k v \bar{v}) = -2\partial_j \Re(\partial_k v \overline{\partial_j v}) + \frac{1}{2}\partial_k \Delta(|v|^2) + \partial_k \left(-\frac{2}{n}|u|^{\frac{2n}{n-2}} \right)$$

With this in mind, we multiply (136) by an appropriate spatial cutoff, in the same spirit as Bourgain [1] and Grillakis [5], to prove a Morawetz-type estimate. We follow closely an argument of Tao [10]: we introduce the weight $a(x) :=$

$\left(\epsilon^2 + \left(\frac{|x|}{A|I|^{\frac{1}{2}} \right)^2 \right)^{\frac{1}{2}} \chi \left(\frac{x}{A|I|^{\frac{1}{2}}} \right)$ where χ is a smooth function, radial such that $\chi(|x|) = 1$ for $|x| \leq 1$ and $\chi(|x|) = 0$ for $|x| \geq 2$. We give here the details since this equation, unlike the energy-critical Schrödinger equation, has no scaling property. Notice that a is convex on $|x| \leq A|I|^{\frac{1}{2}}$ since it is a composition of two convex functions. We multiply (136) by $\partial_k a$ and we integrate by parts

$$(138) \quad \partial_t \int_{\mathbb{R}^n} \partial_k a \Im(\partial_k u \bar{u}) = 2 \int_{\mathbb{R}^n} \partial_j \partial_k a \Re(\partial_k u \overline{\partial_j u}) - \frac{1}{2} \int_{\mathbb{R}^n} \Delta(\Delta a) |u|^2 dx + \int_{\mathbb{R}^n} \Delta a \tilde{F}(u, \bar{u})(t, x) dx$$

A computation shows that for $0 \leq |x| \leq A|I|^{\frac{1}{2}}$

$$(139) \quad \Delta a = \frac{n-1}{(A|I|^{\frac{1}{2}})^2} \left(\epsilon^2 + \frac{|x|^2}{(A|I|^{\frac{1}{2}})^2} \right)^{-\frac{1}{2}} + \frac{\epsilon^2}{(A|I|^{\frac{1}{2}})^2} \left(\epsilon^2 + \frac{|x|^2}{(A|I|^{\frac{1}{2}})^2} \right)^{-\frac{3}{2}}$$

and

$$(140) \quad -\Delta \Delta a = \frac{(n-1)(n-3)}{(A|I|^{\frac{1}{2}})^4} \left(\epsilon^2 + \frac{|x|^2}{(A|I|^{\frac{1}{2}})^2} \right)^{-\frac{3}{2}} + \frac{6(n-3)\epsilon^2}{(A|I|^{\frac{1}{2}})^4} \left(\epsilon^2 + \frac{|x|^2}{(A|I|^{\frac{1}{2}})^2} \right)^{-\frac{5}{2}} + \frac{15\epsilon^4}{(A|I|^{\frac{1}{2}})^4} \left(\epsilon^2 + \frac{|x|^2}{(A|I|^{\frac{1}{2}})^2} \right)^{-\frac{7}{2}}$$

Moreover we have $|\Delta(\Delta a)| \lesssim \frac{1}{(A|I|^{\frac{1}{2}})^4}$, $|\Delta a| \lesssim \frac{1}{(A|I|^{\frac{1}{2}})^2}$ and $|\partial_j \partial_k a| \lesssim \frac{1}{(A|I|^{\frac{1}{2}})^2}$ for $A|I|^{\frac{1}{2}} \leq |x| \leq 2A|I|^{\frac{1}{2}}$ and $|\partial_k a| \lesssim \frac{1}{A|I|^{\frac{1}{2}}}$ for $|x| \leq 2A|I|^{\frac{1}{2}}$. Therefore by the previous estimates, (20), (23) and the inequality $|x|g'(|x|) \lesssim g(|x|)$ we get, after integrating on $I \times \mathbb{R}^n$ and letting ϵ go to zero

$$(141) \quad \frac{1}{A|I|^{\frac{1}{2}}} \int_I \int_{|x| \leq A|I|^{\frac{1}{2}}} \frac{\tilde{F}(u, \bar{u})(t, x)}{|x|} dx dt - C(A|I|^{\frac{1}{2}})^{-2} E|I| - C(A|I|^{\frac{1}{2}})^{-4} E(A|I|^{\frac{1}{2}})^2 |I| \lesssim E$$

for some constant $C \geq 1$. After rearranging we get (54).

5. APPENDIX A

We shall prove the following Leibnitz rule:

Proposition 7. “A fractional Leibnitz rule” Let $0 \leq \alpha < 1$, k and β be integers such that $k \geq 2$ and $\beta > k - 1$, $(r, r_2) \in (1, \infty)^2$, $(r_1, r_3) \in (1, \infty]^2$ be such that $\frac{1}{r} = \frac{\beta}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}$. Let $F : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a C^k -function and let $G := \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a C^k -function such that

$$(142) \quad F^{[i]}(x) = O\left(\frac{F(x)}{x^i}\right), \quad \tau \in [0, 1] : |F(|\tau x + (1 - \tau)y|^2)| \lesssim |F(|x|^2)| + |F(|y|^2)|,$$

and

$$(143) \quad G^{[i]}(x, \bar{x}) = O(|x|^{\beta+1-i})$$

for $0 \leq i \leq k$. Then

$$(144) \quad \|D^{k-1+\alpha}(G(f, \bar{f})F(|f|^2))\|_{L^r} \lesssim \|f\|_{L^{r_1}}^\beta \|D^{k-1+\alpha}f\|_{L^{r_2}} \|F(|f|^2)\|_{L^{r_3}}$$

Here $F^{[i]}$ and $G^{[i]}$ denote the i^{th} -derivatives of F and G respectively. More generally, let $\tilde{F} : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a C^k function. Substitute F with \tilde{F} on the right-hand side of the equality of (142), in the inequality of (142), and on the right-hand side of (144). With these substitutions made, if F , \tilde{F} , and G satisfy (142) and (143), then F and G satisfy (144).

Proof. The proof relies upon an induction process, the usual product rule for fractional derivatives

$$(145) \quad \|D^{\alpha_1}(fg)\|_{L^q} \lesssim \|D^{\alpha_1}f\|_{L^{q_1}} \|g\|_{L^{q_2}} + \|f\|_{L^{q_3}} \|D^{\alpha_1}g\|_{L^{q_4}}$$

and the usual Leibnitz rule for fractional derivatives :

$$(146) \quad \|D^{\alpha_2}H(f)\|_{L^q} \lesssim \|\tilde{H}(f)\|_{L^{q_1}} \|D^{\alpha_2}f\|_{L^{q_2}}$$

if H is C^1 and it satisfies $\tau \in [0, 1] : |H'(\tau x + (1 - \tau)y)| \lesssim \tilde{H}(x) + \tilde{H}(y)$, $0 \leq \alpha_1 < \infty$, $0 < \alpha_2 \leq 1$, $(q, q_4) \in (1, \infty)^2$, $q_3 \in (1, \infty]$, $(q_1, q_2) \in (1, \infty) \times (1, \infty]$ in (145), $(q_1, q_2) \in (1, \infty) \times (1, \infty)$ in (146), $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2}$, and $\frac{1}{q} = \frac{1}{q_3} + \frac{1}{q_4}$ (see e.g. Christ-Weinstein [3], Taylor [8] and references in [8])¹⁷. Moreover we shall use interpolation and the properties of F to control the intermediate terms.

Let $k = 2$. Then

$$(147) \quad \begin{aligned} \|D^{2-1+\alpha}(G(f, \bar{f})F(|f|^2))\|_{L^r} &\sim \|D^\alpha \nabla(G(f, \bar{f})F(|f|^2))\|_{L^r} \\ &\lesssim \|D^\alpha(\partial_z G(f, \bar{f}) \nabla f F(|f|^2))\|_{L^r} + \|D^\alpha(\partial_{\bar{z}} G(f, \bar{f}) \nabla \bar{f} F(|f|^2))\|_{L^r} \\ &\quad + \|D^\alpha(F'(|f|^2)(2\Re(\bar{f} \nabla f) G(f, \bar{f}))\|_{L^r} \\ &\lesssim A_1 + A_2 + A_3 \end{aligned}$$

We estimate A_1 . A_2 is estimated in a similar fashion. By (145), (146) and the assumption (142)

¹⁷Notice that in [3], they add the restriction $0 < \alpha_1 < 1$. It is not difficult to see that this restriction is not necessary: see Taylor [8] for example

$$\begin{aligned}
(148) \quad A_1 &\lesssim \|D^\alpha(\partial_z G(f, \bar{f})F(|f|^2))\|_{L^{r_4}} \|Df\|_{L^{r_5}} + \|\partial_z G(f, \bar{f})F(|f|^2)\|_{L^{r_6}} \|D^{(2-1)+\alpha}f\|_{L^{r_2}} \\
&\lesssim \|f\|_{L^{r_1}}^{\beta-1} \|F(|f|^2)\|_{L^{r_3}} \|D^\alpha f\|_{L^{r_8}} \|Df\|_{L^{r_5}} + \|f\|_{L^{r_1}}^\beta \|D^{(2-1)+\alpha}f\|_{L^{r_2}} \|F(|f|^2)\|_{L^{r_3}}
\end{aligned}$$

with $\frac{1}{r} = \frac{1}{r_4} + \frac{1}{r_5}$, $\frac{1}{r} = \frac{1}{r_6} + \frac{1}{r_2}$, $\frac{1}{r_4} = \frac{\beta-1}{r_1} + \frac{1}{r_3} + \frac{1}{r_8}$, $\frac{1}{r_5} = \frac{1-\theta_1}{r_1} + \frac{\theta_1}{r_2}$ and $\theta_1 = \frac{1}{1+\alpha}$. Notice that these relations imply that $\frac{1}{r_8} = \frac{\theta_1}{r_1} + \frac{1-\theta_1}{r_2}$. Now, by complex interpolation, we have

$$(149) \quad \|D^\alpha f\|_{L^{r_8}} \lesssim \|f\|_{L^{r_1}}^{\theta_1} \|D^{(2-1)+\alpha}f\|_{L^{r_2}}^{1-\theta_1}$$

and

$$(150) \quad \|Df\|_{L^{r_5}} \lesssim \|f\|_{L^{r_1}}^{1-\theta_1} \|D^{(2-1)+\alpha}f\|_{L^{r_2}}^{\theta_1}$$

Plugging (149) and (150) into (148) we get (144).

We estimate A_3 .

$$\begin{aligned}
(151) \quad A_3 &\lesssim \sum_{\tilde{f} \in \{f, \bar{f}\}} \left\| D^\alpha \left(F'(|f|^2) \tilde{f} G(f, \bar{f}) \right) \right\|_{L^{r_4}} \|Df\|_{L^{r_5}} + \|D^{\alpha+1}f\|_{L^{r_2}} \left\| F'(|f|^2) \tilde{f} G(f, \bar{f}) \right\|_{L^{r_6}} \\
&\lesssim A_{3,1} + A_{3,2}
\end{aligned}$$

Using the assumption $F'(|x|^2) = O\left(\frac{F(|x|^2)}{|x|^2}\right)$ we get $A_{3,2} \lesssim \|f\|_{L^{r_1}}^\beta \|D^{1+\alpha}f\|_{L^{r_2}} \|F(|f|^2)\|_{L^{r_3}}$.

Moreover, by (146), the assumptions on F and G , (149) and (150) we get

$$\begin{aligned}
(152) \quad A_{3,1} &\lesssim \|F(|f|^2)|f|^{\beta-1}\|_{L^{r_7}} \|D^\alpha f\|_{L^{r_8}} \|Df\|_{L^{r_5}} \\
&\lesssim \|f\|_{L^{r_1}}^\beta \|D^{1+\alpha}f\|_{L^{r_2}} \|F(|f|^2)\|_{L^{r_3}}
\end{aligned}$$

with $\frac{1}{r_7} + \frac{1}{r_8} = \frac{1}{r_4}$. The more general statement follows exactly the same steps and its proof is left to the reader.

Now let us assume that the result is true for k . Let us prove that it is also true for $k+1$. By (145) we have

$$\begin{aligned}
(153) \quad \|D^{k+\alpha}(G(f, \bar{f})F(|f|^2))\|_{L^r} &\sim \|D^{k-1+\alpha}\nabla(G(f, \bar{f})F(|f|^2))\|_{L^r} \\
&\lesssim \|D^{k-1+\alpha}\partial_z G(f, \bar{f})\nabla f F(|f|^2)\|_{L^r} + \|D^{k-1+\alpha}\partial_{\bar{z}} G(f, \bar{f})\overline{\nabla f} F(|f|^2)\|_{L^r} \\
&\quad + \left\| D^{k-1+\alpha} \left[G(f, \bar{f})F'(|f|^2) (2\Re(\bar{f}\nabla f)) \right] \right\|_{L^r} \\
&\lesssim A'_1 + A'_2 + A'_3
\end{aligned}$$

We estimate A'_1 and A'_3 . A'_2 is estimated in a similar fashion as A'_1 . By (145), (146) and the assumption $|\partial_z G(f, \bar{f})| \lesssim |f|^\beta$ we have

$$\begin{aligned}
(154) \quad A'_1 &\lesssim \|D^{k+\alpha}f\|_{L^{r_2}} \|\partial_z G(f, \bar{f})F(|f|^2)\|_{L^{r_6}} + \|D^{k-1+\alpha}(\partial_z G(f, \bar{f})F(|f|^2))\|_{L^{r'_4}} \|Df\|_{L^{r'_5}} \\
&\lesssim \|f\|_{L^{r_1}}^\beta \|D^{(k+1)-1+\alpha}f\|_{L^{r_2}} \|F(|f|^2)\|_{L^{r_3}} + A'_{1,1}
\end{aligned}$$

with r'_4, r'_5 such that $\frac{1}{r'_4} + \frac{1}{r'_5} = \frac{1}{r}$, $\frac{1}{r'_5} = \frac{1-\theta'_1}{r_1} + \frac{\theta'_1}{r_2}$ and $\theta'_1 = \frac{1}{k+\alpha}$. Notice that, since we assumed that the result is true for k , we get, after checking that $\partial_z G$ satisfies the right assumptions

$$(155) \quad \|D^{k-1+\alpha}(\partial_z G(f, \bar{f})F(|f|^2))\|_{L^{r'_4}} \lesssim \|f\|_{L^{r_1}}^{\beta-1} \|D^{k-1+\alpha}f\|_{L^{r'_8}} \|F(|f|^2)\|_{L^{r_3}}$$

with r'_8 such that $\frac{1}{r'_4} = \frac{\beta-1}{r_1} + \frac{1}{r'_8} + \frac{1}{r_3}$. Notice also that, by complex interpolation

$$(156) \quad \|Df\|_{L^{r'_5}} \lesssim \|f\|_{L^{r_1}}^{1-\theta'_1} \|D^{(k+1)-1+\alpha}f\|_{L^{r_2}}^{\theta'_1}$$

and

$$(157) \quad \|D^{k-1+\alpha}f\|_{L^{r'_8}} \lesssim \|f\|_{L^{r_1}}^{\theta'_1} \|D^{(k+1)-1+\alpha}f\|_{L^{r_2}}^{1-\theta'_1}$$

Combining (155), (156) and (157) we have

$$(158) \quad A'_{1,1} \lesssim \|f\|_{L^{r_1}}^{\beta} \|D^{k+\alpha}f\|_{L^{r_2}} \|F(|f|^2)\|_{L^{r_3}}$$

Plugging this bound into (154) we get the required bound for $A'_{1,1}$.

We turn to A'_3 . Let $\tilde{F}(x) := xF'(x)$. From the induction assumption applied to \tilde{F} we get

$$(159) \quad \begin{aligned} A'_3 &\lesssim \sum_{\tilde{f} \in \{f, \bar{f}\}} \left\| D^{k-1+\alpha} \left[G(f, \bar{f})F'(|f|^2)\tilde{f} \right] \right\|_{L^{r'_4}} \|Df\|_{L^{r'_5}} + \|D^{k+\alpha}f\|_{L^{r_2}} \|G(f, \bar{f})F'(|f|^2)\|_{L^{r_6}} \\ &\lesssim \|f\|_{L^{r_1}}^{\beta-1} \|D^{k-1+\alpha}f\|_{L^{r'_8}} \|F(|f|^2)\|_{L^{r_3}} \|Df\|_{L^{r'_5}} + \|D^{k+\alpha}f\|_{L^{r_2}} \|f\|_{L^{r_1}}^{\beta} \|F(|f|^2)\|_{L^{r_3}} \\ &\lesssim \|f\|_{L^{r_1}}^{\beta} \|D^{k+\alpha}f\|_{L^{r_2}} \|F(|f|^2)\|_{L^{r_3}} \end{aligned}$$

Again the more general statement follows exactly the same steps and its proof is left to the reader. \square

6. APPENDIX B

We shall prove the following proposition:

Proposition 8. *Let $\lambda \in \mathbb{N}^*$ and (Q, R) be such that $\left(\frac{1}{Q}, \frac{1}{R}\right) = \left(\frac{(\lambda-1)(n-2)}{2(n+2)} + \frac{n}{2(n+2)}\right)(1, 1)$.*

Let J be an interval. Let $k > \frac{n}{2}$. Let $\bar{Q}_k(J, u) := \|u\|_{L_t^\infty \dot{H}^k(J)} + \|Du\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)}$

$\|D^k u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)}$. Let $\psi(k)$ be defined as follows: if $\frac{n}{2} < k < \frac{n+2}{n-2}$ then

$\psi(k) := k$ and if $k \geq \frac{n+2}{n-2}$ then $\psi(k) := \frac{n+2}{n-2}$. There exists $\bar{C} > 0$ such that

$$(160) \quad \begin{aligned} \left\| D^k(u^\lambda g(|u|)) \right\|_{L_t^Q L_x^R(J)} &\lesssim \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)}^{\lambda-1} \bar{Q}_k(J, u) \langle \bar{Q}_{\frac{n}{2}+}(J, u) \rangle^{\bar{C}} \\ &\quad \left(g(\bar{Q}_k(J, u)) + \langle \bar{Q}_{\psi(k)}(J, u) \rangle^{\bar{C}} + \langle \bar{Q}_{k-\frac{1}{4}}(J, u) \rangle^{\bar{C}} \right). \end{aligned}$$

The same estimate holds if u^λ is replaced with $u^{\lambda_1}\bar{u}^{\lambda_2}$ with $(\lambda_1, \lambda_2) \in \mathbb{N}^2$ such that $\lambda_1 + \lambda_2 = \lambda$, or if $g(|u|)$ is replaced with $\tilde{g}'(|u|^2)u^{\lambda_3}\bar{u}^{\lambda_4}$ with $(\lambda_3, \lambda_4) \in \mathbb{N}^2$ such that $\lambda_3 + \lambda_4 = 2$ and $\tilde{g}(x) := \log^c \log(10 + x)$.

Proof. Let $k = m + \alpha$ with $0 \leq \alpha < 1$ and m integer. Then by the product rule (see proof in Appendix A) and the Sobolev embedding (15) we have

$$(161) \quad \begin{aligned} \|D^k(u^\lambda g(|u|))\|_{L_t^Q L_x^R(J)} &\lesssim \|D^k u\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)} \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)}^{\lambda-1} g\left(\|u\|_{L_t^\infty \bar{H}^k(J)}\right) \\ &\quad + \|D^k g(|u|)\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)} \|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)}^{\lambda-1} \\ &\quad \|u\|_{L_t^\infty \bar{H}^{\frac{n}{2}+}(J)} \end{aligned}$$

Let RHS' be the right-hand side of (160) multiplied by $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)}^{\lambda-1}$. We have

$$\|D^k g(|u|)\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)} \lesssim \sum_{\gamma \in \mathbb{N}^n: |\gamma|=m} \|D^\alpha \partial^\gamma (g(|u|))\|_{L_t^{\frac{2(n+2)}{n}} L_x^{\frac{2(n+2)}{n}}(J)},$$

Let $X := \partial^\gamma (g(|u|))$. Expanding we see that X is a finite sum of terms of the form $X' := \partial^{\bar{\theta}} \tilde{g}(|u|^2) X'_0 X'_1 \dots X'_m$ with

$$X'_p := (\partial^{\delta_{p,1}} u)^{\theta_{p,1}} \dots (\partial^{\delta_{p,q}} u)^{\theta_{p,q}} (\partial^{\bar{\delta}_{p,1}} \bar{u})^{\bar{\theta}_{p,1}} \dots (\partial^{\bar{\delta}_{p,q}} \bar{u})^{\bar{\theta}_{p,q}}.$$

Here $p \in \{0, 1, \dots, m\}$, $(\delta_{p,j}, \bar{\delta}_{p,j}) \in \mathbb{N}^n \times \mathbb{N}^n$, $\bar{\theta} \in \mathbb{N}^*$, and $(\theta_{p,j}, \bar{\theta}_{p,j}) \in \mathbb{N} \times \mathbb{N}$ are such that $|\delta_{p,j}| = |\bar{\delta}_{p,j}| = p$ and $\sum_{l=1}^m l \theta'_l = m$ with $\theta'_l := \sum_{j=1}^q \theta_{l,j} + \sum_{j=1}^{\bar{q}} \bar{\theta}_{l,j}$.

We prove the following claim:

Claim:

(1) $\theta'_m \in \{0, 1\}$, and if $\theta'_m = 1$ then $\theta'_{m-1} = \dots = \theta'_1 = 0$.

(2) Let $l \in \{1, \dots, m-1\}$. If $\theta'_m = \dots = \theta'_{m-(l-1)} = 0$ and $\theta'_{m-l} \neq 0$ then $\sum_{j=1}^{m-l} \theta'_j \leq l+1$.

The proof of the first statement is left to the reader. Clearly $\theta'_{m-l} \leq \frac{m}{m-l} \leq l+1$.

Hence $\theta'_{m-l} = q$ with $q \in \{1, \dots, l+1\}$. From $(m-l)q + \sum_{j=1}^{m-(l+1)} j \theta'_j \leq m$ we get

$\sum_{j=1}^{m-(l+1)} \theta'_j \leq lq - m(q-1) \leq l+1-q$, which implies that the estimate of the second claim holds.

The following elementary estimates hold:¹⁸

($((n = 4 \text{ and } l \geq 2) \text{ or } (n = 3 \text{ and } l \geq 1))$ and $1 \leq \delta \leq l + 1$ and $1 \leq \bar{m} \leq m - l$ and $0 \leq \bar{\alpha} \leq \alpha$) or
 $((n, l) = (4, 1)$ and $1 \leq \delta \leq 2$ and $1 \leq \bar{m} \leq m - 2$ and $0 \leq \bar{\alpha} \leq \alpha$) :

$$\|D^{\bar{m}+\bar{\alpha}}u\|_{L_t^{\frac{2\delta(n+2)}{n}}L_x^{\frac{2\delta(n+2)}{n}}(J)} \lesssim \|u\|_{L_t^{\frac{2\delta(n+2)}{n}}\tilde{H}^{k-\frac{1}{4},\frac{2(n+2)\delta}{(n+2)\delta-2}}(J)};$$

$$((n = 4 \text{ and } l \in \{0, 1\}) \text{ or } (n = 3 \text{ and } l = 0)) \text{ and } 1 \leq \delta \leq l + 1 \text{ and } 1 \leq \bar{m} \leq m - l \text{ and } 0 \leq \bar{\alpha} \leq \alpha : \\ \|D^{\bar{m}+\bar{\alpha}}u\|_{L_t^{\frac{2\delta(n+2)}{n}}L_x^{\frac{2\delta(n+2)}{n}}(J)} \lesssim \|u\|_{L_t^{\frac{2\delta(n+2)}{n}}\tilde{H}^{k,\frac{2(n+2)\delta}{(n+2)\delta-2}}(J)};$$

$l \geq 2$ and $1 \leq \delta \leq l + 2$ and $0 \leq \bar{m} \leq m - l$:

$$\|D^{\bar{m}}u\|_{L_t^{\frac{2\delta(n+2)}{n}}L_x^{\frac{2\delta(n+2)}{n}}(J)} \lesssim \|u\|_{L_t^{\frac{2\delta(n+2)}{n}}\tilde{H}^{k-\frac{1}{4},\frac{2(n+2)\delta}{(n+2)\delta-2}}(J)};$$

$$n = 4 : \begin{cases} \alpha \leq \bar{\alpha} \leq (\alpha+)+ : \|D^{\bar{\alpha}}P_{\geq 1}u\|_{L_t^\infty L_x^{\infty-}(J)} \lesssim \|D^{\psi(k)}u\|_{L_t^\infty L_x^2(J)}, \\ \alpha > 0, 1 \leq \bar{m} \leq m - 1 : \|D^{\bar{m}}u\|_{L_t^{\frac{4(n+2)}{n}}L_x^{\frac{4(n+2)}{n}+}(J)} \lesssim \|u\|_{L_t^{\frac{4(n+2)}{n}}\tilde{H}^{k,\frac{4(n+2)}{2(n+2)-2}}(J)}, \text{ and} \\ \alpha > 0 : \|D^m u\|_{L_t^{\frac{2(n+2)}{n}}L_x^{\frac{2(n+2)}{n}+}(J)} \lesssim \|u\|_{L_t^{\frac{2(n+2)}{n}}\tilde{H}^{k,\frac{2(n+2)}{n}}(J)}; \end{cases}$$

$$n = 3 : \begin{cases} \|Du\|_{L_t^{\frac{2(n+2)}{n}}L_x^{\frac{15}{2}}(J)} \lesssim \|u\|_{L_t^{\frac{2(n+2)}{n}}\tilde{H}^{k,\frac{2(n+2)}{n}}(J)}, \text{ and} \\ \alpha \leq \bar{\alpha} \leq (\alpha+)+ : \begin{cases} m > 1 : \|D^{\bar{\alpha}}P_{\geq 1}u\|_{L_t^\infty L_x^\infty(J)} \lesssim \|D^{\psi(k)}u\|_{L_t^\infty L_x^2(J)} \\ m = 1 : \|D^{\bar{\alpha}}P_{\geq 1}u\|_{L_t^\infty L_x^{6-}(J)} \lesssim \|D^{\psi(k)}u\|_{L_t^\infty L_x^2(J)} \end{cases} \end{cases}$$

There exists $1 \geq \theta \geq 0$ ¹⁹ such that

$$k' > 0, \delta \in \{1, \dots, l + 2\} : \|u\|_{L_t^{\frac{2\delta(n+2)}{n}}\tilde{H}^{k',\frac{2\delta(n+2)}{\delta(n+2)-2}}(J)} \lesssim \|u\|_{L_t^{\frac{2(n+2)}{n}}\tilde{H}^{k',\frac{2(n+2)}{n}}(J)}^\theta \|u\|_{L_t^\infty \tilde{H}^{k'}(J)}^{1-\theta}$$

$$0 \leq \bar{\alpha} \leq 1 : \|D^{\bar{\alpha}}P_{< 1}u\|_{L_t^\infty L_x^\infty(J)} \lesssim \|Du\|_{L_t^\infty L_x^2(J)}.$$

$\|D^k(g(|u|))\|_{L_t^{\frac{2(n+2)}{n}}L_x^{\frac{2(n+2)}{n}}(J)}$ is bounded by a finite sum of terms of the form²⁰

$$\bar{Y}_p := \|u\|_{L_t^{q_0}L_x^{r_0}(J)}^{\theta'_0} \cdots \|D^{p-1}u\|_{L_t^{q_{p-1}}L_x^{r_{p-1}}(J)}^{\theta'_{p-1}} \\ \|D^p u\|_{L_t^{q_p}L_x^{r_p}(J)}^{\theta'_p-1} \|D^{\alpha+p}u\|_{L_t^{q_p}L_x^{r_p}(J)} \\ \|D^{p+1}u\|_{L_t^{q_{p+1}}L_x^{r_{p+1}}(J)}^{\theta'_{p+1}} \cdots \|D^{m-l}u\|_{L_t^{q_{m-l}}L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|\partial^{\bar{\theta}} \tilde{g}(|u|^2)\|_{L_t^{q'}L_x^{r'}(J)},$$

(with $p \in \{0, \dots, m-l\}$, $\sum_{j=0:j \neq p}^{m-l} \frac{\theta'_j}{q_j} + \frac{\theta'_p-1}{q_p} + \frac{1}{q'} = \frac{n}{2(n+2)}$, and $\sum_{j=0:j \neq p}^{m-l} \frac{\theta'_j}{r_j} + \frac{\theta'_p-1}{r_p} + \frac{1}{r'} = \frac{n}{2(n+2)}$), and of the form

$$\tilde{Y} := \|u\|_{L_t^{q_0}L_x^{r_0}(J)}^{\theta'_0} \cdots \|D^{m-l}u\|_{L_t^{q_{m-l}}L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|D^{\alpha} \partial^{\bar{\theta}} \tilde{g}(|u|^2)\|_{L_t^{q'}L_x^{r'}(J)}$$

¹⁸In the sequel $\tilde{H}^{k,p} := D^{-1}L^p \cap D^{-k}L^p$

¹⁹In the sequel we allow the value of θ to change from one line to the other one. Here $\widehat{P_{<1}f}(\xi) := \phi(\xi)\hat{f}(\xi)$ with ϕ a bump function equal to one for $|\xi| \leq 1$ and supported on $|\xi| \leq 2$ and $\widehat{P_{\geq 1}f}(\xi) := \hat{f}(\xi) - \widehat{P_{<1}f}(\xi)$.

²⁰In the sequel if some terms do not make sense we do not take them into account. Example: if $p = 0$ then one should not take into account the term where D^{p-1} appears. Also if $\theta'_j = 0$ for some $j \in \{0, \dots, m-l\}$ then we ignore all the terms where j appears.

(with $\sum_{j=0}^{m-l} \frac{\theta'_j}{q_j} + \frac{1}{q'} = \frac{n}{2(n+2)}$ and $\sum_{j=0}^{m-l} \frac{\theta'_j}{r_j} + \frac{1}{r'} = \frac{n}{2(n+2)}$). Here $l \in \{0, \dots, m-1\}$ and $\theta'_{m-l} \neq 0$.

We first estimate \bar{Y}_p .
Assume that $p \neq 0$.

Then with $q_0 = r_0 = \infty$ and $q_j = r_j = \bar{q}_p = \bar{r}_p = \frac{2(n+2)}{n} \sum_{s=1}^{m-l} \theta'_s$ for $j \neq 0$ we see from elementary estimates of the derivatives of g , the above claim, the above estimates, and elementary consequences of $\sum_{l=1}^m l\theta'_l = m$ that \bar{Y}_p is bounded by RHS' .

Assume that $p = 0$.

We first consider the case where $(n, l) \neq (3, 0)$, $(n, l) \neq (3, 1)$, $(n, l) \neq (4, 0)$, and $(n, l) \neq (4, 1)$. Letting $q_0 = r_0 = \infty$ and $\bar{q}_0 = \bar{r}_0 = q_j = r_j = \frac{2(n+2)}{n} \left(\sum_{s=1}^{m-l} \theta'_s + 1 \right)$ for $j \neq 0$ we see that \bar{Y}_p is bounded by RHS' .

We then consider the other cases. By decomposition one has to estimate

$$Z_{lo} := \|u\|_{L_t^{q_0} L_x^{r_0}(J)}^{\theta'_0-1} \|D^\alpha P_{<1} u\|_{L_t^{\bar{q}_0} L_x^{\bar{r}_0}(J)} \dots \|D^{m-l} u\|_{L_t^{q_{m-l}} L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|\partial^{\bar{\theta}} \tilde{g}(|u|^2)\|_{L_t^{q'} L_x^{r'}(J)}$$

$$\text{and } Z_{hi} := \|u\|_{L_t^{q_0} L_x^{r_0}(J)}^{\theta'_0-1} \|D^\alpha P_{\geq 1} u\|_{L_t^{\bar{q}_0} L_x^{\bar{r}_0}(J)} \dots \|D^{m-l} u\|_{L_t^{q_{m-l}} L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|\partial^{\bar{\theta}} \tilde{g}(|u|^2)\|_{L_t^{q'} L_x^{r'}(J)}.$$

Assume that $(n, l) = (3, 0)$. Letting $(q_0, r_0) = (\bar{q}_0, \bar{r}_0) = (\infty, \infty)$ and $q_m = r_m = \frac{2(n+2)}{2}$ we see that Z_{lo} is bounded by RHS' . Letting $q_0 = \bar{q}_0 = r_0 = \infty$, $\bar{r}_0 = 6-$ if $m = 1$ (resp. $\bar{r}_0 = \infty$ if $m > 1$), $q_m = \frac{2(n+2)}{n}$, and $r_m = \frac{15}{2} +$ if $m = 1$ (resp. $r_m = \frac{2(n+2)}{n}$ if $m > 1$), we see that Z_{hi} is bounded by RHS' . Assume that $(n, l) = (4, 0)$. Z_{lo} is bounded by RHS' , by assigning the same values to the exponents as for the case $(n, l) = (3, 0)$. Let $(q_0, r_0) = (\infty, \infty)$. If $\alpha > 0$ (resp. $\alpha = 0$) let $(\bar{q}_0, \bar{r}_0) = (\infty, \infty-)$ (resp. $(\bar{q}_0, \bar{r}_0) = (\infty, \infty)$) and $(q_m, r_m) = \left(\frac{2(n+2)}{n}, \frac{2(n+2)}{n} + \right)$ (resp. $\left(\frac{2(n+2)}{n}, \frac{2(n+2)}{n} \right)$): this implies that Z_{hi} is bounded by RHS' . Assume now that $(n, l) = (3, 1)$ or $(4, 1)$. Consider the subcase $(\theta'_1, \dots, \theta'_{m-2}, \theta'_{m-1}) = (0, \dots, 0, 2)$ (resp. $(\theta'_1, \dots, \theta'_{m-2}, \theta'_{m-1}) = (1, \dots, 0, 1)$). Letting $r_{m-1} = \frac{4(n+2)}{n-2}$ (resp. $r_1 = r_{m-1} = \frac{4(n+2)}{n-2}$), $(\bar{q}_0, \bar{r}_0) = (\infty, \infty)$, we see that Z_{lo} is bounded by RHS' . Consider the subcase $(\theta'_1, \dots, \theta'_{m-2}, \theta'_{m-1}) = (0, \dots, 0, 2)$. If $(n, l) = (4, 1)$ and $\alpha > 0$ (resp. $\alpha = 0$) let $(q_{m-1}, r_{m-1}) = \left(\frac{4(n+2)}{n}, \frac{4(n+2)}{n} + \right)$ and $(\bar{q}_0, \bar{r}_0) = (\infty, \infty-)$ (resp. $(q_{m-1}, r_{m-1}) = \left(\frac{4(n+2)}{n}, \frac{4(n+2)}{n} \right)$ and $(\bar{q}_0, \bar{r}_0) = (\infty, \infty)$). If $(n, l) = (3, 1)$ let $(q_{m-1}, r_{m-1}) = \frac{4(n+2)}{n}(1, 1)$ and $(\bar{q}_0, \bar{r}_0) = (\infty, \infty)$. This implies that Z_{hi} is bounded by RHS' . Now consider the subcase $(\theta'_1, \dots, \theta'_{m-2}, \theta'_{m-1}) = (1, \dots, 0, 1)$: this subcase is treated similarly, except that we assign the same value of r_{m-1} (resp. q_{m-1}) of the previous subcase to that of the variable r_1 (resp. q_1).

We then estimate \bar{Y} .

Writing $f = P_{<1} f + P_{\geq 1} f$, we see that given $p \geq 1$, $\|D^\alpha f\|_{L^p(\mathbb{R}^n)} \lesssim \|f\|_{L^p(\mathbb{R}^n)} + \|f\|_{\dot{B}_{p,p}^{\alpha,+}(\mathbb{R}^n)}$ and $\|f\|_{\dot{B}_{p,p}^{\alpha,+}(\mathbb{R}^n)} \lesssim \|f\|_{L^p(\mathbb{R}^n)} + \|D^{(\alpha+)+} f\|_{L^p(\mathbb{R}^n)}$. Here $\dot{B}_{p,p}^{\alpha,+}(\mathbb{R}^n)$ is the standard homogeneous Besov space. Elementary estimates show that $\|\partial^{\bar{\theta}} \tilde{g}(|u|^2)(x+h) - \partial^{\bar{\theta}} \tilde{g}(|u|^2)(x)\|_{L^p(\mathbb{R}^n)} \lesssim \|u(x+h) - u(x)\|_{L^p(\mathbb{R}^n)} \langle \bar{Q}_{\frac{n}{2}+}(J, u) \rangle^C$ for some constant

$C > 0$. Hence from the characterization of the Besov norm by the modulus of continuity, we see that $\|D^\alpha \partial^{\bar{\theta}} \tilde{g}(|u|^2)\|_{L^p(\mathbb{R}^n)} \lesssim \|\partial^{\bar{\theta}} \tilde{g}(|u|^2)\|_{L^p(\mathbb{R}^n)} + \|D^{(\alpha+)+} u\|_{L^p(\mathbb{R}^n)}$.

Hence one has to estimate

$$\tilde{Y}_1 := \|u\|_{L_t^{q_0} L_x^{r_0}(J)}^{\theta'_0} \cdots \|D^{m-l} u\|_{L_t^{q_{m-l}} L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|\partial^{\bar{\theta}} \tilde{g}(|u|^2)\|_{L_t^{q'} L_x^{r'}(J)}, \text{ and}$$

$$\tilde{Y}_2 := \|u\|_{L_t^{q_0} L_x^{r_0}(J)}^{\theta'_0} \cdots \|D^{m-l} u\|_{L_t^{q_{m-l}} L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|D^{(\alpha+)+} u\|_{L_t^{q'} L_x^{r'}(J)}. \text{ We write } \tilde{Y}_2 = \tilde{Y}_{2,lo} + \tilde{Y}_{2,hi} \text{ with}$$

$$\tilde{Y}_{2,lo} := \|u\|_{L_t^{q_0} L_x^{r_0}(J)}^{\theta'_0} \cdots \|D^{m-l} u\|_{L_t^{q_{m-l}} L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|D^{(\alpha+)+} P_{<1} u\|_{L_t^{q'} L_x^{r'}(J)} \text{ and } \tilde{Y}_{2,hi} := \|u\|_{L_t^{q_0} L_x^{r_0}(J)}^{\theta'_0} \cdots \|D^{m-l} u\|_{L_t^{q_{m-l}} L_x^{r_{m-l}}(J)}^{\theta'_{m-l}} \|D^{(\alpha+)+} P_{\geq 1} u\|_{L_t^{q'} L_x^{r'}(J)}.$$

We first consider the case when $(n, l) \neq (3, 0)$, $(n, l) \neq (3, 1)$, $(n, l) \neq (4, 0)$, and $(n, l) \neq (4, 1)$. We can estimate \tilde{Y}_1 by RHS' , assigning the same values for $q_0, r_0, \dots, q_{m-l}, r_{m-l}$ (resp. q', r') as those for the same exponents (resp. \bar{q}_0, \bar{r}_0) when we estimated \tilde{Y}_0 . We can estimate $\tilde{Y}_{2,lo}$ (resp. $\tilde{Y}_{2,hi}$) by RHS' , assigning the same values for $q_0, r_0, \dots, q_{m-l}, r_{m-l}$ (resp. q', r') as those for the same exponents (resp. \bar{q}_0, \bar{r}_0) when we estimated Z_{lo} (resp. Z_{hi} , with $\alpha > 0$). We then consider the other cases. We can estimate $\tilde{Y}_{2,lo}$ and \tilde{Y}_1 (resp. $\tilde{Y}_{2,hi}$) by RHS' , assigning the same values for $q_0, r_0, \dots, q_{m-l}, r_{m-l}$ (resp. q', r') as those for the same exponents (resp. \bar{q}_0, \bar{r}_0) when we estimate Z_{lo} (resp. Z_{hi}).

A straightforward modification of the proof shows that (160) holds if u^λ is replaced with $u^{\lambda_1} \bar{u}^{\lambda_2}$ with $(\lambda_1, \lambda_2) \in \mathbb{N}^2$ such that $\lambda_1 + \lambda_2 = \lambda$.

If we replace $g(|u|)$ with $\tilde{g}'(|u|^2) u^{\lambda_3} \bar{u}^{\lambda_4}$, then (160) also holds by replacing in the proof $\partial^{\bar{\theta}} \tilde{g}(|u|^2)$ with $\partial^{\bar{\theta}+1} \tilde{g}(|u|^2) u^{\lambda_a} \bar{u}^{\lambda_b}$, taking into account that $\left\| \partial^{\bar{\theta}+1} \tilde{g}(|u|^2) u^{\lambda_a} \bar{u}^{\lambda_b} \right\|_{L^r(\mathbb{R}^n)} \lesssim \left\| \partial^{\bar{\theta}} \tilde{g}(|u|^2) \right\|_{L^r(\mathbb{R}^n)}$ and $\left\| D^\alpha \left(\partial^{\bar{\theta}+1} \tilde{g}(|u|^2) u^{\lambda_a} \bar{u}^{\lambda_b} \right) \right\|_{L^r(\mathbb{R}^n)} \lesssim \left\| \partial^{\bar{\theta}+1} \tilde{g}(|u|^2) u^{\lambda_a} \bar{u}^{\lambda_b} \right\|_{L^r(\mathbb{R}^n)} + \|D^{(\alpha+)+} u\|_{L^r(\mathbb{R}^n)}$ (Here $r \in [1, \infty]$ and $(\lambda_a, \lambda_b) \in \mathbb{N}^2$ such that $\lambda_a + \lambda_b = 2$). The proof is left to the reader. \square

7. APPENDIX C

We shall prove the following proposition:

Proposition 9. *Let u be a solution of (1) with data $u_0 \in \tilde{H}^k$, $k > \frac{n}{2}$. Assume that u exists globally in time and that $\|u\|_{L_t^{\frac{2(n+2)}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(\mathbb{R})} < \infty$. Then $Q(\mathbb{R}, u) < \infty$.*

Proof. By symmetry we may WLOG restrict ourselves to \mathbb{R}^+ .

First assume that $\frac{n+2}{n-2} > k > \frac{n}{2}$. Repeating the same steps from ‘ $J := [0, a]$ ’ just below (45) to (49) included and replacing ‘ $[0, T']$ ’ (resp. ‘LHS of (48) \gtrsim ’) with ‘ \mathbb{R}^+ ’ (resp. ‘ $\infty >$ ’) we get $Q(\mathbb{R}^+, u) < \infty$.

Now assume that $k \geq \frac{n+2}{n-2}$. In view of the previous paragraph it is sufficient to show that if for all $1 \leq j \leq k - \frac{1}{4}$, $Q_j(\mathbb{R}^+, u) < \infty$, then $Q(\mathbb{R}^+, u) < \infty$. Let $J := [0, a]$ be an interval. By (16), (15), Proposition 7 and Proposition 8, we get

$$Q(J, u) \lesssim \|u_0\|_{\tilde{H}^k} + Q(J, u) \|u\|_{L_t^{\frac{4}{n-2}} L_x^{\frac{2(n+2)}{n-2}}(J)}^{\frac{2(n+2)}{n-2}} g(Q(J, u)).$$

Again repeating the same steps from ‘ $J := [0, a]$ ’ just below (45) to (49) included, and taking into account the replacements that were pointed out for the case $\frac{n+2}{n-2} > k > \frac{n}{2}$, we get $Q(\mathbb{R}^+, u) < \infty$. □

REFERENCES

- [1] J. Bourgain, *Global well-posedness of defocusing 3D critical NLS in the radial case*, JAMS 12 (1999), 145-171
- [2] T. Cazenave, F.B. Weissler, *The Cauchy problem for the critical nonlinear Schrödinger equation in H^s* , Nonlinear. Anal. 14 (1990), no. 10, 807-836.
- [3] M. Christ and M. Weinstein, *Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation*, J. Funct. Analysis 100 (1991), 87-109
- [4] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, *Global well-posedness and scattering in the energy space for the critical nonlinear Schrödinger equation in \mathbb{R}^3* , Annals of Math. 167 (2008), 767-865
- [5] M. Grillakis, *On nonlinear Schrödinger equations*, Comm. Partial Differential Equations 25 (2000), no 9-10, 1827-1844
- [6] M. Keel and T. Tao, *Endpoint Strichartz estimates*, Amer. Math. J. 120 (1998), 955-980
- [7] E. Rickman and M. Visan, *Global well-posedness and scattering for the defocusing energy-critical nonlinear Schrödinger equation in R^{1+4}* , Amer. J. Math. 129 (2007), 1-60
- [8] M. Taylor, *Tools for PDE. Pseudodifferential operators, paradifferential operators, and layer Potentials*. Mathematical Surveys and Monographs, 81. American Mathematical Society, Providence, RI, 2000.
- [9] T. Roy, *One remark on barely \dot{H}^{s_p} supercritical wave equations*, preprint, arXiv:0906.0044
- [10] T. Tao, *Global well-posedness and scattering for the higher-dimensional energy-critical nonlinear Schrödinger equation for radial data*, New York J. Math., 11, 2005, 57-80
- [11] M. Visan, *The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions*, Duke Math. J. 138 (2007), 281-374.

NAGOYA UNIVERSITY

E-mail address: `tristanroy@math.nagoya-u.ac.jp`