BLACK HOLE (THERMODYNAMICS) - A SIMPLIFIED THEORY FOR QUANTUM GRAVITY NON-SPECIALISTS

Vladan Panković

Department of Physics, Faculty of Sciences, 21000 Novi Sad, Trg Dositeja Obradovića 4., Serbia

PACS number: 04.70.Dy

Abstract

In this work we suggest a simplified "quasi-classical" formalism of the Schwarzschild black hole thermodynamics. We define such small quantum system at Schwarzschild black hole horizon surface whose reduced Compton wavelength equals one circumference of a great circle on the black hole horizon surface. (It is in some way similar to the Bohr orbital momentum quantization postulate for the ground state interpreted via de Broglie relation.) It admits very simple functional connections between all black hole basic thermodynamic characteristics (Unruh temperature, Hawking temperature, Bekenstein-Hawking entropy, Bekenstein entropy horizon surface quantization, evaporation time) and small quantum system mass. All this can be very interesting for the quantum gravity specialists as well as non-specialists.

Key words: black hole, Hawking temperature

In this work we shall suggest a simplified "quasi-classical" formalism of the Schwarzschild black hole thermodynamics. We shall define such small quantum system at Schwarzschild black hole horizon surface whose reduced Compton wavelength equals one circumference of a great circle on the black hole horizon surface. (It is in some way similar to the Bohr orbital momentum quantization postulate for the ground state interpreted via de Broglie relation.) It admits very simple functional connections between all black hole basic thermodynamic characteristics (Unruh temperature, Hawking temperature, Bekenstein-Hawking entropy, Bekenstein entropy horizon surface quantization, evaporation time [1]-[4]) and small quantum system mass. All this can be very interesting for the quantum gravity specialists as well as non-specialists ("without knowing the details of quantum gravity", we paraphrase Fursaev [5]).

Consider a large, massive spherical system with mass M and characteristic radius R.

Define the following reduced Compton wavelength λ_{r} and corresponding small system mass m

(1)
$$\lambda_r = \hbar/(mc) = 2\pi R$$

(2)
$$m = \hbar/(2\pi Rc)$$

for m << M where h represents reduced Planck constant and c – speed of light. Here $2\pi R$ represents the circumference of the large system, i.e. circumference of a great circle at large system surface. Expression (1) simply means that circumference of the large system holds exactly one mentioned reduced Compton wavelength. (It is in some way similar to Bohr angular momentum quantization postulate interpreted via de Broglie relation for ground state.)

Expression (2), after multiplication with GM/R where G represents the Newtonian gravitational constant, implies

(3) GmM/R =
$$\hbar/(2\pi c)$$
 GM/R² = $\hbar/(2\pi c) a = kT$

Here GmM/R represents classical potential energy (absolute value) of the gravitational interaction between large system and small system with mass m at large system surface, $a = GM/R^2$ represents surface gravity (absolute value) and T - Unruh temperature multiplied with Boltzmann constant k.

For Schwarzschild black hole R represents Schwarzschild radius

$$(4) R = 2GM/c^2$$

which, according to (2),(3), implies

(5)
$$m = \hbar c / (4\pi GM)$$

(6)
$$kT = \hbar c^3 / (8\pi GM) = mc^2 / 2$$

Last expression means that T represents Hawking temperature and that it can be simply connected with m. Moreover, expression (5) simply implies

(7)
$$M/m = (4\pi G M^2) / (\hbar c) = (4\pi R^2) / (2 L_P)^2 = A/(2 L_P)^2 = A(2 L_P) / (2 L_P)^3 = S/k$$

where $L_P = (\hbar G/c^3)^{1/2}$ represents the Planck length, $A = (4\pi R^2) - Schwarzschild black hole horizon surface, <math>A(2 L_P) - volume$ of the tiny spherical shell in horizon surface vicinity and S - Bekenstein-Hawking entropy of Schwarzschild black hole. In this way Schwarzschild black hole entropy is simply connected with m too. Moreover, according to (7), Bekenstein-Hawking entropy of Schwarzschild black hole can be considered as the number of different ways at which small quantum system with mass m that occupies surface (2 L_P)² or more precisely volume (2 L_P)³ can be placed at horizon surface or more precisely in spherical shell volume.

Further, as it is well known, Schwarzschild black hole satisfies thermodynamic

$$dMc^2 = T dS$$

or

law

(9)
$$dS = dMc^2/T$$

where Mc^2 represents Schwarzschild black hole total energy. Then, after introduction of (6) in (9) and after simpler integration, it follows expression equivalent to (7).

Suppose, however, that differential form of (9) is approximately changed by corresponding finite difference form

(10)
$$\Delta S = c^2 / T \Delta M$$

and suppose additionally

(11)
$$\Delta M = n m$$
 for $n = 1, 2, ...$

Then, according to (6) and (11), (10) turns out in

(12)
$$\Delta S = 2n k$$
 for $n = 1, 2, ...$

that implies

(13)
$$\Delta A = 2n (2 L_P)^2$$
 for $n = 1, 2, ...$

Expressions (12), (13) represent, obviously, Bekenstein quantization of the Schwarzschild black hole entropy and horizon surface. In this way Bekenstein quantization of the Schwarzschild black hole entropy (12) and horizon surface (13) are simply connected with Schwarzschild black hole mass quantization (11), i.e. with m too.

Further, as it is well-known, time t_e of Schwarzschild black hole total evaporation via Hawking radiation can be obtained after integration of the Stefan-Boltzmann law

(14)
$$dM/dt c^2 = -\sigma T^4 A$$

(where $\sigma = (\pi^2 k^4)/(60 \hbar^3 c^2)$ represents Stefan-Boltzmann constant) under conditions (4), (6)., or after integration of the following expression.

(15)
$$dt = - dM c^2 / (\sigma T^4 A) = 15360\pi (T_P / M^3_P) M^2 dM$$

where $T_P = (\hbar G/c^5)^{1/2}$ represents the Planck time and $M_P = (\hbar c/G)^{1/2}$ represents the Planck mass. It yields

(16)
$$t_e = 5120\pi (M^3/M_P^3) T_P$$

which means that total evaporation time is proportional to the cube of the initial mass of Schwarzschild black hole.

Suppose, however, that differential form of (15) is approximately changed by corresponding finite difference form

(17)
$$\Delta t = 15360\pi (T_{P}/M^{3}_{P}) M^{2} \Delta M$$

under condition (11). It, according to (5), implies

(18)
$$\Delta t = n \ 3820 \ (M/M_P)T_P$$
 for $n = 1, 2, ...$

which means that "horizon surface evaporation time" Δt is proportional to (first degree of) corresponding mass of Schwarzschild black hole. It represents an interesting result simply connected with Schwarzschild black hole mass quantization (11), i.e. with m too.

In conclusion we can shortly repeat and point out the following. In this work we suggest a simplified "quasi-classical" formalism of the Schwarzschild black hole thermodynamics. We define such small quantum system at Schwarzschild black hole horizon surface whose reduced Compton wavelength equals one circumference of a great circle on the black hole horizon surface. (It is in some way similar to the Bohr orbital momentum quantization postulate for the ground state interpreted via de Broglie relation.) It admits very simple functional connections between all black hole basic thermodynamic characteristics (Unruh temperature, Hawking temperature, Bekenstein-Hawking entropy, Bekenstein entropy horizon surface quantization, evaporation time) and small quantum system mass. All this can be very interesting for the quantum gravity specialists as well as non-specialists.

References

[1] R. M. Wald, *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics* (University of Chicago press, Chicago, 1994)

[2] R. M. Wald, The Thermodynamics of Black Holes, gr-qc/9912119

[3] D. N. Page, Hawking Radiation and Black Hole Thermodynamics, hep-th/0409024

[4] P. M. Alsing, P. W. Milloni, *Simplified Derivation of the Hawking-Unruh temperature for an accelerating observer in vacuum*, Am. J. Phys. 72 (2004) 1524 ; quant-ph/0401170 v2

[5] D. V. Fursaev, Can One Understand Black Hole Entropy without Knowing Much about Quantum Gravity?, gr-qc/0404038