High-Order Coupled Cluster M ethod (CCM) Form alism 2 {

\G eneralised" Expectation Values: Spin-Spin Correlation Functions for Frustrated and Unfrustrated 2D Antiferrom agnets

D.J.J.Famell

Health Methodology Research Group, School of Community-Based Medicine, Jean McFarlane Building, University Place, University of Manchester, Manchester M13 9PL, United Kingdom (Dated: February 22, 2024)

Abstract

Recent developments of high-order CCM have been to extend existing formalism and codes to s $\frac{1}{2}$ for both the ground and excited states. In this article, we describe how \generalised" expectation values for a wide range of one- and two-body spin operators may also be determined using existing the CCM code for the ground state. We present new results for the spin-spin correlation functions of the spin-half square-and triangular-lattice antiferrom agnets by using the LSUBm approximation. We show that the absolute values of the spin-spin correlation functions fs(0) s(r) ij converge with increasing approximation level for both lattices. We believe that the LSUBm approxim ation provides reasonable results for the correlation functions for lattice separations roughly m for the square lattice. We compare qualitatively our results for the square lattice of order r to those results of quantum M onte C arlo (QMC) and we see that good correspondence is observed. Indeed as seen by QMC, the spin-spin correlation function initially decays strongly with jrjbefore becoming constant for larger values of jrj. CCM results are also compared to results of exact diagonalisations for both lattices. ED results dem onstrate a strong nite-size e ects at lattice separations r = L=2 (where N = L L) for both lattices. The behaviour of the correlation function for the triangular lattice is qualitatively sim ilar to that of the square lattice, nam ely, that it decays strongly at rst before becoming constant. This is in keeping with the behaviour of both models, which are believed strongly to be Neel-ordered from approximate studies. The CCM has been show n m any times to provide consistently good results for the ground-state energy and sublattice magnetisation of a wide range of quantum spin models. Here we have shown that the CCM also provides good results for the spin-spin correlation function.

I. IN TRODUCTION

The coupled cluster method (CCM) [1, 2, 3, 4, 5, 6, 7, 8, 9] is a well-known method of quantum many-body theory (QMBT). The CCM has been applied with much success in order to study quantum magnetic systems at zero temperature (see Refs. [10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52]). In particular, the use of computer-algebraic in plem entations [22, 26, 32, 52] of the CCM has been found to be very elective with respect to these spin-lattice problems. Recent developments of high-order CCM form alism and codes have been to treat systems with spin quantum number of s $\frac{1}{2}$ for both the ground and excited states [52]. In this article, we show how these ground-state form alism and codes may also be used directly to ind \generalised" expectation values; that is, expectation values for a wide range of one- or two-body spin operator that are de ned prior to the CCM calculation.

We apply the CCM to the spin-half Heisenberg model on the square and triangular lattices. The Hamiltonian is specied as follows,

$$H = \sum_{\substack{i \\ hi; ji}}^{X} s_{i} s_{j}$$
(1)

where the sum on hi; ji counts all nearest-neighbour pairs once. The ground states of all of the cases considered here are classically ordered, albeit by a reduced am ount due to quantum uctuations. Indeed, the best estimates of the am ount of classical ordering of the square lattice case from approximate methods are 61% from CCM [41], 61.4% from quantum M onte Carlo studies [53], 61.4% from series expansions [54], 61.38% from spin-wave theory [55], and 63.4% from exact diagonalisations [56]. G cod results for the spin-spin correlation function of the spin-half square-lattice H eisenberg antiferrom agnet were found using quantum M onte Carlo in R ef. [57] on lattices of size up to N = L L, where L = f16;22;32g. They observed that the correlation functions decayed with separation r. However, they also saw nite-size e ects, indicated by cusp-like behaviour in these correlation functions, at distances given by L=2. Exact diagonalisations have also been carried out for the correlation function of the square lattice [58]. For the spin-half triangular-lattice antiferrom agnet, the best estimates of the am ount of classical ordering are 41% from quantum M onte Carlo studies[59], 40% from series expansions[60], 39% from exact diagonalisations[56], and 40% from previous CCM calculations [41]. Very few results for the spin-spin correlation functions of the triangular lattice seem to exist apart from results of exact diagonalisations (e.g., see [62]).

II. M E T H O D

The details of the practical application of high-order coupled cluster m ethod (CCM) form alism to lattice quantum spin systems are given in Refs. [22, 26, 32, 38, 52] and also in the appendices to this article. However, we point out now that the ket and bra ground-state energy eigenvectors, j i and h \sim j of a generalm any-body system described by a H am iltonian H, are given by

$$Hji = E_{q}ji; h^{H} = E_{q}h^{j};$$
(2)

Furtherm ore, the ket and bra states are parametrized within the single-reference CCM as follows:

$$ji = e^{S}ji ; S = \overset{X}{\underset{I \in 0}{}} S_{I}C_{I}^{+};$$
$$h^{j} = hjSe^{S} ; S = 1 + \overset{X}{\underset{I \in 0}{}} S_{I}C_{I} :$$
(3)

O ne of the m ost in portant features of the CCM is that one uses a single m odel or reference state j i that is norm alized. We note that the param etrisation of the ground state has the norm alization condition for the ground-state bra and ket wave functions (h~j i h j i = 1). The m odel state is required to have the property of being a cyclic vector with respect to two well-de ned A belian subalgebras of multi-con gurational creation operators fC⁺_I g and their Herm itian-adjoint destruction counterparts fC_I (C⁺_I)^yg. The interested reader is referred to the Appendices and to Ref. [52] for more inform ation regarding how the CCM problem is solved for.

Here, we use the classical ground state as the model state. For the square lattice, this is the Neel state in which neighbouring spins are antiparallel, and, for the triangular lattice, this a Neel-lke state in which neighbouring spins on three sublattices are at 120 to each other. For the square lattice, we perform a rotation of the local axes of the up-pointing spins by 180 about the y-axis. The transform ation is described by,

$$s^{x} ! s^{x}; s^{y} ! s^{y}; s^{z} ! s^{z} :$$
 (4)

The model state now appears mathematically to consist of purely down-pointing spins. In terms of the spin raising and lowering operators $s_k = s_k^x = is_k^y$ the Hamiltonian may be written in these local axes as,

$$H = \frac{1}{2} \sum_{i_{1};j_{1}}^{X^{N}} s_{j}^{+} s_{j}^{+} + s_{j} s_{j} + 2s_{j}^{z} s_{j}^{z} ; \qquad (5)$$

where the sum on hi; ji again counts all nearest-neighbour pairs once on the square lattice. A gain, the classical ground-state of the H eisenberg m odel of Eq. (1) for the triangular lattice is the N eel-like state where all spins on each sublattice are separately aligned (all in the xz-plane, say). The spins on sublattice A are oriented along the negative z-axis, and spins on sublattices B and C are oriented at + 120 and 120, respectively, with respect to the spins on sublattice A.W e again rotate the local spin axes of those spins on the di erent sublattices. Speci cally, we leave the spin axes on sublattice A unchanged, and we rotate about the y-axis the spin axes on sublattices B and C by 120 and + 120 respectively,

$$s_{B}^{x} ! \frac{1}{2} s_{B}^{x} \frac{p_{\overline{3}}}{2} s_{B}^{z} ; s_{C}^{x} ! \frac{1}{2} s_{C}^{x} + \frac{p_{\overline{3}}}{2} s_{C}^{z} ;$$

$$s_{B}^{x} ! s_{B}^{y} ! s_{B}^{y} ; s_{C}^{v} ! s_{C}^{v} ; s_{C}^{y} ;$$

$$s_{B}^{z} ! \frac{p_{\overline{3}}}{2} s_{B}^{x} \frac{1}{2} s_{B}^{z} ; s_{C}^{z} ! \frac{q_{\overline{3}}}{2} s_{C}^{x} \frac{1}{2} s_{C}^{z} ;$$
(6)

Once again, the model state now appears to consist of purely down-pointing spins. We may rewrite Eq. (1) in terms of spins de ned in these local quantisation axes for the triangular lattice, such that

$$H = \sum_{\substack{\text{hi! ji} \\ \text{hi! ji}}}^{\text{X} \text{n}} \frac{1}{2} s_{i}^{z} s_{j}^{z} + \frac{p_{\overline{3}}}{4} s_{i}^{z} s_{j}^{+} + s_{i}^{z} s_{j} s_{i}^{+} s_{j}^{z} s_{i} s_{j}^{z} s_{j}^{z} + \frac{1}{8} s_{i}^{+} s_{j} + s_{i} s_{j}^{+} \frac{3}{8} s_{i}^{+} s_{j}^{+} + s_{i} s_{j}^{-} s_{i}^{-} s_{j}^{-}$$

$$(7)$$

We note that the summation in Eq. (7) again runs over nearest-neighbour bonds, but now also with a directionality indicated by hi! ji, which goes from A to B, B to C, and C to A.

The CCM form alism is exact in the limit of inclusion of all possible multi-spin cluster correlations within S and S, although this is usually impossible to achieve practically. Hence, we generally make approximations in both S and S. The three most commonly employed approximation schemes previously utilised have been: (1) the SUBn scheme, in which all correlations involving only n or fewer spins are retained, but no further restriction is made

concerning their spatial separation on the lattice; (2) the SUBn-m sub-approximation, in which all SUBn correlations spanning a range of nomore than m adjacent lattice sites are retained; and (3) the localised LSUBm scheme, in which all multi-spin correlations over all distinct locales on the lattice de ned by m or fewer contiguous sites are retained. Another in portant feature of the method is that the bra and ket states are not always explicitly constrained to be Herm itian conjugates when we make such approximations, although the in portant Helm ann-Feynman theorem is always preserved. We remark that the CCM provides results in the in nite-lattice limit N ! 1 from the outset.

In this article, we wish to determ ine the spin-spin correlation functions js(0) s(r)ij as a function of lattice separation jrj for the (unfrustrated) square-lattice and (frustrated) triangular-lattice antiferrom agnets. We must take into account the rotation of the local spin axes, although this proceeds in exactly the sam emanner as for the Ham iltonian above. A gain, we remark that the manner in which high-order CCM may be solved has been discussed in Ref. [52]. The manner in which the ground-state CCM ket- and bra-state equations are solved is discussed in the Appendix to this article. In particular, the method by which \generalised expectation values" for a variety of one- and two-body spin operators may be obtained is explained in the appendices.

III. RESULTS

The results for the absolute values of the spin-spin correlation functions for the (unfrustrated) spin-half square-lattice antiferrom agnet using the LSUBm approximation with m = f2;4;6;8g are shown in Fig. 1. The LSUBm results are in very good mutual agreement for small lattice separations for dierent values of m. Indeed, LSUBm results are clearly converging with increasing levels of approximation. Furthermore, we see that LSUB6 and LSUB8 correspond reasonably well with each other up to r < 5. From Fig. 1 (as a \nule of thum b" only), we believe LSUBm results ought to provide reasonable results to lattice separations approximately of order m for the square lattice.

CCM results are also compared to those results of exact diagonalisations (ED) [58] in Fig. 1. However, we see that we obtain good correspondence only for very sm all lattice separations r < 3. (W e note that N = 40 is at the current upper limit for ED of computational tractability.) However, the agreement between ED and CCM becomes rapidly worse for

6



FIG.1: CCM results for the absolute values of the spin-spin correlation function for the spin-half square-lattice antiferrom agnet using the LSUBm approximation with m = f2;4;6;8g. CCM results are compared to those results of exact diagonalisations (ED) with N = 36;40 [58].

3. This disparity can be understood by considering results of quantum lattice separation r Monte Carlo (QMC) of Ref. [57] for the spin-spin correlation function, which were carried out for for much larger lattices of size of N = L L, where L = f16;22;32q. Indeed, we nd good correspondence qualitatively (ie., com pared by by eye only) between the LSUB8 results of Fig. 1 and those QMC results for the largest lattice N = 3232 of Fig. 3a in Ref. [57]. Furthermore, the authors of Ref. [57] noted a strong cusp-like behaviour was seen in the spin-spin correlation function at lattice separations of L=2 due to nite-lattice e ects. This is also seen in Fig. 1 for the ED results at r = 3, although this cusp may be exacerbated by a sm all \kink" in the \true" spin-spin correlation function that occurs at this point anyway (as seen in both CCM and QMC results). Interestingly, we see from Fig. 3b of Ref. [57] that the accuracy of QMC results for the spin-spin correlation functions becomes more problematic (again as a \rule of thum b" only) for separations of order r > L=2 for a given lattice size N = L L. Hence, we might also reasonably expect that results of ED m ight only be good for separations of r < 3 for lattices of size N = 36 and n = 40. Indeed, we note that ED agrees well with highly converged CCM results (and incidentally QMC results of Fig. 3a in Ref. [57] { again comparing by eye only) in this region.



FIG.2: CCM results for the absolute values of the spin-spin correlation function for the spin-half triangular-lattice antiferrom agnet using the LSUBm approximation with m = f2;3;4;5;6g.CCM results are compared to those results of exact diagonalisations (ED) [62] with N = 36.

The author of the QMC calculations in Ref. [57] noted that: \to good approximation, the two-point function depends only on jrj." As also noted in Ref. [57], the long-range behaviour of correlation function gives a constant, thus indicating a long-range ordered ground state. We see this also at all levels of approximation as shown in Fig. 1. Results [26] for the sublattice magnetisation have shown that extrapolation of the CCM LSUBm results gives good correspondence to the results of QMC, as noted above, where both methods indicate from the sublattice magnetisation that approximately 61% of the classical Neel order remains in the quantum limit. Finally, the author of Ref. [57] remarked that the approach to the constant for larger separations could be tted by either an exponential or power-law form. We tried both for the LSUB8 data with r 8 and found nothing to contradict this statement, although the power-law decay seemed to work better (i.e., had a lower residual error) that the exponential law. The residual sum-of-squares error for the power-law form was 0.0015 and for the exponential form was 0.0034. Fits of these forms to the LSUB8 data were carried out using the R statistics language: http://cran.r-project.org/.

The results for the (frustrated) spin-half square-lattice antiferrom agnet using the LSUBm approximation with m = f2;3;4;5;6g are shown in Fig. 2. Again, LSUBm results are clearly

8

converging with increasing levels of approximation, and again they are invery good mutual agreement for small lattice separations. CCM results are again compared to results of ED for N = 36 (e.g., see Ref. [62]), and again good agreement is found for very small lattice separations (r < 3). Another strong cusp-like behaviour is seen in ED results for the spin-spin correlation function for r = 3. Again, it is highly likely that this is primarily due to nite-size e ects. Indeed, we see that CCM results for r = 3 are quite different to those of ED.Furthermore, CCM LSUBm results initially decay strongly and then tend to a constant value for larger lattice separations. The behaviour of the correlation function for the triangular lattice is broadly similar to that of the square lattice. Again, extrapolation of LSUBm results [41] for the sublattice magnetisation demonstrate that the spin-half triangular lattice antiferrom agnet is Neel ordered. For LSUB 6 with r = 5.5, we found that the initial decay of the correlation function for CCM data in this case could be tted well by either an exponential or power-law form (for both form s, residual sum -of-squares: 0.0002).

IV. CONCLUSIONS

We have presented new form alism for the CCM in order to form \generalised" expectation values for a wide range of terms in the spin Ham iltonian. This is particularly useful for \high-order" CCM [63]. We have applied this approach in order to determ ine the spin-spin correlation functions for frustrated and unfrustrated 2D antiferrom agnets, namely, the spinhalf triangular- and square-lattice Heisenberg models. We found good correspondence of CCM with results of quantum M onte C arlo (QMC) for the square-lattice antiferrom agnet. A strong cusp-like behaviour was noted in exact diagonalisation (ED) results for the correlation function at a lattice separation L=2 for a lattice of size N = L L for both lattices. This is a nite-size e ect. A cusp in the spin-spin correlation function was also seen in QMC results at r = L=2 for a N = L L lattice.

By contrast, this cusp was not seen in CCM results for either lattice at any value of r at any level of LSUBm approximation. Indeed, we note that CCM results are determined in the in nite-lattice limit (N ! 1) from the outset. We believe that (as a \rule of thum b") we obtain reasonable results for the correlation function up to separations of r m for the square lattice { even despite the fact that we are using a purely \localised" LSUBm approximation scheme. LSUBm results for the triangular lattice are clearly converging, and

9

are in good mutual agreement for small lattice separations. For both lattices, CCM results for the correlation function decay strongly initially, although they become constant for larger values of r. P revious CCM results for the sublattice magnetisation [41] indicate that the spin-half H eisenberg models for the square and triangular lattices are N cel ordered, which agrees with the results of other approximate methods.

It is still fair to say that QMC results still generally provide the most accurate results for the unfrustrated H eisenberg model on lattices of two spatial dimensions. However, QMC is severely restricted by the \sign problem," which is a consequence of frustration at T = 0. The CCM has been shown [41] to provides good results for the ground-state properties of the spin-half triangular lattice antiferrom agnet such as the ground-state energy and sublattice m agnetisation. Here we have shown that the CCM also provides reasonable results for the spin-spin correlation function of the spin-half triangular lattice antiferrom agnet. Finally, we note that CCM results indicate that the behaviour of the correlation function for the triangular lattice is broadly sim ilar to that of the square lattice.

A cknow ledgm ents

We thank and acknowledge Joerg Schulenburg for the continuing development of the CCCM code; excellent advice, help, and support over the years; and nally for providing exact diagonalisation results quoted in this manuscript [58] using the SpinPack code (http://www.ovgu.de/jschulen/spin/) We begin the description of the application of the CCM by noting that it may be proven from Eqs. (2) and (3) in a straightforward manner that the ket- and bra-state equations are thus given by

$$h_{T} e^{S} H e^{S} j i = 0; 8I \neq 0;$$
 (A1)

h j
$$\hat{s}e^{s}$$
 [H ;C₁⁺]e^s j i = 0; 8I \notin 0 : (A2)

The index I refers to a particular choice of cluster from the set of (N_F) fundam ental clusters that are distinct under the symmetries of the crystallographic lattice and the Ham iltonian and for a given approximation scheme at a given level of approximation. We note that these equations are equivalent to the minimization of the expectation value of $H = h^{-}H ji$ with respect to the CCM bra- and ket-state correlation coecients f $S_{I};S_{I}g$. We note that Eq. (A1) is equivalent to $H = S_{I} = 0$, whereas Eq. (A2) is equivalent to $H = S_{I} = 0$. Furthermore, we note that Eq. (A1) leads directly to simple form for the ground-state energy given by

$$E_{g} = E_{g} (fS_{I}g) = h je^{S} H e^{S} ji :$$
(A3)

The full set fS_1 ; S_1 g provides a complete description of the ground state. For instance, an arbitrary operator A will have a ground-state expectation value given as

The sim ilarity transform of A is given by,

$$A^{\sim} = {}^{S}Ae^{S} = A + [A;S] + \frac{1}{2!}[A;S];S] + : (A5)$$

APPENDIX B:HIGH-ORDER CCM

The manner is which the ground-state problem is solved to high-orders of approximation via computational is provided in Ref. [52]. However, we note that use new \high-order CCM operators" that are formed purely of spin-raising operators with respect to the model state. The model state is taken to be a state in which all spins point in the downwards z-direction after some appropriate rotation of the local axes of the spins. This allows us to write one-and two-body spin operators in term softhese new high-order CCM operators. The problem

inherent in Eq. (A1) becomes one of pattern-matching the fundamental set of clusters in C_{I} to the terms in the similarity transformed version of the Hamiltonian e^SH e^S j i, where the I-th such equation is given by

$$E_{I} h j C_{T} e^{S} H e^{S} j i = 0 ; 8I \neq 0 :$$
(B1)

(Note that we assume that h $jC_{I}C_{I}^{+}ji = 1$ in the above equation). Specic terms in the H am iltonian that m ay be used in the CCM code are: $s^{z}s^{z}; s^{+}s^{z}; s s^{z}; s^{z}s^{+}; s^{z}s; s^{+}s^{z}; s s^{+}; s^{z}s; s^{+}s^{z}; s^{+}s^{z}; s^{+}s^{z}; s^{+}s^{+}; s^{+}s^{+}; s^{+}s; s$

We now de ne the following new set of CCM bra-state correlation coe cients given by $x_I = S_I$ and $x_I = N_B = N$ (l!) $_IS_I$ and we assume again that $h \downarrow C_I C_I^+ j i = 1$. Note that N_B is the number of B ravais lattice sites. Note also that for a given cluster I then $_I$ is a symmetry factor which is dependent purely on the point-group symmetries (and not the translational symmetries) of the crystallographic lattice and that 1 is the number of spin operators. We note that the factors $_I$, N, N_B , and (l!) never need to be explicitly determined. The CCM bra-state operator may thus be rewritten as

$$S = 1 + N = \underset{I=1}{\overset{X_F}{\times} C_I} ;$$
 (B2)

such that we have a particularly simple form for H , given by

$$H = N \underset{I=0}{\overset{X_{F}}{\times}} x_{I} E_{I} ; \qquad (B3)$$

where $\mathbf{x}_0 = 1$. We note that the \mathbf{E}_0 is defined by $\mathbf{E}_0 = \frac{1}{N}$ h je ^S H e^S j i (and, thus, $\mathbf{E}_0 = \frac{1}{N} \mathbf{E}_g$) and that \mathbf{E}_I is the I-th CCM ket-state equation defined by Eq. (B1). The CCM ket-state equations are easily re-derived by taking the partial derivative of H = N with respect to \mathbf{x}_I , where

$$\frac{(H = N)}{\kappa_{I}} (0) = E_{I} :$$
(B4)

We now take the partial derivative of H = N with respect to x_I such that the bra-state equations take on a particularly simple form, given by

$$\frac{(H = N)}{x_{I}} = \frac{E_{0}}{x_{I}} + \frac{X_{F}}{x_{J}} \times \frac{E_{J}}{x_{I}} (0) = E_{I} :$$
(B5)

The expectation value of a \generalized" spin operator that we shall call A m ay be treated in an analogous m anner to that of the expectation value of the H am iltonian, given by H . W e write:

$$A_{I} = h j C_{I} e^{S} A e^{S} j i$$
 (C1)

and with $C_0 = 1$. The similarity transform of A is dened by Eq. (A4). We may treat a wide range of one- and two-body spin operations. However, unlike the Heisenberg Hamiltonian of Eq. (1), we do not constrain i and j in the two-body terms to be only nearest-neighbors or next-nearest-neighbors. For example, we consider here the spin-spin correlation functions. The expectation value of the generalized (spin) operator may again be written in a particularly simple form as:

$$A = h^{*}A^{j}i$$

$$= N^{*}K_{I}h^{*}C_{I}e^{S}Ae^{S}ji$$

$$I = 0$$

$$A = N^{*}K_{I}A_{I};$$
(C2)

where $\mathbf{x}_0 = 1$ also and $A_0 = \frac{1}{N}h$ je ^SAe^S j i. The same code used to nd ground-state equations may be used to nd the generalized expectation values. Again we note that the index I in Eq. (C2) runs from zero to N_F. Again, we note that factors such as N_B or _I etc. do not need to be determined explicitly because they cancel because of the de nition of $f\mathbf{x}_{Ig}$ given above. The techniques need to achieve a computational solution have been discussed extensively elsewhere, and the interested reader is referred to R efs. [22, 26, 32, 38, 52] for m ore information. However, clearly we see that the sum mation over all fundamental clusters involved in evaluating A is carried out readily once the ket- and bra-state equations of Eqs. (B 4) and (B 5) have been solved for.

^[1] F.Coester, Nucl. Phys. 7, 421 (1958); F.Coester and H.Kummel, ibid. 17, 477 (1960).

^[2] J.Cizek, J.Chem. Phys. 45, 4256 (1966); Adv.Chem. Phys. 14, 35 (1969).

^[3] R F.Bishop and K H.Luhim ann, Phys. Rev. B 17, 3757 (1978); ibid. 26, 5523 (1982).

^[4] H.Kummel, K.H.Luhm ann, and J.G.Zabolitzky, PhysRep. 36C, 1 (1978).

- [5] J.S.Arponen, Ann. Phys. (N.Y.) 151, 311 (1983).
- [6] R F. Bishop and H. Kummel, Phys. Today 40(3), 52 (1987).
- [7] J.S. Arponen, R.F. Bishop, and E. Pajanne, Phys. Rev. A 36, 2519 (1987); ibid. 36, 2539 (1987); in: Condensed Matter Theories, Vol. 2, P. Vashishta, R.K. Kalia, and R.F. Bishop, eds. (Plenum, New York, 1987), p. 357.
- [8] R.J.Bartlett, J.Phys.Chem. 93, 1697 (1989).
- [9] R.F.Bishop, Theor. Chim. Acta 80, 95 (1991).
- [10] M.Roger and J.H.Hetherington, Phys.Rev.B 41, 200 (1990); Europhys.Lett.11, 255 (1990).
- [11] R.F.Bishop, J.B. Parkinson, and Y. Xian, Phys. Rev. B 44, 9425 (1991).
- [12] R.F.Bishop, J.B. Parkinson, and Y. Xian, Phys. Rev. B 46, 880 (1992).
- [13] R F.Bishop, JB.Parkinson, and Y.Xian, J.Phys.: Condens. Matter 5, 9169 (1993).
- [14] D JJ.Famell and JB.Parkinson, J.Phys.: Condens.M atter 6, 5521 (1994).
- [15] R F. Bishop, R G. Hale, and Y. Xian, Phys. Rev. Lett. 73, 3157 (1994).
- [16] Y.Xian, J.Phys.: Condens.M atter 6, 5965 (1994).
- [17] R. Bursill, G A. Gehring, D J.J. Famell, J.B. Parkinson, T. Xiang, and C. Zeng, J. Phys.: Condens. Matter 7, 8605 (1995).
- [18] R.G. Hale. Ph.D. Thesis, UM IST, Manchester, United Kingdom (1995).
- [19] R F.Bishop, D J.J.Farnell, and J.B.Parkinson, J.Phys.: Condens. Matter 8, 11153 (1996).
- [20] D JJ. Famell, SA. Kruger, and JB. Parkinson, J. Phys.: Condens. Matter 9, 7601 (1997).
- [21] R F.Bishop, Y.Xian, and C.Zeng, in: Condensed Matter Theories, Vol. 11, E.V. Ludena, P. Vashishta, and R F.Bishop, eds. (Nova Science, Commack, New York, 1996), p. 91.
- [22] C. Zeng, D J.J. Famell, and R F. Bishop, J. Stat. Phys. 90, 327 (1998).
- [23] RF.Bishop, DJJ.Famell, and JB.Parkinson, Phys.Rev.B 58, 6394 (1998).
- [24] R F. Bishop in M icroscopic M any-Body Theories and Their Applications, Lecture Notes in Physics 510, J.N avarro and A.Polls, eds.Lecture Notes in Physics Vol.510 (Springer-Verlag, Berlin, 1998), p.1.
- [25] J.Rosenfeld, N.E.Ligterink, and R.F.Bishop, Phys. Rev. B 60, 4030 (1999).
- [26] R F. Bishop, D J.J. Famell, S E. K ruger, J B. Parkinson, J. Richter, and C. Zeng, J. Phys.: Condens. M atter 12, 6887 (2000).
- [27] R F.Bishop, D J.J.Famell, and M L.Ristig, Int. J.M od. Phys. B 14, 1517 (2000).
- [28] SE.Kruger, J.Richter, J.Schulenberg, DJJ.Famell, and RF.Bishop, Phys.Rev.B 61,

14607 (2000).

- [29] D.J.J.Famell, R.F.Bishop, and K.A.Gemoth, Phys. Rev. B 63, 220402R (2001).
- [30] D J.J. Famell, K A. Gemoth, and R F. Bishop, Phys. Rev. B 64, 172409 (2001).
- [31] S.E.K ruger and J.R ichter, Phys. Rev. B 64, 024433 (2001).
- [32] D JJ.Famell, R F.Bishop, and K A.Gemoth, J.Stat. Phys. 108, 401 (2002).
- [33] N.B. Ivanov, J.Richter, and D.J.J. Famell, Phys. Rev. B 66, 014421 (2002).
- [34] D JJ.Famelland R F.Bishop, arxiv.org/abs/cond-m at/0311126.
- [35] S.E.K ruger, D.J.J.Famell, and J.R ichter, Int.J.M od. Phys. B 17, 5347 (2003).
- [36] R.Darradi, J.Richter, and D.J.J.Famell, Phys. Rev. B. 72, 104425 (2005).
- [37] R.Darradi, J.Richter, and D.J.J.Famell, J.Phys.: Condens.Matter 17, 341 (2005).
- [38] D.J.J.Famell, J.Schulenberg, J.Richter, and K.A.Gemoth, Phys. Rev. B. 72, 172408 (2005).
- [39] S.E.Kruger, R.Darradi, J.Richter, and D.J.J.Famell, Phys. Rev. B 73, 094404 (2006)
- [40] D.Schmalfu, R.Darradi, J.Richter, J.Schulenburg, and D. Ihle, Phys. Rev. Lett. 97, 157201 (2006).
- [41] D JJ.Famell and R F.Bishop, arxiv.org/abs/cond-m at/0606060.
- [42] J.Richter, R.Darradi, R.Zinke, and R.F.Bishop, Int. J.M od. Phys. B 21, 2273 (2007).
- [43] R.Zinke, J.Schulenburg, and J.Richter, Eur. Phys. J.B 61, 147 (2008).
- [44] R F.Bishop, P.H.Y. Li, R.Darradi, and J.Richter, J.Phys.: Condens. Matt. 20 255251 (2008).
- [45] R.F.Bishop, P.H.Y.Li, R.Darradi, J.Schulenburg, and J.Richter, Phys. Rev. B 78, 054412 (2008).
- [46] R F. Bishop, P H. Y. Li, R. Darradi, and J.Richter, Europhys. Lett. 83, 47004 (2008).
- [47] R F. Bishop, P H.Y. Li, R. Darradi, J.Richter, and C E. Campbell, J. Phys.: Condens. M att. 20, 415213 (2008).
- [48] R.Darradi, O.Derzhko, R.Zinke, J.Schulenburg, SE.K ruger, and J.Richter, Phys. Rev. B 78, 214415 (2008).
- [49] D JJ.Famell and R F.Bishop, Int. J.M od. Phys. B. 22, 3369 (2008).
- [50] D J.J.Famell, J.Richter, R.Zinke, and R.F.Bishop, J.Stat. Phys. 135, 175 (2009).
- [51] P.Li, D.J.J.Famell, and R.F.Bishop, Phys. Rev. B 79, 174405 (2009).
- [52] D JJ.Famell, arX iv:0909.1226
- [53] A.W. Sandvik, Phys. Rev. B 56, 11678 (1997).
- [54] W .Zeng, J.O itm aa, and C.J.Ham er, Phys.Rev.B 43, 8321 (1991).

- [55] C.J.Hamer, Z.Weihong, P.Amdt, Phys. Rev. B 46, 6276 (1992).
- [56] J. Richter, J. Schulenburg, and A. Honecker, in Lecture Notes in Physics 645 edited by U. Schollwock, J. Richter, D J.J. Famell and R.F. Bishop (Springer Verlag Berlin Heidelberg 2004) p. 85{153.
- [57] S.Liang, Phys. Rev. B 42, 6555 (1990).
- [58] J.Richter, J.Schulenburg, arX iv:0909.3723
- [59] L.Capriotti, A.E.Trum per, S.Sorella, Phys.Rev.Lett. 82, 3899 (1999).
- [60] R P. Singh and D A. Huse, Phys. Rev. Lett. 68, 1766 (1992).
- [61] S.J.M iyake, J.Phys.Soc.Jpn.61, 983 (1992).
- [62] PW.Leung, and K.Runge, Phys. Rev. B 47, 5861 (1993).
- [63] A GPL licensed version of the Crystallographic Coupled Cluster M ethod' (CCCM) code of DJJ.Famell and J.Schulenburg is available online at: http://www.ovgu.de/jschulen/ccm/