

A Classical Background for the Wave Function Prediction in the Infinite System Density Matrix Renormalization Group Method

Hiroshi Ueda¹⁾, Andrej Gendiar²⁾, and Tomotoshi Ishino³⁾

¹Department of Material Engineering Science, Graduate School of Engineering Science, Osaka University, Osaka 560-8531, Japan

²Institute of Electrical Engineering, Slovak Academy of Sciences, Dubravská cesta 9, SK-841 04, Bratislava, Slovakia

³Department of Physics, Graduate School of Science, Kobe University, Kobe 657-8501, Japan

We report a physical background of the wave function prediction in the infinite system density matrix renormalization group (DMRG) method, from the view point of two-dimensional vertex model, a typical lattice model in statistical mechanics. Singular value decomposition applied to rectangular corner transfer matrices naturally draws matrix product representation for the maximal eigenvector of the row-to-row transfer matrix. The wave function prediction can be expressed as the insertion of an approximate half-column transfer matrix. This insertion process is in accordance with the scheme proposed by McCulloch recently.

KEYWORDS: DMRG, PWRG, CTMRG, Renormalization

1. Introduction

The density matrix renormalization group (DMRG) method is one of the efficient numerical methods, which has been applied extensively to one-dimensional (1D) quantum systems and two-dimensional (2D) classical systems.¹⁽⁴⁾ The method is variational in the sense that it assumes a trial state, the matrix product state (MPS), which is written as a product of local tensors.⁵⁽¹⁵⁾ Orthogonality of each matrix ensures the numerical stability.

One of the bottlenecks in the computation of the DMRG method is the diagonalization of super block Hamiltonian. The construction of a good initial vector for this diagonalization is very important. For the finite-system DMRG method, the so-called wave function renormalization scheme provides the answer.^{16,17)} For the infinite-system DMRG method, Baxter's method of corner transfer matrix (CTM),¹⁸⁽²⁰⁾ which can be reinterpreted from the view point of the DMRG method,^{21,22)} essentially solves the problem of initial vector. Based on Baxter's CTM method, the product wave function renormalization group (PWRG) method was proposed,²³⁽²⁵⁾ and has been applied to the study of 1D spin chains.²⁶⁽⁴⁰⁾

Recently McCulloch proposed a way of precise wave function prediction, which works better than the PWRG method especially when the system size is small compared with the correlation length.⁴¹⁾ In this paper we present a physical background for McCulloch's scheme from the view point of 2D vertex model, one of the typical lattice models in statistical mechanics.²⁰⁾ Although we employ classical lattice model, most of the obtained results can be applicable for 1D quantum systems through the quantum-classical correspondence.

Structure of the paper is as follows. In the next section we explain the symmetric vertex model, and express the maximal eigenstate of the row-to-row transfer matrix by use of CTMs. In §3 we consider the area extension of CTMs, introducing an approximate half-column trans-




Fig. 1. A finite size vertex model of width $N = 8$. Cross marks show boundary spins and open circles show row spins, which are the variables of σ in Eq. (2.1). The system consists of its left-half and the right half, where the vertical stitch corresponds to the half-column spin $\sigma_{1/2}; \sigma_{3/2}; \sigma_{5/2}; \dots$ between them.

fer matrix. We show the connection between MPS and CTM formulation in §4, where the system size extension scheme by McCulloch is obtained naturally. We summarize the obtained result in the last section, and discuss the remaining problem on the MPS obtained by the finite-system DMRG method.

2. Eigenstate of row-to-row Transfer Matrix Approximated by Corner Transfer Matrices

Throughout this article we consider a square-lattice symmetric vertex model,²⁰⁾ as an example of 2D classical lattice models. There is a d -state spin variable on each bond, which connects neighboring lattice points. Four spins around a lattice point determine the local Boltzmann weight W , which is called as the vertex weight. We assume that the vertex weight is position independent, and therefore the system is uniform. We also assume that each vertex weight is invariant under exchange of left and right spin variables, and those of up and down spin variables. In other words, we consider the symmetric vertex

model in order to simplify the following formulation.

As shown on the left side of Fig. 1, we treat a finite size system that has a rectangular shape. This system corresponds to the stack of row-to-row transfer matrices T_N , whose width is N , multiplied by an initial vector V_N . We choose V_N so that it corresponds to the boundary condition at the bottom of the system, where there is a row of boundary spins shown by the cross marks. Those cross marks aligned vertically also represent boundary spins, that are located at the both ends of T_N . The row of open circles represents spins on top of the rectangular system. We consider a d^N -dimensional vector

$$| \psi \rangle = T_N T_N T_N \dots T_N V_N; \quad (2.1)$$

where the number of the row-to-row transfer matrix T_N is sufficiently large. Under this assumption we can expect that $| \psi \rangle$ is a good approximation of the maximal eigenvector of T_N if V_N is not orthogonal to that.

For a while let us consider the case $N = 8$; generalization to arbitrary N is straightforward. We label the top spins as $q_1; q_2; q_3; q_4; p_4; p_3; p_2; p_1$ from left to right. The vector elements of $| \psi \rangle$ are then written as $| \psi \rangle (q_1; q_2; q_3; q_4; p_4; p_3; p_2; p_1)$. Since we have assumed the left-right symmetry for the vertex weight, it is convenient to divide the row-spin into the left half $q_1; q_2; q_3; q_4$, where we have counted them from left to right, and the right half $p_1; p_2; p_3; p_4$, where we have counted them from right to left. (See right bottom of Fig. 1.) According to this division, we can interpret $| \psi \rangle$ as a d^4 -dimensional real symmetric matrix, whose elements can be expressed as $| \psi \rangle (q_1 q_2 q_3 q_4 | p_1 p_2 p_3 p_4)$. We have used the vertical bar $| \cdot \rangle$ to separate the left and the right indices, and dropped the commas between the spin variables for the book keeping. If necessary, we further abbreviate the matrix notation as $| \psi \rangle (q|p)$.

We express the left half of the rectangular system by use of the CTM, whose elements are written as $C_4 (q_1 q_2 q_3 q_4 | j_1 j_2 j_3 \dots)$, where $j_1 j_2 j_3 \dots$ represent the half-column spins at the center of the system. In the same manner we can express the right half by the transpose of C_4 , i.e., C_4^T . Joining process of these halves by stitching C_4 and C_4^T via the contraction of the half-column spins can be expressed simply by the product of matrices $| \psi \rangle = C_4 C_4^T$. More precisely, there is a relation

$$| \psi \rangle (q|p) = \sum_{j_1, j_2, j_3, \dots} C_4 (q|j) C_4 (j|p); \quad (2.2)$$

where we have used the abbreviations $q = q_1 q_2 q_3 q_4$, $p = p_1 p_2 p_3 p_4$, and $j = j_1 j_2 j_3 \dots$. Since we have assumed that the number of T_8 in Eq. (2.1) is sufficiently large, the same for the number of column-spin j . Although we treat j , we do not think of them as spins directly treated in numerical calculations, unlike q and p .

One of the fundamental mathematical tool in the DMRG method is the singular value decomposition (SVD).^{1,2)} Let us apply it to the CTM

$$C_4 (q|p) = \sum_{j_1, j_2, j_3, \dots} A_4 (q|j) \lambda_4(j) U_4 (j|p); \quad (2.3)$$

where λ_4 is a d^4 -state block-spin (or an auxiliary) variable,




Fig. 2. The singular value decomposition applied to C_4 . The black square and circle corresponds to the block spin and the singular values $\lambda_4(j)$, respectively. The left and the right triangles represent A_4 and U_4 , respectively.

and $\lambda_4(j)$ represents the singular values. The matrix A_4 is d^4 -dimensional, and it satisfies the orthogonal relations

$$\begin{aligned} \sum_{q_1, q_2, q_3, q_4} A_4 (q^0|j) A_4 (q|j) &= \delta_4^0(j); \\ \sum_{q_1, q_2, q_3, q_4} A_4 (q|j^0) A_4 (q|j) &= \delta_4^0(j); \end{aligned} \quad (2.4)$$

where $\delta_4^0(j)$ is Kronecker's delta, and where $\delta_4^0(j)$ is defined as

$$\delta_4^0(j) = \prod_{i=1}^4 \delta_4^0(j_i); \quad (2.5)$$

The above orthogonal relation can be written shortly as $A_4 A_4^T = A_4^T A_4 = I_4$. Column vectors of the rectangular matrix U_4 are also orthogonal with each other,

$$\sum_{q_1, q_2, q_3, q_4} U_4 (j^0|q) U_4 (j|q) = \delta_4^0(j); \quad (2.6)$$

but the row vectors are not

$$\sum_{q_1, q_2, q_3, q_4} U_4 (j^0|q) U_4 (j|q) \neq \delta_4^0(j); \quad (2.7)$$

This is because the degree of freedom of j is far larger than that of q or p . Figure 2 is the pictorial representation of SVD applied to C_4 .

We often regard the singular values λ_4 as the diagonal matrix $\lambda_4(j) = \lambda_4(j) \delta_4^0(j)$, and write Eq. (2.3) shortly as $C_4 = A_4 \lambda_4 U_4^T$. For the latter convenience, let us introduce the generalized inverse of the CTM

$$C_4^{-1} = U_4 \lambda_4^{-1} A_4^T; \quad (2.8)$$

which satisfies the relation

$$\begin{aligned} C_4 C_4^{-1} &= A_4 \lambda_4 U_4^T U_4 \lambda_4^{-1} A_4^T \\ &= A_4 A_4^T = I_4; \end{aligned} \quad (2.9)$$

It should be noted that $C_4^{-1} C_4$ is a projection operator

$$U_4 \lambda_4^{-1} A_4^T A_4 U_4^T = U_4 U_4^T \quad (2.10)$$

in the left hand side of Eq. (2.7), where $(C_4^{-1} C_4)^2 = C_4^{-1} C_4$ holds.

In the context of the DMRG method, small singular values are neglected when it is impossible to store matrix




Fig. 3. The pictorial representation of $P_4 = C_3^{-1} C_4$ in Eq. (3.1). The circle with the cross mark shows C_3^{-1} . Since P_4 has a function of extending the area of CTM, it can be regarded as an approximation for the half-column transfer matrix.




Fig. 4. The area of CTM can be extended by applying the approximate half-column transfer matrices.

elements during the numerical calculation. This truncation is a kind of decimation in the renormalization group (RG) theory. Under the truncation, the matrices A_4 work as the RG transformation that controls numerical precision. In the next section we do not truncate singular values, in order to avoid complications in notations, and the introduction of truncation is straightforward.

3. Half Column Transfer Matrix and Matrix Product State

We introduce a new notation between matrices, the dot product, which contract variables according to Einstein rule. As an example, let us consider

$$P_4 = C_3^{-1} C_4; \tag{3.1}$$

where q_1, q_2 , and q_3 are contracted but q_4 is not, since the first three spins are shared by C_3^{-1} and C_4 . Figure 3 shows this rule graphically. Substituting Eq. (2.3) and (2.8) to $C_3^{-1} C_4$, we obtain

$$\begin{aligned} P_4 &= (U_3^{-1} A_3^T) (A_4 U_4^T) \\ &= U_3^{-1} (A_3^T A_4) U_4^T \\ &= U_3^{-1} \tilde{A}_4 U_4^T; \end{aligned} \tag{3.2}$$

To avoid any confusion, let us write down element of P_4

$$\begin{aligned} P_4 ({}^0 j_4 j) &= \sum_{q_1, q_2, q_3} U_3 ({}^0 j) U_3^{-1} (\tilde{A}_4 (q_1 q_2 q_3 q_4 j)) U_4 (j); \end{aligned} \tag{3.3}$$

where the new matrix $\tilde{A}_4 = A_3^T A_4$ is the renormalized orthogonal matrix

$$\tilde{A}_4 (q_4 j) = \sum_{q_1, q_2, q_3} A_3 (q_1 q_2 q_3 j) A_4 (q_1 q_2 q_3 q_4 j); \tag{3.4}$$

which satisfies the relation

$$\sum_{q_4} \tilde{A}_4 (q_4 j) \tilde{A}_4 (q_4 j) = ({}^0 j): \tag{3.5}$$

In Eq. (3.4) the group of spins q_1, q_2 , and q_3 are mapped onto the block spin by the RG transformation A_3 . The obtained \tilde{A}_4 corresponds to the matrix that constructs MPS, which is constructed by the infinite system DMRG




Fig. 5. Pictorial representation of Eq. (3.8).

method, as shown later.

The P_4 thus obtained has a function of half-column transfer matrix (HCTM), since it extends the width of C_3 by one by way of the dot product

$$C_3 P_4 = C_3 (C_3^{-1} C_4) = (C_3 C_3^{-1}) C_4 = C_4 \tag{3.6}$$

as shown in the left side of Fig. 4. Applying SVD to C_3 and substituting Eq. (3.2), $C_3 P_4$ is calculated as

$$(A_3^{-1} U_3^T) (U_3^{-1} \tilde{A}_4 U_4^T) = A_3 \tilde{A}_4 U_4^T; \tag{3.7}$$

Since C_3 is again constructed from C_2 and P_3 , as shown in the right side of Fig. 4, we can further decompose C_4 as

$$C_4 = C_2 P_3 P_4 = A_2 \tilde{A}_3 \tilde{A}_4 U_4^T; \tag{3.8}$$

The contraction process by the dot products are shown in the right side of Fig. 5. It should be noted that $C_2 P_4$ is not C_3 , since U_2^T contained in C_2 and U_3 contained in P_4 do not matches to give an identity. In this sense, P_4 is an approximation for the half column transfer matrix, optimized for the area extension of C_3 only.

Using the decomposition of C_4 in Eq. (3.8), we obtain the matrix product representation of $\tilde{\rho}_8 = C_4 C_4^T$. We have

$$\begin{aligned} \tilde{\rho}_8 &= A_2 \tilde{A}_3 \tilde{A}_4 ({}^4)^2 \tilde{A}_4^T \tilde{A}_3^T A_2^T \\ &= A_2 \tilde{A}_3 \tilde{A}_4 \tilde{\lambda}_4^T \tilde{A}_3^T A_2^T; \end{aligned} \tag{3.9}$$

where $\tilde{\lambda}_4 = ({}^4)^2$ is the singular value of $\tilde{\rho}_8$. (See Fig. 6.) Such a construction of $\tilde{\rho}_8$ is equivalent to the MPS




Fig. 6. Matrix product expression of $|\psi_8\rangle$ in Eq. (3.9). Double circle represent $\tilde{C}_4 = (C_4)^2$.




Fig. 7. Approximate area extension process $C_5^{A_{PP}} = C_4 P_4$.

considered in the context of the infinite system DMRG method.

4. Approximate Area Extension

Let us consider a problem of obtaining an approximation of $C_5 = C_4 P_5$ without using P_5 . This attempt is equivalent to construct an approximation for C_5 using C_2, C_3 , or C_4 . One might think that $P_4 = C_3^{-1} C_4 = U_3 C_3^{-1} \tilde{A}_4 U_4^T$ can be of use as an approximation for P_5 . But this idea should be rejected since $U_4^T U_3$, which appears in the calculation of $C_4 P_4$, is not an identity. A way to avoid this mismatching is to introduce a spatial reflection of P_4 , which is defined as

$$P_4 = U_4 \tilde{A}_4^T C_3^{-1} U_3^T; \quad (4.1)$$

and use it as an approximation for P_5 . Leaving the validity of the approximation scheme by the latter discussion, let us calculate the approximate extension $C_5^{A_{PP}} = C_4 P_4$ and write it into the matrix product representation. (See Fig. 7.) We obtain

$$\begin{aligned} C_5^{A_{PP}} &= A_2 \tilde{A}_3 \tilde{A}_4 U_4^T U_4 \tilde{A}_4^T C_3^{-1} U_3^T \\ &= A_2 \tilde{A}_3 \tilde{A}_4 (C_4)^2 \tilde{A}_4^T C_3^{-1} U_3^T \\ &= A_2 \tilde{A}_3 \tilde{A}_4 \tilde{A}_4^T C_3^{-1} U_3^T; \end{aligned} \quad (4.2)$$

and from this approximation we can construct

$$|\psi_{10}^{A_{PP}}\rangle = C_5^{A_{PP}} (C_5^{A_{PP}})^T \quad (4.3)$$




Fig. 8. Reorthogonalization process in Eq. (4.3). The rectangular in the lower diagram corresponds to \tilde{C}_4 in Eq. (4.4).

$$= A_2 \tilde{A}_3 \tilde{A}_4 \tilde{A}_4^T C_3^{-1} \tilde{A}_4 \tilde{A}_4^T \tilde{A}_3^T A_2^T;$$

which may approximate $|\psi_{10}\rangle$. Applying Schmidt orthogonalization for $\tilde{A}_4 \tilde{A}_4^T$ from the left side

$$\tilde{A}_4 \tilde{A}_4^T = B^T \quad (4.4)$$

we obtain the new orthogonal matrix B^T and the right triangular matrix \tilde{C}_4 . (See Fig. 8.) Substituting Eq. (4.4) into Eq. (4.3), we get the matrix product expression

$$|\psi_{10}^{A_{PP}}\rangle = A_2 \tilde{A}_3 \tilde{A}_4 B^T C_3^{-1} B^T \tilde{A}_4^T \tilde{A}_3^T A_2^T; \quad (4.5)$$

The extension from $|\psi_8\rangle$ to $|\psi_{10}^{A_{PP}}\rangle$ is the same as the wave function extension scheme proposed by McCulloch,⁴¹⁾ where the approximation for the renormalized wave function is given by

$$|\tilde{\psi}_{10}^{A_{PP}}\rangle = B^T C_3^{-1} B^T; \quad (4.6)$$

We have thus obtained a natural explanation for McCulloch's extension scheme from the view point of 2D vertex model. Up to now we have not considered the effect of basis truncation, which is used in numerical calculation of the infinite system DMRG method. First of all, the extension in Eqs. (4.4) and (4.5) is still efficient under the truncation, as it was shown numerically.⁴¹⁾ We then consider the extension from N to $N+2$ in the large system size limit $N \rightarrow \infty$. For simplicity, let us assume that the MPS in this limit is uniform, and the system is away from criticality. In this limit we can drop the site index from Eq. (4.1), and can express the approximate transfer matrix as

$$P = U^{-1} \tilde{A} U^T = U S U^T; \quad (4.7)$$

where $S = U^{-1} \tilde{A}$. From the assumed symmetry of the vertex model, both P and S are symmetric

$$\begin{aligned} P(|j\rangle\langle j|) &= P(|j\rangle\langle j|) \\ S(|j\rangle\langle j|) &= S(|j\rangle\langle j|); \end{aligned} \quad (4.8)$$

This symmetry is also expressed in short form as $P = P$ and $S = S$. Thus at least when the system size $N = 2i$ is large enough, typically several times larger than the correlation length, one can justify the usage of $C_i P_i$ as the approximation for $C_{i+1} P_{i+1}$.

Before closing this section, we consider the MPS expression for $|\psi_N\rangle$ that is optimized by way of the sweeping




Fig. 9. Graphical representation of Eq. (4.14).

process in the finite system DMRG method. The matrix product structure

$$N = A_2 \tilde{A}_3 \tilde{A}_{\frac{N}{2}} \tilde{A}_{\frac{N}{2}} \tilde{A}_{\frac{N}{2}}^T \tilde{A}_3^T A_2^T \quad (4.9)$$

is similar to that obtained by the infinite system DMRG method, but in this case the matrices satisfies the additional relation

$$\tilde{A}_{i-1} = \sqrt{\lambda_i} \tilde{A}_i^T; \quad (4.10)$$

where both \tilde{A}_i and $\sqrt{\lambda_i}$ differ from those obtained by the infinite system DMRG method. Taking the square root of λ_i , we formally obtain a diagonal matrix $\sqrt{\lambda_i} = \sqrt{\lambda_i}$. It should be noted that this $\sqrt{\lambda_i}$ is different from that obtained from the SVD applied to C_i . Defining

$$S_i = \sqrt{\lambda_i} \tilde{A}_i = \sqrt{\lambda_i} \tilde{A}_i^T \sqrt{\lambda_i} \quad (4.11)$$

and substituting it to Eq. (4.9), we obtain a new standard form for MPS

$$N = \lambda_1 S_2 S_3 S_{\frac{N}{2}} S_{\frac{N}{2}}^T S_3^T S_2^T \lambda_1; \quad (4.12)$$

where λ_1 is just a constant and is not essential. It is then straightforward to obtain the approximation $\frac{\text{APP}}{N+2}$ just by putting $S_{\frac{N}{2}} S_{\frac{N}{2}}^T$ at the center of the above MPS, where this insertion is a variant of Eq. (4.5). In the thermodynamic limit $N \rightarrow \infty$ the matrix S_i in Eq. (4.11) is independent on the site index i , and therefore it coincides with S in Eq. (4.8). This symmetric representation of uniform MPS is often of use.

5. Conclusions and Discussions

We have considered the wave function prediction in the finite system DMRG method, when it is applied to the 2D vertex model. Through the singular value decomposition of CTM C_i , we obtained the approximate half-column transfer matrix P_i . The insertion of P_i naturally explains the wave function prediction proposed by McCulloch,⁴¹⁾ which works better than the product wave function renormalization group (PWFRG) method,²³⁽²⁵⁾ especially when the system size is small. The difference between these two prediction methods can be explained by the shape of finite size system. The PWFRG method treats growing triangular cluster,²⁴⁾ whereas McCulloch's scheme always treat half-infinite stripe.

The relation between CTM and MPS in the finite system DMRG method is not so clear. For example, \mathcal{S}_8 can be expressed as $C_3 C_5^T$, but the MPS representation of the optimized \mathcal{S}_8 by the finite system DMRG cannot

be obtained from the SVD applied to C_3 and C_5 independently. This puzzle is something to do with the targeting scheme for asymmetric vertex model, and also with the determination of optimal RG transformation in the real-time DMRG method, where the density matrix is time dependent.

Acknowledgement

We thank I. McCulloch for valuable comments and discussions. H. U. thanks Dr. Okunishi for helpful comments on the DMRG method and continuous encouragement. A. G. Acknowledge QUTE and VEGA 1/0633/09 grants.

- 1) S. R. White: Phys. Rev. Lett. 69 (1992) 2863; Phys. Rev. B 48 (1993) 10345.
- 2) Density-Matrix Renormalization - A New Numerical Method in Physics -, eds. I. Peschel, X. Wang, M. Kaulke, and K. Hallberg (Springer, Berlin, 1999) and references therein.
- 3) T. Nishino, T. Hikihara, K. Okunishi, and Y. Hieida: Int. J. Mod. Phys. B 13 (1999) 1.
- 4) U. Schollwöck: Rev. Mod. Phys. 77 (2005) 259.
- 5) I. Aeck, T. Kennedy, E. H. Lieb, and H. Tasaki: Phys. Rev. Lett. 59 (1987) 799.
- 6) M. Fannes, B. Nachtergale, and R. F. Werner: Europhys. Lett. 10 (1989) 633.
- 7) M. Fannes, B. Nachtergale, and R. F. Werner: Commun. Math. Phys. 144 (1992) 443.
- 8) M. Fannes, B. Nachtergale, and R. F. Werner: Commun. Math. Phys. 174 (1995) 477.
- 9) A. Klumper, A. Schadschneider, and J. Zittartz: Z. Phys. B 87 (1992) 281.
- 10) H. Niggemann, A. Klumper, and J. Zittartz: Z. Phys. B 104 (1997) 103.
- 11) S. Ostlund and S. Rommer: Phys. Rev. Lett. 75 (1995) 3537.
- 12) S. Rommer and S. Ostlund: Phys. Rev. B 55 (1997) 2164.
- 13) M. Andersson, M. Bonan, and S. Ostlund: Phys. Rev. B 59 (1999) 10493.
- 14) H. Takasaki, T. Hikihara, and T. Nishino: J. Phys. Soc. Jpn. 68 (1999) 1537.
- 15) J. Dukelsky, M. A. Martı́n-Delgado, T. Nishino, and G. Sierra: Europhys. Lett. 43 (1998) 457.
- 16) S. R. White and I. Aeck: Phys. Rev. B 54 (1996) 9862.
- 17) S. R. White: Phys. Rev. Lett. 77 (1996) 3633.
- 18) R. J. Baxter: J. Math. Phys. 9 (1968) 650.
- 19) R. J. Baxter: J. Stat. Phys. 19 (1978) 461.
- 20) R. J. Baxter: Exactly Solved Models in Statistical Mechanics (Academic Press, London, 1982).
- 21) T. Nishino and K. Okunishi: J. Phys. Soc. Jpn. 65 (1996) 891.
- 22) T. Nishino and K. Okunishi: J. Phys. Soc. Jpn. 66 (1997) 3040.
- 23) T. Nishino and K. Okunishi: J. Phys. Soc. Jpn. 64 (1995) 4084.
- 24) K. Ueda, T. Nishino, K. Okunishi, Y. Hieida, R. Derian, and A. Gendiar: J. Phys. Soc. Jpn. 75 (2006) 014003.
- 25) H. Ueda, T. Nishino, K. Kusakabe: J. Phys. Soc. Jpn. 77 (2008) 114002.
- 26) N. A. Kutsu and Y. A. Kutsu: Phys. Rev. B 57 (1998) R4233.
- 27) N. A. Kutsu and Y. A. Kutsu: Prog. Theor. Phys. 105 (2001) 123.
- 28) N. A. Kutsu, Y. A. Kutsu, and T. Yamamoto: Prog. Theor. Phys. 105 (2001) 361.
- 29) N. A. Kutsu, Y. A. Kutsu, and T. Yamamoto: Phys. Rev. B 64 (2001) 085415.
- 30) N. A. Kutsu, Y. A. Kutsu, and T. Yamamoto: J. of Cryst. Growth 237-239 (2002) 14.
- 31) N. A. Kutsu, Y. A. Kutsu, and T. Yamamoto: Phys. Rev. B 67 (2003) 125407.
- 32) Y. Hieida, K. Okunishi, and Y. A. Kutsu: Phys. Lett. A 233 (1997) 464.
- 33) M. Hagiwara, Y. Narumi, K. Kindo, M. Kohno, H. Nakano, R. Sato, and M. Takahashi: Phys. Rev. Lett. 80 (1998) 1312.
- 34) K. Okunishi, Y. Hieida, and Y. A. Kutsu: Phys. Rev. B 59

- (1999) 6806.
- 35) K.Okunishi, Y.Hieida, and Y.Akutsu Phys.Rev.E 59 (1999) R 6227.
- 36) Y.Hieida, K.Okunishi, and Y.Akutsu: New J.of Phys. 1 (1999) 7.1.
- 37) K.Okunishi, Y.Hieida, and Y.Akutsu: Phys.Rev. B 60 (1999) R 6953.
- 38) Y.Hieida, K.Okunishi, and Y.Akutsu: Phys.Rev. B 64 (2001) 224422.
- 39) Y.Nanumi, K.Kindo, M.Hagiwara, H.Nakano, A.Kawaguchi, K.Okunishi, and M.Kohno: Phys.Rev. B 69 (2004) 174405.
- 40) S.Yoshikawa, K.Okunishi, M.Senda, and S.Miyashita: J.Phys.Soc.Jpn. 73 (2004) 1798.
- 41) I.McCulloch: arXiv:0804.2509.