Variable X-ray Absorption in mini-BAL QSOs

Margherita Giustini, 1,2,3 Massimo Cappi, George Chartas, Michael Eracleous, Giorgio G.C. Palumbo, and Cristian Vignali,

Abstract. We present the results of X-ray spectral analysis of two mini-BAL QSOs, PG 1126-041 and PG 1351+640, aimed at getting insights into the physics of quasar outflows. We find strong X-ray spectral variability on timescales of years. These variations can be well reproduced by variations of physical properties as the covering factor and column density of ionized gas along the line of sight, compatible with radiatively-driven accretion disk wind models.

Introduction

X-ray observations are crucial in getting insights into the physical mechanism responsible for launching and accelerating quasar outflows.

For example, radiatively-driven accretion disk wind models predict the presence of high column densities of X-ray absorbing gas which prevents the overionization of the UV-driven wind, allowing for its effective acceleration (see e.g. Proga et al. 2004). Furthermore, the physical properties of this X-ray absorbing gas are predicted to be variable with time (see e.g. Schurch et al. 2009). However, high signal-to-noise ratio X-ray observations of BAL QSOs are rare, because of their known "X-ray weakness" – e.g. Gallagher et al. 2006; but see Giustini et al. 2008.

We present the results of multiple XMM-Newton observations of two low redshift mini-BAL QSOs, PG 1126-041 (z=0.06) and PG 1351+640 (z=0.09).

Data Analysis and Results

We analyzed five EPIC datasets, three for PG 1351+640 (observed in 2004/12/31, 2008/06/06 and 2008/06/08) and two for PG 1126-041 (observed in 2008/06/15and 2008/12/13). We reduced all the datasets following standard XMM-Newton analysis threads.

PG 1351+640: The source flux dropped by a factor of ~ 10 in the soft (E <1 keV) band, while remaining constant at E > 7 keV during 4 years. No changes are detected during the two days elapsed between the two observations taken in June 2008, so we analyzed them together. The best fit model

¹Department of Astronomy & Astrophysics, Penn State University, University Park, PA 16802

²Dipartimento di Astronomia, Università di Bologna, via Ranzani 1, 40127 Bologna, Italy

³INAF/Istituto di Astrofisica Spaziale e Fisica cosmica, via Gobetti 101, 40138 Bologna, Italy




Figure 1. Left: EPIC-pn spectra of PG 1351+640 taken in December 2004 (black dots) and in June 2008 (red triangles). Right: EPIC-pn spectra of PG 1126-041 taken in June 2008 (black squares) and in December 2008 (red triangles) epochs. Bottom: χ^2 residuals in units of σ .

 $(\chi^2/\text{d.o.f.}=875/877, \text{ Fig.1, left})$ is a powerlaw continuum with $\Gamma \sim 2.3$ absorbed by two layers of ionized gas; one is totally covering the continuum source, has a column density $N_H \sim 1.5 \times 10^{21} \text{ cm}^{-2}$ and an ionization parameter $\xi \sim 20 \text{ erg cm s}^{-1}$ in both states. The other ionized absorber has an ionization parameter $\xi \sim 300 \text{ erg cm s}^{-1}$ and is covering 50%(90%) of the source, with a column density $N_H \sim 9(1) \times 10^{23} \text{ cm}^{-2}$ in the 2004(2008) observation.

PG 1126-041: The 0.2-10 keV source flux increased by a factor of $\sim 60\%$ in the six months elapsed between the two observations presented here. The best fit model (χ^2 /d.o.f.=96/102, Fig.1, right) is a powerlaw continuum with $\Gamma \sim 2.2$ absorbed by a layer of ionized gas covering $\sim 95\%$ of the source, with a column density $N_H \sim 8(5) \times 10^{22}$ cm⁻² and an ionization parameter $\xi \sim 30(40)$ erg cm s⁻¹ in the 2008 June(December) observation.

3. Conclusions

Our observations are consistent with radiatively-driven accretion disk wind models in that they find high column densities of X-ray absorbing gas along the line of sight, and strong variability in the physical properties of these ionized absorbers. Future, deeper X-ray observations of quasars which show powerful UV-driven winds are needed to confirm this scenario: for example, the adopted models predict the presence of strong iron absorption lines which cannot be detected with the present moderate signal-to-noise ratio spectra.

Acknowledgments. We acknowledge financial support from NASA grant NNX08AB71G and ASI contract I/088/06/0.

References

Gallagher, S. C., Brandt, W. N., Chartas, G., Priddey, R., Garmire, G. P., Sambruna, R. M. 2006, ApJ, 644, 709
Giustini, M., Cappi, M., Vignali, C. 2008, A&A, 491, 425
Proga, D., & Kallman, T. R. 2004, ApJ, 616, 688
Schurch, N. J., Done, C., Proga, D. 2009, ApJ, 694, 1