

NEW IDENTITIES INVOLVING q -EULER POLYNOMIALS OF HIGHER ORDER

T. KIM AND Y. H. KIM

ABSTRACT. In this paper, we present new generating functions which are related to q -Euler numbers and polynomials of higher order. From these generating functions, we give new identities involving q -Euler numbers and polynomials of higher order.

§1. Introduction/ Preliminaries

Let \mathbb{C} be the complex number field. We assume that $q \in \mathbb{C}$ with $|q| < 1$ and the q -number is defined by $[x]_q = \frac{1-q^x}{1-q}$ in this paper. The q -factorial is given by $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$ and the q -binomial formulae are known that

$$(x : q)_n = \prod_{i=1}^n (1 - xq^{i-1}) = \sum_{i=0}^n \binom{n}{i}_q q^{\binom{i}{2}} (-x)^i, \quad (\text{see [3, 14, 15]}),$$

and

$$\frac{1}{(x : q)_n} = \prod_{i=1}^n \left(\frac{1}{1 - xq^{i-1}} \right) = \sum_{i=0}^{\infty} \binom{n+i-1}{i}_q x^i, \quad (\text{see [3, 5, 14, 15]}),$$

where $\binom{n}{i}_q = \frac{[n]_q!}{[n-i]_q![i]_q!} = \frac{[n]_q[n-1]_q \cdots [n-i+1]_q}{[i]_q!}$.

The Euler polynomials are defined by $\frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}$, for $|t| < \pi$. In the special case $x = 0$, $E_n (= E_n(0))$ are called the n -th Euler numbers. In this paper, we consider the q -extensions of Euler numbers and polynomials of higher order. Barnes' multiple Bernoulli polynomials are also defined by

$$(1) \quad \frac{t^r}{\prod_{j=1}^r (e^{a_j t} - 1)} e^{xt} = \sum_{n=0}^{\infty} B_n(x, r | a_1, \dots, a_r) \frac{t^n}{n!}, \quad \text{where } |t| < \max_{1 \leq i \leq r} \frac{2\pi}{|a_i|}, \quad (\text{see [1, 14]}).$$

Key words and phrases. : multiple q -zeta function, q -Euler numbers and polynomials, higher order q -Euler numbers, Laurent series, Cauchy integral.

2000 AMS Subject Classification: 11B68, 11S80

The present Research has been conducted by the research Grant of Kwangwoon University in 2010

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

In one of an impressive series of papers (see [1, 6, 14]), Barnes developed the so-called multiple zeta and multiple gamma function. Let a_1, \dots, a_N be positive parameters. Then Barnes' multiple zeta function is defined by

$$\zeta_N(s, w|a_1, \dots, a_N) = \sum_{m_1, \dots, m_N=0} (w + m_1 a_1 + \dots + m_N a_N)^{-s}, \quad (\text{see [1]}),$$

where $\Re(s) > N$, $\Re(w) > 0$. For $m \in \mathbb{Z}_+$, we have

$$\zeta_N(-m, w|a_1, \dots, a_N) = \frac{(-1)^m m!}{(N+m)!} B_{N+m}(w, N|a_1, \dots, a_N).$$

In this paper, we consider Barnes' type multiple q -Euler numbers and polynomials. The purpose of this paper is to present new generating functions which are related to q -Euler numbers and polynomials of higher order. From the Mellin transformation of these generating functions, we derive the q -extensions of Barnes' type multiple zeta functions, which interpolate the q -Euler polynomials of higher order at negative integer. Finally, we give new identities involving q -Euler numbers and polynomials of higher order.

§2. q -Euler numbers and polynomials of higher order

In this section, we assume that $q \in \mathbb{C}$ with $|q| < 1$. Let x, a_1, \dots, a_r be complex numbers with positive real parts. Barnes' type multiple Euler polynomials are defined by

$$(2) \quad \frac{2^r}{\prod_{j=1}^r (e^{a_j t} + 1)} e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x|a_1, \dots, a_r) \frac{t^n}{n!}, \quad \text{for } |t| < \max_{1 \leq i \leq r} \frac{\pi}{|w_i|}, \quad (\text{see [6]}),$$

and $E_n^{(r)}(a_1, \dots, a_r) (= E_n^{(r)}(0|a_1, \dots, a_r))$ are called the n -th Barnes' type multiple Euler numbers. First, we consider the q -extension of Euler polynomials. The q -Euler polynomials are defined by

$$(3) \quad F_q(t, x) = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!} = [2]_q \sum_{m=0}^{\infty} (-q)^m e^{[m+x]_q t}, \quad (\text{see [8, 11, 13, 14, 15]}).$$

From (3), we have

$$E_{n,q}(x) = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l q^{lx}}{(1+q^{l+1})}.$$

In the special case $x = 0$, $E_{n,q} (= E_{n,q}(0))$ are called the n -th q -Euler numbers. From (3), we can easily derive the following relation.

$$E_{0,q} = 1, \quad \text{and } q(qE + 1)^n + E_{n,q} = 0 \text{ if } n \geq 1, \quad (\text{see [8, 16, 17]}),$$

where we use the standard convention about replacing E^k by $E_{k,q}$. It is easy to show that

$$\lim_{q \rightarrow 1} F_q(t, x) = \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (\text{see [2, 3, 19-23]}),$$

where $E_n(x)$ are the n -th Euler polynomials. For $r \in \mathbb{N}$, the Euler polynomials of order r is defined by

$$(4) \quad \left(\frac{2}{e^t + 1} \right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}, \quad \text{for } |t| < \pi.$$

Now we consider the q -extension of (4).

$$(5) \quad F_q^{(r)}(t, x) = [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} (-q)^{m_1 + \dots + m_r} e^{[m_1 + \dots + m_r + x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x) \frac{t^n}{n!},$$

where $E_{n,q}^{(r)}(x)$ are called the n -th q -Euler polynomials of order r (see [10-15]). From (5), we can derive

$$(6) \quad E_{n,q}^{(r)}(x) = \frac{[2]_q^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l q^{lx}}{(1+q^{l+1})^r}.$$

By (5) and (6), we see that

$$(7) \quad F_q^{(r)}(t, x) = [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-q)^m e^{[m+x]_q t}.$$

Thus, we note that $\lim_{q \rightarrow 1} F_q^{(r)}(t, x) = \left(\frac{2}{e^t + 1} \right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}$. In the special case $x = 0$, $E_{n,q}^{(r)} (= E_{n,q}^{(r)}(0))$ are called the n -th q -Euler numbers of order r . By (5), (6) and (7), we obtain the following proposition.

Proposition 1. For $r \in \mathbb{N}$, let

$$F_q^{(r)}(t, x) = [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} (-q)^{m_1 + \dots + m_r} e^{[m_1 + \dots + m_r + x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x) \frac{t^n}{n!}.$$

Then we have

$$E_{n,q}^{(r)}(x) = \frac{[2]_q^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l q^{lx}}{(1+q^{l+1})^r} = [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-q)^m [m+x]_q^n.$$

From the Mellin transformation of $F_q^{(r)}(t, x)$, we can derive the following equation.

$$(8) \quad \begin{aligned} \frac{1}{\Gamma(s)} \int_0^\infty F_q^{(r)}(-t, x) t^{s-1} dt &= [2]_q^r \sum_{m_1, \dots, m_r=0}^\infty \frac{(-q)^{m_1+\dots+m_r}}{[m_1+\dots+m_r+x]_q^s} \\ &= [2]_q^r \sum_{m=0}^\infty \binom{m+r-1}{m} (-q)^m \frac{1}{[m+x]_q^s}, \end{aligned}$$

where $s \in \mathbb{C}$, $x \neq 0, -1, -2, \dots$. By (8), we can define the multiple q -zeta function related to q -Euler polynomials.

Definition 2. For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we define the multiple q -zeta function related to q -Euler polynomials as

$$\zeta_{q,r}(s, x) = [2]_q^r \sum_{m_1, \dots, m_r=0}^\infty \frac{(-q)^{m_1+\dots+m_r}}{[m_1+\dots+m_r+x]_q^s}.$$

Note that $\zeta_{q,r}(s, x)$ is a meromorphic function in whole complex s -plane. From (8), we also note that

$$\zeta_{q,r}(s, x) = [2]_q^r \sum_{m=0}^\infty \binom{m+r-1}{m} (-q)^m \frac{1}{[m+x]_q^s}.$$

By Laurent series and the Cauchy residue theorem in (5) and (8), we see that

$$\zeta_q(-n, x) = E_{n,q}^{(n)}(x), \text{ for } n \in \mathbb{Z}_+.$$

Therefore, we obtain the following theorem.

Theorem 3. For $r \in \mathbb{N}$, $n \in \mathbb{Z}_+$, and $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we have

$$\zeta_q(-n, x) = E_{n,q}^{(r)}(x).$$

Let χ be the Dirichlet's character with conductor $f \in \mathbb{N}$ with $f \equiv 1 \pmod{2}$. Then the generalized q -Euler polynomial attached to χ are considered by

$$F_{q,\chi}(x) = \sum_{n=0}^\infty E_{n,\chi,q}(x) \frac{t^n}{n!} = [2]_q \sum_{m=0}^\infty (-q)^m \chi(m) e^{[m+x]_q t}.$$

From (3) and (9), we have

$$E_{n,\chi,q}(x) = \frac{[2]_q}{[2]_{q^f}} \sum_{a=0}^{f-1} (-q)^a \chi(a) E_{n,q^f}\left(\frac{x+a}{f}\right).$$

In the special case $x = 0$, $E_{n,\chi,q} = E_{n,\chi,q}(0)$ are called the n -th generated q -Euler number attached to χ .

It is known that the generalized Euler polynomials of order r are defined by

$$(10) \quad \left(\frac{2 \sum_{a=0}^{f-1} (-1)^a \chi(a) e^{at}}{e^{ft} + 1} \right)^r e^{xt} = \sum_{n=0}^{\infty} E_{n,\chi}^{(r)}(x) \frac{t^n}{n!},$$

for $|t| < \frac{\pi}{f}$.

We consider the q -extension of (10). The generalized q -Euler polynomials of order r attached to χ are defined by

$$(11) \quad \begin{aligned} F_{q,\chi}^{(r)}(t, x) &= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} (-q)^{m_1+\dots+m_r} \left(\prod_{i=1}^r \chi(m_i) \right) e^{[m_1+\dots+m_r+x]_q t} \\ &= \sum_{n=0}^{\infty} E_{n,\chi,q}^{(r)}(x) \frac{t^n}{n!}, \quad (\text{see [14, 15]}). \end{aligned}$$

Note that

$$\lim_{q \rightarrow 1} F_{q,\chi}^{(r)}(t, x) = \left(\frac{2 \sum_{a=0}^{f-1} (-1)^a \chi(a) e^{at}}{e^{ft} + 1} \right)^r.$$

By (11), we easily see that

$$\begin{aligned} E_{n,\chi,q}^{(r)}(x) &= \frac{[2]_q^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-q^x)^l \sum_{a_1, \dots, a_r=0}^{f-1} \left(\prod_{j=1}^r \chi(a_j) \right) \frac{(-q^{l+1})^{\sum_{i=1}^r a_i}}{(1+q^{(l+1)f})^r} \\ &= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} (-q)^{m_1+\dots+m_r} \left(\prod_{i=1}^r \chi(m_i) \right) [m_1 + \dots + m_r + x]_q^n. \end{aligned}$$

For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we have

$$(12) \quad \begin{aligned} &\frac{1}{\Gamma(s)} \int_0^{\infty} F_{q,\chi}^{(r)}(-t, x) t^{s-1} dt \\ &= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-q)^{m_1+\dots+m_r} \left(\prod_{i=1}^r \chi(m_i) \right)}{[m_1 + \dots + m_r + x]_q^s}, \quad (\text{see [15]}). \end{aligned}$$

From (12), we can consider the Dirichlet's type multiple q - l -function as follows :

Definition 4. For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we define the Dirichlet's type multiple q - l -function as

$$l_q(s, x|\chi) = [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-q)^{m_1+\dots+m_r} \left(\prod_{i=1}^r \chi(m_i) \right)}{[m_1 + \dots + m_r + x]_q^s}, \quad (\text{see [15]}).$$

By Laurent series and the Cauchy residue theorem in (11) and (12), we obtain the following theorem.

Theorem 5. For $n \in \mathbb{Z}_+$, we have

$$l_q(-n, x|\chi) = E_{n,\chi,q}^{(r)}(x).$$

For $h \in \mathbb{Z}$ and $r \in \mathbb{N}$, we consider the extended r -ple q -Euler polynomials.

$$\begin{aligned} F_q^{(h,r)}(t, x) &= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} q^{\sum_{j=1}^r (h-j+1)m_j} (-1)^{\sum_{j=1}^r m_j} e^{[m_1 + \dots + m_r + x]_q t} \\ (13) \quad &= \sum_{n=0}^{\infty} E_{n,q}^{(h,r)}(x) \frac{t^n}{n!}. \end{aligned}$$

Note that

$$\lim_{q \rightarrow 1} F_q^{(h,r)}(t, x) = \left(\frac{2}{e^t + 1}\right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}.$$

From (13), we note that

$$\begin{aligned} E_{n,q}^{(h,r)}(x) &= \frac{[2]_q^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-q^x)^l}{(-q^{h-r+l+1}; q)_r} \\ (14) \quad &= [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-q^{h-r+1})^m [m+x]_q^n. \end{aligned}$$

By (14), we easily see that

$$(15) \quad F_q^{(h,r)}(t, x) = [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-q^{h-r+1})^m e^{[m+x]_q t}, \quad (\text{see [11, 13, 14]}).$$

Using the Mellin transform for $F_q^{(h,r)}(t, x)$, we have

$$\begin{aligned} (16) \quad &\frac{1}{\Gamma(s)} \int_0^{\infty} F_q^{(r)}(-t, x) t^{s-1} dt \\ &= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-1)^{m_1 + \dots + m_r} q^{\sum_{j=1}^r (h-j+1)m_j}}{[m_1 + \dots + m_r + x]_q^s}, \quad (\text{see [13, 14, 15]}), \end{aligned}$$

for $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$. Now we can define the extended q -zeta function associated with $E_{n,q}^{(h,r)}(x)$.

Definition 6. For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we define the (h, q) -zeta function as

$$\zeta_{q,r}^{(h)}(s, x) = [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-1)^{m_1+\dots+m_r} q^{\sum_{j=1}^r (h-j+1)m_j}}{[m_1 + \dots + m_r + x]_q^s}.$$

Note that $\zeta_{q,r}^{(h)}(s, x)$ is also a meromorphic function in whole complex s -plane. From (16) and (15), we note that

$$(17) \quad \zeta_{q,r}^{(h)}(s, x) = [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-q^{h-j+1})^m \frac{1}{[m+x]_q^s}.$$

Using the Cauchy residue theorem and Laurent series in (16), we obtain the following theorem.

Theorem 7. For $n \in \mathbb{Z}_+$, we have

$$\zeta_{q,r}^{(h)}(-n, x) = E_{n,q}^{(h,r)}(x).$$

We consider the extended r -ple generalized q -Euler polynomials as follows :

$$(18) \quad \begin{aligned} & F_{q,\chi}^{(h,r)}(t, x) \\ &= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} q^{\sum_{j=1}^r (h-j+1)m_j} (-1)^{\sum_{j=1}^r m_j} \left(\prod_{j=1}^r \chi(m_j) \right) e^{[m_1+\dots+m_r+x]_q t} \\ &= \sum_{n=0}^{\infty} E_{n,\chi,q}^{(h,r)}(x) \frac{t^n}{n!}. \end{aligned}$$

By (18), we see that

$$(19) \quad \begin{aligned} E_{n,\chi,q}^{(h,r)}(x) &= \frac{[2]_q^r}{(1-q)^n} \sum_{a_1, \dots, a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} \left(\prod_{j=1}^r \chi(a_j) \right) \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l q^{lx} q^{(h-j+l+1)a_j}}{(-q^{(h-r+l+1)f} : q^f)_r} \\ &= \frac{[2]_q^r}{[2]_{q^f}^r} [f]_q^n \sum_{a_1, \dots, a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} \left(\prod_{j=1}^r \chi(a_j) \right) q^{\sum_{j=1}^r (h-j+1)a_j} \zeta_{q^f,r}^{(h)}\left(-n, \frac{x + \sum_{j=1}^r a_j}{f}\right). \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 8. For $n \in \mathbb{Z}_+$, we have

$$E_{n,\chi,q}^{(h,r)}(x) = \frac{[2]_q^r}{[2]_{q^f}^r} [f]_q^n \sum_{a_1, \dots, a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} \left(\prod_{j=1}^r \chi(a_j) \right) q^{\sum_{j=1}^r (h-j+1)a_j} \zeta_{q^f,r}^{(h)} \left(-n, \frac{x + \sum_{j=1}^r a_j}{f} \right).$$

From (18), we note that

$$(20) \quad \frac{1}{\Gamma(s)} \int_0^\infty F_{q,\chi}^{(h,r)}(-t, x) t^{s-1} dt = [2]_q^r \sum_{m_1, \dots, m_r=0}^\infty \frac{q^{\sum_{j=1}^r (h-j+1)m_j} (\prod_{j=1}^r \chi(m_j)) (-1)^{m_1+\dots+m_r}}{[m_1 + \dots + m_r + x]_q^s},$$

where $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$.

From (20), we define the Dirichlet's type multiple (h, q) - l -function associated with the generalized multiple q -Euler polynomials attached to χ .

Definition 9. For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we define the Dirichlet's type multiple q - l -function as follows :

$$l_q^{(h)}(s, x|\chi) = [2]_q^r \sum_{m_1, \dots, m_r=0}^\infty \frac{q^{\sum_{j=1}^r (h-j+1)m_j} (\prod_{i=1}^r \chi(m_i)) (-1)^{m_1+\dots+m_r}}{[m_1 + \dots + m_r + x]_q^s}.$$

Note that $l_q^{(h)}(s, x|\chi)$ is a meromorphic function in whole complex plane. It is easy to show that

$$l_q^{(h)}(s, x|\chi) = \frac{[2]_q^r}{[2]_{q^f}^r} \frac{1}{[f]_q^s} \sum_{a_1, \dots, a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} \left(\prod_{j=1}^r \chi(a_j) \right) q^{\sum_{j=1}^r (h-j+1)a_j} \zeta_{q^f,r}^{(h)} \left(s, \frac{x + \sum_{j=1}^r a_j}{f} \right).$$

By (19) and (20), we obtain the following theorem.

Theorem 10. For $n \in \mathbb{Z}_+$, we have

$$l_q^{(h)}(-n, x|\chi) = E_{n,\chi,q}^{(h,r)}(x).$$

Finally, we give the q -extension of Barnes' type multiple Euler polynomials in (2). For $x, a_1, \dots, a_r \in \mathbb{C}$ with positive real part, let us define the Barnes' type multiple

q -Euler polynomials in \mathbb{C} as follows :

$$\begin{aligned}
(21) \quad & F_q^{(r)}(t, x|a_1, \dots, a_r; b_1, \dots, b_r) \\
&= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} (-1)^{m_1+\dots+m_r} q^{(b_1+1)m_1+\dots+(b_r+1)m_r} e^{[a_1 m_1+\dots+a_r m_r+x]t} \\
&= \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r) \frac{t^n}{n!},
\end{aligned}$$

where $b_1, \dots, b_r \in \mathbb{Z}$. By (21), we see that

$$\begin{aligned}
& E_{n,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r) \\
&= \frac{[2]_q^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l q^{lx}}{(1+q^{la_1+b_1+1}) \dots (1+q^{la_r+b_r+1})} \\
&= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} (-1)^{m_1+\dots+m_r} q^{(b_1+1)m_1+\dots+(b_r+1)m_r} [a_1 m_1 + \dots + a_r m_r + x]_q^n.
\end{aligned}$$

From (21), we note that

$$\begin{aligned}
(22) \quad & \frac{1}{\Gamma(s)} \int_0^\infty F_q^{(r)}(-t, x|a_1, \dots, a_r; b_1, \dots, b_r) t^{s-1} dt \\
&= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-q)^{m_1+\dots+m_r} q^{b_1 m_1+\dots+b_r m_r}}{[a_1 m_1 + \dots + a_r m_r + x]_q^s}.
\end{aligned}$$

By (22), we define the Barnes' type multiple q -zeta function as follows :

$$\begin{aligned}
& \zeta_{q,r}(s, x|a_1, \dots, a_r; b_1, \dots, b_r) \\
&= [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-q)^{m_1+\dots+m_r} q^{b_1 m_1+\dots+b_r m_r}}{[a_1 m_1 + \dots + a_r m_r + x]_q^s},
\end{aligned}$$

where $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$. By (21), (22) and (23), we obtain the following theorem.

Theorem 11. For $n \in \mathbb{Z}_+$, we have

$$\zeta_{q,r}(s, x|a_1, \dots, a_r; b_1, \dots, b_r) = E_{n,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r).$$

Let χ be the Dirichlet's character with conductor $f \in \mathbb{N}$ with $f \equiv 1 \pmod{2}$. Then the generalized Barnes' type multiple q -Euler polynomials attached to χ are defined

by

$$\begin{aligned}
& F_{q,\chi}^{(r)}(t, x|a_1, \dots, a_r; b_1, \dots, b_r) \\
(24) \quad & = [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} (-q)^{m_1+\dots+m_r} q^{b_1 m_1+\dots+b_r m_r} \left(\prod_{i=1}^r \chi(m_i) \right) e^{[a_1 m_1+\dots+a_r m_r+x]_q t} \\
& = \sum_{n=0}^{\infty} E_{n,\chi,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r) \frac{t^n}{n!},
\end{aligned}$$

From (24), we note that

$$\begin{aligned}
& \frac{1}{\Gamma(s)} \int_0^{\infty} F_{q,\chi}^{(r)}(-t, x|a_1, \dots, a_r; b_1, \dots, b_r) t^{s-1} dt \\
(25) \quad & = [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-q)^{m_1+\dots+m_r} q^{b_1 m_1+\dots+b_r m_r} \left(\prod_{i=1}^r \chi(m_i) \right)}{[a_1 m_1 + \dots + a_r m_r + x]_q^s}.
\end{aligned}$$

By (25), we can define Barnes' type multiple q - l -function in \mathbb{C} . For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, let us define the Barnes' type multiple q - l -function as follows :

$$\begin{aligned}
& l_q^{(r)}(s, x|a_1, \dots, a_r; b_1, \dots, b_r) \\
(26) \quad & = [2]_q^r \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(-q)^{m_1+\dots+m_r} q^{b_1 m_1+\dots+b_r m_r} \left(\prod_{i=1}^r \chi(m_i) \right)}{[a_1 m_1 + \dots + a_r m_r + x]_q^s}.
\end{aligned}$$

Note that $l_q^{(r)}(s, x|a_1, \dots, a_r; b_1, \dots, b_r)$ is a meromorphic function in whole complex s -plane. By (24), (25) and (26), we easily see that

$$l_q^{(r)}(-n, x|a_1, \dots, a_r; b_1, \dots, b_r) = E_{n,\chi,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r)$$

for $n \in \mathbb{Z}_+$, (see [1-18]).

REFERENCES

- [1] E. W. Barnes, *On the theory of multiple gamma function*, Trans. Camb. Philos. Soc. A **196** (1904), 374-425.
- [2] I. N. Cangul, V. Kurt, H. Ozden, Y. Simsek, *On the higher-order w - q -Genocchi numbers*, Adv. Stud. Contemp. Math. **19** (2009), 39-57.
- [3] N. K. Govil, V. Gupta, *Convergence of q -Meyer-Konig-Zeller-Durrmeyer operators*, Adv. Stud. Contemp. Math. **19** (2009), 97-108.
- [4] T. Kim, *On a q -analogue of the p -adic log gamma functions and related integrals*, J. Number Theory **76** (1999), 320-329.
- [5] T. Kim, *q -Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
- [6] T. Kim, *On Euler-Barnes multiple zeta functions*, Russ. J. Math. Phys. **10** (2003), 261-267.

- [7] T. Kim, *Analytic continuation of multiple q -zeta functions and their values at negative integers*, Russ. J. Math. Phys. **11** (2004), 71–76.
- [8] T. Kim, *The modified q -Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **16** (2008), 161–170.
- [9] T. Kim, *Note on the q -Euler numbers of higher order*, Adv. Stud. Contemp. Math. **19** (2009), 25–29.
- [10] T. Kim, *Note on Dedekind type DC sums*, Adv. Stud. Contemp. Math. **18** (2009), 249–260.
- [11] T. Kim, *Note on the Euler q -zeta functions*, J. Number Theory **129** (2009), 1798–1804.
- [12] T. Kim, *A note on the generalized q -Euler numbers*, Proc. Jangjeon Math. Soc. **12** (2009), 45–50.
- [13] T. Kim, *Some identities on the q -Euler polynomials of higher order and q -stirling numbers by the fermionic p -adic integral on \mathbb{Z}_p* , Russ. J. Math. Phys. **16** (2009), 1061–9208.
- [14] T. Kim, *Barnes type multiple q -zeta functions and q -Euler polynomials*, arXiv:0912.5119v1.
- [15] T. Kim, *Note on multiple q -zeta functions*, to be appeared in Russ. J. Math. Phys., arXiv:0912.5477v1.
- [16] T. Kim, *On the q -extension of Euler and Genocchi numbers*, J. Math. Anal. Appl. **326**, 1458–1465.
- [17] T. Kim, *On p -adic q - l -functions and sums of powers*, J. Math. Anal. Appl. **329**, 1472–1481.
- [18] T. Kim, Y. Simsek, *Analytic continuation of the multiple Daehee q - l -functions associated with Daehee numbers*, Russ. J. Math. Phys. **15** (2008), 58–65.
- [19] Y. H. Kim, W. Kim, C. S. Ryoo, *On the twisted q -Euler zeta function associated with twisted q -Euler numbers*, Proc. Jangjeon Math. Soc. **12** (2009), 93–100.
- [20] H. Ozden, I. N. Cangul, Y. Simsek, *Remarks on q -Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. **18** (2009), 41–48.
- [21] K. Shiratani, S. Yamamoto, *On a p -adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci., Kyushu University Ser. A **39** (1985), 113–125.
- [22] Y. Simsek, *Theorems on twisted L -function and twisted Bernoulli numbers*, Advan. Stud. Contemp. Math. **11** (2005), 205–218.
- [23] Z. Zhang, Y. Zhang, *Summation formulas of q -series by modified Abel’s lemma*, Adv. Stud. Contemp. Math. **17** (2008), 119–129.

Taekyun Kim

Division of General Education-Mathematics, Kwangwoon University,
Seoul 139-701, S. Korea e-mail: tkkim@kw.ac.kr

Young-Hee Kim

Division of General Education-Mathematics,
Kwangwoon University,
Seoul 139-701, S. Korea e-mail: yhkim@kw.ac.kr