

WHAT IS THE ROLE OF CONTINUITY IN CONTINUOUS LINEAR FORMS REPRESENTATION?

PAUL PONCET

ABSTRACT. The recent extensions of domain theory have proved particularly efficient to study lattice-valued maxitive measures, when the target lattice is continuous. Maxitive measures are defined analogously to classical measures with the supremum operation in place of the addition. Building further on the links between domain theory and idempotent analysis highlighted by Lawson (2004), we investigate the concept of domain-valued *linear forms* on an idempotent (semi)module. In addition to proving representation theorems for continuous linear forms, we address two applications: the idempotent Radon–Nikodym theorem and the idempotent Riesz representation theorem. To unify similar results from different mathematical areas, our analysis is carried out in the general $\mathbf{Z}$ framework of domain theory.

1. INTRODUCTION

Maxitive measures are defined analogously to classical (additive) measures with the supremum operation $\oplus$ in place of the addition $+$. These measures were first introduced by Shilkret [66], and rediscovered many times. This explains why similar notions and results coexist in the literature, that we tried to survey, unify, and surpass in [56, Chapter I].

Maslov’s monograph [48], in which maxitive measures with values in ordered semirings were considered, testifies to deep connections between idempotent analysis and *order theory* or *lattice theory*. Similar initiatives have been undertaken in the framework of fuzzy set theory,

Date: February 4, 2022.

2010 *Mathematics Subject Classification.* 06A12, 06A15, 06F99, 12K10, 18A99, 28B15.

Key words and phrases. domains, continuous lattices, $\mathbf{Z}$ -theory, max-plus algebra, idempotent analysis, idempotent semifields, idempotent semimodules, Dedekind–MacNeille completion, residuation, maxitive measures, Radon–Nikodym theorem, Riesz representation theorem.

where $[0, 1]$ -valued possibility measures have been replaced by lattice-valued possibility measures (see Greco [29], Liu and Zhang [41], de Cooman et al. [17], Kramosil [37]). More recently, the branch of order theory dealing with *continuous lattices* and *domains* turned out to play a crucial role in the study of lattice-valued maxitive measures; see the work of Heckmann and Huth [31, 32], treating fuzzy set theory, category theory and continuous lattices, and of Akian [2], who favoured applications to idempotent analysis and large deviations of random processes. Connections between idempotent mathematics and continuous lattices (or domain theory) also arose in the work of Akian and Singer [3], and were surveyed by Lawson [39]. See also the early developments of Norberg [53, 54] on domain-valued random variables and the use of continuous (semi)lattices in random set theory.

The article [57] was another contribution to the strengthening of these links; we considered maxitive measures with values in a domain rather than in $\overline{\mathbb{R}}_+$. In the present paper we shall build further on the role of domain theory in idempotent analysis. We shall be especially interested in linear forms on a module over an idempotent semifield $\mathbb{k}$.

Our motivation partly comes from the following apparent paradox. Let ν be a completely maxitive measure defined on the open subsets $\mathcal{G}(E)$ of a topological space E , and taking its values in a complete lattice $\mathbb{k}$. It is known since Heckmann and Huth [31, 32] and Akian [2] that, if $\mathbb{k}$ is a continuous lattice (hence a domain), then ν admits a cardinal density, i.e. is of the form

$$(1.1) \quad \nu(\cdot) = \bigoplus_{x \in \cdot} c^+(x),$$

for some map $c^+ : E \rightarrow \mathbb{k}$. See also [56, Corollary II-5.9]. But Heckmann and Huth proved a stronger result, for they *characterized* continuity of $\mathbb{k}$ as follows: if $\mathbb{k}$ is a given complete lattice, then it is continuous *if and only if*, for every topological space E , each completely maxitive measure $\nu : \mathcal{G}(E) \rightarrow \mathbb{k}$ admits a cardinal density [32, Theorem 5].

Surprisingly, the work of Litvinov et al. [40] and Cohen et al. [16] seems to contradict this result. Indeed, these authors proved a representation theorem for *continuous* linear forms v defined on a complete $\mathbb{k}$ -module M , with $\mathbb{k}$ a complete idempotent semifield: one can write

$$(1.2) \quad v(\cdot) = \langle c, \cdot \rangle,$$

for some $c \in M$, where $\langle c, x \rangle$ denotes a $\mathbb{k}$ -valued operation defined on a subset of $M \times M$. This representation has formal and theoretical affinities with that of Equation (1.1). However, its terms require no kind of continuity assumption on $\mathbb{k}$! Coherently no reference to domain

theory appears in the last-mentioned papers. How can one understand this paradox?

To unravel it, we need to go beyond the tools of classical domain theory, and use instead the general $\mathbf{Z}$ framework of domain theory (see Bandelt and Ern  [7]). This is about selecting other subsets than the usual filtered subsets, e.g. singletons or nonempty subsets. This is done by a functor $\mathbf{Z} : \mathbf{Po} \rightarrow \mathbf{Set}$ from the category of posets to the category of sets. Then one can redefine the notions of way-above relation and continuous poset. In the case where $\mathbf{Z}$ selects nonempty subsets, a continuous poset is nothing but a *completely distributive* poset or *supercontinuous* poset in the sense of Ern  et al. [21]. And if $\mathbf{Z}$ selects singletons, it happens that the way-above relation coincides with the partial order $\geq$ and that every poset is continuous! This functor is implicitly used in [40] and [16], and this explains why these articles apparently do not ask for continuity of $\mathbb{k}$.

We warn the reader that the following notions will depend on a given functor $\mathbf{Z}$:

- way-above relation,
- continuous idempotent semifield,
- smooth linear map (or form),
- continuous linear map (or form),
- completable and complete modules,
- cuts and normal completion of a completable module,
- strongly archimedean element of a module.

In works related to $\mathbf{Z}$ -theory, it is common practice to constantly recall the dependency on $\mathbf{Z}$ ($\mathbf{Z}$ -complete poset, $\mathbf{Z}$ -way-above relation, $\mathbf{Z}$ -continuous poset, etc.); we believe however that it makes the text heavy and is not really useful if the context is clear.

A linear form $v : M \rightarrow \mathbb{k}$ is *smooth* if v commutes with infima of $\mathbf{Z}$ -sets, and *continuous* if v is smooth and commutes with arbitrary existing suprema. Under appropriate hypotheses, smooth linear forms can be represented by an ideal of the module M ; for continuous linear forms, this ideal becomes principal, i.e. is generated by an element c , and one obtains Equation (1.2) as stated by the following theorem.

Theorem 1.1. *Suppose that M is a complete module over a continuous complete idempotent semifield $\mathbb{k} \neq \{0, 1\}$, and let $v : M \rightarrow \mathbb{k}$. Then v is a non-degenerate continuous linear form on M if and only if there is an strongly archimedean element $c \in M$ such that $v(\cdot) = \langle c, \cdot \rangle$. In this case, c is unique and equals the supremum of the set $\{1 \geq v\}$.*

This result generalizes [40, Theorems 5.1 and 5.2] and [16, Corollary 39]. The implicit functor $\mathbf{Z}$ is supposed to be *union-complete*, so

that in every continuous poset the way-above relation is interpolating, i.e. such that $t \gg r$ implies $t \gg s \gg r$ for some s .

Using Theorem 1.1 we reprove the idempotent Radon–Nikodym theorem (or Sugeno–Murofushi theorem, see [68] and [56, Chapter I]). For this purpose we choose for $\mathbf{Z}$ the functor that selects singletons. If τ (resp. ν) denotes the dominating (resp. dominated) σ -maxitive measure, the module we work with is not $L_+^1(\tau)$ but a module $\mathbf{M}$ that depends on both τ and ν . We show that τ is localizable (resp. σ -principal) if and only if $\mathbf{M}$ is a complete module (resp. a σ -principal module). Moreover, if τ is σ -principal, then every σ -continuous linear form on $\mathbf{M}$ is continuous. With this result, the idempotent Radon–Nikodym theorem can be deduced easily.

Unfortunately, Theorem 1.1 is not sufficient for proving an idempotent version of the Riesz representation theorem. The idempotent Riesz theorem usually applies to a linear form $V : M \rightarrow \mathbb{R}_+$ defined on the module M of nonnegative bounded continuous maps of a Tychonoff space. It asserts that V can be expressed as a Shilkret integral with respect to some regular maxitive measure that is finite on compact subsets. Since such a measure always admits a finite cardinal density c^+ , this amounts to writing V as

$$V(f) = \bigoplus_{x \in E} \frac{f(x)}{c(x)},$$

for some map $c : E \rightarrow \mathbb{R}_+^*$ (and in fact $c = 1/c^+$), where E is the underlying topological space. But c does not need to be continuous, it is only lower-semicontinuous in general. This means that c is *outside* M , a case that is not treated by Theorem 1.1.

To take account of this situation, we introduce *module extensions*, i.e. pairs $\overline{M}/M$ with M a submodule of a complete module $\overline{M}$. For instance the module of nonnegative lower-semicontinuous maps is an extension of the module of nonnegative bounded continuous maps. We obtain the following result.

Theorem 1.2. *Suppose that $\overline{M}/M$ is a extension over a complete idempotent semifield $\mathbb{k}$, and let $v : M \rightarrow \mathbb{k}$ be a linear form on M . Assume that the extension is meet-continuous. Then v is non-degenerate continuous on $\overline{M}/M$ if and only if there is an archimedean element c in $\overline{M}/M$ such that $v(\cdot) = \langle c, \cdot \rangle$. In this case, the supremum of $\{1 \geq v\}$ in $\overline{M}$ is the least c satisfying $v(\cdot) = \langle c, \cdot \rangle$.*

For simplification purposes this theorem is limited to the case where $\mathbf{Z}$ selects singletons. A novel assumption is introduced: we ask for the extension $\overline{M}/M$ to be *meet-continuous*. This specifies that finite

infima distribute over directed suprema. This result enables one to tackle the idempotent Riesz theorem. We reprove, with a few improvements, a version of this theorem given by Choquet [15] and proved by Kolokoltsov and Maslov [36] in the locally-compact case, and a version due to Breyer and Gulinsky [12] and also reproved by Puhalskii [59]. We also prove a Riesz like theorem in the case where the topological space E is separable metrizable.

The paper is organized as follows. Section 2 recalls basics of domains and continuous posets, in the categorical framework of $\mathbf{Z}$ -theory. Section 3 deals with the concepts of idempotent semifields and modules over semirings. In Section 4 we introduce the notion of linear forms defined on a $\mathbb{k}$ -module, where $\mathbb{k}$ is an idempotent semifield. We propose a generic way of constructing such maps using ideals of the underlying module. When continuity assumptions on $\mathbb{k}$ are required, we use the tools of $\mathbf{Z}$ -theory introduced in Section 2. In Section 5 our main theorem on representation of continuous linear forms on a complete module is proved. In Section 6 we go through some applications to maxitive measures and the idempotent Radon–Nikodym theorem. Section 7 provides necessary and sufficient conditions for a module to be embeddable into a complete module. In Section 8 we prove a representation theorem for residuated forms on a module extension. In Section 9 the idempotent Riesz representation theorem is proved.

2. A PRIMER ON $\mathbf{Z}$ -THEORY FOR CONTINUOUS POSETS AND DOMAINS

A *poset* or *partially ordered set* $(P, \leq)$ is a set P equipped with a reflexive, antisymmetric and transitive binary relation $\leq$. Let us denote by $\mathbf{Po}$ the category of all posets with order-preserving maps as morphisms. A *subset selection* is a function that assigns to each poset P a certain collection $\mathbf{Z}[P]$ of subsets of P called the *$\mathbf{Z}$ -sets* of P . A *subset system* is a subset selection $\mathbf{Z}$ such that

- i) at least one $\mathbf{Z}[P]$ has a nonempty element,
- ii) for each order-preserving map $f : P \rightarrow Q$, $f(Z) \in \mathbf{Z}[Q]$ for every $Z \in \mathbf{Z}[P]$,

the point ii) meaning that $\mathbf{Z}$ is a covariant functor from $\mathbf{Po}$ to $\mathbf{Set}$ (the category of sets) with $\mathbf{Z}[f]$ defined by $\mathbf{Z}[f](Z) = f(Z)$ if $Z \in \mathbf{Z}[P]$, for every order-preserving map $f : P \rightarrow Q$. To this definition, first given by Wright et al. [72], we add a third (unusual but useful in the framework of this paper) condition:

- iii) the empty set is not in $\mathbf{Z}[P]$, for all posets P .

The suggestion of [72] to apply subset systems to the theory of continuous posets was followed by Nelson [51], Novak [55], Bandelt [6], Bandelt and Ern  [7], [8], and this research was carried on by Venugopalan [70], [71], Xu [73], Baranga [9], Menon [49], Shi and Wang [65], Ern  [20], [23] among others. Conditions *i*) and *ii*) together ensure that each $Z[P]$ contains all singletons.

The basic example of subset system is the set of directed subsets of P . This subset system is behind the classical theory of continuous posets and domains, see the monograph by Gierz et al. [27]. Here are some further examples:

- (1) Taking $Z[P]$ as the set of all nonempty subsets of P works well for investigating completely distributive lattices, see Ern  et al. [21]. Completely distributive lattices were initially examined by Raney [60], [61].
- (2) The case where $Z[P]$ is the set of filtered subsets of P was used for instance by G. Gerritse [26], Jonasson [35], Akian and Singer [3]. See also [57].
- (3) If $Z[P]$ is the set of all singletons of P , then Z is also a subset selection.
- (4) A series of papers deals with the case where $Z[P]$ is the set of chains of P , see Markowsky and Rosen [46], and Markowsky [43], [44], [45]. Using the Hausdorff maximality theorem, relations between directed subsets and chains were explored by Iwamura [34], Bruns [13], and Markowsky [42]. See also Ern  [20, p. 54].
- (5) The case where $Z[P]$ is the set of nonempty finite subsets of P was investigated by Martinez [47]. See also Frink [24] and Ern  [18].

Rather than Z , we shall often deal with the subset selection F , defined by $F[P] = \{\uparrow Z : Z \in Z[P]\}$, where $\uparrow Z$ is the upper subset generated by Z , i.e. $\uparrow Z := \{y \in P : \exists x \in Z, x \leq y\}$. The elements of $F[P]$ are the *F-sets*, or the *(Z-)filters*, of P . Although F is not a subset system in general, it satisfies the following conditions:

- i*) at least one $F[P]$ has a nonempty element,
- ii'*) for each order-preserving map $f : P \rightarrow Q$, $\uparrow f(F) \in F[Q]$ for every $F \in F[P]$,
- iii*) an F -set is never empty.

A subset selection F derived from a subset system Z as above will be called a *filter selection*. Note that, like Z , F is functorial, i.e. $F[g \circ f] = F[g] \circ F[f]$ for all order-preserving maps $f : P \rightarrow Q$ and $g : Q \rightarrow R$, if one naturally defines $F[f](F) = \uparrow f(F)$ for all $F \in F[P]$.

Translation 2.1 (Filter selections). The first three examples of subset systems given above lead to the following filter selections, respectively:

- (1) $F[P]$ is the set $\text{Up}^*[P]$ of *nonempty upper subsets* of P ,
- (2) $F[P]$ is the set $\text{Fi}[P]$ of *filters* (in the sense of [27]) of P ,
- (3) $F[P]$ is the set $\text{PFi}[P]$ of *principal filters* of P .

We now introduce the *way-above relation*, which in our context is more relevant than the usual *way-below relation*. Thus, our notions of continuous posets and domains are dual to the traditional definitions. The way-above relation has already been used to study lattice-valued upper-semicontinuous functions, see for instance [26] and [35]; see also [57]. We say that $y \in P$ is *way-above* $x \in P$, written $y \gg x$, if, for every F -set F with infimum, $x \geq \bigwedge F$ implies $y \in F$. We use the notations $\uparrow x = \{y \in P : y \gg x\}$, and for $A \subset P$, $\uparrow A = \{y \in P : \exists x \in A, y \gg x\}$. The poset P is *continuous* if every element is the F infimum of elements way-above it, i.e. $\uparrow x \in F[P]$ and $x = \bigwedge \uparrow x$ for all $x \in P$. A *domain* is a continuous poset in which every F -set has an infimum.

Translation 2.2 (Continuous posets). For our three examples of subset systems, the notion of continuous posets translates respectively as follows:

- (1) if $F = \text{Up}^*$, then a poset is continuous if and only if it is *completely distributive* (complete distributivity is sometimes called *supercontinuity*),
- (2) if $F = \text{Fi}$, then a poset is continuous if and only if it is *continuous in the sense of [57]*,
- (3) if $F = \text{PFi}$, then the way-above relation $y \gg x$ reduces to the partial order $y \geq x$, and every poset is continuous.

For a poset P , the way-above relation is *additive* if, for all $x \in P$, the subset $\{y \in P : x \gg y\}$ is either empty or directed, i.e. if whenever $x \gg y$ and $x \gg y'$, we have $x \gg z$ for some $z \in P$ such that $z \geq y$ and $z \geq y'$. A continuous poset with an additive way-above relation is *stably-continuous*. *With respect to the filter selection PFi, every poset is stably-continuous.*

A poset P has the *interpolation property* if, for all $x, y \in P$ with $y \gg x$, there exists some $z \in P$ such that $y \gg z \gg x$. For continuous posets in the classical sense, it is well known that the interpolation property holds, see e.g. [27, Theorem I-1.9]. This is a crucial feature that is behind many important results of the theory. For an arbitrary choice of Z , however, this needs no longer to be true. Deriving sufficient conditions on Z to recover the interpolation property is the goal of the following theorem. The subset selection F is *union-complete* if, for

every $V \in \mathbf{F}[\mathbf{F}[P]]$ (where $\mathbf{F}[P]$ is considered as a poset ordered by reverse inclusion $\supset$), $\bigcup V \in \mathbf{F}[P]$. As explained in [20], this condition embodies the fact that finite unions of finite sets are finite, $\supset$ -filtered unions of filtered sets are filtered, etc. The following theorem restates a result due to [55] and [7] in its dual form. We give the proof here for the sake of completeness.

Theorem 2.3. [55, 7] *If $\mathbf{F}$ is a union-complete filter selection, then every continuous poset has the interpolation property.*

Remark 2.4. In the context of $\mathbf{Z}$ -theory, many authors (see [55], [7], [70]) call *strongly continuous* a continuous poset with the interpolation property.

Proof. Let P be a continuous poset, and let $x \in P$. We need to show that $F \subset \hat{\uparrow}F$, where F denotes the $\mathbf{F}$ -set $F = \hat{\uparrow}x$. For this purpose we first prove that $\hat{\uparrow}F$ is an $\mathbf{F}$ -set. Write $\hat{\uparrow}F = \bigcup_{y \in F} \hat{\uparrow}y = \bigcup V$, where V is the collection of subsets contained in some $\hat{\uparrow}y$, $y \in F$. Considering the order-preserving map $f : P \ni y \mapsto \hat{\uparrow}y \in \mathbf{F}[P]$ (recall that $\mathbf{F}[P]$ is ordered by reverse inclusion) and using Property *ii'*) above, we have $V = \uparrow f(F) \in \mathbf{F}[\mathbf{F}[P]]$. Since $\mathbf{F}$ is union-complete, one has $\hat{\uparrow}F = \bigcup V \in \mathbf{F}[P]$. Since P is continuous,

$$x = \bigwedge \hat{\uparrow}x = \bigwedge F = \bigwedge_{y \in F} y = \bigwedge_{y \in F} (\bigwedge \hat{\uparrow}y) = \bigwedge (\bigcup_{y \in F} \hat{\uparrow}y) = \bigwedge \hat{\uparrow}F.$$

The definition of the way-above relation and the fact that $\hat{\uparrow}F \in \mathbf{F}[P]$ give $y \in \hat{\uparrow}F = \hat{\uparrow}(\hat{\uparrow}x)$, for all $y \in \hat{\uparrow}x$. This proves that P has the interpolation property. $\square$

All subset systems mentioned above are union-complete. It remains an open problem to exhibit a continuous poset with respect to some subset system that does not satisfy the interpolation property.

We should stress the fact that the machinery of category theory is justified as long as relations between posets are examined. If a single poset P is at stake, having just a collection of subsets of P at disposal could be sufficient, as in the works [6], [8], [73] (where the letter $\mathfrak{M}$ is used for the collection of selected subsets). In the present work, we hope that the relevance of using functorial (filter) selections will be made clear.

3. SEMIRINGS, SEMIFIELDS, MODULES OVER A SEMIRING

3.1. Semirings, semifields. A *semiring* is an abelian monoid $(\mathbb{k}, \oplus, 0)$ endowed with an additional binary relation $\times$ (the multiplication) that

is associative, has a unit $1 \neq 0$, distributes over $\oplus$, and admits 0 as absorbing element. A semiring is *idempotent* (or is a *doid*, see Baccelli et al. [5] or Gondran and Minoux [28]) if $\oplus$ is idempotent, i.e. $t \oplus t = t$ for all t , and *commutative* if the multiplication is commutative. An (*idempotent*) *semifield* is an (idempotent) semiring in which every non-zero element has a multiplicative inverse. We do not assume a semifield to be commutative in general (see however Remark 3.2). Notice that, if $\mathbb{k}$ is an idempotent semifield, then $\mathbb{k} \setminus \{0\}$ is a *lattice-group*. This implies that $\mathbb{k}$ is a distributive lattice; in particular, every nonempty finite subset of $\mathbb{k}$ has an infimum, and we have

$$(3.1) \quad s \wedge t = (s^{-1} \oplus t^{-1})^{-1},$$

for all $s, t \in \mathbb{k} \setminus \{0\}$.

A doid has a natural structure of partially ordered set with $s \leq t \Leftrightarrow s \oplus t = t$, whose bottom element is 0. With this point of view $s \oplus t$ is nothing but the supremum of $\{s, t\}$, hence every doid is a commutative idempotent monoid, i.e. a *semilattice*. Given a filter selection F , a doid is *complete* if every upper-bounded subset T has a supremum such that

$$(3.2) \quad s(\bigoplus T) = \bigoplus_{t \in T} st, \quad (\bigoplus T)s = \bigoplus_{t \in T} ts,$$

for all s , and if every F -set F has an infimum such that

$$(3.3) \quad s(\bigwedge F) = \bigwedge_{f \in F} sf, \quad (\bigwedge F)s = \bigwedge_{f \in F} fs,$$

for all s . With respect to the filter selection $\mathbf{PFi}$ that selects principal ideals, Equations (3.3) are trivial.

In an idempotent semifield, Equations (3.2) and (3.3) are satisfied for all subsets T (resp. F) with supremum (resp. with infimum). Thus, an idempotent semifield is complete if and only if every upper-bounded subset has a supremum. This makes the notion of complete idempotent semifield independent of the filter selection F .

The following lemma, which will be used many times in this paper, mimics a result by Akian and Singer [3, Lemma 2.1].

Lemma 3.1 (Extends [3, Lemma 2.1]). *Let F be a filter selection, and let $\mathbb{k}$ be an idempotent semifield. For all $r, s, t \in \mathbb{k}$ with $r \neq 0$, $t \gg s$ implies $tr \gg sr$.*

Proof. Let F be an F -set of $\mathbb{k}$ with infimum such that $sr \geq \bigwedge F$. The map $f : \mathbb{k} \rightarrow \mathbb{k}$ defined by $f(u) = ur^{-1}$ is order-preserving, hence the set $\uparrow f(F) = f(F)$ is an F -set. Since $t \gg s \geq (\bigwedge F)r^{-1} = \bigwedge f(F)$, we have $t \in f(F)$, so $tr \in F$. This shows that $tr \gg sr$. $\square$

From now on we use the acronym *cis* for a complete idempotent semifield distinct from $\{0, 1\}$. A *cis* is never a complete lattice; if it were, there would be a greatest element $\top$, and we would have $\top \geq 1 \Rightarrow \top^2 \geq \top \Rightarrow \top^2 = \top \Rightarrow \top = 1$, while $\top = 1$ is only possible if the *cis* coincides with $\{0, 1\}$ (a case that is excluded in the definition of a *cis*).

Remark 3.2. It is worth recalling that, by the Iwasawa theorem, every *cis* is commutative (see e.g. Birkhoff [11, Theorem 28]).

Remark 3.3 (On quasifields). Litvinov et al. [40] defined a *quasifield* as a dioid in which every non-zero element is the supremum of invertible elements and such that $t \leq 1$ whenever the subset $\{t^n : n = 1, 2, \dots\}$ is upper-bounded. They showed that every quasifield distinct from $\{0, 1\}$ can be embedded into a *cis*, and asserted that, conversely, every *cis* is a quasifield. This latter point indeed holds, for if, for some $t \neq 0$, the subset $\{t^n : n = 1, 2, \dots\}$ is upper-bounded, and if s is its supremum, then with Equations (3.2) we have $ts \leq s$; since $s \neq 0$ we deduce that $t \leq 1$.

3.2. Modules over a semiring. We now turn our attention to modules over a semiring.

Definition 3.4. Let $\mathbb{k}$ be a semiring. A *right $\mathbb{k}$ -module* is a commutative monoid $(M, \oplus, 0)$ equipped with a right action $M \times \mathbb{k} \ni (x, t) \mapsto x.t \in M$ such that, for all $x \in M$, $x.0 = 0$, $x.1 = x$, and for all $y \in M$, $s, t \in \mathbb{k}$,

$$\begin{aligned} x.(st) &= (x.s).t, \\ (x \oplus y).t &= x.t \oplus y.t, \\ x.(s \oplus t) &= x.s \oplus x.t. \end{aligned}$$

A subset of M is a *submodule* if it contains 0 and is closed under addition and external multiplication.

In the sequel we shall say *$\mathbb{k}$ -module* or *module over $\mathbb{k}$* for *right $\mathbb{k}$ -module*, and we shall only deal with modules over an idempotent semifield $\mathbb{k}$. Then the previous axioms imply that $x \oplus x = x$ for all $x \in M$, and $0.t = 0$ for all $t \in \mathbb{k}$, so the addition $x \oplus y$ of two elements x, y of M is the supremum of $\{x, y\}$ with respect to the induced partial order $x \leq y \Leftrightarrow x \oplus y = y$. In other words, $(M, \oplus, 0)$ is a semilattice.

For background on -or applications of- modules over dioids or quantales, see Zimmermann [74], Samborskiĭ and Shpiz [64], Abramsky and Vickers [1], Rosenthal [62], [33], Kruml [38], Cohen et al. [16], Litvinov et al. [40], Shpiz [67], Shpiz and Litvinov [67], Gondran and Minoux [28], Russo [63], Castella [14].

Remark 3.5. Some authors, especially in the area of idempotent analysis and max-plus algebra, prefer to call *semimodule* a module over a semiring, and *idempotent semimodule* a module over a dioid or over an idempotent semifield. However, one can see that the axioms given in Definition 3.4 do not differ from the axioms defining a classical module (over a ring), and distinctions only appear in the choice of the base semiring. The same remark can be made for axioms defining a *morphism* between modules (see Section 4 for the precise definition). Hence, from a categorical (and also from a historical) point of view, we see no reason not to keep on with the term *module*.

The following example is inspired by extreme value theory.

Example 3.6. We equip the set $\mathbb{R}_+$ of nonnegative real numbers with its idempotent semifield structure, i.e. with the maximum operation for $\oplus$, and the usual multiplication. We write $\mathbb{R}_+^{\max} = (\mathbb{R}_+, \oplus, \times)$. Let μ, σ, ξ be real numbers with $\sigma > 0$, and consider

$$M_{\mu, \sigma, \xi} = \left\{ x \in \mathbb{R} \cup \{-\infty\} : 1 + \xi \frac{x - \mu}{\sigma} > 0 \right\}.$$

Then $(M_{\mu, \sigma, \xi}, \oplus, \mathbf{0})$ is an $\mathbb{R}_+^{\max}$ -module if $\oplus$ denotes the usual maximum operation, if $\mathbf{0}$ denotes $\mu - \sigma/\xi$ if ξ is positive, $-\infty$ otherwise, and if we consider the external multiplication defined by

$$x.t = \mu - \frac{\sigma}{\xi} + \frac{\sigma t^\xi}{\xi} \left(1 + \xi \frac{x - \mu}{\sigma}\right),$$

if ξ is non-zero, and

$$x.t = x + \sigma \log(t)$$

otherwise, for all $x \in M_{\mu, \sigma, \xi}$, $t \in \mathbb{R}_+$.

4. MORPHISMS AND LINEAR FORMS

In this section, F is a union-complete filter selection, and $\mathbb{k}$ is an idempotent semifield. When continuity assumptions on $\mathbb{k}$ are required, we use the tools of Z -theory introduced in Section 2.

A *morphism* (or *linear map*) between two $\mathbb{k}$ -modules M and N is a map $f : M \rightarrow N$ satisfying both following conditions:

- *homogeneity*: $f(x.t) = f(x).t$,
- *maxitivity*: $f(x \oplus y) = f(x) \oplus f(y)$,

for all $x, y \in M$, $t \in \mathbb{k}$. Or equivalently, $f(0) = 0$ and $f(x \oplus y.t) = f(x) \oplus f(y).t$, for all $x, y \in M$ and $t \in \mathbb{k}$. A morphism f is *smooth* if, for all F -sets F of M with infimum, $f(F)$ has an infimum in N such that $f(\bigwedge F) = \bigwedge f(F)$. A (*smooth*) *linear form* on a $\mathbb{k}$ -module M is a (*smooth*) morphism $v : M \rightarrow \mathbb{k}$, where $\mathbb{k}$ is considered as a $\mathbb{k}$ -module.

Translation 4.1 (Smoothness).

- (1) If $F = \mathbf{Up}^*$, then a morphism is smooth if and only if it preserves all nonempty existing infima.
- (2) If $F = \mathbf{Fi}$, then a morphism is smooth if and only if it is Scott-continuous.
- (3) If $F = \mathbf{PFI}$, then every morphism is smooth.

Example 4.2 (Example 3.6 continued). The map $v : M_{\mu, \sigma, \xi} \rightarrow \mathbb{R}_+$ defined by $v(x) = (1 + \xi \frac{x-\mu}{\sigma})^{1/\xi}$ if ξ is non-zero, $v(x) = \exp(\frac{x-\mu}{\sigma})$ otherwise, is a linear form on $M_{\mu, \sigma, \xi}$, smooth with respect to $\mathbf{Up}^*$.

Example 4.3. The set $\mathbb{R}_+$ is still equipped with its idempotent semi-field structure. A *maxitive measure* on a σ -algebra $\mathcal{B}$ is a map $\nu : \mathcal{B} \rightarrow \overline{\mathbb{R}}_+$ such that $\nu(\emptyset) = 0$ and

$$\nu(B_1 \cup B_2) = \nu(B_1) \oplus \nu(B_2),$$

for all $B_1, B_2 \in \mathcal{B}$. It is σ -*maxitive* if it commutes with unions of nondecreasing sequences of elements of $\mathcal{B}$. The *Shilkret integral* (or *idempotent integral*) of some measurable map $f : E \rightarrow \overline{\mathbb{R}}_+$ with respect to a (σ) -maxitive measure ν on $\mathcal{B}$ is defined by

$$\int_E f.d\nu = \bigoplus_{t \in \mathbb{R}_+} t.\nu(f > t).$$

Such a map f is ν -*integrable* if its Shilkret integral is finite. Then the set M of ν -integrable maps is an $\mathbb{R}_+^{\max}$ -module, and the Shilkret integral is a linear form on M .

Notations 4.4. Let I be a subset of a $\mathbb{k}$ -module M .

- For all $t \in \mathbb{k} \setminus \{0\}$, we write $I.t = \{x.t : x \in I\}$, and $I.0 = \bigcap_{t \neq 0} I.t$. The reader is warned that $I.0$ does not coincide with $\{0\}$ in general.
- For all $x \in M$, we denote by $\langle I, x \rangle$ the set $\{t \in \mathbb{k} : x \in I.t\}$.

A subset X is *lower* if $X = \downarrow X$, where $\downarrow X := \{y : \exists x \in X, y \leq x\}$. An *ideal* I of M is a lower subset of M such that $x \oplus y \in I$, for all $x, y \in I$. An ideal I is *smooth* if, for all F -sets F of M with infimum, $\bigwedge F \in I$ implies $F \cap I \neq \emptyset$. An ideal I is *right-continuous* if $I.t = \bigcap_{s \gg t} I.s$ for all $t \in \mathbb{k}$, and *left-continuous* if $I.t = \bigcup_{t \gg s} I.s$ for all $t \in \mathbb{k}$.

The next proposition, which is inspired by the concept of *Minkowski functional* (or *gauge*) in convex analysis and by a remark of Nguyen et al. [52] on maxitive measures, provides a generic way of constructing a linear form from an ideal. We first prove a useful lemma.

Lemma 4.5 (Compare with [40, Lemma 5.1]). *Let $\mathbb{k}$ be an idempotent semifield. Then $\mathbb{k} \setminus \{0\}$ has an infimum, and*

- $\bigwedge \mathbb{k} \setminus \{0\} = 1$ if and only if $\mathbb{k} = \{0, 1\}$,
- $\bigwedge \mathbb{k} \setminus \{0\} = 0$ if and only if $\mathbb{k} \neq \{0, 1\}$.

Proof. If $\mathbb{k} = \{0, 1\}$, the result is clear, so suppose that $\mathbb{k} \neq \{0, 1\}$. This implies the existence of some $t \in \mathbb{k}, t > 0$ and $t \neq 1$. To show that 0 is the infimum of $\mathbb{k} \setminus \{0\}$, we pick some lower bound s of $\mathbb{k} \setminus \{0\}$, and we assume that $s > 0$. If $s = 1$, the definition of s gives $t > 1$, hence $0 < t^{-1} < 1 = s$, a contradiction. As a consequence, $s \neq 1$. Since $1 \in \mathbb{k} \setminus \{0\}$, we have $s < 1$, by definition of s . Hence, $0 < s^2 < s$, another contradiction. We conclude that $s = 0$, which proves that 0 is the infimum of $\mathbb{k} \setminus \{0\}$ whenever $\mathbb{k} \neq \{0, 1\}$. $\square$

Proposition 4.6. *Let $\mathbb{k}$ be an idempotent semifield, $\mathbb{k} \neq \{0, 1\}$, and let I be an ideal (resp. a smooth ideal) of M such that, for all $x \in M$, $\langle I, x \rangle$ is an F-set with infimum. Define $v : M \rightarrow \mathbb{k}$ by*

$$(4.1) \quad v(x) = \bigwedge \langle I, x \rangle,$$

for all $x \in M$. If I is right-continuous, then $I = \{1 \geq v\}$ and v is a linear form (resp. a smooth linear form) on M .

Proof. If v is given by Equation (4.1) with a right-continuous ideal I , then v is order-preserving, for if $x \leq y$, $y \in I.t$, and $t \neq 0$, then $x.t^{-1} \leq y.t^{-1} \in I$, so that $x = (x.t^{-1}).t \in I.t$. Thus, one has $\{t \in \mathbb{k} : x \in I.t\} \supset \{t \in \mathbb{k} : y \in I.t\}$, so that $v(x) \leq v(y)$.

Now let us show that $v(x) \oplus v(x') \geq v(x \oplus x')$. So let $s \gg v(x) \oplus v(x')$. Then $s \gg v(x)$, so there exists some $t \in \mathbb{k}$ such that $s \geq t$ and $x \in I.t$. There is also some t' with the corresponding properties with respect to x' . Note that $x, x' \in I.s$, which implies $x \oplus x' \in I.s$. Since I is right-continuous, we have $x \oplus x' \in I.s_0$, where $s_0 := v(x) \oplus v(x')$. If $s_0 = 0$, then $x \oplus x' \in I.0$, so that $v(x \oplus x') = 0 = s_0 = v(x) \oplus v(x')$ by definition of v . Otherwise, we can write $x \oplus x' = y.s_0$, with $y \in I$. We thus have $v(x \oplus x') = v(y).s_0$. Since $y \in I$, $v(y) \leq 1$. This leads to $v(x \oplus x') \leq s_0$, i.e. $v(x \oplus x') \leq v(x) \oplus v(x')$.

For v to be a linear form, it remains to show that $v(x.t) = v(x).t$, for all $x \in M, t \in \mathbb{k}$. This step is not difficult and left to the reader.

It is clear that $I \subset \{1 \geq v\}$. For the reverse inclusion, let $x \in \{1 \geq v\}$, i.e. $1 \geq v(x)$. To prove that $x \in I$, we use the right-continuity of I , i.e. we show that $x \in I.s$ for all $s \gg 1$. We have $s \gg v(x) = \bigwedge \langle I, x \rangle$. The subset $\langle I, x \rangle$ is assumed to be an F-set, so $s \in \langle I, x \rangle$, i.e. $x \in I.s$, which is the desired result.

Suppose in addition that I is smooth, and let us show that v is smooth. First recall that, if F is an F-set of M with infimum f_0 and $t \in \mathbb{k} \setminus \{0\}$, then $F.t^{-1}$ is an F-set such that $\bigwedge (F.t^{-1}) = f_0.t^{-1}$. We

obtain

$$\begin{aligned}
v(f_0) &= \bigwedge \{t \in \mathbb{k} \setminus \{0\} : f_0 \in I.t\} \\
&= \bigwedge \{t \in \mathbb{k} \setminus \{0\} : \bigwedge (F.t^{-1}) \in I\} \\
&= \bigwedge \bigcup_{f \in F} \{t \in \mathbb{k} \setminus \{0\} : f.t^{-1} \in I\},
\end{aligned}$$

since I is smooth. We deduce that $v(f_0) = \bigwedge_{f \in F} \bigwedge \{t \in \mathbb{k} \setminus \{0\} : f \in I.t\} = \bigwedge_{f \in F} v(f)$, so v is smooth. $\square$

When the range $\mathbb{k}$ of the map v is continuous, one can remove the assumption of right-continuity of I . This leads to the converse statement as follows.

Proposition 4.7. *Assume that $\mathbb{k}$ is a (stably-)continuous cis. A map $v : M \rightarrow \mathbb{k}$ is a (smooth) linear form on M if and only if there is some (smooth) ideal I of M such that $\langle I, x \rangle$ is an $\mathbf{F}$ -set and*

$$v(x) = \bigwedge \langle I, x \rangle,$$

for all $x \in M$. In this case:

- (1) I is right-continuous if and only if $I = \{x \in M : 1 \geq v(x)\}$;
- (2) I is left-continuous if and only if $I = \{x \in M : 1 \gg v(x)\}$.

Proof. At first we consider the case where $\mathbb{k}$ is a continuous cis. If v is a linear form, define $I := \{1 \geq v\}$. This is an ideal such that, for all $t \neq 0$, $I.t = \{t \geq v\}$, and, by Lemma 4.5, $I.0 = \{x \in M : v(x) = 0\}$. Since $\mathbb{k}$ is continuous, I is right-continuous. Moreover, $\langle I, x \rangle$ equals the principal filter generated by $v(x)$, hence is an $\mathbf{F}$ -set, and $v(x) = \bigwedge \langle I, x \rangle$ for all x .

If $\mathbb{k}$ is stably-continuous and v is smooth, we can rather define $I := \{1 \gg v\}$. By hypothesis the way-above relation is additive, so that I is an ideal. Also, $I.t = \{t \gg v\}$ by Lemma 3.1, and $I.0 = \{x \in M : v(x) = 0\}$ since $\mathbb{k}$ is continuous. Left-continuity of I holds by the interpolation property. Moreover, for all x , $\langle I, x \rangle = \uparrow v(x)$, which is an $\mathbf{F}$ -set whose infimum is $v(x)$ since $\mathbb{k}$ is continuous. Smoothness of I is a consequence of the smoothness of v and of the fact that $\langle I, x \rangle$ is nonempty.

Conversely, assume that Equation (4.1) is satisfied, and let us show that v is a linear form. Let $J = \bigcap_{s \gg 1} I.s$. Then J is an ideal of M containing I . We prove that, for all $t \in \mathbb{k}$,

$$(4.2) \quad t \geq v(x) \Leftrightarrow x \in J.t.$$

Using Lemma 4.5, it suffices to prove Equivalence (4.2) for $t \neq 0$. If $t \geq v(x)$ and $s \gg 1$, then $st \gg t$ by Lemma 3.1, so $st \gg v(x)$. This

gives $x \in I.st$, hence $x.t^{-1} \in I.s$, for all $s \gg 1$. By definition of J we get $x.t^{-1} \in J$, i.e. $x \in J.t$. Now we suppose that $x \in J.t$, and we want to show that $t \geq v(x)$. So let $s \gg t$. If $u = st^{-1}$, then $u \gg 1$ (see again Lemma 3.1), so $xt^{-1} \in I.u$. Thus, $x \in I.s$. The definition of v implies $s \geq v(x)$, for all $s \gg t$. By continuity of $\mathbb{k}$ we have $t \geq v(x)$. So Equivalence (4.2) is proved. This also shows that J is right-continuous and that $\langle J, x \rangle$ is an F-set whose infimum is $v(x)$, for all $x \in M$. By Proposition 4.6, v is a linear form, and, as in the proof of Proposition 4.6, v is smooth if I is smooth.

To finish the proof, suppose again that v is a linear form defined by Equation (4.1). If I is right-continuous, then the previous point implies $I = J$ and $I.t = \{t \geq v\}$, for all $t \in \mathbb{k}$, so Item (1) is proved. If I is left-continuous, the inclusion $I \supset \{1 \gg v\}$ is clear, by definition of v and $\gg$. If $x \in I$, then $x \in I.s$ for some $s \in \mathbb{k}$ such that $1 \gg s$, by left-continuity of I . This implies that $s \geq v(x)$, so that $1 \gg v(x)$, and Item (2) is proved. $\square$

Translation 4.8. Back to the three main instances of filter selections, the assumptions of Proposition 4.7 translate as follows.

- (1) $\langle I, x \rangle$ is an Up*-set if and only if $\langle I, x \rangle$ is nonempty. This condition is satisfied for all $x \in M$ as soon as $I \neq \{0\}$ and, for all $x, y \in M$ with $y \neq 0$, there exists some $t \in \mathbb{k}$ with $x \leq y.t$.
- (2) $\langle I, x \rangle$ is a Fi-set if and only if $\langle I, x \rangle$ is nonempty. As above, this condition is satisfied for all $x \in M$ as soon as $I \neq \{0\}$ and, for all $x, y \in M$ with $y \neq 0$, there exists some $t \in \mathbb{k}$ with $x \leq y.t$.
- (3) $\langle I, x \rangle$ is a PFi-set if and only if $\langle I, x \rangle$ has a least element.

The last case leads to the following corollary.

Corollary 4.9. *Let $\mathbb{k}$ be an idempotent semifield, $\mathbb{k} \neq \{0, 1\}$. A map $v : M \rightarrow \mathbb{k}$ is a linear form on M if and only if there is some ideal I of M such that*

$$x \in I.t \iff t \geq v(x),$$

for all $x \in M$, $t \in \mathbb{k}$. In this case, I equals $\{1 \geq v\}$.

Proof. Let F be the filter selection PFi that selects principal filters. With this choice, $\mathbb{k}$ is continuous, and the way-above relation coincides with $\geq$, so that I is necessarily right-continuous. To conclude, use Proposition 4.7. $\square$

At this early stage, the reader may already understand, from the previous proof, where the “paradox” evoked in the Introduction comes from: no continuity assumption seems to be needed in the terms of Corollary 4.9, but this is simply due to the fact that, with respect to

the filter selection $\mathbf{PFi}$, the $\text{cis } \mathbb{k}$ is always continuous. This will be made even clearer in Section 5, where we shall deal with the representation of continuous linear forms.

5. CONTINUOUS LINEAR FORMS ON A COMPLETE MODULE

5.1. Continuity, residuation. In this section, we prove a representation theorem for continuous linear forms on a complete module. Let $\mathbf{F}$ be a union-complete filter selection, and let M, N be modules over an idempotent semifield $\mathbb{k}$. A morphism $f : M \rightarrow N$ is *continuous* if it is smooth and such that, for every subset $X \subset M$ with a supremum in M , $f(X)$ has a supremum in N satisfying $f(\bigoplus X) = \bigoplus f(X)$.

Translation 5.1 (Continuity).

- (1) If $\mathbf{F} = \mathbf{Up}^*$, then a morphism is continuous if and only if it preserves all existing infima and suprema.
- (2) If $\mathbf{F} = \mathbf{Fi}$, then a morphism is continuous if and only if it is bi-Scott-continuous (this notion was called *wo-continuity* by Shpiz [67]).
- (3) If $\mathbf{F} = \mathbf{PFi}$, then a morphism is continuous if and only if it preserves all existing suprema (this notion of continuity is the one adopted by Cohen et al. [16]; Litvinov et al. [40] and Shpiz [67] called such a morphism a *b-morphism*).

We say that M is *completable* if, for all $x \in M$, the map $\mathbb{k} \rightarrow M, t \mapsto x.t$ is a continuous morphism, i.e. if, for all $x \in M$ and all $T \subset \mathbb{k}$ with supremum,

$$(5.1) \quad x. \bigoplus T = \bigoplus_{t \in T} x.t,$$

and if for all $x \in M$ and all $\mathbf{F}$ -sets F of $\mathbb{k}$ with infimum,

$$(5.2) \quad x. \bigwedge F = \bigwedge_{f \in F} x.f.$$

Remark 5.2. Note that, if Equation (5.1) is satisfied for every $T \subset \mathbb{k}$ with supremum, then Equation (5.2) is also satisfied for every $F \subset \mathbb{k}$ with *non-zero infimum* (one does not need F be to an $\mathbf{F}$ -set in this case). Hence in the definition of completability the role of Equation (5.2) is to control the behaviour of $\mathbb{k} \rightarrow M, t \mapsto x.t$ around zero. Therefore, the case $\mathbf{F} = \mathbf{Up}^*$ is demanding, while with $\mathbf{F} = \mathbf{PFi}$ this behaviour is unconstrained.

Also, M is *complete* if it is completable and such that every upper-bounded subset (resp. every $\mathbf{F}$ -set) has a supremum (resp. an infimum). In Section 7, Theorem 7.1 will define the concept of *normal completion*

of a completable module and show that completability is equivalent to embeddability into a complete module.

A map $f : M \rightarrow N$ between $\mathbb{k}$ -modules M, N is *residuated* if there exists a (necessarily unique) map $f^\# : N \rightarrow M$, called the *adjoint* of f , and satisfying

$$x \leq f^\#(y) \iff y \geq f(x),$$

for all $x \in M, y \in N$. Residuated maps are related to Galois connections, see Ern  et al. [22]. A *residuated form* on M is a homogeneous residuated map from M to $\mathbb{k}$. A map $v : M \rightarrow \mathbb{k}$ is *non-degenerate* if $\{x \in M : 1 \geq v(x)\}$ is upper-bounded. For instance the map $M \ni x \mapsto 0 \in \mathbb{k}$ is a non-degenerate (continuous) linear form if and only if M has a greatest element.

Lemma 5.3. *Let M be a complete module over a cis $\mathbb{k}$. Then a map $v : M \rightarrow \mathbb{k}$ is a smooth residuated form if and only if it is a non-degenerate continuous linear form.*

Proof. Necessity is clear. For sufficiency, let v be a non-degenerate continuous linear form, and let $w : \mathbb{k} \rightarrow M, t \mapsto \bigoplus\{x \in M : t \geq v(x)\}$. This map is well-defined since v is non-degenerate and homogeneous. If $t \geq v(x)$, then $x \leq w(t)$ by definition of w . Conversely, if $x \leq w(t)$, then $v(x) \leq v(w(t))$, and since v preserves arbitrary existing suprema, $v(x) \leq \bigoplus\{v(x') : x' \in M, t \geq v(x')\} \leq t$. This proves that w is the adjoint of v , hence v is a (smooth) residuated form. $\square$

Remark 5.4. If $\mathbf{F} = \mathbf{PFi}$, the previous lemma identifies residuated forms with non-degenerate *b-linear functionals* in the sense of Litvinov et al. [40].

5.2. Archimedean elements and scalar product. As explained in the paragraph after Lemma 3.1, a cis $\mathbb{k}$ has no greatest element. However, we can conventionally add a top $\top$ to $\mathbb{k}$ and define $\bar{\mathbb{k}} = \mathbb{k} \cup \{\top\}$. Naturally extending $\oplus$ and $\times$ to $\bar{\mathbb{k}}$ by $t \oplus \top = \top \oplus t = \top, t \cdot \top = \top \cdot t = \top$ if $t \neq 0$, and $0 \cdot \top = \top \cdot 0 = 0$, we see that $\bar{\mathbb{k}}$ has the structure of a complete dioid. We also define $\top^{-1} = 0$ and $0^{-1} = \top$. If $x, c \in M$, we let

$$x \backslash c = \bigoplus\{t \in \mathbb{k} : c \geq x \cdot t\}$$

whenever this set is upper-bounded, and $x \backslash c = \top$ otherwise. Also, the infimum of the subset $F_c(x) = \{t \in \mathbb{k} : c \cdot t \geq x\}$ in $\bar{\mathbb{k}}$ is denoted by $\langle c, x \rangle$, and one can easily check that

$$\langle c, x \rangle = (x \backslash c)^{-1},$$

for all $x, c \in M$. We are interested in conditions on c ensuring that the map $x \mapsto \langle c, x \rangle$ is a continuous or a residuated linear form on M . So we need the

Definition 5.5. Let M be a module over an idempotent semifield $\mathbb{k}$. An element $c \in M$ is called *archimedean* if both following conditions are satisfied:

- the subset $F_c(x)$ is nonempty for all $x \in M$,
- if the infimum of $F_c(x)$ is zero, then $F_c(x) \supset \mathbb{k} \setminus \{0\}$,

where $F_c(x)$ denotes the subset $\{t \in \mathbb{k} : c.t \geq x\}$.

The first condition implies that the bracket $\langle c, x \rangle$ is well-defined in $\mathbb{k}$ for all $x \in M$. The second condition may seem unnatural and so deserves some explanation. First observe that it will automatically be satisfied in any of the following standard situations:

- if $F \in \{\text{Up}^*, \text{Fi}\}$ and M is completable,
- if the cis $\mathbb{k}$ is continuous with respect Fi ,
- in particular if $\mathbb{k}$ is a totally ordered cis (e.g. $\mathbb{k} = \mathbb{R}_+^{\max}$).

To see how it works, let us suppose in the following lines that M is a completable module over a cis $\mathbb{k}$, and that c is such that $F_c(x)$ is nonempty for all $x \in M$. How far is then $x \mapsto \langle c, x \rangle$ from being a residuated map?

If $\langle c, x \rangle$ is non-zero, then the infimum of the subset $F_c(x)$ is reached (see Remark 5.2), so that $c.\langle c, x \rangle \geq x$ or, in other words,

$$(5.3) \quad x \leq c.t \iff t \geq \langle c, x \rangle,$$

for all $t \in \mathbb{k}$. However, this equivalence is no longer guaranteed if $\langle c, x \rangle = 0$. This is where a “smooth” behaviour of $\mathbb{k} \rightarrow M, t \mapsto x.t$ around zero is needed, in accordance with Remark 5.2; so at this stage we must distinguish between the different filter selections.

The important fact is that the subset $F_c(x)$ is always filtered (if nonempty) by Equation (3.1). *This makes Fi the most natural filter selection to use on $\mathbb{k}$ -modules.* As a consequence, if $F \in \{\text{Up}^*, \text{Fi}\}$, Equivalence (5.3) is satisfied by completeness of M , even if $\langle c, x \rangle = 0$. Thus, the second condition in Definition 5.5 is fulfilled, and the map $x \mapsto \langle c, x \rangle$ is residuated. Moreover, $\langle c, x \rangle = 0$ implies $x = 0$.

The case $F = \text{PFi}$ is more delicate. We might include in the definition of c that $F_c(x)$ be an F -set, so here a principal filter; but this would imply that $\langle c, x \rangle \neq 0$ whenever $x \neq 0$, a property that is not desirable for applications (see e.g. the case of the Riesz representation theorem in Section 9). That is why we introduced a second *ad hoc* condition in the definition of an archimedean element. The following proposition gives sufficient conditions on $\mathbb{k}$ for this condition to hold.

Proposition 5.6. *Let $\mathbb{k}$ be an idempotent semifield. Consider the following conditions:*

- (1) *1 is way-above 0 with respect to $\mathbf{Fi}$;*
- (2) *there is a $t \in \mathbb{k}$ way-above 0 with respect to $\mathbf{Fi}$;*
- (3) *every filter with a zero infimum contains $\mathbb{k} \setminus \{0\}$;*
- (4) *every unbounded ideal coincides with $\mathbb{k}$;*
- (5) *$\mathbb{k}$ is continuous with respect to $\mathbf{Fi}$;*
- (6) *$\mathbb{k}$ is totally ordered;*

Then (6) $\Rightarrow$ (5) $\Rightarrow$ (4) $\Leftrightarrow$ (3) $\Leftrightarrow$ (2) $\Leftrightarrow$ (1). If any of these conditions is satisfied, then

- *$\mathbb{k}$ is join-continuous (with respect to $\mathbf{Fi}$), i.e. satisfies $s \oplus \bigwedge F = \bigwedge (s \oplus F)$ for all filters F and $s \in \mathbb{k}$;*
- *the second condition of Definition 5.5 always holds;*
- *for all $t \in \mathbb{k} \setminus \{0\}$ and $s < 1$, there is some $n \in \mathbb{N}$ such that $t \geq s^n$.*

An archimedean element c of M is *strongly archimedean* if $t \gg s$ implies $c.t \gg c.s$ for all $s, t \in \mathbb{k}$. For an archimedean element c , the map $x \mapsto \langle c, x \rangle$ is smooth only if c is strongly archimedean; the converse statement holds as soon as $\mathbb{k}$ is continuous.

The following result justifies the term *scalar product* for the bracket $\langle \cdot, \cdot \rangle$ (see also Cohen et al. [16, Section 3] for more on this topic; note that these authors preferred to call scalar product the bracket $(\cdot \setminus \cdot)$ rather than $\langle \cdot, \cdot \rangle$).

Lemma 5.7. *Let M, N be modules over a cis $\mathbb{k}$, and let $f : M \rightarrow N$ be a (smooth) residuated linear map. If c is a (strongly) archimedean element of N then $f^\#(c)$ is a (strongly) archimedean element of M , and we have*

$$\langle c, f(x) \rangle = \langle f^\#(c), x \rangle,$$

for all $x \in M$, where $f^\#$ denotes the upper adjoint of f .

Proof. If $t \in \mathbb{k} \setminus \{0\}$, we have $c.t \geq f(x) \Leftrightarrow c \geq f(x.t^{-1}) \Leftrightarrow f^\#(c) \geq x.t^{-1} \Leftrightarrow f^\#(c).t \geq x$. Since c is archimedean, $F_c(f(x))$ contains some non-zero element, so that $F_{f^\#(c)}(x)$ is nonempty, so the first condition for $f^\#(c)$ to be archimedean is checked. Now if the subset $F_{f^\#(c)}(x)$ has zero infimum, then either it contains 0 (and in this case $x = 0$ so that $F_{f^\#(c)}(x) = \mathbb{k}$) or it does not. In the latter case, the series of equivalence at the beginning of the proof shows that $F_{f^\#(c)}(x) = F_c(f(x)) \setminus \{0\}$. This implies that $F_c(f(x))$ has zero infimum, thus contains $\mathbb{k} \setminus \{0\}$ since c is archimedean. Therefore, $F_{f^\#(c)}(x)$ also contains $\mathbb{k} \setminus \{0\}$, and we have proved that $f^\#(c)$ is archimedean. Using again the equivalence

$c.t \geq f(x) \Leftrightarrow f^\#(c).t \geq x$ for all $t \in \mathbb{k} \setminus \{0\}$, we deduce that $t \geq \langle c, f(x) \rangle \Leftrightarrow t \geq \langle f^\#(c), x \rangle$ for all $t \in \mathbb{k} \setminus \{0\}$, so

$$\langle c, f(x) \rangle = \langle f^\#(c), x \rangle.$$

For the rest of the proof, assume that f is smooth and that c is strongly archimedean. Let $t \gg s$, and let us show that $f^\#(c).t \gg f^\#(c).s$. For this purpose, let F be an F-set of M with infimum such that $f^\#(c).s \geq \bigwedge F$. Then $f^\#(c.s) \geq \bigwedge F$, hence $c.s \geq f(\bigwedge F)$. Since f is smooth, this implies that $c.s \geq \bigwedge f(F)$. Since $\uparrow f(F)$ is an F-set of $\mathbb{k}$ and $c.t \geq c.s$, we obtain $c.t \geq f(x)$ for some $x \in F$. This gives $f^\#(c).t = f^\#(c.t) \geq x$, and the result is proved. $\square$

We are now in a position to prove the main result of this section.

Theorem 5.8 (Compare [40, Theorems 5.1-5.2], [16, Corollary 39]). *Suppose that M is a complete module over a continuous cis $\mathbb{k}$, and let $v : M \rightarrow \mathbb{k}$. The following conditions are equivalent:*

- (1) v is a smooth residuated form on M ,
- (2) v is a non-degenerate continuous linear form on M ,
- (3) $v(\cdot) = \langle c, \cdot \rangle$, for some strongly archimedean element $c \in M$.

If these conditions are satisfied, then c is unique and equals the supremum of the set $\{x \in M : 1 \geq v(x)\}$.

Proof. Equivalence between (1) and (2) is given by Lemma 5.3, and the implication (3) $\Rightarrow$ (1) was the purpose of Paragraph 5.2. So let us prove that (1) implies (3). Let v be a smooth residuated form on M . Define c as the supremum of the set $\{x \in M : 1 \geq v(x)\}$, i.e. $c = v^\#(1)$, where $v^\#$ is the adjoint of v . If one notices that 1 is a strongly archimedean element in $\mathbb{k}$, then c is a strongly archimedean element in M by Lemma 5.7, and one has

$$\langle 1, v(x) \rangle = \langle v^\#(1), x \rangle,$$

for all $x \in M$, that is $v(x) = \langle c, x \rangle$, for all $x \in M$. Uniqueness is deduced from the fact that $x \leq c \Leftrightarrow 1 \geq \langle c, x \rangle$. $\square$

Example 5.9 (Example 4.2 continued). We introduced the linear form v on $M_{\mu, \sigma, \xi}$ defined by $v(x) = (1 + \xi \frac{x-\mu}{\sigma})^{1/\xi}$ if ξ is non-zero, $v(x) = \exp(\frac{x-\mu}{\sigma})$ otherwise. An easy computation shows that μ is the supremum of $\{x \in M_{\mu, \sigma, \xi} : 1 \geq v(x)\}$, and that $v(x) = \langle \mu, x \rangle$ for all $x \in M_{\mu, \sigma, \xi}$. Moreover, μ is strongly archimedean (with respect to $F = \text{Up}^*$), for if $t > s$, then $\mu.t = \mu + \sigma \frac{t^\xi - 1}{\xi} > \mu.s$ if ξ is non-zero, and $\mu.t = \mu + \sigma \log(t) > \mu.s$ otherwise. Hence, v is a smooth residuated form on $M_{\mu, \sigma, \xi}$ (where smoothness is understood with respect to $F = \text{Up}^*$).

This theorem upgrades a result by Cohen et al. [16, Corollary 39] deduced from a geometric Hahn–Banach type theorem [16, Theorem 34]. A different formulation will be proved in Section 8, in the framework of *module extensions*. As for now, we use Theorem 5.8 to reprove the Radon–Nikodym theorem for the Shilkret integral.

6. THE RADON–NIKODYM THEOREM: A DIFFERENT PERSPECTIVE

We come back to the Radon–Nikodym theorem for the Shilkret integral surveyed in [56, Chapter I]; here we deduce this result from the order-theoretical developments of the previous section. See [56, Chapters I–II] for definitions and notations related to maxitive measures. The filter selection used throughout this section is **PFi**, i.e. the one that selects principal filters.

6.1. Complements on σ -complete modules. The next result prepares applications to the Radon–Nikodym theorem. It gives sufficient conditions on a module M over a *cis* $\mathbb{k}$ in order that every linear form on M be continuous. The module M is *σ -complete* if every upper-bounded countable subset has a supremum and if M is completable, i.e. if

$$x. \bigoplus_{t \in T} T = \bigoplus_{t \in T} x.t,$$

for all $x \in M$, $T \subset \mathbb{k}$ with supremum. We say that M is *σ -principal* if every upper-bounded σ -ideal is principal, i.e. of the form $\downarrow x$ for some $x \in M$. A subset G of M is *generating* if, for all $x \in M$, $x = \bigoplus \downarrow x \cap G$. Also, the module is *countably generated* if there exists a generating subset G such that $\downarrow x \cap G$ is countable, for all $x \in M$. A linear form v on M is *σ -continuous* if, for every countable subset $X \subset M$ admitting a supremum in M , $v(X)$ has a supremum in $\mathbb{k}$ satisfying $v(\bigoplus X) = \bigoplus v(X)$.

Proposition 6.1. *Let M be a σ -complete module over a *cis* $\mathbb{k}$.*

- (1) *If M is countably generated, then M is σ -principal.*
- (2) *If M is σ -principal, then M is complete and every σ -continuous linear form is continuous.*

Proof. (1) Assume that M is countably generated by some subset G . Let I be an upper-bounded σ -ideal of M . If u is an upper-bound of I , the subset $I \cap G$ is included in the countable subset $\downarrow u \cap G$, hence is countable. So let $x := \bigoplus G \cap I \in I$. It is easily seen that $x = \bigoplus I$, hence $I = \downarrow x$, i.e. I is a principal ideal.

(2) Assume that M is σ -principal, and let X be an upper-bounded subset of M . The σ -ideal I generated by X is made up of elements

lower than joins of countable subsets of X . Since I is upper-bounded, it is principal, so we have $I = \downarrow x$ for some $x \in I$, and x is of the form $x = \bigoplus G$ for some countable subset G of X . Thus, we have $x = \bigoplus G = \bigoplus I = \bigoplus X$, so that M is complete. Moreover, if $v : M \rightarrow \mathbb{k}$ is a σ -continuous linear form, then $v(\bigoplus X) = v(\bigoplus G) = \bigoplus v(G) \leq \bigoplus v(X)$, hence v is continuous. $\square$

6.2. Maxitive measures as linear forms. Let $\mathcal{E}$ be a semi- σ -algebra on some nonempty set E and ν, τ be σ -maxitive measures on $\mathcal{E}$. We shall assume that ν is finite and absolutely continuous with respect to τ , in symbols $\nu \dashv \tau$. In order to apply the results of Section 5, we merely want to get rid of the collection of τ -negligible subsets. We could consider the quotient space $\mathcal{E}/\tau$, but this would not give us the structure of module over the idempotent semifield $\mathbb{R}_+^{\max} = (\mathbb{R}_+, \max, \times)$ that we need. A better idea is the following.

Let $\mathcal{E}_+^\nu = \mathcal{L}_+^1(E, \mathcal{E}, \nu)$ be the set of all ν -integrable lsm maps $g : E \rightarrow \overline{\mathbb{R}}_+$. A map n in $\mathcal{E}_+^\nu$ is τ -negligible if the subset $\{n > 0\}$ is τ -negligible. We define on $\mathcal{E}_+^\nu$ the equivalence relation $\diamond$ by $f \diamond g$ if and only if, for some τ -negligible map n , we have $f \oplus n = g \oplus n$. We denote by $\langle g \rangle$ the equivalence class of a $g \in \mathcal{E}_+^\nu$. Then the quotient set $\mathbf{M} := \mathcal{E}_+^\nu / \tau := \mathcal{E}_+^\nu / \diamond$ is a σ -complete module over $\mathbb{R}_+^{\max}$ with external multiplication $\mathbf{f} \cdot t := \langle t \cdot f \rangle$ and countable addition $\bigoplus_{j=1}^\infty \mathbf{g}_j = \langle \bigoplus_{j=1}^\infty g_j \rangle$, for all $t \in \mathbb{R}_+$ and $\mathbf{f} = \langle f \rangle, \mathbf{g}_j = \langle g_j \rangle \in \mathbf{M}$. The induced partial order is $\mathbf{f} \leq \mathbf{g}$ if and only if $\{f > g\}$ is τ -negligible. The reader can check that the previous definitions do not depend on the choice of the representatives f, g , etc.

Recall that τ on $\mathcal{E}$ is *localizable* if, for each σ -ideal $\mathcal{J}$ of $\mathcal{E}$, there exists some $L \in \mathcal{E}$ such that

- $S \setminus L$ is τ -negligible, for all $S \in \mathcal{J}$,
- if there is some $G \in \mathcal{E}$ such that $S \setminus G$ is τ -negligible for all $S \in \mathcal{J}$, then $L \setminus G$ is τ -negligible.

In this case, $\mathcal{J}$ is said to be *localized* in L .

Proposition 6.2. *Let ν, τ be σ -maxitive measures on $\mathcal{E}$. Assume that ν is finite and such that $\nu \dashv \tau$. Then τ is localizable (resp. σ -principal) if and only if $\mathcal{E}_+^\nu / \tau$ is a complete module (resp. a σ -principal module).*

Proof. A preliminary remark is that, since ν is finite, the lsm map 1_G is ν -integrable for all $G \in \mathcal{E}$. Assume that $\mathbf{M} = \mathcal{E}_+^\nu / \tau$ is a complete module, and let $\mathcal{J}$ be a σ -ideal of $\mathcal{E}$. Then the σ -ideal I generated by $\{\langle 1_S \rangle : S \in \mathcal{J}\}$ is upper-bounded (by $\langle 1_E \rangle$) in $\mathbf{M}$. Hence there is some $f \in \mathcal{E}_+^\nu$ such that $\langle f \rangle$ is the supremum of I . In particular, if $S \in \mathcal{J}$, there is some τ -negligible lsm map n such that $1_S \leq f \oplus n$, so that $S \subset L \cup \{n > 0\}$, where $L := \{f > 2^{-1}\}$. As a consequence, $S \setminus L$ is

τ -negligible for all $S \in \mathcal{J}$. To show that $\mathcal{J}$ is localized in L , let $G \in \mathcal{E}$ such that $S \setminus G$ is τ -negligible for all $S \in \mathcal{J}$. Then $\langle 1_G \rangle$ is an upper-bound of I , so that $\langle f \rangle \leq \langle 1_G \rangle$ by definition of f . Since $2^{-1} \cdot 1_L \leq f$, we deduce that $L \setminus G$ is τ -negligible, hence that τ is localizable.

If $\mathbf{M}$ is σ -principal, we can impose L to belong to $\mathcal{J}$ and to be such that $\langle 1_L \rangle$ generates I . Then L generates $\mathcal{J}$, and this proves that τ is σ -principal.

Conversely, suppose that τ is localizable, and let I be an upper-bounded σ -ideal of $\mathbf{M}$. If $q \in \mathbb{Q}_+$, let $\mathcal{J}_q = \{\{f > q\} : \langle f \rangle \in I\}$. This is a σ -ideal, hence it is localized in some $L_q \in \mathcal{E}$. Let $\langle g \rangle$ be an upper-bound of I . Then $S \setminus \{g > q\}$ is τ -negligible, for all $S \in \mathcal{J}_q$ and all $q \in \mathbb{Q}_+$. Since $\mathcal{J}_q$ is localized in L_q we deduce that $L_q \setminus \{g > q\}$ is τ -negligible. This implies that the map ℓ defined by $\ell = \bigoplus_{q \in \mathbb{Q}_+} q \cdot 1_{L_q}$ is ν -integrable and satisfies $\langle \ell \rangle \leq \langle g \rangle$. To show that $\langle \ell \rangle$ is the supremum of I , it suffices to prove that $\langle \ell \rangle$ is an upper-bound of I . If $\langle f \rangle \in I$, there exists some τ -negligible subset $N_q \in \mathcal{E}$ such that $\{f > q\} \subset L_q \cup N_q$. If $n = \bigoplus_{q \in \mathbb{Q}_+} q \cdot 1_{N_q}$, then $\{n > 0\} \subset \bigcup_{q \in \mathbb{Q}_+} N_q$, so n is τ -negligible. We have $f \leq \ell \oplus n$, so that $\langle f \rangle \leq \langle \ell \rangle$. This proves that $\langle \ell \rangle$ is the supremum of I , and that $\mathbf{M}$ is complete.

If τ is σ -principal, then the set L_q can be chosen of the form $\{\ell_q > q\}$, where $\langle \ell_q \rangle \in I$. It can be seen that ℓ and $\bigoplus_{q \in \mathbb{Q}_+} \ell_q$ are equivalent, so that $\langle \ell \rangle = \langle \bigoplus_{q \in \mathbb{Q}_+} \ell_q \rangle \in I$. This shows that I is principal, so that $\mathbf{M}$ is a σ -principal module. $\square$

We denote by $\mathbf{v}$ the map induced by ν on $\mathbf{M}$, i.e.

$$\mathbf{v}(\mathbf{f}) = \int^\infty f \, d\nu,$$

for all $\mathbf{f} = \langle f \rangle \in \mathbf{M}$. Since $\mathcal{E}_+^\nu$ demands ν -integrable maps, we have $\mathbf{v}(\mathbf{f}) < \infty$, so $\mathbf{v}$ is a σ -continuous linear form on $\mathbf{M}$. We shall say that ν is τ -continuous if $\mathbf{v}$ is continuous. As a corollary of Theorem 5.8 we have the following result. Recall that a map $g : E \rightarrow \overline{\mathbb{R}}_+$ is *upper-semimeasurable* or *usm* if $\{g < t\} \in \mathcal{E}$ for all $t \in \mathbb{R}_+$.

Theorem 6.3. *Let ν, τ be σ -maxitive measures on $\mathcal{E}$. Assume that ν is finite and τ is localizable. Then the following assertions are equivalent:*

- $\nu \dashv \tau$ and ν is τ -continuous,
- ν has a usm relative density with respect to τ .

Proof. Since ν is finite and τ is localizable, $\mathbf{M} = \mathcal{E}_+^\nu / \tau$ is a complete module by Proposition 6.2. From the identity $\langle \mathbf{c}, \mathbf{f} \rangle = \bigoplus_{x \in E}^\tau \frac{f(x)}{c(x)}$, which holds for all $\mathbf{f} = \langle f \rangle, \mathbf{c} = \langle c \rangle \in \mathbf{M}$, we deduce that ν has a usm relative density with respect to τ if and only if there is some $\mathbf{c} \in \mathbf{M}$ such that

$\mathbf{v}(\cdot) = \langle \mathbf{c}, \cdot \rangle$. This situation implies that $\nu \dashv \tau$ and ν is τ -continuous by Theorem 5.8.

For the converse statement, assume that $\nu \dashv \tau$ and ν is τ -continuous. We only have to prove that $\mathbf{v}$ is non-degenerate, for then Theorem 5.8 gives the desired result. So let $f \in \mathcal{E}_+^\nu$ such that $\mathbf{v}(\mathbf{f}) \leq 1$, where $\mathbf{f} = \langle f \rangle$. Then, for all rational numbers $q > 0$, the subset $\{f > q\}$ is in the σ -ideal $\mathcal{I}_q = \{G \in \mathcal{E} : \nu(G) \leq q^{-1}\}$. Since τ is localizable, $\mathcal{I}_q$ is localized in some $L_q \in \mathcal{E}$, and since ν is τ -continuous, we have $\nu(L_q) \leq q^{-1}$. As a consequence, the map $g = \bigoplus_{q \in \mathbb{Q}_+} q \cdot 1_{L_q}$ is lsm, ν -integrable (with $\int^\infty g d\nu \leq 1$), and such that $\langle f \rangle \leq \langle g \rangle$. This proves that the subset $\{\mathbf{f} \in \mathbf{M} : 1 \geq \mathbf{v}(\mathbf{f})\}$ is upper-bounded in $\mathbf{M}$, i.e. that $\mathbf{v}$ is non-degenerate. $\square$

Corollary 6.4. *Let ν, τ be σ -maxitive measures on $\mathcal{E}$. Assume that τ is σ -principal. Then $\nu \dashv \tau$ if and only if ν has a usm relative density with respect to τ .*

Proof. We first suppose that ν is finite. If τ is σ -principal, then $\mathcal{E}_+^\nu/\tau$ is a σ -principal module by Proposition 6.2. By Proposition 6.1, ν is τ -continuous and τ is localizable. So Theorem 6.3 applies, and ν admits a usm relative density with respect to τ . In the case where ν is non-finite, we replace ν by $\nu_1 : B \mapsto \arctan \nu(B)$, which has a usm relative density c_1 with respect to τ . Therefore, $\tan c_1$ is a usm relative density of ν with respect to τ . $\square$

Example 6.5. Let E be a topological space, $\mathcal{E}$ be the collection $\mathcal{G}$ of open subsets of E , and $\tau = \delta_\#$. Then $\delta_\#$ is localizable and $\nu \dashv \delta_\#$, for all maxitive measures ν on $\mathcal{G}$. Moreover, ν is $\delta_\#$ -continuous if and only if ν is completely maxitive if and only if ν has a usc cardinal density. Also, $\delta_\#$ is σ -principal if and only if every subset of E is Lindelöf (then E is usually said to be *hereditarily Lindelöf*, a property that is implied by second-countability), in which case every ν on $\mathcal{G}$ has a cardinal density.

To conclude this section we propose a new proof of the Sugeno–Murofushi theorem, which is a Radon–Nikodym like theorem for the Shilkret integral (see [56, Theorem I-6.4]).

Theorem 6.6 (Sugeno–Murofushi). *Let ν, τ be σ -maxitive measures on a σ -algebra $\mathcal{B}$. Assume that τ is σ -finite and σ -principal. Then $\nu \dashv \tau$ if and only if there exists some $\mathcal{B}$ -measurable map $c : E \rightarrow \mathbb{R}_+$ such that*

$$\nu(B) = \int_B^\infty c d\tau,$$

for all $B \in \mathcal{B}$. If these conditions are satisfied, then c is unique τ -almost everywhere.

Proof. If $\nu \dashv \tau$, then by Corollary 6.4 there are $\mathcal{B}$ -measurable maps $c_1, c_2 : E \rightarrow \overline{\mathbb{R}}_+$ such that $\nu(B) = \bigoplus_{x \in B}^\tau c_1(x)$ and $\tau(B) = \bigoplus_{x \in B}^\tau c_2(x)$, for all $B \in \mathcal{B}$. Since τ is σ -finite, one can choose a map c_2 that takes only finite values (see [56, Proposition I-6.1]). Using the fact that $\tau(\{c_2 = 0\}) = 0$, it is easy to show that $\nu(\cdot) = \int^\infty c d\tau$ on $\mathcal{B}$, where c is the measurable map defined by $c(x) = c_1(x)/c_2(x)$ if $c_2(x) \neq 0$, $c(x) = 0$ otherwise. $\square$

7. COMPLETABLE MODULES AND THE NORMAL COMPLETION

7.1. The normal completion of a completable module. The following theorem defines the concept of *normal completion* of a completable module, alias *Dedekind–MacNeille completion* or *completion by cuts*. See e.g. Ern  [19] for the normal completion of quasiordered sets.

Theorem 7.1. *Let $\mathbb{k}$ be an idempotent semifield. A $\mathbb{k}$ -module is completable if and only if it can be continuously embedded into a complete $\mathbb{k}$ -module.*

Sketch of the proof. Sufficiency is obvious. For necessity, let M be a completable $\mathbb{k}$ -module. We follow the usual Dedekind–MacNeille completion method for partially ordered sets. If $X \subset M$, we write $X^\downarrow$ (resp. $X^\uparrow$) for the subset of lower (resp. upper) bounds of X in M , and we write $X^{\uparrow\downarrow}$ instead of $(X^\uparrow)^\downarrow$. A subset X of M is *closed* if $X^{\uparrow\downarrow} = X$, and *proper* if either $X \neq M$ or M has a greatest element. Let $\mathcal{N}(M)$ be the collection of all proper closed subsets X of M . If $X \oplus X' := (X \cup X')^{\uparrow\downarrow}$ for all proper closed subsets X, X' , then $X \oplus X'$ is closed, proper (to prove this, note that a closed subset is proper if and only if it is upper-bounded) and $(\mathcal{N}(M), \oplus, \{0\})$ is a commutative idempotent monoid. The partial order induced by $\oplus$ on $\mathcal{N}(M)$ is the inclusion, i.e. $X \leq X' \Leftrightarrow X \subset X'$. For the external multiplication we let $X.t := \{x.t : x \in X\}$ if $t \neq 0$ and $X.0 = \{0\}$, and one can check that $X.t$ is proper closed for all proper closed subsets X . Also, since M is completable, the following relations hold:

$$\begin{aligned} (X \oplus X').t &= X.t \oplus X'.t, \\ X. \bigoplus_{t \in T} T &= \bigoplus_{t \in T} X.t, \end{aligned}$$

for all $X, X' \in \mathcal{N}(M)$, $t \in \mathbb{k}$ and $T \subset \mathbb{k}$ with supremum, and

$$X. \bigwedge F = \bigwedge_{f \in F} X.f,$$

for all $X \in \mathcal{N}(M)$ and $\mathbf{F}$ -sets F in $\mathbb{k}$ with infimum. Thus, $\mathcal{N}(M)$ is a completable $\mathbb{k}$ -module, which is actually complete for

$$\bigoplus_{j \in J} X_j = (\bigcup_{j \in J} X_j)^{\uparrow\downarrow},$$

for all upper-bounded families $(X_j)_{j \in J}$ of proper closed subsets. Note that the infimum in $\mathcal{N}(M)$ satisfies

$$\bigwedge_{j \in J} X_j = \bigcap_{j \in J} X_j,$$

for all families $(X_j)_{j \in J}$ of proper closed subsets.

To embed M into $\mathcal{N}(M)$, let $i_M : M \rightarrow \mathcal{N}(M), x \mapsto \downarrow x$. This map i_M is well defined, for $\downarrow x$ is proper closed for all $x \in M$. Clearly, we have $i_M(x.t) = i_M(x).t$ and $i_M(x \oplus y) = i_M(x) \oplus i_M(y)$ for all $x, y \in M$ and $t \in \mathbb{k}$, so that i_M is an injective morphism. Moreover, for all subsets X of M with supremum (resp. with infimum), we have $i_M(\bigoplus X) = \bigoplus i_M(X)$ (resp. $i_M(\bigwedge X) = \bigwedge i_M(X)$), so that i_M is continuous. $\square$

Remark 7.2. If $\mathbf{F} = \mathbf{PFi}$, then a module is completable if and only if it is *b-regular* in the sense of Litvinov et al. [40, Definition 3.9].

Remark 7.3. Identifying M and $i_M(M)$, every element of $\mathcal{N}(M)$ can be expressed as a supremum (resp. an infimum) of elements of M . In particular, every element of $\mathcal{N}(M)$ is upper-bounded by some element of M .

Remark 7.4. Every idempotent semifield considered as a module over itself is completable. However, an idempotent semifield can be embedded into a complete idempotent semifield if and only if it is commutative (see Remark 3.2).

7.2. Cut-stability and extensions. In this paragraph we give two categorical results on the normal completion. Since every $\{0, 1\}$ -module is completable, they extend that of Ern  [19].

Ern  introduced the concept of cut-stability, that we modify as follows. A map $f : M \rightarrow N$ is *lower cut-stable* if

$$f(X^\uparrow)^\downarrow = f(X)^{\uparrow\downarrow},$$

for all subsets X of M , and *cut-stable* if it is lower cut-stable and such that

$$f(F^\downarrow)^\uparrow = f(F)^\downarrow{}^\uparrow,$$

for all $\mathbf{F}$ -sets F of M . For instance, the map i_M that embeds a completable module into its normal completion (see the proof of Theorem 7.1) is cut-stable. Note that every cut-stable morphism is continuous.

Proposition 7.5 (Compare with [19, Theorem 3.1]). *Let $\mathbb{k}$ be an idempotent semifield. A morphism f between completable $\mathbb{k}$ -modules M and N is lower cut-stable if and only if there exists a (unique) morphism $\mathcal{N}(f)$ between the normal completions $\mathcal{N}(M)$ and $\mathcal{N}(N)$ that preserves arbitrary existing suprema and extending f , i.e. such that the following diagram commutes:*

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ i_M \downarrow & & \downarrow i_N \\ \mathcal{N}(M) & \xrightarrow{\mathcal{N}(f)} & \mathcal{N}(N) \end{array}$$

Moreover, in the cases $\mathbf{F} = \mathbf{Up}^*$ and $\mathbf{F} = \mathbf{PFi}$, the morphism f is cut-stable if and only if $\mathcal{N}(f)$ is continuous, and the normal completion extends to a functor $\mathcal{N}$ on the category of completable $\mathbb{k}$ -modules with continuous morphisms.

Proof. Analogous to that of [19, Theorem 3.1]. □

Remark 7.6. In the case $\mathbf{F} = \mathbf{Fi}$, it is only possible to say that $\mathcal{N}(f)$ preserves infima of filters of $\mathcal{N}(M)$ of the form $\mathcal{F}_F = \{X \in \mathcal{N}(M) : X \cap F \neq \emptyset\}$, with F a filter of M . This is not enough to make $\mathcal{N}(f)$ smooth (i.e. continuous) in general.

The following universal property of the normal completion is deduced immediately.

Corollary 7.7 (Compare with [19, Corollary 3.2]). *Let $\mathbb{k}$ be an idempotent semifield. A morphism f from a completable $\mathbb{k}$ -module M into a complete $\mathbb{k}$ -module N is lower cut-stable if and only if there exists a (unique) morphism from $\mathcal{N}(M)$ into N preserving arbitrary existing suprema*

and extending f , i.e. such that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ i_M \downarrow & \nearrow \mathcal{N}(f) & \\ \mathcal{N}(M) & & \end{array}$$

Moreover, in the cases $\mathbf{F} = \mathbf{Up}^*$ and $\mathbf{F} = \mathbf{PFi}$, the morphism f is cut-stable if and only if $\mathcal{N}(f)$ is continuous.

We call a pair $\overline{M}/M$ an *extension* over $\mathbb{k}$ (we shall also speak of the extension $\overline{M}$ of M over $\mathbb{k}$) if $\overline{M}$ is a complete module over $\mathbb{k}$ and M is a submodule of $\overline{M}$. In this situation, M is necessarily completable. The extension is *short* if, for all $y \in \overline{M}$, there is some $x \in M$ such that $y \leq x$. This condition restricts the “size” of $\overline{M}$ and will reveal its importance in the next section. Also, the extension is *cut-stable* if the map $i : M \ni x \mapsto x \in \overline{M}$ is cut-stable; in this case, if a subset (resp. an $\mathbf{F}$ -set) of M has a supremum (resp. an infimum) in M , then it coincides with its supremum (resp. its infimum) in $\overline{M}$. For a completable module M , the normal completion leads to a short and cut-stable extension $\mathcal{N}(M)/i_M(M)$ (see Remark 7.3).

Example 7.8. Let E be a Hausdorff topological space, let C_c^+ be the set of compactly-supported continuous maps from E to $\mathbb{R}_+$, and let L^+ be the set of lower-semicontinuous maps from E to $\overline{\mathbb{R}}_+$. Then L^+/C_c^+ is an extension over $\mathbb{R}_+^{\max}$ that is neither short nor cut-stable in general. Figure 1 gives a sequence of continuous functions on $E = [0, 1]$, whose supremum is $x \mapsto 1$ in C_c^+ , and is $x \mapsto 1_{(0,1)}(x)$ in L^+ .

8. RESIDUATED FORMS ON A MODULE EXTENSION

This section is expressed in the language $\mathbf{PFi}$ of principal filters, and $\overline{M}/M$ is an extension over an idempotent semifield $\mathbb{k}$. Henceforth, all suprema of subsets of M or $\overline{M}$ are taken in $\overline{M}$. A map $v : M \rightarrow \mathbb{k}$ is *residuated on $\overline{M}/M$* if there exists a map $w : \mathbb{k} \rightarrow \overline{M}$ satisfying

$$(8.1) \quad x \leq w(t) \iff t \geq v(x),$$

for all $x \in M$, $t \in \mathbb{k}$. In this case, there exists a least map w such that Equivalence (8.1) holds, called the *adjoint* of v with respect to $\overline{M}/M$, denoted by $v^\#$, and defined by

$$v^\#(t) = \bigoplus \{x \in M : t \geq v(x)\},$$

for all $t \in \mathbb{k}$, where the supremum is taken in $\overline{M}$.




FIGURE 1. A nondecreasing sequence of continuous functions on $[0, 1]$.

Lemma 8.1. *A map that is residuated on a short extension of M is residuated on each extension of M .*

Proof. Let $\overline{M}/M$ be a short extension. Consider the following commutative diagram:

$$\begin{array}{ccc}
 M & \xrightarrow{i} & \overline{M} \\
 i_M \downarrow & \nearrow \bar{i} & \\
 \mathcal{N}(M) & &
 \end{array}$$

where $\bar{i}$ is defined by $\bar{i}(X) = \bigoplus X$. Let $v : M \rightarrow \mathbb{k}$ be a residuated map on $\overline{M}/M$. Then v admits an adjoint $v^\# : \mathbb{k} \rightarrow \overline{M}$. We first show that v is residuated on $\mathcal{N}(M)/M$. If $t \in \mathbb{k}$, the subset $I_t = \{x \in M : t \geq v(x)\}$ is upper-bounded (by $v^\#(t)$) in $\overline{M}$, hence also in M since $\overline{M}/M$ is short. Thus, I_t admits a supremum in $\mathcal{N}(M)$, that we denote by $w(t)$. Since $\bar{i}$ preserves arbitrary existing suprema, $\bar{i}(w(t)) = \bigoplus \{x \in M : t \geq v(x)\}$, where the supremum is taken in $\overline{M}$. Thus, $\bar{i} \circ w = v^\#$. We show that Equivalence (8.1) holds. Clearly, $t \geq v(x)$ implies $x \leq w(t)$. Conversely, assume that $x \leq w(t)$. Composing by $\bar{i}$, we get $x \leq v^\#(t)$, so that $t \geq v(x)$. This proves that v is residuated on $\mathcal{N}(M)/M$.

Now let $\tilde{M}/M$ be some extension of M , and consider the related commutative diagram:

$$\begin{array}{ccc} M & \xrightarrow{\quad} & \tilde{M} \\ i_M \downarrow & \nearrow \tilde{j} & \\ \mathcal{N}(M) & & \end{array}$$

where $j : M \ni x \mapsto x \in \tilde{M}$ and $\tilde{j} : X \mapsto \bigoplus X$. Then one can show that

$$x \leq \tilde{j} \circ w(t) \Leftrightarrow t \geq v(x),$$

for all $x \in M, t \in \mathbb{k}$, so v is residuated on $\tilde{M}/M$. $\square$

We define a *residuated form* on $\overline{M}/M$ as a homogeneous residuated map on $\overline{M}/M$. A map $v : M \rightarrow \mathbb{k}$ is *non-degenerate* on $\overline{M}/M$ if $\{x \in M : 1 \geq v(x)\}$ has an upper-bound in $\overline{M}$.

Lemma 8.2. *Let $\overline{M}/M$ be a short extension over a cis $\mathbb{k}$. We suppose that every element of $\overline{M}$ can be expressed as the supremum in $\overline{M}$ of elements of M . Then a map $v : M \rightarrow \mathbb{k}$ is a residuated form on $\overline{M}/M$ if and only if v is non-degenerate on $\overline{M}/M$ and extends to a residuated form on $\overline{M}$.*

$$\begin{array}{ccc} M & \xrightarrow{\quad} & \mathbb{k} \\ i \downarrow & \nearrow \bar{v} & \\ \overline{M} & & \end{array}$$

Proof. Let v be a residuated form on $\overline{M}/M$ with adjoint $v^\# : \mathbb{k} \rightarrow \overline{M}$. Then $\{x \in M : 1 \geq v(x)\}$ is upper-bounded (by $v^\#(1)$) in $\overline{M}$, so v is non-degenerate on $\overline{M}/M$. Moreover, if $y \in \overline{M}$, then y is upper-bounded by some $x \in M$ since $\overline{M}/M$ is short, so the subset $\{t \in \mathbb{k} : y \leq v^\#(t)\}$ of $\mathbb{k}$, which contains $t = v(x)$, is nonempty. Thus, we can define the map $\bar{v} : \overline{M} \rightarrow \mathbb{k}$ by $\bar{v}(y) = \bigwedge \{t \in \mathbb{k} : y \leq v^\#(t)\}$. Since every element of $\overline{M}$ can be expressed as the supremum in $\overline{M}$ of elements of M , we have

$$y \leq v^\#(t) \Leftrightarrow t \geq \bar{v}(y),$$

for all $y \in \overline{M}, t \in \mathbb{k}$. So we obtain that $\bar{v}$ is a residuated form on $\overline{M}$.

Conversely, assume that a map $v : M \rightarrow \mathbb{k}$ is non-degenerate on $\overline{M}/M$ and extends to a residuated form $\bar{v}$ on $\overline{M}$. If w is the adjoint of

$\bar{v}$ (with respect to $\bar{M}/\bar{M}$), then

$$x \leq w(t) \Leftrightarrow t \geq v(x),$$

for all $x \in M, t \in \mathbb{k}$, so v is a residuated form on $\bar{M}/M$. $\square$

With respect to the filter selection **PFi**, we say that an element $c \in \bar{M}$ is *archimedean in $\bar{M}/M$* if

- the subset $F_c(x)$ is nonempty for all $x \in M$,
- if the infimum of $F_c(x)$ is zero, then $F_c(x) \supset \mathbb{k} \setminus \{0\}$,

where $F_c(x)$ denotes the subset $\{t \in \mathbb{k} : c.t \geq x\}$. The next lemma can be proved along the same lines as Lemma 5.7.

Lemma 8.3. *Let $\bar{M}/M$ be an extension over a cis $\mathbb{k}$, and let $v : M \rightarrow \mathbb{k}$ be a residuated form on $\bar{M}/M$. Then the supremum of $\{x \in M : 1 \geq v(x)\}$ is archimedean in $\bar{M}/M$.*

The innovation of this section mainly relies on highlighting the role of the following concept in the representation of continuous linear forms. An extension $\bar{M}/M$ is *meet-continuous* if

$$x \wedge \bigoplus I = \bigoplus \downarrow x \cap I,$$

for all $x \in M$ and all ideals I of M with an upper-bound in $\bar{M}$.

Example 8.4 (Example 7.8 continued). Let E be a topological space. We still denote by L^+ the set of lower-semicontinuous maps from E to $\mathbb{R}_+$. If M is some submodule of the set of continuous maps from E to $\mathbb{R}_+$, then the extension L^+/M is meet-continuous. Before proving this assertion, note that the supremum in L^+ coincides with the pointwise supremum. Now let $f \in M$ and let I be an ideal in M . We want to show that $f \wedge \bigoplus I \leq \bigoplus \downarrow f \cap I$, i.e. that $f(x) \wedge \bigoplus_{g \in I} g(x) \leq \bigoplus_{h \in I, h \leq f} h(x)$. For this purpose, let $s < f(x) \wedge \bigoplus_{g \in I} g(x)$. There is some $g \in I$ such that $s < f(x) \wedge g(x)$. Then the map $h = f \wedge g$ is continuous, is in I and satisfies $h \leq f$ and $s < h(x)$, so the claim follows.

A map $f : M \rightarrow \mathbb{k}$ is *continuous* on $\bar{M}/M$ if, for every subset $X \subset M$ such that $\bigoplus X \in M$, $f(X)$ has a supremum in $\mathbb{k}$ satisfying $f(\bigoplus X) = \bigoplus f(X)$. If $x \in M$ and $c \in \bar{M}$, we write $\langle c, x \rangle$ for the infimum of $\{t \in \mathbb{k} : c.t \geq x\}$ whenever this set is nonempty, and $\langle c, x \rangle = \top$ otherwise. The following theorem shows that, under conditions different than those of Theorem 5.8, the representation $v(\cdot) = \langle c, \cdot \rangle$ still holds, at the price that c no longer needs to belong to M . But it is actually important to authorize c to be outside M , in order to encompass the (idempotent) Riesz representation theorem (see Theorem 9.3 below).

Theorem 8.5. *Suppose that $\overline{M}/M$ is a meet-continuous extension over a cisl $\mathbb{k}$, and let $v : M \rightarrow \mathbb{k}$ be a linear form on M . Then the following conditions are equivalent:*

- (1) *v is residuated on $\overline{M}/M$,*
- (2) *v is non-degenerate continuous on $\overline{M}/M$,*
- (3) *$v(\cdot) = \langle c, \cdot \rangle$, for some archimedean element c in $\overline{M}/M$.*

If any of these conditions is satisfied, then the supremum of $\{1 \geq v\}$ in $\overline{M}$ is the least c satisfying $v(\cdot) = \langle c, \cdot \rangle$.

Proof. Assume that v is non-degenerate and continuous. Let $t \in \mathbb{k}$ and $I_t = \{x \in M : t \geq v(x)\}$. Clearly I_t is an ideal, let $w(t)$ denote its supremum in $\overline{M}$. To prove that v is residuated on $\overline{M}/M$, we have to show that, if $x \in M$ and $x \leq w(t)$, then $t \geq v(x)$, i.e. $x \in I_t$. If $\overline{M}/M$ is meet-continuous, then $x = x \wedge w(t) = x \wedge \bigoplus I_t = \bigoplus \downarrow x \cap I_t \in M$, and using the fact that v is continuous we get $v(x) = \bigoplus_{y \in \downarrow x \cap I_t} v(y) \leq t$, i.e. $x \in I_t$. This proves that (2) implies (1), and the converse implication is clear. The equivalence between (1) and (3) can be proved along the same lines as in the proof of Theorem 5.8. $\square$

Remark 8.6. If every element of $\overline{M}$ can be expressed as the supremum in $\overline{M}$ of elements of M , there is a *unique* such element c in the previous theorem. Indeed, assume that, for some archimedean elements b, c in $\overline{M}/M$, we have $\langle b, x \rangle = \langle c, x \rangle$ for all $x \in M$. Then $x \leq b \Leftrightarrow 1 \geq \langle b, x \rangle \Leftrightarrow 1 \geq \langle c, x \rangle \Leftrightarrow x \leq c$, for all $x \in M$, so that $\downarrow b \cap M = \downarrow c \cap M$. This implies that $b = \bigoplus \downarrow b \cap M = \bigoplus \downarrow c \cap M = c$.

As a direct application, we shall reprove an idempotent version of the Riesz representation theorem in Section 9.

9. THE RIESZ REPRESENTATION THEOREM

In this section, we aim at proving Riesz representation theorems for the Shilkret integral with the help of Theorem 8.5. The filter selection implicitly used here is **PFi**, i.e. the one that selects principal filters.

Let E be a Hausdorff topological space, $\mathcal{G}$ (resp. $\mathcal{B}$) be the collection of open subsets (resp. Borel subsets) of E , and C_c^+ be the set of nonnegative compactly-supported continuous maps from E to $\mathbb{R}_+$. The next theorem is of historical importance, for (part of) it was originally stated by Choquet [15, Paragraph 53.1] without proof; the first proof is due to Kolokoltsov and Maslov [36, Theorem 1]. The reader can also refer to Puhalskii [59, Theorem 1.7.21].

Lemma 9.1 (Urysohn). *Let E be a locally-compact Hausdorff space. If $K \subset U \subset E$ with K compact and U open, then there exists a compactly-supported continuous map $f : E \rightarrow [0, 1]$ such that $f(x) = 1$ for all $x \in K$ and $\overline{\{x \in E : f(x) > 0\}} \subset U$.*

Proof. This is customarily proved by using the fact that the one-point compactification of E is a normal space, see e.g. Aliprantis and Border [4, Corollary 2.74]. $\square$

Lemma 9.2. *Let E be a locally-compact Hausdorff space. Every lower-semicontinuous map $g : E \rightarrow \overline{\mathbb{R}}_+$ is a supremum of elements of C_c^+ .*

Proof. Let $s \in \mathbb{R}_+$ and $x \in E$ be such that $s < g(x)$. Since $\{g > s\}$ is an open subset, there is some $f_1 \in C_c^+$ such that $s < f_1(x) < g(x)$ and $f_1 = 0$ on $\{g \leq s\}$, by Urysohn's lemma. Now $\{g > f_1\}$ is also open, so there is some $f_2 \in C_c^+$ such that $f_2(x) = f_1(x)$ and $f_2 = 0$ on $\{g \leq f_1\}$. This proves that the map $f_{s,x} = f_1 \wedge f_2$ is in C_c^+ and satisfies $f_{s,x} \leq g$ and $s < f_{s,x}(x)$. As a consequence, one can see that g is the pointwise supremum of all such maps $f_{s,x}$. $\square$

Theorem 9.3 (Improves [59, Theorem 1.7.21]). *Let E be a locally-compact Hausdorff space, and let $V : C_c^+ \rightarrow \mathbb{R}_+$ be a linear form on C_c^+ . Then there exists a unique regular maxitive measure ν on $\mathcal{B}$ such that*

$$V(f) = \int^\infty f \, d\nu,$$

for all $f \in C_c^+$. Moreover, ν takes finite values on compact subsets of E .

Proof. The functional V is a linear form on the $\mathbb{R}_+^{\max}$ -module C_c^+ . If $\overline{M} = L^+$ is the module of $\overline{\mathbb{R}}_+$ -valued lower-semicontinuous maps and $M = C_c^+$, then $\overline{M}/M$ is a meet-continuous extension by Example 8.4, so Theorem 8.5 applies if we show that V is continuous and non-degenerate on $\overline{M}/M$. Non-degeneracy of V on $\overline{M}/M$ is ensured by the existence of arbitrary suprema in $\overline{M}$. For the continuity of V , let $(f_j)_{j \in J}$ be a nondecreasing net of elements of C_c^+ such that $f := \bigoplus_{j \in J} f_j \in M$, where the supremum is taken in $\overline{M}$. In particular, $(f_j)_{j \in J}$ converges pointwise to f , see Example 8.4. We want to prove that $V(f) = \bigoplus_{j \in J} V(f_j)$. So let $1 > \varepsilon > 0$, let K_ε be the compact set $\{f \geq \varepsilon\}$, and define h_j on K_ε by $h_j(x) = f_j(x)/f(x)$. Then $h_j \in C_c^+(K_\varepsilon)$ and $(h_j)_{j \in J}$ is a nondecreasing net converging to 1 pointwise. Applying Dini's Theorem, the convergence is uniform on K_ε , hence there is some $j_0 \in J$ such that $1 \leq \varepsilon + h_{j_0}$ on K_ε . Thus, $f \leq \varepsilon \cdot f + f_{j_0}$ on K_ε . Let K be a compact set containing $\{0 < f < 1\}$. By Urysohn's

lemma, we may find a compactly-supported continuous map h such that $h = 1$ on K . Then $f \leq (\frac{1}{1-\varepsilon}f_{j_0}) \oplus (\varepsilon h)$ on E . This implies $V(f) \leq (\frac{1}{1-\varepsilon} \bigoplus_{j \in J} V(f_j)) \oplus (\varepsilon V(h))$, for all $1 > \varepsilon > 0$, so V is continuous. By Theorem 8.5, there exists some archimedean element c in $\overline{M}/M$ such that $V(f) = \langle c, f \rangle$, for all $f \in M$. Defining the usc map c^+ by $c^+(x) = 1/c(x)$, we have $V(f) = \bigoplus_{x \in E} f(x).c^+(x)$, for all $f \in M$. The maxitive measure ν defined on $\mathcal{B}$ by $\nu(B) = \bigoplus_{x \in B} c^+(x)$ for all $B \in \mathcal{B}$ is regular by [57, Theorem 3.22], and we have

$$V(f) = \int^\infty f \, d\nu,$$

for all $f \in M$. If K is a compact subset of E , Urysohn's lemma provides a map $f \in M$ such that $f(x) = 1$ for all $x \in K$, so that $c^+(x) \leq V(f)$ for all $x \in K$. This ensures that $\nu(K) = \bigoplus_{x \in K} c^+(x)$ is finite.

Uniqueness of ν is a direct consequence of the uniqueness of c , which itself derives from Lemma 9.2 and Remark 8.6. $\square$

In the same line, one can formulate Riesz like theorems for a functional V defined on the set C_b^+ of nonnegative bounded continuous maps instead of C_c^+ . Breyer and Gulinsky [12] proved the next theorem¹, see also Puhalskii [59, Theorem 1.7.25] and Gulinsky [30, Theorem 3.4]. A functional $V : C_b^+ \rightarrow \mathbb{R}_+$ is *tight* if, for all $\varepsilon > 0$, there is some compact subset K of E such that $V(f) \leq \varepsilon \|f\|$, for each $f \in C_b^+$ that equals 0 on K .

Theorem 9.4. [12] *Assume that E is a Tychonoff space, and let $V : C_b^+ \rightarrow \mathbb{R}_+$ be a tight linear form on C_b^+ that preserves countable pointwise suprema. Then there exists a unique (finite) tight regular maxitive measure ν on $\mathcal{B}$ such that*

$$V(f) = \int^\infty f \, d\nu,$$

for all $f \in C_b^+$.

Proof. See e.g. Puhalskii [59, Theorem 1.7.25]. The idea of the proof is to use the Stone-Ćech compactification of E and to apply Theorem 9.3. $\square$

In order to treat the case of a non-tight linear form on C_b^+ , we shall assume that the Tychonoff space is also second-countable, i.e. is a separable metrizable space. A functional $V : C_b^+ \rightarrow \mathbb{R}_+$ is *optimal* if, for all nonincreasing sequences $(f_n)_{n \in \mathbb{N}}$ of elements of C_b^+ tending pointwise to 0, the sequence $(V(f_n))_{n \in \mathbb{N}}$ tends to 0.

¹We were not in a position to access this article.

Theorem 9.5. *Assume that E is a separable metrizable space, and let $V : C_b^+ \rightarrow \mathbb{R}_+$ be a linear form on C_b^+ that preserves countable pointwise suprema. Then there exists a unique (finite) regular maxitive measure ν on $\mathcal{B}$ such that*

$$V(f) = \int^\infty f \, d\nu,$$

for all $f \in C_b^+$. Moreover, if E is Polish, then the following conditions are equivalent:

- V is optimal,
- V is tight,
- ν is tight,
- c^+ is upper-compact,

where c^+ is the maximal cardinal density of ν .

Proof. We denote L^+ by $\overline{M}$ and C_b^+ by M . As in the proof of Theorem 9.3, $\overline{M}/M$ is a meet-continuous extension by Example 8.4, V is a non-degenerate linear form on $\overline{M}/M$, and we want to prove that

$$(9.1) \quad V(f) = \bigoplus_{j \in J} V(f_j),$$

for all nondecreasing nets $(f_j)_{j \in J}$ in M such that $f := \bigoplus_{j \in J} f_j \in M$, where the supremum is taken in $\overline{M}$. Let q be a nonnegative rational number. The open subset $\{f > q\}$ is covered by the family of open subsets $\{f_j > q\}$, $j \in J$. Since E is separable metrizable, it is second-countable, so we can extract a countable subcover and write $\{f > q\} = \bigcup_{j \in N_q} \{f_j > q\}$, where N_q is a countable subset of J . Defining N as the union of all N_q , which is countable, we see that $f = \bigoplus_{j \in N} f_j$. Since V preserves countable pointwise suprema in M , Equation (9.1) holds, so V is continuous on $\overline{M}/M$. The existence of ν now follows from the same argument as in the proof of Theorem 9.3. Since E is separable metrizable, E is normal; using Urysohn's lemma for normal spaces (see e.g. [4, Theorem 2.46]), we can show that every $\mathbb{R}_+$ -valued lower-semi-continuous map is a supremum of elements of C_b^+ , as a perfect analogue of Lemma 9.2, and this leads to the uniqueness of ν .

Now suppose that E is Polish. Assume that $\{c^+ \geq t\}$ is not compact, for some $t > 0$. Then there exists some $\varepsilon > 0$ and some sequence (x_n) of elements of $\{c^+ \geq t\}$ such that $d(x_m, x_n) > \varepsilon$, for all $m \neq n$. Since E is Polish, one can find some countable family of open balls with radius $\varepsilon/2$ covering E . Let $B_k \ni x_k$ be one of these balls containing x_k . Let $f_k \in C_b^+$ such that $f_k(x_k) = 1/t$ and $f_k = 0$ on $E \setminus B_k$. Then

$g_n := \bigoplus_{k \geq n} f_k$ tends pointwise to 0. But $V(g_n) \geq g_n(x_n) \cdot c^+(x_n) \geq 1$, so V is not optimal.

Conversely, assume that c^+ is upper-compact and that V is not optimal. So let (f_n) be a nonincreasing sequence of elements of C_b^+ that tends pointwise to zero, and assume that $\bigwedge_{n \in \mathbb{N}} V(f_n) > 0$. Then there exists some $t > 0$ such that $V(f_n) > t$, for all n . We deduce the existence of some $x_n \in E$ such that $f_n(x_n) \cdot c^+(x_n) > t$. Since the sequence (f_n) is nonincreasing, $f_n \leq f_0$. Also, f_0 is upper-bounded by some $u > 0$, so that x_n is in the compact subset $\{c^+ \geq t/u\}$. This implies that (x_n) clusters to some x , but this contradicts the fact that $f_m(x_n) \geq f_m(x_n) \cdot c^+(x_n) \geq f_n(x_n) \cdot c^+(x_n) > t$ for all $n \geq m$.

By [57, Theorem 3.25], we know that ν is tight if and only if c^+ is upper-compact. If V is tight, then ν is tight by uniqueness of ν in Theorem 9.4; the converse statement is obvious. $\square$

Bell and Bryc also investigated the case where E is Polish, their result [10, Theorem 2.1] is encompassed in the previous theorem.

Akian proved a slightly different result for normal spaces that we merely recall for the sake of completeness.

Theorem 9.6. [2, Theorem 4.8] *Assume that E is a normal space, and let $V : C_b^+ \rightarrow \mathbb{R}_+$ be a linear form on C_b^+ that preserves countable pointwise suprema. Then there exists a unique σ -maxitive measure ν on $\mathcal{B}$ such that*

$$V(f) = \int^\infty f \, d\nu,$$

for all $f \in C_b^+$.

Remark 9.7 (On large deviations). The idempotent Riesz representation theorem partly originates from large deviation questionings. Varadhan [69] was interested in the functional defined on the set $C_b^+(E)$ of nonnegative bounded continuous maps, for some Polish space E , by

$$V(f) = \lim_{n \rightarrow \infty} \left(\int_E f^{1/\alpha_n} \, d\mu_n \right)^{\alpha_n},$$

whenever the limit exists, where (μ_n) is a sequence of probability measures on E satisfying a large deviation principle, and $\alpha_n \downarrow 0$ when $n \uparrow \infty$. He proved the representation

$$(9.2) \quad V(f) = \bigoplus_{x \in E} (f(x) e^{-I(x)}),$$

where $I : E \rightarrow \overline{\mathbb{R}}_+$ is the (lower-semicontinuous) *rate function*, that governs large deviations. For more on the links between large deviation

principles and maxitive measures, we refer the reader to Puhalskii [58, 59], B. Gerritse [25], Akian [2], Nedović et al. [50].

10. CONCLUSION AND PERSPECTIVES

Following Cohen et al. [16], we could certainly have pushed on the generalization to the use of *reflexive* idempotent semirings² instead of idempotent semifields, but our main interest here, at least in the first part of the paper, was to stress the role of $\mathbf{Z}$ theory in the gathering of similar but a priori distinct results from different mathematical areas.

Some results on modules and continuous linear forms are of topological flavour; this aspect will be sharpened in a future work with the examination of topological $\mathbb{R}_+^{\max}$ -modules.

Acknowledgments. I would like to thank Marc Leandri for his advice on a preliminary version of the manuscript. I am also indebted in Pr. Jimmie D. Lawson who made numerous comments and pointed out a mistake in an ealier version of Section 7.

REFERENCES

- [1] Samson Abramsky and Steve J. Vickers. Quantales, observational logic and process semantics. *Math. Structures Comput. Sci.*, 3(2):161–227, 1993.
- [2] Marianne Akian. Densities of idempotent measures and large deviations. *Trans. Amer. Math. Soc.*, 351(11):4515–4543, 1999.
- [3] Marianne Akian and Ivan Singer. Topologies on lattice ordered groups, separation from closed downward sets and conjugations of type Lau. *Optimization*, 52(6):629–672, 2003.
- [4] Charalambos D. Aliprantis and Kim C. Border. *Infinite dimensional analysis*. Springer, Berlin, third edition, 2006. A hitchhiker’s guide.
- [5] François L. Baccelli, Guy Cohen, Geert Jan Olsder, and Jean-Pierre Quadrat. *Synchronization and linearity*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons Ltd., Chichester, 1992. An algebra for discrete event systems.
- [6] Hans-J. Bandelt. M -distributive lattices. *Arch. Math. (Basel)*, 39(5):436–442, 1982.
- [7] Hans-J. Bandelt and Marcel Ern . The category of $\mathbf{Z}$ -continuous posets. *J. Pure Appl. Algebra*, 30(3):219–226, 1983.
- [8] Hans-J. Bandelt and Marcel Ern . Representations and embeddings of M -distributive lattices. *Houston J. Math.*, 10(3):315–324, 1984.
- [9] Andrei Baranga. $\mathbf{Z}$ -continuous posets. *Discrete Math.*, 152(1-3):33–45, 1996.
- [10] Harold Bell and Wlodzimierz Bryc. Variational representations of Varadhan functionals. *Proc. Amer. Math. Soc.*, 129(7):2119–2125 (electronic), 2001.

²Or even to the use of idempotent semirings in which every element is the supremum of reflexive elements, so as to generalize both Cohen et al.’s and Litvinov et al.’s approaches.

- [11] Garrett Birkhoff. *Lattice theory*. American Mathematical Society Colloquium Publications, vol. 25, third edition. American Mathematical Society, Providence, RI, 1967.
- [12] V. Breyer and Oleg V. Gulinsky. Large deviations on infinite dimensional spaces (in Russian). Technical report, Moscow Institute of Physics and Technology, Russia, 1996.
- [13] Günter Bruns. A lemma on directed sets and chains. *Arch. Math. (Basel)*, 18:561–563, 1967.
- [14] Dominique Castella. L’algèbre tropicale comme algèbre de la caractéristique 1 : Polynômes rationnels et fonctions polynômiales; arXiv:0809.0231v1. 2008.
- [15] Gustave Choquet. Theory of capacities. *Ann. Inst. Fourier, Grenoble*, 5:131–295, 1953–1954.
- [16] Guy Cohen, Stéphane Gaubert, and Jean-Pierre Quadrat. Duality and separation theorems in idempotent semimodules. *Linear Algebra Appl.*, 379:395–422, 2004. Tenth Conference of the International Linear Algebra Society.
- [17] Gert de Cooman, Guangquan Zhang, and Etienne E. Kerre. Possibility measures and possibility integrals defined on a complete lattice. *Fuzzy Sets and Systems*, 120(3):459–467, 2001.
- [18] Marcel Ern . A completion-invariant extension of the concept of continuous lattices. In *Continuous lattices, Proceedings of the Conference on Topological and Categorical Aspects of Continuous Lattices (Workshop IV) held at the University of Bremen, Bremen, November 9–11, 1979*, volume 871 of *Lecture Notes in Mathematics*, pages 45–60, Berlin, 1981. Springer-Verlag. Edited by Bernhard Banaschewski and Rudolf-E. Hoffmann.
- [19] Marcel Ern . The Dedekind–MacNeille completion as a reflector. *Order*, 8(2):159–173, 1991.
- [20] Marcel Ern . Z -continuous posets and their topological manifestation. *Appl. Categ. Structures*, 7(1-2):31–70, 1999. Applications of ordered sets in computer science (Braunschweig, 1996).
- [21] Marcel Ern , Mai Gehrke, and Al s Pultr. Complete congruences on topologies and down-set lattices. *Appl. Categ. Structures*, 15(1-2):163–184, 2007.
- [22] Marcel Ern , J rgen Koslowski, Austin Melton, and George E. Strecker. A primer on Galois connections. In *Papers on general topology and applications (Madison, WI, 1991)*, volume 704, pages 103–125, 1993.
- [23] Marcel Ern  and Dongsheng Zhao. Z -join spectra of Z -supercompactly generated lattices. *Appl. Categ. Structures*, 9(1):41–63, 2001.
- [24] Orrin Frink. Ideals in partially ordered sets. *Amer. Math. Monthly*, 61:223–234, 1954.
- [25] Bart Gerritse. Varadhan’s theorem for capacities. *Comment. Math. Univ. Carolin.*, 37(4):667–690, 1996.
- [26] Gerard Gerritse. Lattice-valued semicontinuous functions. In *Probability and lattices*, volume 110 of *CWI Tract*, pages 93–125. Math. Centrum Centrum Wisk. Inform., Amsterdam, 1997.
- [27] Gerhard Gierz, Karl Heinrich Hofmann, Klaus Keimel, Jimmie D. Lawson, Michael W. Mislove, and Dana S. Scott. *Continuous lattices and domains*, volume 93 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2003.

- [28] Michel Gondran and Michel Minoux. *Graphs, dioids and semirings*, volume 41 of *Operations Research/Computer Science Interfaces Series*. Springer, New York, 2008. New models and algorithms.
- [29] Gabriele H. Greco. Fuzzy integrals and fuzzy measures with their values in complete lattices. *J. Math. Anal. Appl.*, 126(2):594–603, 1987.
- [30] Oleg V. Gulinsky. The principle of the largest terms and quantum large deviations. *Kybernetika (Prague)*, 39(2):229–247, 2003. Special issue on max-plus algebras (Prague, 2001).
- [31] Reinhold Heckmann and Michael Huth. A duality theory for quantitative semantics. In *Computer science logic (Aarhus, 1997)*, volume 1414 of *Lecture Notes in Comput. Sci.*, pages 255–274. Springer, Berlin, 1998.
- [32] Reinhold Heckmann and Michael Huth. Quantitative semantics, topology, and possibility measures. *Topology Appl.*, 89(1-2):151–178, 1998. Domain theory.
- [33] Michael Huth. A powerdomain of possibility measures. *Electronic Notes in Theoretical Computer Science*, 6:192–202, 1997.
- [34] T. Iwamura. A lemma on directed sets. *Zenkoku Shijo Sugaku Danwakai*, 262:107–111, 1944. In Japanese.
- [35] Johan Jonasson. On positive random objects. *J. Theoret. Probab.*, 11(1):81–125, 1998.
- [36] Vassili N. Kolokoltsov and Victor P. Maslov. The general form of endomorphisms in the space of continuous functions with values in a numerical commutative semiring (with the operation $\oplus = \max$). *Dokl. Akad. Nauk SSSR*, 295(2):283–287, 1987.
- [37] Ivan Kramosil. Generalizations and extensions of lattice-valued possibilistic measures, part I. Technical Report 952, Institute of Computer Science, Academy of Sciences of the Czech Republic, 2005.
- [38] David Kruml. Spatial quantales. *Appl. Categ. Structures*, 10(1):49–62, 2002.
- [39] Jimmie D. Lawson. Idempotent analysis and continuous semilattices. *Theoret. Comput. Sci.*, 316(1-3):75–87, 2004.
- [40] Grigory L. Litvinov, Victor P. Maslov, and Grigori B. Shpiz. Idempotent functional analysis. An algebraic approach. *Mat. Zametki*, 69(5):758–797, 2001.
- [41] Xue Cheng Liu and Guangquan Zhang. Lattice-valued fuzzy measure and lattice-valued fuzzy integral. *Fuzzy Sets and Systems*, 62(3):319–332, 1994.
- [42] George Markowsky. Chain-complete posets and directed sets with applications. *Algebra Universalis*, 6(1):53–68, 1976.
- [43] George Markowsky. Categories of chain-complete posets. *Theoret. Comput. Sci.*, 4(2):125–135, 1977.
- [44] George Markowsky. A motivation and generalization of Scott’s notion of a continuous lattice. In *Continuous lattices, Proceedings of the Conference on Topological and Categorical Aspects of Continuous Lattices (Workshop IV) held at the University of Bremen, Bremen, November 9–11, 1979*, volume 871 of *Lecture Notes in Mathematics*, pages 298–307, Berlin, 1981. Springer-Verlag. Edited by Bernhard Banaschewski and Rudolf-E. Hoffmann.
- [45] George Markowsky. Propaedeutic to chain-complete posets with basis. In *Continuous lattices, Proceedings of the Conference on Topological and Categorical*

- Aspects of Continuous Lattices (Workshop IV) held at the University of Bremen, Bremen, November 9–11, 1979*, volume 871 of *Lecture Notes in Mathematics*, pages 308–314, Berlin, 1981. Springer-Verlag. Edited by Bernhard Banaschewski and Rudolf-E. Hoffmann.
- [46] George Markowsky and Barry K. Rosen. Bases for chain-complete posets. *IBM J. of Res. and Development*, 20:138–147, 1976.
 - [47] Jorge Martinez. Unique factorization in partially ordered sets. *Proc. Amer. Math. Soc.*, 33:213–220, 1972.
 - [48] Victor P. Maslov. *Méthodes opératorielles*. Éditions Mir, Moscow, 1987. Translated from the Russian by Djilali Embarek.
 - [49] Venu G. Menon. A note on topology of Z -continuous posets. *Comment. Math. Univ. Carolin.*, 37(4):821–824, 1996.
 - [50] Ljubo Nedović, Nebojša M. Ralević, and Tatjana Grbić. Large deviation principle with generated pseudo measures. *Fuzzy Sets and Systems*, 155(1):65–76, 2005.
 - [51] Evelyn Nelson. Z -continuous algebras. In *Continuous lattices, Proceedings of the Conference on Topological and Categorical Aspects of Continuous Lattices (Workshop IV) held at the University of Bremen, Bremen, November 9–11, 1979*, volume 871 of *Lecture Notes in Mathematics*, pages 315–334, Berlin, 1981. Springer-Verlag. Edited by Bernhard Banaschewski and Rudolf-E. Hoffmann.
 - [52] Hung T. Nguyen, Nhu T. Nguyen, and Tonghui Wang. On capacity functionals in interval probabilities. *Internat. J. Uncertain. Fuzziness Knowledge-Based Systems*, 5(3):359–377, 1997.
 - [53] Tommy Norberg. Existence theorems for measures on continuous posets, with applications to random set theory. *Math. Scand.*, 64(1):15–51, 1989.
 - [54] Tommy Norberg. On the convergence of probability measures on continuous posets. In *Probability and lattices*, volume 110 of *CWI Tract*, pages 57–92. Math. Centrum Centrum Wisk. Inform., Amsterdam, 1997.
 - [55] Dan Novak. On a duality between the concepts “finite” and “directed”. *Houston J. Math.*, 8(4):545–563, 1982.
 - [56] Paul Poncet. *Infinite-dimensional idempotent analysis: the role of continuous posets*. PhD thesis, École Polytechnique, Palaiseau, France, 2011.
 - [57] Paul Poncet. How regular can maxitive measures be? *Submitted*, 2012.
 - [58] Anatolii A. Puhalskii. Large deviations of semimartingales via convergence of the predictable characteristics. *Stochastics Stochastics Rep.*, 49(1-2):27–85, 1994.
 - [59] Anatolii A. Puhalskii. *Large deviations and idempotent probability*, volume 119 of *Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics*. Chapman & Hall/CRC, Boca Raton, FL, 2001.
 - [60] George N. Raney. Completely distributive complete lattices. *Proc. Amer. Math. Soc.*, 3:677–680, 1952.
 - [61] George N. Raney. A subdirect-union representation for completely distributive complete lattices. *Proc. Amer. Math. Soc.*, 4:518–522, 1953.
 - [62] Kimmo I. Rosenthal. Modules over a quantale and models for the operator ! in linear logic. *Cahiers Topologie Géom. Différentielle Catég.*, 35(4):329–333, 1994.

- [63] Ciro Russo. Quantale modules and their operators, with applications. *J. Logic Comput.*, 20(4):917–946, 2010.
- [64] Serguei N. Samborskiĭ and Grigori B. Shpiz. Convex sets in the semimodule of bounded functions. volume 13 of *Lecture Notes in Control and Information Sciences*, pages 135–138. American Mathematical Society, Providence, RI, 1992. Edited by Victor P. Maslov and Serguei N. Samborskiĭ.
- [65] G.-B. Shi and G.-P. Wang. Z -mappings and a classification theorem. *Semigroup Forum*, 52(3):349–355, 1996.
- [66] Niel Shilkret. Maxitive measure and integration. *Nederl. Akad. Wetensch. Proc. Ser. A 74 = Indag. Math.*, 33:109–116, 1971.
- [67] Grigori B. Shpiz. An eigenvector existence theorem in idempotent analysis. *Mat. Zametki*, 82(3):459–468, 2007.
- [68] Michio Sugeno and Toshiaki Murofushi. Pseudo-additive measures and integrals. *J. Math. Anal. Appl.*, 122(1):197–222, 1987.
- [69] S. R. Srinivasa Varadhan. Asymptotic probabilities and differential equations. *Comm. Pure Appl. Math.*, 19:261–286, 1966.
- [70] Poothampilly Venugopalan. Z -continuous posets. *Houston J. Math.*, 12(2):275–294, 1986.
- [71] Poothampilly Venugopalan. Union complete subset systems. *Houston J. Math.*, 14(4):583–600, 1988.
- [72] Jesse B. Wright, Eric G. Wagner, and James W. Thatcher. A uniform approach to inductive posets and inductive closure. *Theoret. Comput. Sci.*, 7(1):57–77, 1978.
- [73] Xiao-Quan Xu. Construction of homomorphisms of M -continuous lattices. *Trans. Amer. Math. Soc.*, 347(8):3167–3175, 1995.
- [74] Karel Zimmermann. A general separation theorem in extremal algebras. *Ekonom.-Mat. Obzor*, 13(2):179–201, 1977.

CMAP, ÉCOLE POLYTECHNIQUE, ROUTE DE SACLAY, 91128 PALAISEAU CEDEX,
 FRANCE, AND INRIA, SACLAY-ÎLE-DE-FRANCE
E-mail address: poncet@cmap.polytechnique.fr