

Deformations of Poisson structures by closed 3-forms¹

O. I. Mokhov

Abstract

We prove that an arbitrary Poisson structure $\omega^{ij}(u)$ and an arbitrary closed 3-form $T_{ijk}(u)$ generate the local Poisson structure $A^{ij}(u, u_x) = M_s^i(u, u_x)\omega^{sj}(u)$, where $M_s^i(u, u_x)(\delta_j^s + \omega^{sp}(u)T_{pjk}(u)u_x^k) = \delta_j^i$, on the corresponding loop space. We obtain also a special graded ε -deformation of an arbitrary Poisson structure $\omega^{ij}(u)$ by means of an arbitrary closed 3-form $T_{ijk}(u)$.

In this paper we prove that an arbitrary Poisson structure $\omega^{ij}(u)$ and an arbitrary closed 3-form $T_{ijk}(u)$ generate the local Poisson structure

$$A^{ij}(u, u_x) = B_s^i(u, u_x)\omega^{sj}(u), \quad (1)$$

where

$$B_s^i(u, u_x)M_j^s(u, u_x) = \delta_j^i, \quad M_j^s(u, u_x) = \delta_j^s + \omega^{sp}(u)T_{pjk}(u)u_x^k, \quad (2)$$

i.e., the matrix operator $A^{ij}(u, u_x)$ gives the Poisson bracket

$$\{I, J\} = \int \frac{\delta I}{\delta u^i(x)} A^{ij}(u, u_x) \frac{\delta J}{\delta u^j(x)} dx \quad (3)$$

on the space of functionals on the corresponding loop space.

Let M^N be an arbitrary smooth N -dimensional manifold with the local coordinates $u = (u^1, \dots, u^N)$. By the *loop space* ΩM of the manifold M^N we mean, in this paper, the space of all smooth parametrized mappings of the circle S^1 into M^N , $\gamma : S^1 \rightarrow M^N$, $\gamma(x) = \{u^i(x)\}$, $x \in S^1$. The tangent space $T_\gamma \Omega M$ of the loop space ΩM at the point γ consist of all smooth vector fields $\xi = \{\xi^i, 1 \leq i \leq N\}$, defined along the loop γ with $\xi(\gamma(x)) \in T_{\gamma(x)} M$, $\forall x \in S^1$, where $T_{\gamma(x)} M$ is a tangent space of the manifold M at the

¹The work was supported by the Max-Planck-Institut für Mathematik (Bonn, Germany), by the Russian Foundation for Basic Research (project no. 08-01-00464) and by the programme “Leading Scientific Schools” (project no. NSH-1824.2008.1).

point $\gamma(x)$. All closed 2-forms (presymplectic structures) on the loop space ΩM that are given by matrix operators of the form $\omega_{ij}(u, u_x, \dots, u_{(k)})$, i.e., all closed 2-forms of the form

$$\omega(\xi, \eta) = \int_{S^1} \xi^i \omega_{ij}(u, u_x, \dots, u_{(k)}) \eta^j dx, \quad (4)$$

where $\xi, \eta \in T_\gamma \Omega M$, were completely described in [1] (see also descriptions of various differential-geometric classes of symplectic (presymplectic) and Poisson structures in [2]–[9]).

Theorem 1 [1]. *A bilinear form (4) is a closed skew-symmetric 2-form (a presymplectic structure) on the loop space ΩM if and only if*

$$\omega_{ij}(u, u_x, \dots, u_{(k)}) = T_{ijk}(u) u_x^k + \Omega_{ij}(u), \quad (5)$$

where $T_{ijk}(u)$ is an arbitrary closed 3-form on the manifold M^N and $\Omega_{ij}(u)$ is an arbitrary closed 2-form on M^N .

If the matrix $\omega_{ij}(u, u_x, \dots, u_{(k)})$ is nondegenerate, $\det(\omega_{ij}(u, u_x, \dots, u_{(k)})) \neq 0$, then the corresponding presymplectic form (4), (5) is symplectic and the inverse matrix $\omega^{ij}(u, u_x, \dots, u_{(k)})$, $\omega^{is}(u, u_x, \dots, u_{(k)}) \omega_{sj}(u, u_x, \dots, u_{(k)}) = \delta_j^i$, gives the Poisson structure

$$\{I, J\} = \int \frac{\delta I}{\delta u^i(x)} \omega^{ij}(u, u_x, \dots, u_{(k)}) \frac{\delta J}{\delta u^j(x)} dx \quad (6)$$

on the loop space ΩM , i.e., the bracket (6) is skew-symmetric and satisfy the Jacobi identity. Therefore Theorem 1 gives the complete description of all nondegenerate Poisson structures on the loop space ΩM that are given by matrix operators of the form $\omega^{ij}(u, u_x, \dots, u_{(k)})$, i.e., all the nondegenerate Poisson brackets of the form (6), $\det(\omega^{ij}(u, u_x, \dots, u_{(k)})) \neq 0$ (such nondegenerate Poisson structures were studied by Ashtashov and Vinogradov in [9], see also [7]–[8] and [1]–[6]). We note that if the closed 2-form $\Omega_{ij}(u)$ is nondegenerate, $\det(\Omega_{ij}(u)) \neq 0$, i.e, the form $\Omega_{ij}(u)$ is symplectic on M^N , then the 2-form (5) is a nondegenerate form on ΩM for any closed 3-form $T_{ijk}(u)$ on the manifold M^N since it is obvious that in this case $\det(T_{ijk}(u) u_x^k + \Omega_{ij}(u)) \neq 0$. Thus we can define, on the loop space of an arbitrary symplectic manifold M^N , the Poisson bracket

$$\{I, J\} = \int \frac{\delta I}{\delta u^i(x)} \omega^{ij}(u, u_x) \frac{\delta J}{\delta u^j(x)} dx, \quad (7)$$

where

$$\omega^{li}(u, u_x)(T_{ijk}(u) u_x^k + \Omega_{ij}(u)) = \delta_j^l, \quad (8)$$

I and J being arbitrary functionals on ΩM . The Poisson bracket (7), (8) is a partial case of the bracket (1)–(3), namely, the case when the Poisson structure $\omega^{ij}(u)$ is nondegenerate, $\det(\omega^{ij}(u)) \neq 0$, $\omega^{ij}(u) = \Omega^{ij}(u)$, $\Omega^{is}(u) \Omega_{sj}(u) = \delta_j^i$, since

$$\omega^{ij}(u, u_x) = C_s^i(u, u_x) \Omega^{sj}(u), \quad (9)$$

where

$$C_s^l(u, u_x)(\delta_j^s + \Omega^{si}(u)T_{ijk}(u)u_x^k) = \delta_j^l. \quad (10)$$

The case of degenerate Poisson structures $\omega^{ij}(u)$, $\det(\omega^{ij}(u)) = 0$, is much more complicated. We note that in contrast to the case of all closed 2-forms (presymplectic structures) of the form (4) (Theorem 1) the problem of description of all degenerate Poisson structures of the form (6) is a very complicated and unsolved problem.

Theorem 2. *An arbitrary Poisson structure $\omega^{ij}(u)$ and an arbitrary closed 3-form $T_{ijk}(u)$ give the local Poisson bracket (1)–(3).*

First of all, we note that obviously the matrix operator $A^{ij}(u, u_x)$ (1), (2) is skew-symmetric.

Lemma. *A skew-symmetric matrix operator $A^{ij}(u, u_x)$ (1) gives a Poisson bracket (3) if and only if the following relations hold:*

$$\omega^{ij}(u)\omega^{rp}(u)\frac{\partial M_r^s}{\partial u_x^i} = \omega^{is}(u)\omega^{rj}(u)\frac{\partial M_r^p}{\partial u_x^i}, \quad (11)$$

$$\begin{aligned} & \omega^{ij}(u)\omega^{rp}(u)\frac{\partial M_r^s}{\partial u^i} - \omega^{ij}(u)\frac{d}{dx}\left(\frac{\partial M_r^s}{\partial u_x^i}\omega^{rp}(u)\right) + \frac{\partial \omega^{ij}}{\partial u^r}\omega^{rp}(u)M_i^s(u) + \\ & + \omega^{is}(u)\omega^{rj}(u)\frac{\partial M_r^p}{\partial u^i} + \frac{d}{dx}\left(\omega^{is}(u)\right)\frac{\partial M_r^p}{\partial u_x^i}\omega^{rj}(u) + \frac{\partial \omega^{is}}{\partial u^r}\omega^{rj}(u)M_i^p(u) + \\ & + \omega^{ip}(u)\omega^{rs}(u)\frac{\partial M_r^j}{\partial u^i} + \frac{d}{dx}\left(\omega^{ip}(u)\right)\frac{\partial M_r^j}{\partial u_x^i}\omega^{rs}(u) + \frac{\partial \omega^{ip}}{\partial u^r}\omega^{rs}(u)M_i^j(u) = 0. \end{aligned} \quad (12)$$

If $M_s^i(u, u_x) = \delta_j^s + \omega^{sp}(u)T_{pj k}(u)u_x^k$, then relations (11), (12) hold for an arbitrary Poisson structure $\omega^{ij}(u)$ and an arbitrary closed 3-form $T_{ijk}(u)$.

Let us add an arbitrary parameter ε in the formula for our Poisson structure:

$$A^{ij}(\varepsilon, u, u_x) = B_s^i(\varepsilon, u, u_x)\omega^{sj}(u), \quad (13)$$

where

$$B_s^i(\varepsilon, u, u_x)(\delta_j^s + \varepsilon\omega^{sp}(u)T_{pj k}(u)u_x^k) = \delta_j^i. \quad (14)$$

We can now expand the Poisson structure $A^{ij}(\varepsilon, u, u_x)$ in series in ε :

$$A^{ij}(\varepsilon, u, u_x) = \omega^{ij}(u) - \varepsilon\omega^{is}(u)T_{sr k}(u)\omega^{rj}(u)u_x^k + \dots \quad (15)$$

This expansion give an ε -deformation of an arbitrary Poisson structure $\omega^{ij}(u)$ by means of an arbitrary closed 3-form $T_{ijk}(u)$.

We note that this ε -deformation of an arbitrary Poisson structure $\omega^{ij}(u)$ belongs to a special class of graded ε -deformations of Poisson structures (see, for example, [10], [11]).

Acknowledgements. The work was supported by the Max-Planck-Institut für Mathematik (Bonn, Germany), by the Russian Foundation for Basic Research (project no. 08-01-00464) and by a grant of the President of the Russian Federation (project no. NSh-1824.2008.1).

References

- [1] O. I. Mokhov, “Symplectic and Poisson geometry on loop spaces of manifolds and nonlinear equations”, In: *Topics in Topology and Mathematical Physics*, Ed. S.P.Novikov, Amer. Math. Soc., Providence, RI, 1995, pp. 121–151;
<http://arXiv.org/hep-th/9503076> (1995).
- [2] O. I. Mokhov, “Poisson and symplectic geometry on loop spaces of smooth manifolds”, In: *Geometry from the Pacific Rim, Proceedings of the Pacific Rim Geometry Conference held at National University of Singapore, Republic of Singapore, December 12–17, 1994*, Eds. A.J.Berrick, B.Loo, H.-Y.Wang, Walter de Gruyter, Berlin, 1997, pp. 285–309.
- [3] O. I. Mokhov, “Differential geometry of symplectic and Poisson structures on loop spaces of smooth manifolds, and integrable systems”, *Trudy Matem. Inst. Akad. Nauk*, Vol. 217, Moscow, Nauka, 1997, pp. 100–134; English translation in *Proceedings of the Steklov Institute of Mathematics (Moscow)*, Vol. 217, 1997, pp. 91–125.
- [4] O. I. Mokhov, “Symplectic and Poisson structures on loop spaces of smooth manifolds, and integrable systems”, *Uspekhi Matematicheskikh Nauk*, Vol. 53, No. 3, 1998, pp. 85–192; English translation in *Russian Mathematical Surveys*, Vol. 53, No. 3, 1998, pp. 515–622.
- [5] O. I. Mokhov, “Symplectic and Poisson geometry on loop spaces of smooth manifolds and integrable equations”, Moscow–Izhevsk, Institute of Computer Studies, 2004 (In Russian); English version: *Reviews in Mathematics and Mathematical Physics*, Vol. 11, Part 2, Harwood Academic Publishers, 2001; Second Edition: *Reviews in Mathematics and Mathematical Physics*, Vol. 13, Part 2, Cambridge Scientific Publishers, 2009.
- [6] O. I. Mokhov, “Symplectic forms on loop space and Riemannian geometry”, *Funkts. Analiz i Ego Prilozh.*, Vol. 24, No. 3, 1990, pp. 86–87; English translation in *Functional Analysis and its Applications*, Vol. 24, No. 3, 1990, pp. 247–249.
- [7] A. M. Astashov, “Normal forms of Hamiltonian operators in field theory”, *Dokl. Akad. Nauk SSSR*, Vol. 270, No. 5, 1983, pp. 363–368; English translation in *Soviet Math. Dokl.*, Vol. 27, No. 3, 1983, pp. 685–689.
- [8] A. M. Vinogradov, “Hamiltonian structures in field theory”, *Dokl. Akad. Nauk SSSR*, Vol. 241, No. 1, 1978, pp. 18–21; English translation in *Soviet Math. Dokl.*, Vol. 19, No. 4, 1978, pp. 790–794.

- [9] A. M. Astashov and A. M. Vinogradov, “On the structure of Hamiltonian operators in field theory”, J. Geom. Phys., Vol. 3, No. 2, 1986, pp. 781–785; English translation in Soviet Math. Dokl., Vol. 27, 1983, pp. 263–287.
- [10] B. Dubrovin and Y. Zhang, “Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov - Witten invariants”; arXiv:math/0108160 (2001).
- [11] B. Dubrovin, S.-Q. Liu and Y. Zhang, “On Hamiltonian perturbations of hyperbolic systems of conservation laws, I: Quasi-triviality of bi- Hamiltonian perturbations,” Comm. Pure Appl. Math., Vol. 59, 2006, 559 - 615; arXiv:math/0410027 (2004).

O. I. Mokhov

Centre for Nonlinear Studies,
 L.D.Landau Institute for Theoretical Physics,
 Russian Academy of Sciences,
 Kosygina str., 2,
 Moscow, 117940, Russia;
 Department of Geometry and Topology,
 Faculty of Mechanics and Mathematics,
 M.V.Lomonosov Moscow State University,
 Moscow, 119992, Russia
E-mail: mokhov@mi.ras.ru; mokhov@landau.ac.ru; mokhov@bk.ru