

Nonlinear stability of periodic traveling wave solutions of systems of viscous conservation laws in the generic case

MATHEW A. JOHNSON* KEVIN ZUMBRUN†

February 13, 2022

Keywords: Periodic traveling waves; Bloch decomposition; modulated waves.

2000 MR Subject Classification: 35B35.

Abstract

Extending previous results of Oh–Zumbrun and Johnson–Zumbrun, we show that spectral stability implies linearized and nonlinear stability of spatially periodic traveling-wave solutions of viscous systems of conservation laws for systems of generic type, removing a restrictive assumption that wave speed be constant to first order along the manifold of nearby periodic solutions.

1 Introduction

Nonclassical viscous conservation laws arising in multiphase fluid and solid mechanics exhibit a rich variety of traveling wave phenomena, including homoclinic (pulse-type) and periodic solutions along with the standard heteroclinic (shock, or front-type) solutions [GZ, Z6, OZ1, OZ2]. Here, we investigate stability of spatially periodic traveling waves: specifically, sufficient conditions for stability of the wave.

In previous work [OZ4, JZ3], we showed that strong spectral stability in the sense of Schneider [S1, S2, S3] implies linearized and nonlinear $L^1 \cap H^K \rightarrow L^\infty$ stability in all dimensions $d \geq 1$. However, as pointed out in [OZ1, Se1], the conditions of Schneider are *nongeneric in the conservation law setting*, implying the restrictive condition that wave speed be constant to first order along the manifold of nearby periodic solutions. Indeed, it was shown in [OZ2] that failure of this condition implies a degradation in the decay rates of the Green function of the linearized equations about the periodic wave, suggesting

*Indiana University, Bloomington, IN 47405; matjohn@indiana.edu: Research of M.J. was partially supported by an NSF Postdoctoral Fellowship under NSF grant DMS-0902192.

†Indiana University, Bloomington, IN 47405; kzumbrun@indiana.edu: Research of K.Z. was partially supported under NSF grants no. DMS-0300487 and DMS-0801745.

that nonlinear stability would be unlikely in the general (nonstationary wave speed) case in dimension $d = 1$.

In this paper, we show that these difficulties are only apparent, and that, somewhat surprisingly, spectral stability implies nonlinear stability even if this additional condition on wave speeds is dropped. More precisely, we show that small $L^1 \cap H^s$ perturbations of a planar periodic solution $u(x, t) \equiv \bar{u}(x_1)$ (without loss of generality taken stationary) converge at Gaussian rate in L^p , $p \geq 2$ to a modulation

$$(1.1) \quad \bar{u}(x_1 - \psi(x, t))$$

of the unperturbed wave, where $x = (x_1, \tilde{x})$, $\tilde{x} = (x_2, \dots, x_d)$, and ψ is a scalar function whose x - and t -gradients decay at Gaussian rate in all L^p , $p \geq 2$, but which itself decays more slowly by a factor $t^{1/2}$; in particular, ψ is merely bounded in L^∞ for dimension $d = 1$.

In proving this result, we make crucial use of the tools developed in [OZ4, JZ3], in particular, a key nonlinear cancellation argument of [JZ3]. The key new observation making possible the treatment of the generic case is a careful study of the Bloch perturbation expansion about frequency $\xi = 0$, motivated by relations to the Whitham averaged system observed in [Se1, OZ3, JZ1, JZB].

It was shown in [Se1, OZ3] that the low-frequency dispersion relation near zero of the linearized operator about a periodic solution \bar{u} agrees to first order with that of the linearization about a constant state of the Whitham averaged system

$$(1.2) \quad \begin{aligned} \partial_t M + \sum_j \partial_{x_j} F^j &= 0, \\ \partial_t(\Omega N) + \nabla_x(\Omega S) &= 0, \end{aligned}$$

where $M \in \mathbb{R}^n$ denotes the average over one period, F^j the average of an associated flux, $\Omega = |\nabla_x \Psi| \in \mathbb{R}^1$ the frequency, $S = -\Psi_t / |\nabla_x \Psi| \in \mathbb{R}^1$ the speed s , and $N = \nabla_x \Psi / |\nabla_x \Psi| \in \mathbb{R}^d$ the normal ν associated with nearby periodic waves, with an additional constraint

$$(1.3) \quad \text{curl}(\Omega N) = \text{curl} \nabla_x \Psi \equiv 0.$$

As noted in [Se1, OZ3], this implies both that the eigenvalues $\lambda_j(\xi)$ bifurcating from $\lambda = 0$ at $\xi = 0$ are C^1 along rays through the origin, and that weak hyperbolicity (reality of characteristics of (1.2)–(1.3)) is necessary for spectral or linearized stability.

As noted in [JZB], there is a deeper analogy between the low-frequency linearized dispersion relation and the Whitham averaged system at the structural level, suggesting a useful rescaling of the low-frequency perturbation problem. It is this intuition that motivates our derivation of sharp low-frequency estimates crucial to the analysis of nonlinear stability. With these estimates in place, the rest of the argument goes exactly as in [JZ3, OZ4].

1.1 Equations and assumptions

Consider a parabolic system of conservation laws

$$(1.4) \quad u_t + \sum_j f^j(u)_{x_j} = \Delta_x u,$$

$u \in \mathcal{U}(\text{open}) \in \mathbb{R}^n$, $f^j \in \mathbb{R}^n$, $x \in \mathbb{R}^d$, $d \geq 1$, $t \in \mathbb{R}^+$, and a periodic traveling wave solution

$$(1.5) \quad u = \bar{u}(x \cdot \nu - st),$$

of period X , satisfying the traveling-wave ODE $\bar{u}'' = (\sum_j \nu_j f^j(\bar{u}))' - s\bar{u}'$ with boundary conditions $\bar{u}(0) = \bar{u}(X) =: u_0$. Integrating, we obtain a first-order profile equation

$$(1.6) \quad \bar{u}' = \sum_j \nu_j f^j(\bar{u}) - s\bar{u} - q,$$

where $(u_0, q, s, \nu, X) \equiv \text{constant}$. Without loss of generality take $\nu = e_1$, $s = 0$, so that $\bar{u} = \bar{u}(x_1)$ represents a stationary solution depending only on x_1 .

Following [Se1, OZ3, OZ4], we assume:

(H1) $f^j \in C^{K+1}$, $K \geq [d/2] + 4$.

(H2) The map $H : \mathbb{R} \times \mathcal{U} \times \mathbb{R} \times S^{d-1} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ taking $(X; a, s, \nu, q) \mapsto u(X; a, s, \nu, q) - a$ is full rank at $(\bar{X}; \bar{u}(0), 0, e_1, \bar{q})$, where $u(\cdot; \cdot)$ is the solution operator of (1.6).

Conditions (H1)–(H2) imply that the set of periodic solutions in the vicinity of \bar{u} form a smooth $(n+d+1)$ -dimensional manifold $\{\bar{u}^\alpha(x \cdot \nu(a) - \alpha - s(a)t)\}$, with $\alpha \in \mathbb{R}$, $a \in \mathbb{R}^{n+d}$.

1.1.1 Linearized equations

Linearizing (1.4) about $\bar{u}(\cdot)$, we obtain

$$(1.7) \quad v_t = Lv := \Delta_x v - \sum (A^j v)_{x_j},$$

where coefficients $A^j := Df^j(\bar{u})$ are now periodic functions of x_1 . Taking the Fourier transform in the transverse coordinate $\tilde{x} = (x_2, \dots, x_d)$, we obtain

$$(1.8) \quad \hat{v}_t = L_{\tilde{\xi}} \hat{v} = \hat{v}_{x_1, x_1} - (A^1 \hat{v})_{x_1} - i \sum_{j \neq 1} A^j \xi_j \hat{v} - \sum_{j \neq 1} \xi_j^2 \hat{v},$$

where $\tilde{\xi} = (\xi_2, \dots, \xi_d)$ is the transverse frequency vector.

1.1.2 Bloch–Fourier decomposition and stability conditions

Following [G, S1, S2, S3], we define the family of operators

$$(1.9) \quad L_\xi = e^{-i\xi_1 x_1} L_{\tilde{\xi}} e^{i\xi_1 x_1}$$

operating on the class of L^2 periodic functions on $[0, X]$; the (L^2) spectrum of $L_{\tilde{\xi}}$ is equal to the union of the spectra of all L_ξ with ξ_1 real with associated eigenfunctions

$$(1.10) \quad w(x_1, \tilde{\xi}, \lambda) := e^{i\xi_1 x_1} q(x_1, \xi_1, \tilde{\xi}, \lambda),$$

where q , periodic, is an eigenfunction of L_ξ . By continuity of spectrum, and discreteness of the spectrum of the elliptic operators L_ξ on the compact domain $[0, X]$, we have that the spectra of L_ξ may be described as the union of countably many continuous surfaces $\lambda_j(\xi)$.

Without loss of generality taking $X = 1$, recall now the *Bloch–Fourier representation*

$$(1.11) \quad u(x) = \left(\frac{1}{2\pi}\right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} e^{i\xi \cdot x} \hat{u}(\xi, x_1) d\xi_1 d\tilde{\xi}$$

of an L^2 function u , where $\hat{u}(\xi, x_1) := \sum_k e^{2\pi i k x_1} \hat{u}(\xi_1 + 2\pi k, \tilde{\xi})$ are periodic functions of period $X = 1$, $\hat{u}(\tilde{\xi})$ denoting with slight abuse of notation the Fourier transform of u in the full variable x . By Parseval’s identity, the Bloch–Fourier transform $u(x) \rightarrow \hat{u}(\xi, x_1)$ is an isometry in L^2 :

$$(1.12) \quad \|u\|_{L^2(x)} = \|\hat{u}\|_{L^2(\xi; L^2(x_1))},$$

where $L^2(x_1)$ is taken on $[0, 1]$ and $L^2(\xi)$ on $[-\pi, \pi] \times \mathbb{R}^{d-1}$. Moreover, it diagonalizes the periodic-coefficient operator L , yielding the *inverse Bloch–Fourier transform representation*

$$(1.13) \quad e^{Lt} u_0 = \left(\frac{1}{2\pi}\right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} e^{i\xi \cdot x} e^{L\xi t} \hat{u}_0(\xi, x_1) d\xi_1 d\tilde{\xi}$$

relating behavior of the linearized system to that of the diagonal operators L_ξ .

Loosely following [OZ4], we assume along with (H1)–(H2) the *strong spectral stability* conditions:

(D1) $\sigma(L_\xi) \subset \{\text{Re}\lambda < 0\}$ for $\xi \neq 0$.

(D2) $\text{Re}\sigma(L_\xi) \leq -\theta|\xi|^2$, $\theta > 0$, for $\xi \in \mathbb{R}^d$ and $|\xi|$ sufficiently small.

(D3’) $\lambda = 0$ is an eigenvalue of L_0 of multiplicity exactly $n + 1$.¹

As shown in [OZ3], (H1)–(H2) and (D1)–(D3’) imply that there exist $n + 1$ smooth eigenvalues

$$(1.14) \quad \lambda_j(\xi) = -ia_j(\xi) + o(|\xi|)$$

of L_ξ bifurcating from $\lambda = 0$ at $\xi = 0$, where $-ia_j$ are homogeneous degree one functions; see Lemma 2.1 below.

As in [OZ4], we make the further nondegeneracy hypothesis:

(H3) The functions $a_j(\xi)$ in (1.14) are distinct.

The functions a_j may be seen to be the characteristics associated with the Whitham averaged system (1.2)–(1.3) linearized about the values of M, S, N, Ω associated with the background wave \bar{u} ; see [OZ3, OZ4]. Thus, (D1) implies weak hyperbolicity of (1.2)–(1.3) (reality of a_j), while (H1) corresponds to strict hyperbolicity.

Remark 1.1. Condition (D3’) is a weakened version of the condition (D3) of [OZ4, JZ3] that $\lambda = 0$ be a *semisimple* eigenvalue of L_0 of minimal multiplicity $n + 1$, which implies [OZ1, OZ2, Se1] the special property that wave speed be stationary at \bar{u} along the manifold of nearby periodic solutions. The stronger conditions (D1)–(D3) are exactly the spectral assumptions of [S1, S2, S3] introduced by Schneider in the reaction-diffusion case. Conditions (D1)–(D3) (resp. (D1)–(D3’)) correspond to “dissipativity” of the large-time behavior of the linearized system [S1, S2, S3].

¹ The zero eigenspace of L_0 is at least $(n + 1)$ -dimensional by linearized existence theory and (H2), and hence $n + 1$ is the minimal multiplicity; see [Se1].

1.2 Main result

With these preliminaries, we can now state our main result.

Theorem 1.1. *Assuming (H1)–(H3) and (D1)–(D3’), for some $C > 0$ and $\psi \in W^{K,\infty}(x, t)$,*

$$(1.15) \quad \begin{aligned} |\tilde{u} - \bar{u}(\cdot - \psi)|_{L^p}(t) &\leq C(1+t)^{-\frac{d}{2}(1-1/p)} |\tilde{u} - \bar{u}|_{L^1 \cap H^K}|_{t=0}, \\ |\tilde{u} - \bar{u}(\cdot - \psi)|_{H^K}(t) &\leq C(1+t)^{-\frac{d}{4}} |\tilde{u} - \bar{u}|_{L^1 \cap H^K}|_{t=0}, \\ |(\psi_t, \psi_x)|_{W^{K+1,p}} &\leq C(1+t)^{-\frac{d}{2}(1-1/p)} |\tilde{u} - \bar{u}|_{L^1 \cap H^K}|_{t=0} \end{aligned}$$

for all $t \geq 0$, $p \geq 2$, $d \geq 1$, and

$$(1.16) \quad |\tilde{u} - \bar{u}|_{L^p}(t), |\psi(t)|_{L^p} \leq C(1+t)^{-\frac{d}{2}(1-\frac{1}{p})+\frac{1}{2}} |\tilde{u} - \bar{u}|_{L^1 \cap H^K}|_{t=0}$$

for all $t \geq 0$ and $p = \infty$ or $p \geq 2$ and $d \geq 3$, for solutions \tilde{u} of (1.4) with $|\tilde{u} - \bar{u}|_{L^1 \cap H^K}|_{t=0}$ sufficiently small. In particular, \bar{u} is nonlinearly bounded $L^1 \cap H^K \rightarrow L^\infty$ stable for $d \geq 1$, asymptotically $L^1 \cap H^K \rightarrow L^\infty$ stable for $d \geq 2$, and asymptotically $L^1 \cap H^K \rightarrow H^K$ stable for $d \geq 3$.

Remark 1.2. *In Theorem 1.1, derivatives in $x \in \mathbb{R}^d$ for $d \geq 2$ refer to total derivatives. Moreover, unless specified by an appropriate index, throughout this paper derivatives in spatial variable x will always refer to the total derivative of the function.*

In dimension one, Theorem 1.1 asserts only bounded $L^1 \cap H^K \rightarrow L^\infty$ stability, a very weak notion of stability. The bounds (1.15)–(1.16) agree for dimension $d = 1$ with those obtained in [JZ3] in the stationary wave speed case that (D3) holds in place of (D3’), but for higher dimensions are weaker by roughly factor $t^{1/2}$.

Remark 1.3. In dimension $d = 1$, it is straightforward to show that the results of Theorem 1.1 extend to all $1 \leq p \leq \infty$ using the pointwise techniques of [OZ2]; see Remark 3.6.

1.3 Discussion and open problems

The proof of Theorem 1.1 largely completes the line of investigation carried out in [OZ2, Se1, OZ3, OZ4, JZ3], showing that spectral stability implies linear and nonlinear stability of planar spatially periodic traveling waves. The corresponding spectral stability problem has been studied analytically in [OZ1, Se1, OZ3], yielding various necessary conditions, and by numerical Evans function investigation in [OZ1]. An interesting direction for further study would be more systematic numerical investigation along the lines of [BLZ, HLYZ1, HLYZ2, BHZ, BLZ] in the viscous shock wave case. A second interesting open problem would be to extend the results for planar waves to the case of solutions with multiple periods, as considered in the reaction–diffusion setting in [S1, S2, S3].

The key to the nonlinear analysis in critical dimensions $d = 1, 2$, as in [JZ3, S1, S2, S3], is to subtract out a slower-decaying part of the solution described by an appropriate modulation equation and show that the residual decays sufficiently rapidly to close a nonlinear

iteration. Note that the modulated approximation $\bar{u}(x_1 - \psi(x, t))$ of (1.1) is not the full Ansatz $\bar{u}^a(\Psi(x, t))$, $\Psi(x, t) := x_1 - \psi(x, t)$, associated with the Whitham averaged system (1.2)–(1.3), where \bar{u}^a is the manifold of periodic solutions near \bar{u} introduced below (H2), but only the translational part not involving perturbations a in the profile. (See [OZ3] for the derivation of Ansatz and (1.2)–(1.3).) That is, we don't need to separate out all variations along the manifold of periodic solutions, but only the special variations connected with translation invariance.

This can be understood heuristically by the observation that (1.2) indicates that variables a , $\nabla_x \Psi$ are roughly comparable, which would suggest, by the diffusive behavior $\Psi \gg \nabla_x \Psi$, that a is negligible with respect to Ψ . Indeed, this heuristic argument translates rigorously to our ultimate computation of linearized behavior leading to the final result; see Section 2 and Remark 2.2. In this respect, the connection to the Whitham system is somewhat clearer in the generic case considered here than in the quasi-Hamiltonian case treated previously in [OZ2, OZ4, JZ3].²

It would be interesting to better understand the connection between the Whitham averaged system (or suitable higher-order correction) and behavior at the nonlinear level, as explored at the linear level in [OZ3, OZ4, JZ1, JZB]. As discussed further in [OZ3], another interesting problem would be to try to rigorously justify the WKB expansion for the related vanishing viscosity problem, in the spirit of [GMWZ1, GMWZ2].

2 Spectral preparation

We begin by a careful study of the Bloch perturbation expansion at $\xi = 0$.

Lemma 2.1. *Assuming (H1)–(H3), (D1)–(D3'), the eigenvalues $\lambda_j(\xi/|\xi|, \xi)$ of L_ξ are analytic functions of $\xi/|\xi|$ and $|\xi|$. Suppose further that 0 is a nonsemisimple eigenvalue of L_0 , i.e., (D3') holds, but not (D3). Then, the Jordan structure of the zero eigenspace of L_0 consists of an n -dimensional kernel and a single Jordan chain of height 2, where the left kernel of L_0 is the n -dimensional subspace of constant functions, and \bar{u}' spans the right eigendirection lying at the base of the Jordan chain. Moreover, for $|\xi|$ sufficiently small, there exist right and left eigenfunctions $q_j(\xi/|\xi|, \xi, \cdot)$ and $\tilde{q}_j(\xi/|\xi|, \xi, \cdot)$ of L_ξ associated with λ_j of form $q_j = \sum_k \beta_{j,k} v_k$ and $\tilde{q}_j = \sum_k \tilde{\beta}_{j,k} \tilde{v}_k$ where $\{v_j\}$ and $\{\tilde{v}_j\}$ are dual bases of the total eigenspace of L_ξ associated with sufficiently small eigenvalues, analytic in $\omega = \xi/|\xi|$ and $|\xi|$, with $\tilde{v}_j(\omega; 0)$ constant for $j \neq n$ and $v_n(\omega; 0) \equiv \bar{u}'(\cdot)$; $\tilde{\beta}_{j,1}, \dots, \tilde{\beta}_{j,n-1}, |\xi|^{-1} \tilde{\beta}_{j,n}, \tilde{\beta}_{j,n+1}$ and $\beta_{j,1}, \dots, \beta_{j,n-1}, |\xi| \beta_{j,n}, \beta_{j,n+1}$ are analytic in $\xi/|\xi|, |\xi|$; and $\langle \tilde{q}_j, q_k \rangle = \delta_j^k$.*

Proof. Recall that L_ξ as an elliptic second-order operator on bounded domain has spectrum consisting of isolated eigenvalues of finite multiplicity. Expanding

$$(2.1) \quad L_\xi = L_0 + |\xi| L_{\xi/|\xi|}^1 + |\xi|^2 L_{\xi/|\xi|}^2$$

² In the degenerate case that the stronger condition (D3) holds, i.e., wave speed is stationary at \bar{u} , the situation is somewhat more complicated, and these relations break down; see [JZ3] for further discussion.

for each fixed angle $\hat{\xi} := \xi/|\xi|$, consider the continuous family of spectral perturbation problems in $|\xi|$ indexed by angle $\omega = \xi/|\xi|$ about the eigenvalue $\lambda = 0$ of L_0 .

Because 0 is an isolated eigenvalue of L_0 , the associated total right and left eigenprojections P_0 and \tilde{P}_0 perturb analytically in both ω and $|\xi|$, giving projection P_ξ and \tilde{P}_ξ [K]. These yield in standard fashion (for example, by projecting appropriately chosen fixed subspaces) locally analytic right and left bases $\{v_j\}$ and $\{\tilde{v}_j\}$ of the associated total eigenspaces given by the range of P_ξ, \tilde{P}_ξ .

Defining $V = (v_1, \dots, v_{n+1})$ and $\tilde{V} = (\tilde{v}_1, \dots, \tilde{v}_{n+1})^*$, $*$ denoting adjoint, we may convert the infinite-dimensional perturbation problem (2.1) into an $(n+1) \times (n+1)$ matrix perturbation problem

$$(2.2) \quad M_\xi = M_0 + |\xi|M_1 + |\xi|^2M_2,$$

where $M_\xi(\omega, |\xi|) := \langle \tilde{V}_\xi^*, L_\xi V_\xi \rangle$ and $\langle \cdot, \cdot \rangle$ refers to the $L^2(x_1)$ inner product on $[0, X]$. That is, the eigenvalues $\lambda_j(\xi)$ lying near 0 of L_ξ are the eigenvalues of M_ξ , and the associated right and left eigenfunctions of L_ξ are

$$(2.3) \quad f_j = Vw_j \text{ and } \tilde{f}_j = \tilde{w}_j\tilde{V}^*,$$

where w_j and \tilde{w}_j are the associated right and left eigenvectors of M_ξ .

Case (i). If $\lambda = 0$ is a semisimple eigenvalue of L_0 , then $M_0 = 0$, and (2.2) reduces to the simpler perturbation problem $\tilde{M}_\xi := |\xi|^{-1}M_\xi = M_1 + |\xi|M_2$ studied in [OZ4, JZ3], which $\lambda_j(\xi) = |\xi|\tilde{\lambda}_j(\xi)$, $\tilde{\lambda}_j(\xi)$ denoting the eigenvalues of \tilde{M}_ξ . Since $\tilde{\lambda}_j$ are continuous, λ_j are differentiable at $|\xi| = 0$ in the parameter $|\xi|$ as asserted in the introduction. Moreover, by (H3), the eigenvalues $\tilde{\lambda}_j(0)$ of $M_1 = \tilde{M}_0$ are distinct, and so they perturb analytically in $\omega, |\xi|$, as do the associated right and left eigenvectors.

Case (ii). Hereafter, assume that $\lambda = 0$ is a nonsemisimple eigenvalue of L_0 , so that M_0 is nilpotent but nonzero, possessing a nontrivial associated Jordan chain. Moreover, as the n -dimensional subspace of constant functions by direct computation lie in the kernel of $L_0^* = (\partial_{x_1}^2 + A_1^*\partial_{x_1})$, where $A_1(x_1) := df^1(\bar{u}(x_1))$, we have that the $(n+1)$ -dimensional zero eigenspace of L_0 consists precisely of an n -dimensional kernel and a single Jordan chain of height two. Moreover, by translation-invariance (differentiate in x_1 the profile equation (1.6)), we have $L_0\bar{u}' = 0$, so that \bar{u}' lies in the right kernel of L_0 .

Now, recall the assumption (H2) that $H : \mathbb{R} \times \mathcal{U} \times \mathbb{R} \times S^{d-1} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ taking $(X; a, s, \nu, q) \mapsto u(X; a, s, \nu, q) - a$ is full rank at $(\bar{X}; \bar{u}(0), 0, e_1, \bar{q})$, where $u(\cdot; \cdot)$ is the solution operator of profile ODE (1.6). The fact that $\ker L_0$ is n -dimensional implies that the restriction \tilde{H} taking $(a, q) \mapsto u(X; a, s, \nu, q) - a$ for fixed (X, ν, s) is also full rank, i.e., H is full rank with respect to the specific parameters (X, s, ν) . Applying the Implicit Function Theorem and counting dimensions, we find that the set of periodic solutions, i.e., the inverse image of zero under map H local to \bar{u} is a smooth $(n+d+1)$ -dimensional manifold $\{\bar{u}^a(x \cdot \nu(a) - \alpha - s(a)t)\}$, with $\alpha \in \mathbb{R}$, $a \in \mathbb{R}^{n+d}$. Moreover, $d+1$ dimensions may be parametrized by (X, s, ν) , or without loss of generality $(a_1, \dots, a_{d+1}) = (X, s, \nu)$.

Fixing (X, ν) and $(a_{d+2}, \dots, a_{n+d+1})$, and varying s , we find by differentiation of (1.6) that $f_* := -\partial_s \bar{u}$ satisfies³ the generalized eigenfunction equation

$$L_0 f_* = \bar{u}'.$$

Thus, \bar{u}' spans the eigendirection lying at the base of the Jordan chain, with the generalized zero-eigenfunction of L_0 corresponding to variations in speed along the manifold of periodic solutions about \bar{u} . Without loss of generality, therefore, we may take $\tilde{v}_1, \dots, \tilde{v}_{n-1}$ and \tilde{v}_{n+1} to be constant at $|\xi| = 0$, i.e., depending only on $\omega = \xi/|\xi|$ and not x_1 , and $v_n \equiv \bar{u}'$ at $|\xi| = 0$ independent of ω .

Recalling from [JZ3] the fact that

$$\langle c, L^1 \bar{u}' \rangle = \langle c, (\omega_1(2\partial_{x_1} - A_1) - \sum_{j \neq 1} \omega_j A_j) \bar{u}' \rangle = \langle c, \omega_1 \partial_{x_1}^2 \bar{u} - \sum_{j \neq 1} \omega_j \partial_{x_1} f^j(\bar{u}) \rangle \equiv 0$$

for any constant functions c , where again $\langle \cdot, \cdot \rangle$ denotes $L^2(x_1)$ inner product on the interval $x_1 \in [0, X]$, and $A_j := df^j(\bar{u}(\cdot))$, we find under this normalization that (2.2) has the special structure

$$(2.4) \quad M_0 = \begin{pmatrix} 0_{(n-1) \times (n-1)} & 0_{n-1} & 0_{n-1} \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad M_1 = \begin{pmatrix} * & 0_{n-1} & * \\ * & * & * \\ * & 0 & * \end{pmatrix}.$$

Now, rescaling (2.2) as

$$(2.5) \quad \check{M}_\xi := |\xi|^{-1} S(\xi) M_\xi S(\xi)^{-1},$$

where

$$(2.6) \quad S := \begin{pmatrix} I_{n-1} & 0 & 0 \\ 0 & |\xi| & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

we obtain

$$(2.7) \quad \check{M}_\xi = \check{M}_0 + |\xi| \check{M}_1 + O(|\xi|^2),$$

where $\check{M}_j = \check{M}_j(\omega)$ like the original M_j are analytic matrix-valued functions of ω , and the eigenvalues $m_j(\xi) = m_j(\omega; |\xi|)$ of \check{M}_ξ are $|\xi|^{-1} \lambda_j(\xi)$.

As the eigenvalues m_j of \check{M}_ξ are continuous, the eigenvalues $\lambda_j(\xi) = |\xi| m_j$ are differentiable at $|\xi| = 0$ as asserted in the introduction. Moreover, by (H3), the eigenvalues $\check{\lambda}_j(0)$ of \check{M}_0 are distinct, and so they perturb analytically in $\omega, |\xi|$, as do the associated right and left eigenvectors z_j and \tilde{z}_j . Undoing the rescaling (2.5), and recalling (2.3), we obtain the result. \square

Remark 2.2. Note that the n th coordinate of vectors $w \in \mathbb{C}^{n+1}$ in the perturbation problem (2.2) corresponds as the coefficient of \bar{u}' to variations Ψ in displacement. Thus, rescaling (2.5) amounts to substituting for Ψ the variable $|\xi| \Psi \sim \Psi_x$ of the Whitham averaged system (1.2).

³Note the function f_* is X -periodic, and hence in the domain of L_0 since we have fixed the period X .

3 Linearized stability estimates

By standard spectral perturbation theory [K], the total eigenprojection $P(\xi)$ onto the eigenspace of L_ξ associated with the eigenvalues $\lambda_j(\xi)$, $j = 1, \dots, n+1$ described in the previous section is well-defined and analytic in ξ for ξ sufficiently small, since these (by discreteness of the spectra of L_ξ) are separated at $\xi = 0$ from the rest of the spectrum of L_0 . Introducing a smooth cutoff function $\phi(\xi)$ that is identically one for $|\xi| \leq \varepsilon$ and identically zero for $|\xi| \geq 2\varepsilon$, $\varepsilon > 0$ sufficiently small, we split the solution operator $S(t) := e^{Lt}$ into low- and high-frequency parts

$$(3.1) \quad S^I(t)u_0 := \left(\frac{1}{2\pi}\right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} e^{i\xi \cdot x} \phi(\xi) P(\xi) e^{L\xi t} \hat{u}_0(\xi, x_1) d\xi_1 d\tilde{\xi}$$

and

$$(3.2) \quad S^{II}(t)u_0 := \left(\frac{1}{2\pi}\right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} e^{i\xi \cdot x} (I - \phi P(\xi)) e^{L\xi t} \hat{u}_0(\xi, x_1) d\xi_1 d\tilde{\xi}.$$

3.1 High-frequency bounds

By standard sectorial bounds [He, Pa] and spectral separation of $\lambda_j(\xi)$ from the remaining spectra of L_ξ , we have trivially the exponential decay bounds

$$(3.3) \quad \begin{aligned} \|e^{L\xi t} (I - \phi P(\xi)) f\|_{L^2([0, X])} &\leq C e^{-\theta t} \|f\|_{L^2([0, X])}, \\ \|e^{L\xi t} (I - \phi P(\xi)) \partial_{x_1}^l f\|_{L^2([0, X])} &\leq C t^{-\frac{l}{2}} e^{-\theta t} \|f\|_{L^2([0, X])}, \\ \|\partial_{x_1}^l e^{L\xi t} (I - \phi P(\xi)) f\|_{L^2([0, X])} &\leq C t^{-\frac{l}{2}} e^{-\theta t} \|f\|_{L^2([0, X])}, \end{aligned}$$

for $\theta, C > 0$, and $0 \leq m \leq K$ (K as in (H1)). Together with (1.12), these give immediately the following estimates.

Proposition 3.1 ([OZ4]). *Under assumptions (H1)–(H3) and (D1)–(D2), for some $\theta, C > 0$, and all $t > 0$, $2 \leq p \leq \infty$, $0 \leq l \leq K+1$, $0 \leq m \leq K$,*

$$(3.4) \quad \begin{aligned} \|\partial_x^l S^{II}(t) f\|_{L^2(x)}, \|S^{II}(t) \partial_x^l f\|_{L^2(x)} &\leq C t^{-\frac{l}{2}} e^{-\theta t} \|f\|_{L^2(x)}, \\ \|\partial_x^m S^{II}(t) f\|_{L^p(x)}, \|S^{II}(t) \partial_x^m f\|_{L^p(x)} &\leq C t^{-\frac{d}{2}(\frac{1}{2} - \frac{1}{p}) - \frac{m}{2}} e^{-\theta t} \|f\|_{L^2(x)}, \end{aligned}$$

where, again, derivatives in $x \in \mathbb{R}^d$ refers to total derivatives.

Proof. The first inequalities follow immediately by (1.12). The second follows for $p = \infty$, $m = 0$ by Sobolev embedding from

$$\|S^{II}(t) f\|_{L^\infty(\tilde{x}; L^2(x_1))} \leq C t^{-\frac{d-1}{4}} e^{-\theta t} \|f\|_{L^2([0, X])}$$

and

$$\|\partial_{x_1} S^{II}(t) f\|_{L^\infty(\tilde{x}; L^2(x_1))} \leq C t^{-\frac{d-1}{4} - \frac{1}{2}} e^{-\theta t} \|f\|_{L^2([0, X])},$$

which follow by an application of (1.12) in the x_1 variable and the Hausdorff–Young inequality $\|f\|_{L^\infty(\tilde{x})} \leq \|\hat{f}\|_{L^1(\tilde{\xi})}$ in the variable \tilde{x} . The result for derivatives in x_1 and general $2 \leq p \leq \infty$ then follows by L^p interpolation. Finally, the result for derivatives in \tilde{x} follows from the inverse Fourier transform, equation (3.2), and the large $|\xi|$ bound

$$|e^{Lt}f|_{L^2(x_1)} \leq e^{-\theta|\tilde{\xi}|^2t}|f|_{L^2(x_1)}, \quad |\xi| \text{ sufficiently large,}$$

which easily follows from Parseval and the fact that L_ξ is a relatively compact perturbation of $\partial_x^2 - |\xi|^2$. Thus, by the above estimate we have

$$\begin{aligned} \|e^{Lt}\partial_{\tilde{x}}f\|_{L^2(x)} &\leq C\|e^{L\xi t}|\tilde{\xi}|\hat{f}\|_{L^2(x_1,\xi)} \\ &\leq C\sup\left(e^{-\theta|\tilde{\xi}|^2t}|\xi|\right)\|\hat{f}\|_{L^2(x_1,\xi)} \\ &\leq Ct^{-1/2}\|f\|_{L^2(x)}. \end{aligned}$$

A similar argument applies for $1 \leq m \leq K$. □

3.2 Low-frequency bounds

Denote by

$$(3.5) \quad G^I(x, t; y) := S^I(t)\delta_y(x)$$

the Green kernel associated with S^I , and

$$(3.6) \quad [G_\xi^I(x_1, t; y_1)] := \phi(\xi)P(\xi)e^{L\xi t}[\delta_{y_1}(x_1)]$$

the corresponding kernel appearing within the Bloch–Fourier representation of G^I , where the brackets on $[G_\xi^I]$ and $[\delta_y]$ denote the periodic extensions of these functions onto the whole line. Then, we have the following descriptions of G^I , $[G_\xi^I]$, deriving from the spectral expansion (1.14) of L_ξ near $\xi = 0$.

Proposition 3.2 ([OZ4]). *Under assumptions (H1)–(H3) and (D1)–(D3'),*

$$(3.7) \quad \begin{aligned} [G_\xi^I(x_1, t; y_1)] &= \phi(\xi) \sum_{j=1}^{n+1} e^{\lambda_j(\xi)t} q_j(\xi, x_1) \tilde{q}_j(\xi, y_1)^*, \\ G^I(x, t; y) &= \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} e^{i\xi \cdot (x-y)} [G_\xi^I(x_1, t; y_1)] d\xi \\ &= \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} e^{i\xi \cdot (x-y)} \phi(\xi) \sum_{j=1}^{n+1} e^{\lambda_j(\xi)t} q_j(\xi, x_1) \tilde{q}_j(\xi, y_1)^* d\xi, \end{aligned}$$

where $*$ denotes matrix adjoint, or complex conjugate transpose, $q_j(\xi, \cdot)$ and $\tilde{q}_j(\xi, \cdot)$ are right and left eigenfunctions of L_ξ associated with eigenvalues $\lambda_j(\xi)$ defined in (1.14), normalized so that $\langle \tilde{q}_j, q_j \rangle \equiv 1$.

Proof. Relation (3.7)(i) is immediate from the spectral decomposition of elliptic operators on finite domains, and the fact that λ_j are distinct for $|\xi| > 0$ sufficiently small, by (H3). Substituting (3.5) into (3.1) and computing

$$(3.8) \quad \widehat{\delta}_y(\xi, x_1) = \sum_k e^{2\pi i k x_1} \widehat{\delta}_y(\xi + 2\pi k e_1) = \sum_k e^{2\pi i k x_1} e^{-i\xi \cdot y - 2\pi i k y_1} = e^{-i\xi \cdot y} [\delta_{y_1}(x_1)],$$

where the second and third equalities follow from the fact that the Fourier transform either continuous or discrete of the delta-function is unity, we obtain

$$\begin{aligned} G^I(x, t; y) &= \left(\frac{1}{2\pi}\right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} e^{i\xi \cdot x} \phi P(\xi) e^{L\xi t} \widehat{\delta}_y(\xi, x_1) d\xi \\ &= \left(\frac{1}{2\pi}\right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} e^{i\xi \cdot (x-y)} \phi P(\xi) e^{L\xi t} [\delta_{y_1}(x_1)] d\xi, \end{aligned}$$

yielding (3.7)(ii) by (3.6)(i) and the fact that ϕ is supported on $[-\pi, \pi]$. \square

Proposition 3.3. *Under assumptions (H1)-(H3) and (D1)-(D3'), the low-frequency Green function $G^I(x, t; y)$ of (3.5) decomposes as $G^I = E + \tilde{G}^I$,*

$$(3.9) \quad E = \bar{u}'(x) e(x, t; y),$$

where, for some $C > 0$, all $t > 0$,

$$(3.10) \quad \begin{aligned} \sup_y \|\tilde{G}^I(\cdot, t; y)\|_{L^p(x)} &\leq C(1+t)^{-\frac{d}{2}(1-\frac{1}{p})} \\ \sup_y \|\partial_y^r \tilde{G}^I(\cdot, t; y)\|_{L^p(x)}, \quad \sup_y \|\partial_t^r \tilde{G}^I(\cdot, t; y)\|_{L^p(x)} &\leq C(1+t)^{-\frac{d}{2}(1-\frac{1}{p})-\frac{1}{2}} \end{aligned}$$

for $p \geq 2$, $1 \leq r \leq 2$,

$$(3.11) \quad \sup_y \|\partial_x^j \partial_t^l \partial_y^r e(\cdot, t; y)\|_{L^p(x)} \leq C(1+t)^{-\frac{d}{2}(1-\frac{1}{p})-\frac{(j+l)}{2}-\frac{1}{2}}$$

for $p \geq 2$, $0 \leq j, k, l$, $j+l \leq K$, $1 \leq r \leq 2$, and

$$(3.12) \quad \sup_y \|\tilde{\partial}_x^j \partial_t^l e(\cdot, t; y)\|_{L^p(x)} \leq C(1+t)^{-\frac{d}{2}(1-\frac{1}{p})-\frac{(j+l)}{2}}$$

for $0 \leq j, k, l$, $j+l \leq K$, provided that $p \geq 2$ and $j+l \geq 1$ or $d \geq 3$, or $p = \infty$ and $d \geq 1$. Moreover, $e(x, t; y) \equiv 0$ for $t \leq 1$.

Remark 3.4. *In Proposition 3.3, and throughout the remainder of the paper, derivatives in $y \in \mathbb{R}^d$ refer to total derivatives, just as with the variable $x \in \mathbb{R}^d$.*

Proof. In the degenerate case (D3) that 0 is a semisimple eigenvalue of L_0 , these estimates have been established in [OZ4, JZ3]. Without loss of generality, therefore, we hereafter assume that 0 is a nonsemisimple eigenvalue of L_0 , with the consequences described in Lemma 2.1. Recalling that

$$(3.13) \quad \begin{aligned} G^I(x, t; y) &= \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} e^{i\xi \cdot (x-y)} \phi(\xi) \sum_{j=1}^{n+1} e^{\lambda_j(\xi)t} q_j(\xi, x_1) \tilde{q}_j(\xi, y_1)^* d\xi \\ &= \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} e^{i\xi \cdot (x-y)} \phi(\xi) \sum_{j,k,l=1}^{n+1} e^{\lambda_j(\xi)t} \beta_{j,k} v_k(\xi, x_1) \tilde{\beta}_{j,l} \tilde{v}_l(\xi, y_1)^* d\xi, \end{aligned}$$

define

$$(3.14) \quad \tilde{e}(x, t; y) = \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} e^{i\xi \cdot (x-y)} \phi(\xi) \sum_{j,l} e^{\lambda_j(\xi)t} \beta_{j,n} \tilde{\beta}_{j,l} \tilde{v}_l(\xi, y_1)^* d\xi$$

so that

$$(3.15) \quad \begin{aligned} G^I(x, t; y) - \bar{u}'(x_1) \tilde{e}(x, t; y) &= \\ &= \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} e^{i\xi \cdot (x-y)} \phi(\xi) \sum_{j,k \neq n,l} e^{\lambda_j(\xi)t} \beta_{j,k} \tilde{\beta}_{j,l} v_k(\xi, x_1) \tilde{v}_l(\xi, y_1)^* d\xi \\ &\quad + \left(\frac{1}{2\pi}\right)^d \int_{\mathbb{R}^d} e^{i\xi \cdot (x-y)} \phi(\xi) \sum_{j,l} e^{\lambda_j(\xi)t} \beta_{j,n} \tilde{\beta}_{j,l} \left(v_n(\xi, x_1) - \bar{u}'(x_1)\right) \tilde{v}_l(\xi, y_1)^* d\xi, \end{aligned}$$

where, by analyticity of v_n , $v_n(\xi, x_1) - \bar{u}'(x_1) = O(|\xi|)$, and so, by Lemma 2.1,

$$(3.16) \quad \beta_{j,n} \tilde{\beta}_{j,l} \left(v_n(\xi, x_1) - \bar{u}'(x_1)\right) \tilde{v}_l(\xi, y_1)^* = O(1)$$

and

$$(3.17) \quad \beta_{j,k} \tilde{\beta}_{j,l} v_k(\xi, x_1) \tilde{v}_l(\xi, y_1)^* = O(1) \text{ for } k \neq n.$$

Note further that $\tilde{v}_l \equiv \text{constant}$ unless $l = n$, in which case $\tilde{\beta}_{j,l} = O(|\xi|)$ by Lemma 2.1; hence

$$(3.18) \quad \partial_{y_1} \left(\beta_{j,n} \tilde{\beta}_{j,l} \left(v_n(\xi, x_1) - \bar{u}'(x_1)\right) \tilde{v}_l(\xi, y_1)^* \right) = O(|\xi|)$$

and

$$(3.19) \quad \partial_{y_1} \left(\beta_{j,k} \tilde{\beta}_{j,l} v_k(\xi, x_1) \tilde{v}_l(\xi, y_1)^* \right) = O(|\xi|) \text{ for } k \neq n.$$

From representation (3.15), bounds (3.16)–(3.17), and $\Re \lambda_j(\xi) \leq -\theta|\xi|^2$, we obtain by the triangle inequality

$$(3.20) \quad \|G^I - \bar{u}' \tilde{e}\|_{L^\infty(x,y)} \leq C \|e^{-\theta|\xi|^2 t} \phi(\xi)\|_{L^1(\xi)} \leq C(1+t)^{-\frac{d}{2}}.$$

Derivative bounds follow similarly, since x_1 -derivatives falling on v_{jk} are harmless, whereas, by (3.18)–(3.19), y_1 - or t -derivatives falling on \tilde{v}_{jl} or on $e^{i\xi \cdot (x-y)}$ bring down a factor of $|\xi|$ improving the decay rate by factor $(1+t)^{-1/2}$. (Note that $|\xi|$ is bounded because of the cutoff function ϕ , so there is no singularity at $t = 0$.)

To obtain bounds for $p = 2$, we note that (3.10 may be viewed itself as a Bloch–Fourier decomposition with respect to variable $z := x - y$, with y appearing as a parameter. Recalling (1.12), we may thus estimate

$$\begin{aligned}
(3.21) \quad & \sup_y \|G^I(\cdot, t; y) - \bar{u}'\tilde{e}(\cdot, t; y)\|_{L^2(x)} \leq \\
& C \sum_{j,k \neq n,l} \sup_y \|\phi(\xi) e^{\lambda_j(\xi)t} v_k(\cdot, z_1) \tilde{v}_l^*(\cdot, y_1) \tilde{v}_l(\cdot, y_1)^*\|_{L^2(\xi; L^2(z_1 \in [0, X]))} \\
& + C \sum_{j,l} \sup_y \|\phi(\xi) e^{\lambda_j(\xi)t} \left(\frac{v_n(\cdot, x_1) - \bar{u}'(x_1)}{|\cdot|} \right) \tilde{v}_l(\cdot, y_1)^*\|_{L^2(\xi; L^2(z_1 \in [0, X]))} \\
& \leq C \sum_{j,k \neq n,l} \sup_y \|\phi(\xi) e^{-\theta|\xi|^2 t}\|_{L^2(\xi)} \sup_{\xi} \|v_k(\cdot, z_1)\|_{L^2(0, X)} \|\tilde{v}_l(\cdot, y_1)^*\|_{L^\infty(0, X)} \\
& + C \sum_{j,l} \sup_y \|\phi(\xi) e^{-\theta|\xi|^2 t}\|_{L^2(\xi)} \sup_{\xi} \left\| \left(\frac{v_n(\xi, x_1) - \bar{u}'(x_1)}{|\xi|} \right) \right\|_{L^2(0, X)} \|\tilde{v}_l(\cdot, y_1)^*\|_{L^\infty(0, X)} \\
& \leq C(1+t)^{-\frac{d}{4}},
\end{aligned}$$

where we have used in a crucial way the boundedness of \tilde{v}_l in L^∞ ,⁴ and also the boundedness of

$$\left(\frac{v_n(\xi, x_1) - \bar{u}'(x_1)}{|\xi|} \right) \sim \partial_{|\xi|} v_n(\omega; r)$$

in L^2 , where $0 < r < |\xi|$. Derivative bounds follow similarly as above, noting that y - or t -derivatives bring down a factor $|\xi|$, while x -derivatives are harmless, to obtain an additional factor of $(1+t)^{-1/2}$ decay. Finally, bounds for $2 \leq p \leq \infty$ follow by L^p -interpolation.

Defining

$$(3.22) \quad e(x, t; y) := \chi(t)\tilde{e}(x, t; y),$$

where χ is a smooth cutoff function such that $\chi(t) \equiv 1$ for $t \geq 2$ and $\chi(t) \equiv 0$ for $t \leq 1$, and setting $\tilde{G} := G - \bar{u}'(x_1)e(x, t; y)$, we readily obtain the estimates (3.25) by combining the above estimates on $G^I - \bar{u}\tilde{e}$ with bound (3.4) on G^{II} .

⁴This is clear for $\xi = 0$, since v_j are linear combinations of genuine and generalized eigenfunctions, which are solutions of the homogeneous or inhomogeneous eigenvalue ODE. More generally, note that resolvent of $L_\xi - \gamma$ gains one derivative, hence the total eigenprojection, as a contour integral of the resolvent, does too— now, use the one-dimensional Sobolev inequality for periodic boundary conditions to bound the L^∞ difference from the mean by the (bounded) H^1 norm, then bound the mean by the L^1 norm, which is controlled by the L^2 norm.

Finally, recalling, by Lemma 2.1, that $\tilde{v}_l \equiv \text{constant}$ for $l \neq n$ while $\tilde{\beta}_{j,n} = O(|\xi|)$, we have

$$\partial_{y_1} \left(\beta_{j,n} \tilde{\beta}_{j,l} \tilde{v}_l(\xi, y_1)^* \right) = o(|\xi|).$$

Bounds (3.11) thus follow from (3.14) by the argument used to prove (3.10), together with the observation that x - or t -derivatives bring down factors of $|\xi|$.

Bounds (3.12) follow similarly for $p = \infty$ if $e^{-\theta|\xi|^2 t}/|\xi|$ is integrable in \mathbb{R}^d , and for $p \geq 2$ if $e^{-\theta|\xi|^2 t}/|\xi|^2$ is integrable, thus yielding the stated results for all $d \geq 2$. In the special case $d = 1$, $p = \infty$, (2.1) becomes a simpler one-parameter perturbation in ξ , and the $|\xi|^{-1}$ contributions become analytic multiples of ξ^{-1} , whose principal value integrals may be carried out explicitly to give a sum of traveling error functions that is bounded in L^∞ ; see the proof of Proposition 1.5, [OZ2] in the one-dimensional case. We omit this calculation as largely outside our analysis. (However, note that we need this bound to conclude L^∞ bounded stability in the one-dimensional case.) \square

Remark 3.5. Underlying our analysis, and that of [OZ2, JZ3], is the fundamental relation

$$(3.23) \quad G(x, t; y) = \left(\frac{1}{2\pi} \right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} e^{i\xi \cdot (x-y)} [G_\xi(x_1, t; y_1)] d\xi.$$

3.3 Final linearized bounds

Corollary 3.1. *Under assumptions (H1)–(H3), (D1)–(D3'), the Green function $G(x, t; y)$ of (1.7) decomposes as $G = E + \tilde{G}$,*

$$(3.24) \quad E = \bar{u}'(x)e(x, t; y),$$

where, for some $C > 0$, all $t > 0$, $1 \leq q \leq 2 \leq p \leq \infty$, $0 \leq j, k, l$, $j + l \leq K$, $1 \leq r \leq 2$,

$$(3.25) \quad \begin{aligned} \left| \int_{-\infty}^{+\infty} \tilde{G}(x, t; y) f(y) dy \right|_{L^p(x)} &\leq C(1+t)^{-\frac{d}{2}(1/2-1/p)} t^{-\frac{1}{2}(1/q-1/2)} |f|_{L^q \cap L^2}, \\ \left| \int_{-\infty}^{+\infty} \partial_y^r \tilde{G}(x, t; y) f(y) dy \right|_{L^p(x)} &\leq C(1+t)^{-\frac{d}{2}(1/2-1/p) - \frac{1}{2} + \frac{r}{2}} \\ &\quad \times t^{-\frac{d}{2}(1/q-1/2) - \frac{r}{2}} |f|_{L^q \cap L^2}, \\ \left| \int_{-\infty}^{+\infty} \partial_t^r \tilde{G}(x, t; y) f(y) dy \right|_{L^p(x)} &\leq C(1+t)^{-\frac{d}{2}(1/2-1/p) - \frac{1}{2} + r} \\ &\quad \times t^{-\frac{d}{2}(1/q-1/2) - r} |f|_{L^q \cap L^2}. \end{aligned}$$

$$(3.26) \quad \begin{aligned} \left| \int_{-\infty}^{+\infty} \partial_x^j \partial_t^k e(x, t; y) f(y) dy \right|_{L^p} &\leq (1+t)^{-\frac{d}{2}(1/q-1/p) - \frac{(j+k)}{2} + \frac{1}{2}} |f|_{L^q}, \\ \left| \int_{-\infty}^{+\infty} \partial_x^j \partial_t^k \partial_y^r e(x, t; y) f(y) dy \right|_{L^p} &\leq (1+t)^{-\frac{d}{2}(1/q-1/p) - \frac{(j+k)}{2}} |f|_{L^q}. \end{aligned}$$

Moreover, $e(x, t; y) \equiv 0$ for $t \leq 1$.

Proof. (Case $q = 1$). From (3.10) and the triangle inequality we obtain

$$\left\| \int_{\mathbb{R}^d} \tilde{G}^I(x, t; y) f(y) dy \right\|_{L^p(x)} \leq \int_{\mathbb{R}^d} \sup_y \|\tilde{G}^I(\cdot, t; y)\|_{L^p} |f(y)| dy \leq C(1+t)^{-\frac{d}{2}(1-1/p)} \|f\|_{L^1}$$

and similarly for y - and t -derivative estimates, which, together with (3.4), yield (3.25). Bounds (3.26) follow similarly by the triangle inequality and (3.11)–(3.12).

(Case $q = 2$). From (3.16)–(3.17), and analyticity of v_j , \tilde{v}_j , we have boundedness from $L^2[0, X] \rightarrow L^2[0, X]$ of the projection-type operators

$$(3.27) \quad f \rightarrow \beta_{j,n} \tilde{\beta}_{j,l} \left(v_n(\xi, x_1) - \bar{u}'(x_1) \right) \langle \tilde{v}_l, f \rangle$$

and

$$(3.28) \quad f \rightarrow \beta_{j,k} \tilde{\beta}_{j,l} v_k(\xi, x_1) \langle \tilde{v}_l, f \rangle \text{ for } k \neq n,$$

uniformly with respect to ξ , from which we obtain by (3.15), (3.22), and (1.12) the bound

$$(3.29) \quad \left| \int_{-\infty}^{+\infty} \tilde{G}^I(x, t; y) f(y) dy \right|_{L^2(x)} \leq C \|f\|_{L^2(x)},$$

for all $t \geq 0$, yielding together with (3.4) the result (3.25) for $p = 2$, $r = 1$. Similarly, by boundedness of \tilde{v}_j , v_j , \bar{u}' in all $L^p[0, X]$, we have

$$\begin{aligned} |e^{\lambda_j(\xi)t} \beta_{j,n} \tilde{\beta}_{j,l} \left(v_n(\xi, x_1) - \bar{u}'(x_1) \right) \langle \tilde{v}_l, \hat{f} \rangle|_{L^\infty(x_1)} &\leq C e^{-\theta|\xi|^2 t} |\hat{f}(\xi, \cdot)|_{L^2(x_1)}, \\ |e^{\lambda_j(\xi)t} \beta_{j,k} \tilde{\beta}_{j,l} v_k(\xi, x_1) \langle \tilde{v}_l, \hat{f} \rangle|_{L^\infty(x_1)} &\leq C e^{-\theta|\xi|^2 t} |\hat{f}(\xi, \cdot)|_{L^2(x_1)}, \text{ for } k \neq n, \end{aligned}$$

$C, \theta > 0$, yielding by definitions (3.15), (3.22) the bound

$$(3.30) \quad \begin{aligned} \left| \int_{-\infty}^{+\infty} \tilde{G}^I(x, t; y) f(y) dy \right|_{L^\infty(x)} &\leq \left(\frac{1}{2\pi} \right)^d \int_{-\pi}^{\pi} \int_{\mathbb{R}^{d-1}} C \phi(\xi) e^{-\theta|\xi|^2 t} |\hat{f}(\xi, \cdot)|_{L^2(x_1)} d\xi_1 d\tilde{\xi} \\ &\leq C |\phi(\xi) e^{-\theta|\xi|^2 t}|_{L^2(\xi)} |\hat{f}|_{L^2(\xi, x_1)} \\ &= C(1+t)^{-\frac{d}{4}} \|f\|_{L^2([0, X])}, \end{aligned}$$

hence giving the result for $p = \infty$, $r = 0$. The result for $r = 0$ and general $2 \leq p \leq \infty$ then follows by L^p interpolation between $p = 2$ and $p = \infty$. Derivative bounds $1 \leq r \leq 2$ follow by similar arguments, using (3.18)–(3.19). Bounds (3.26) follow similarly.

(Case $1 \leq q \leq 2$). By Riesz–Thorin interpolation between the cases $q = 1$ and $q = 2$, we obtain the bounds asserted in the general case $1 \leq q \leq 2$, $2 \leq p \leq \infty$. \square

Remark 3.6. The bounds on \tilde{G} , e_t , e_x may be recognized as the standard diffusive bounds satisfied for the heat equation [Z7]. For dimension $d = 1$, it may be shown using pointwise techniques as in [OZ2] that the bounds of Corollary 3.1 extend to all $1 \leq q \leq p \leq \infty$.

We note a striking analogy between the Green function decomposition of Corollary 3.1 and that of [MaZ3, Z4] in the viscous shock case; compare Proposition 3.3, [Z7].

4 Nonlinear stability in dimension one

With the bounds of Corollary (3.1), nonlinear stability follows by exactly the same argument as in [JZ3], included here for completeness. We carry out the nonlinear stability analysis only in the most difficult, one-dimensional, case. The extension to the multi-dimensional case is straightforward [JZ3, OZ4]. (Recall that the nonlinear iteration is easier to close in multi-dimensions, since the linearized behavior is faster decaying [OZ4, JZ3, S1, S2, S3].)

Hereafter, take $x \in \mathbb{R}^1$, dropping the indices on f^j and x_j and writing $u_t + f(u)_x = u_{xx}$.

4.1 Nonlinear perturbation equations

Given a solution $\tilde{u}(x, t)$ of (1.4), define the nonlinear perturbation variable

$$(4.1) \quad v = u - \bar{u} = \tilde{u}(x + \psi(x, t)) - \bar{u}(x),$$

where

$$(4.2) \quad u(x, t) := \tilde{u}(x + \psi(x, t))$$

and $\psi : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is to be chosen later.

Lemma 4.1. *For v, u as in (4.1),(4.2),*

$$(4.3) \quad u_t + f(u)_x - u_{xx} = (\partial_t - L)\bar{u}'(x_1)\psi(x, t) + \partial_x R + (\partial_t + \partial_x^2)S,$$

where

$$R := v\psi_t + v\psi_{xx} + (\bar{u}_x + v_x)\frac{\psi_x^2}{1 + \psi_x} = O(|v|(|\psi_t| + |\psi_{xx}|) + \left(\frac{|\bar{u}_x| + |v_x|}{1 - |\psi_x|}\right)|\psi_x|^2)$$

and

$$S := -v\psi_x = O(|v|(|\psi_x|).$$

Proof. To begin, notice from the definition of u in (4.2) we have by a straightforward computation

$$\begin{aligned} u_t(x, t) &= \tilde{u}_x(x + \psi(x, t), t)\psi_t(x, t) + \tilde{u}_t(x + \psi, t) \\ f(u(x, t))_x &= df(\tilde{u}(x + \psi(x, t), t))\tilde{u}_x(x + \psi, t) \cdot (1 + \psi_x(x, t)) \end{aligned}$$

and

$$\begin{aligned} u_{xx}(x, t) &= (\tilde{u}_{xx}(x + \psi(x, t), t) \cdot (1 + \psi_x(x, t)))_x \\ &= \tilde{u}_{xxx}(x + \psi(x, t), t) \cdot (1 + \psi_x(x, t)) + (\tilde{u}_{xx}(x + \psi(x, t), t) \cdot \psi_x(x, t))_x. \end{aligned}$$

Using the fact that $\tilde{u}_t + df(\tilde{u})\tilde{u}_x - \tilde{u}_{xx} = 0$, it follows that

$$(4.4) \quad \begin{aligned} u_t + f(u)_x - u_{xx} &= \tilde{u}_x\psi_t + df(\tilde{u})\tilde{u}_x\psi_x - \tilde{u}_{xx}\psi_x - (\tilde{u}_x\psi_x)_x \\ &= \tilde{u}_x\psi_t - \tilde{u}_t\psi_x - (\tilde{u}_x\psi_x)_x \end{aligned}$$

where it is understood that derivatives of \tilde{u} appearing on the righthand side are evaluated at $(x + \psi(x, t), t)$. Moreover, by another direct calculation, using the fact that $L(\bar{u}'(x)) = 0$ by translation invariance, we have

$$(\partial_t - L)\bar{u}'(x)\psi = \bar{u}_x\psi_t - \bar{u}_t\psi_x - (\bar{u}_x\psi_x)_x.$$

Subtracting, and using the facts that, by differentiation of $(\bar{u} + v)(x, t) = \tilde{u}(x + \psi, t)$,

$$(4.5) \quad \begin{aligned} \bar{u}_x + v_x &= \tilde{u}_x(1 + \psi_x), \\ \bar{u}_t + v_t &= \tilde{u}_t + \tilde{u}_x\psi_t, \end{aligned}$$

so that

$$(4.6) \quad \begin{aligned} \tilde{u}_x - \bar{u}_x - v_x &= -(\bar{u}_x + v_x)\frac{\psi_x}{1 + \psi_x}, \\ \tilde{u}_t - \bar{u}_t - v_t &= -(\bar{u}_x + v_x)\frac{\psi_t}{1 + \psi_x}, \end{aligned}$$

we obtain

$$u_t + f(u)_x - u_{xx} = (\partial_t - L)\bar{u}'(x)\psi + v_x\psi_t - v_t\psi_x - (v_x\psi_x)_x + \left((\bar{u}_x + v_x)\frac{\psi_x^2}{1 + \psi_x} \right)_x,$$

yielding (4.3) by $v_x\psi_t - v_t\psi_x = (v\psi_t)_x - (v\psi_x)_t$ and $(v_x\psi_x)_x = (v\psi_x)_{xx} - (v\psi_{xx})_x$. \square

Corollary 4.2. *The nonlinear residual v defined in (4.1) satisfies*

$$(4.7) \quad v_t - Lv = (\partial_t - L)\bar{u}'(x_1)\psi - Q_x + R_x + (\partial_t + \partial_x^2)S,$$

where

$$(4.8) \quad Q := f(\tilde{u}(x + \psi(x, t), t)) - f(\bar{u}(x)) - df(\bar{u}(x))v = \mathcal{O}(|v|^2),$$

$$(4.9) \quad R := v\psi_t + v\psi_{xx} + (\bar{u}_x + v_x)\frac{\psi_x^2}{1 + \psi_x},$$

and

$$(4.10) \quad S := -v\psi_x = \mathcal{O}(|v|(|\psi_x|).$$

Proof. Taylor expansion comparing (4.3) and $\bar{u}_t + f(\bar{u})_x - \bar{u}_{xx} = 0$. \square

4.2 Cancellation estimate

Our strategy in writing (4.7) is motivated by the following basic cancellation principle.

Proposition 4.3 ([HoZ]). *For any $f(y, s) \in L^p \cap C^2$ with $f(y, 0) \equiv 0$, there holds*

$$(4.11) \quad \int_0^t \int G(x, t-s; y) (\partial_s - L_y) f(y, s) dy ds = f(x, t).$$

Proof. Integrating the left hand side by parts, we obtain

$$(4.12) \quad \int G(x, 0; y) f(y, t) dy - \int G(x, t; y) f(y, 0) dy + \int_0^t \int (\partial_t - L_y)^* G(x, t-s; y) f(y, s) dy ds.$$

Noting that, by duality,

$$(\partial_t - L_y)^* G(x, t-s; y) = \delta(x-y) \delta(t-s),$$

$\delta(\cdot)$ here denoting the Dirac delta-distribution, we find that the third term on the righthand side vanishes in (4.12), while, because $G(x, 0; y) = \delta(x-y)$, the first term is simply $f(x, t)$. The second term vanishes by $f(y, 0) \equiv 0$. \square

4.3 Nonlinear damping estimate

Proposition 4.1. *Let $v_0 \in H^K$ (K as in (H1)), and suppose that for $0 \leq t \leq T$, the H^K norm of v and the $H^K(x, t)$ norms of ψ_t and ψ_x remain bounded by a sufficiently small constant. There are then constants $\theta_{1,2} > 0$ so that, for all $0 \leq t \leq T$,*

$$(4.13) \quad |v(t)|_{H^K}^2 \leq C e^{-\theta_1 t} |v(0)|_{H^K}^2 + C \int_0^t e^{-\theta_2(t-s)} \left(|v|_{L^2}^2 + |(\psi_t, \psi_x)|_{H^K(x,t)}^2 \right) (s) ds.$$

Proof. Subtracting from the equation (4.4) for u the equation for \bar{u} , we may write the nonlinear perturbation equation as

$$(4.14) \quad v_t + (df(\bar{u})v)_x - v_{xx} = Q(v)_x + \tilde{u}_x \psi_t - \tilde{u}_t \psi_x - (\tilde{u}_x \psi_x)_x,$$

where it is understood that derivatives of \tilde{u} appearing on the righthand side are evaluated at $(x + \psi(x, t), t)$. Using (4.6) to replace \tilde{u}_x and \tilde{u}_t respectively by $\bar{u}_x + v_x - (\bar{u}_x + v_x) \frac{\psi_x}{1+\psi_x}$ and $\bar{u}_t + v_t - (\bar{u}_x + v_x) \frac{\psi_t}{1+\psi_x}$, and moving the resulting $v_t \psi_x$ term to the lefthand side of (4.14), we obtain

$$(4.15) \quad \begin{aligned} (1 + \psi_x) v_t - v_{xx} = & -(df(\bar{u})v)_x + Q(v)_x + \bar{u}_x \psi_t \\ & - ((\bar{u}_x + v_x) \psi_x)_x + \left((\bar{u}_x + v_x) \frac{\psi_x^2}{1 + \psi_x} \right)_x. \end{aligned}$$

Taking the L^2 inner product in x of $\sum_{j=0}^K \frac{\partial_x^{2j} v}{1+\psi_x}$ against (4.15), integrating by parts, and rearranging the resulting terms, we arrive at the inequality

$$\partial_t |v|_{H^K}^2(t) \leq -\theta |\partial_x^{K+1} v|_{L^2}^2 + C \left(|v|_{H^K}^2 + |(\psi_t, \psi_x)|_{H^K(x,t)}^2 \right),$$

for some $\theta > 0$, $C > 0$, so long as $|\tilde{u}|_{H^K}$ remains bounded, and $|v|_{H^K}$ and $|(\psi_t, \psi_x)|_{H^K(x,t)}$ remain sufficiently small. Using the Sobolev interpolation $|v|_{H^K}^2 \leq |\partial_x^{K+1} v|_{L^2}^2 + \tilde{C} |v|_{L^2}^2$ for $\tilde{C} > 0$ sufficiently large, we obtain $\partial_t |v|_{H^K}^2(t) \leq -\tilde{\theta} |v|_{H^K}^2 + C \left(|v|_{L^2}^2 + |(\psi_t, \psi_x)|_{H^K(x,t)}^2 \right)$ from which (4.13) follows by Gronwall's inequality. \square

4.4 Integral representation/ ψ -evolution scheme

By Proposition 4.3, we have, applying Duhamel's principle to (4.7),

$$(4.16) \quad \begin{aligned} v(x, t) &= \int_{-\infty}^{\infty} G(x, t; y) v_0(y) dy \\ &+ \int_0^t \int_{-\infty}^{\infty} G(x, t-s; y) (-Q_y + R_x + S_t + S_{yy})(y, s) dy ds + \psi(t) \bar{u}'(x). \end{aligned}$$

Defining ψ implicitly as

$$(4.17) \quad \begin{aligned} \psi(x, t) &= - \int_{-\infty}^{\infty} e(x, t; y) u_0(y) dy \\ &- \int_0^t \int_{-\infty}^{+\infty} e(x, t-s; y) (-Q_y + R_x + S_t + S_{yy})(y, s) dy ds, \end{aligned}$$

following [ZH, Z4, MaZ2, MaZ3], where e is defined as in (3.24), and substituting in (4.16) the decomposition $G = \bar{u}'(x)e + \tilde{G}$ of Corollary 3.1, we obtain the *integral representation*

$$(4.18) \quad \begin{aligned} v(x, t) &= \int_{-\infty}^{\infty} \tilde{G}(x, t; y) v_0(y) dy \\ &+ \int_0^t \int_{-\infty}^{\infty} \tilde{G}(x, t-s; y) (-Q_y + R_x + S_t + S_{yy})(y, s) dy ds, \end{aligned}$$

and, differentiating (4.17) with respect to t , and recalling that $e(x, s; y) \equiv 0$ for $s \leq 1$,

$$(4.19) \quad \begin{aligned} \partial_t^j \partial_x^k \psi(x, t) &= - \int_{-\infty}^{\infty} \partial_t^j \partial_x^k e(x, t; y) u_0(y) dy \\ &- \int_0^t \int_{-\infty}^{+\infty} \partial_t^j \partial_x^k e(x, t-s; y) (-Q_y + R_x + S_t + S_{yy})(y, s) dy ds. \end{aligned}$$

Equations (4.18), (4.19) together form a complete system in the variables $(v, \partial_t^j \psi, \partial_x^k \psi)$, $0 \leq j \leq 1$, $0 \leq k \leq K$, from the solution of which we may afterward recover the shift ψ via (4.17). From the original differential equation (4.7) together with (4.19), we readily obtain short-time existence and continuity with respect to t of solutions $(v, \psi_t, \psi_x) \in H^K$ by a standard contraction-mapping argument based on (4.13), (4.17), and (3.26).

4.5 Nonlinear iteration

Associated with the solution (u, ψ_t, ψ_x) of integral system (4.18)–(4.19), define

$$(4.20) \quad \zeta(t) := \sup_{0 \leq s \leq t} |(v, \psi_t, \psi_x)|_{HK}(s)(1+s)^{1/4}.$$

Lemma 4.2. *For all $t \geq 0$ for which $\zeta(t)$ is finite, some $C > 0$, and $E_0 := |u_0|_{L^1 \cap HK}$,*

$$(4.21) \quad \zeta(t) \leq C(E_0 + \zeta(t)^2).$$

Proof. By (4.9)–(4.10) and definition (4.20),

$$(4.22) \quad |(Q, R, S)|_{L^1 \cap L^\infty} \leq |(v, v_x, \psi_t, \psi_x)|_{L^2}^2 + |(v, v_x, \psi_t, \psi_x)|_{L^\infty}^2 \leq C\zeta(t)^2(1+t)^{-\frac{1}{2}},$$

so long as $|\psi_x| \leq |\psi_x|_{HK} \leq \zeta(t)$ remains small, and likewise (using the equation to bound t derivatives in terms of x -derivatives of up to two orders)

$$(4.23) \quad |(\partial_t + \partial_x^2)S|_{L^1 \cap L^\infty} \leq |(v, \psi_x)|_{H^2}^2 + |(v, \psi_x)|_{W^{2,\infty}}^2 \leq C\zeta(t)^2(1+t)^{-\frac{1}{2}}.$$

Applying Corollary 3.1 with $q = 1$, $d = 1$ to representations (4.18)–(4.19), we obtain for any $2 \leq p < \infty$

$$(4.24) \quad \begin{aligned} |v(\cdot, t)|_{L^p(x)} &\leq C(1+t)^{-\frac{1}{2}(1-1/p)} E_0 \\ &\quad + C\zeta(t)^2 \int_0^t (1+t-s)^{-\frac{1}{2}(1/2-1/p)} (t-s)^{-\frac{3}{4}} (1+s)^{-\frac{1}{2}} ds \\ &\leq C(E_0 + \zeta(t)^2)(1+t)^{-\frac{1}{2}(1-1/p)} \end{aligned}$$

and

$$(4.25) \quad \begin{aligned} |(\psi_t, \psi_x)(\cdot, t)|_{W^{K,p}} &\leq C(1+t)^{-\frac{1}{2}} E_0 + C\zeta(t)^2 \int_0^t (1+t-s)^{-\frac{1}{2}(1-1/p)-1/2} (1+s)^{-\frac{1}{2}} ds \\ &\leq C(E_0 + \zeta(t)^2)(1+t)^{-\frac{1}{2}(1-1/p)}. \end{aligned}$$

Using (4.13) and (4.24)–(4.25), we obtain $|v(\cdot, t)|_{HK(x)} \leq C(E_0 + \zeta(t)^2)(1+t)^{-\frac{1}{4}}$. Combining this with (4.25), $p = 2$, rearranging, and recalling definition (4.20), we obtain (4.2). \square

Proof of Theorem 1.1. By short-time H^K existence theory, $\|(v, \psi_t, \psi_x)\|_{HK}$ is continuous so long as it remains small, hence η remains continuous so long as it remains small. By (4.2), therefore, it follows by continuous induction that $\eta(t) \leq 2C\eta_0$ for $t \geq 0$, if $\eta_0 < 1/4C$, yielding by (4.20) the result (1.15) for $p = 2$. Applying (4.24)–(4.25), we obtain (1.15) for $2 \leq p \leq p_*$ for any $p_* < \infty$, with uniform constant C . Taking $p_* > 4$ and estimating

$$|Q|_{L^2}, |R|_{L^2}, |S|_{L^2}(t) \leq |(v, \psi_t, \psi_x)|_{L^4}^2 \leq CE_0(1+t)^{-\frac{3}{4}}$$

in place of the weaker (4.22), then applying Corollary 3.1 with $q = 2$, $d = 1$, we obtain finally (1.15) for $2 \leq p \leq \infty$, by a computation similar (4.24)–(4.25); we omit the details of this final bootstrap argument. Estimate (1.16) then follows using (3.26) with $q = d = 1$, by

$$(4.26) \quad \begin{aligned} |\psi(t)|_{L^p} &\leq CE_0(1+t)^{\frac{1}{2p}} + C\zeta(t)^2 \int_0^t (1+t-s)^{-\frac{1}{2}(1-1/p)}(1+s)^{-\frac{1}{2}} ds \\ &\leq C(1+t)^{\frac{1}{2p}}(E_0 + \zeta(t)^2), \end{aligned}$$

together with the fact that $\tilde{u}(x, t) - \bar{u}(x) = v(x - \psi, t) + (\bar{u}(x) - \bar{u}(x - \psi))$, so that $|\tilde{u}(\cdot, t) - \bar{u}|$ is controlled by the sum of $|v|$ and $|\bar{u}(x) - \bar{u}(x - \psi)| \sim |\psi|$. This yields stability for $|u - \bar{u}|_{L^1 \cap H^K}|_{t=0}$ sufficiently small, as described in the final line of the theorem. \square

References

- [BHZ] B. Barker, J. Humpherys, and K. Zumbrun. *One-dimensional stability of parallel shock layers in isentropic magnetohydrodynamics*, Preprint (2007).
- [BLZ] B. Barker, O. Lafitte, and K. Zumbrun, *Existence and stability of viscous shock profiles for 2-D isentropic MHD with infinite electrical resistivity*, preprint (2009).
- [G] R. Gardner, *On the structure of the spectra of periodic traveling waves*, J. Math. Pures Appl. 72 (1993), 415-439.
- [GZ] R. Gardner and K. Zumbrun, *The Gap Lemma and geometric criteria for instability of viscous shock profiles*, Comm. Pure Appl. Math. 51 (1998), no. 7, 797–85.
- [GMWZ1] Guès, O., Métivier, G., Williams, M., and Zumbrun, K., *Existence and stability of multidimensional shock fronts in the vanishing viscosity limit*, Arch. Rat. Mech. Anal. 175. (2004), 151-244.
- [GMWZ2] O. Gues, G. Métivier, M. Williams, and K. Zumbrun, *Navier–Stokes regularization of multidimensional Euler shocks*. Ann. Sci. École Norm. Sup. (4) 39 (2006), no. 1, 75–175.
- [He] D. Henry, *Geometric theory of semilinear parabolic equations*, Lecture Notes in Mathematics, Springer–Verlag, Berlin (1981).
- [HoZ] D. Hoff and K. Zumbrun *Asymptotic behavior of multidimensional scalar viscous shock fronts*, Indiana Univ. Math. Journal, Vol. 49, No. 2 (2000).
- [HLZ] J. Humpherys, O. Lafitte, and K. Zumbrun. *Stability of viscous shock profiles in the high Mach number limit*, Comm. Math. Phys, to appear, 2009.

- [HLYZ1] J. Humpherys, G. Lyng, and K. Zumbrun. *Spectral stability of ideal-gas shock layers*, Arch. Ration. Mech. Anal., to appear, 2009.
- [HLYZ2] J. Humpherys, G. Lyng, and K. Zumbrun. *Multidimensional spectral stability of large-amplitude Navier–Stokes shocks*, In preparation.
- [JZ1] M. Johnson and K. Zumbrun, *Rigorous Justification of the Whitham Modulation Equations for the Generalized Korteweg-de Vries Equation*, preprint (2009).
- [JZ3] M. Johnson and K. Zumbrun, *Nonlinear stability and asymptotic behavior of periodic traveling waves of multidimensional viscous conservation laws in dimensions one and two*, preprint (2009).
- [JZB] M. Johnson, K. Zumbrun, and J. Bronski, *Bloch wave expansion vs. Whitham Modulation Equations for the Generalized Korteweg-de Vries Equation*, in preparation.
- [K] T. Kato, *Perturbation theory for linear operators*, Springer–Verlag, Berlin Heidelberg (1985).
- [MaZ2] C. Mascia and K. Zumbrun, *Stability of small-amplitude shock profiles of symmetric hyperbolic-parabolic systems*, Comm. Pure Appl. Math. 57 (2004), no. 7, 841–876.
- [MaZ3] C. Mascia and K. Zumbrun, *Pointwise Green function bounds for shock profiles of systems with real viscosity*. Arch. Ration. Mech. Anal. 169 (2003), no. 3, 177–263.
- [MaZ4] C. Mascia and K. Zumbrun, *Stability of large-amplitude viscous shock profiles of hyperbolic-parabolic systems*, Arch. Ration. Mech. Anal. 172 (2004), no. 1, 93–131.
- [OZ1] M. Oh and K. Zumbrun, *Stability of periodic solutions of viscous conservation laws with viscosity- 1. Analysis of the Evans function*, Arch. Ration. Mech. Anal. 166 (2003), no. 2, 99–166.
- [OZ2] M. Oh and K. Zumbrun, *Stability of periodic solutions of viscous conservation laws with viscosity- Pointwise bounds on the Green function*, Arch. Ration. Mech. Anal. 166 (2003), no. 2, 167–196.
- [OZ3] M. Oh, and K. Zumbrun, *Low-frequency stability analysis of periodic traveling-wave solutions of viscous conservation laws in several dimensions*, Journal for Analysis and its Applications, 25 (2006), 1–21.
- [OZ4] M. Oh, and K. Zumbrun, *Stability and asymptotic behavior of traveling-wave solutions of viscous conservation laws in several dimensions*, to appear, Arch. Ration. Mech. Anal.

- [Pa] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, Applied Mathematical Sciences, 44, Springer-Verlag, New York-Berlin, (1983) viii+279 pp. ISBN: 0-387-90845-5.
- [S1] G. Schneider, *Nonlinear diffusive stability of spatially periodic solutions– abstract theorem and higher space dimensions*, Proceedings of the International Conference on Asymptotics in Nonlinear Diffusive Systems (Sendai, 1997), 159–167, Tohoku Math. Publ., 8, Tohoku Univ., Sendai, 1998.
- [S2] G. Schneider, *Diffusive stability of spatial periodic solutions of the Swift-Hohenberg equation*, (English. English summary) Comm. Math. Phys. 178 (1996), no. 3, 679–702.
- [S3] G. Schneider, *Nonlinear stability of Taylor vortices in infinite cylinders*, Arch. Rat. Mech. Anal. 144 (1998) no. 2, 121–200.
- [Se1] D. Serre, *Spectral stability of periodic solutions of viscous conservation laws: Large wavelength analysis*, Comm. Partial Differential Equations 30 (2005), no. 1-3, 259–282.
- [Z4] K. Zumbrun, *Stability of large-amplitude shock waves of compressible Navier–Stokes equations*, with an appendix by Helge Kristian Jenssen and Gregory Lyng, in Handbook of mathematical fluid dynamics. Vol. III, 311–533, North-Holland, Amsterdam, (2004).
- [Z6] K. Zumbrun, *Dynamical stability of phase transitions in the p-system with viscosity-capillarity*, SIAM J. Appl. Math. 60 (2000), 1913-1929.
- [Z7] K. Zumbrun, *Instantaneous shock location and one-dimensional nonlinear stability of viscous shock waves*, preprint (2009).
- [ZH] K. Zumbrun and P. Howard, *Pointwise semigroup methods and stability of viscous shock waves*. Indiana Mathematics Journal V47 (1998), 741–871; Errata, Indiana Univ. Math. J. 51 (2002), no. 4, 1017–1021.