

Inverse problem for wave equation with sources and observations on disjoint sets

Matti Lassas and Lauri Oksanen

Department of Mathematics and Statistics, University of Helsinki, P.O. Box 68
FI-00014, Finland

E-mail: matti.lassas@helsinki.fi, lauri.oksanen@helsinki.fi

Abstract. We consider an inverse problem for a hyperbolic partial differential equation on a compact Riemannian manifold. Assuming that Γ_1 and Γ_2 are two disjoint open subsets of the boundary of the manifold we define the restricted Dirichlet-to-Neumann operator $\Lambda_{\Gamma_1, \Gamma_2}$. This operator corresponds the boundary measurements when we have smooth sources supported on Γ_1 and the fields produced by these sources are observed on Γ_2 . We show that when Γ_1 and Γ_2 are disjoint but their closures intersect at least at one point, then the restricted Dirichlet-to-Neumann operator $\Lambda_{\Gamma_1, \Gamma_2}$ determines the Riemannian manifold and the metric on it up to an isometry. In the Euclidian space, the result yields that an anisotropic wave speed inside a compact body is determined, up to a natural coordinate transformations, by measurements on the boundary of the body even when wave sources are kept away from receivers. Moreover, we show that if we have three arbitrary non-empty open subsets Γ_1, Γ_2 , and Γ_3 of the boundary, then the restricted Dirichlet-to-Neumann operators $\Lambda_{\Gamma_j, \Gamma_k}$ for $1 \leq j < k \leq 3$ determine the Riemannian manifold to an isometry. Similar result is proven also for the finite-time boundary measurements when the hyperbolic equation satisfies an exact controllability condition.

AMS classification scheme numbers: 35R30

Submitted to: *Inverse Problems*

Keywords: Inverse problems, wave equation, partial data.

1. Introduction and main results

Let M be a compact and connected C^∞ -smooth manifold of dimension n and let g be a C^∞ -smooth Riemannian metric on M . Let q be a real-valued C^∞ -smooth function on M , and denote by Δ_g the Laplace-Beltrami operator on M . We consider a hyperbolic inverse problem corresponding to the 2nd order elliptic operator

$$a(x, D) := -\Delta_g + q(x).$$

In local coordinates g is a positive-definite C^∞ -smooth matrix $(g_{jk}(x))_{j,k=1}^n$ with the inverse $(g^{jk}(x))_{j,k=1}^n$ and

$$a(x, D)u = -|g|^{-1/2} \sum_{j,k=1}^n \frac{\partial}{\partial x^j} \left(g^{jk} |g|^{1/2} \frac{\partial}{\partial x^k} u \right) + qu, \quad (1)$$

where $|g| := \det(g_{jk})$. Hence our results cover the setting, where $M \subset \mathbb{R}^n$ is an open domain with smooth boundary and $a(x, D)$ is an elliptic operator of the form (1).

Let $H^s(M)$ be the Sobolev space of $s \in \mathbb{N}$ times weakly differentiable functions on M , and let $H_0^1(M)$ be the $H^1(M)$ closure of $C_0^\infty(M)$, the space of smooth compactly supported functions. The operator

$$Au(x) := a(x, D)u, \quad D(A) := H^2(M) \cap H_0^1(M) \quad (2)$$

is self-adjoint in $L^2(M) = L^2(M, dV_g)$, where dV_g is the Riemannian volume measure. In local coordinates $dV_g = |g|^{1/2} dx$.

Denote by $v^f(x, t) = v(x, t)$ the solution of the initial boundary value problem

$$\begin{aligned} \partial_t^2 v + a(x, D)v &= 0 \quad \text{in } M \times (0, \infty), \\ v|_{\partial M \times (0, \infty)} &= f, \\ v|_{t=0} = \partial_t v|_{t=0} &= 0, \end{aligned} \quad (3)$$

for $f \in C_0^\infty(\partial M \times (0, \infty))$, and define the hyperbolic Dirichlet-to-Neumann operator

$$\Lambda : C_0^\infty(\partial M \times (0, \infty)) \rightarrow C^\infty(\partial M \times (0, \infty)), \quad \Lambda f := \partial_\nu v^f|_{\partial M \times \mathbb{R}_+},$$

where ∂_ν is the normal derivative on ∂M . In local coordinates the exterior conormal ν is the covector (ν_1, \dots, ν_n) with

$$\sum_{j,k=1}^n \nu_j(x) g^{jk}(x) \nu_k(x) = 1, \quad x \in \partial M,$$

and

$$\partial_\nu = \sum_{j,k=1}^n \nu_j g^{jk} \frac{\partial}{\partial x^k}.$$

Denote by $\Lambda_{\Gamma_1, \Gamma_2}^T$ the restriction of the Dirichlet-to-Neumann operator

$$\Lambda_{\Gamma_1, \Gamma_2}^T : C_0^\infty(\Gamma_1 \times (0, T)) \rightarrow C^\infty(\Gamma_2 \times (0, T)),$$

where $\Gamma_1, \Gamma_2 \subset \partial M$ are open. Furthermore, denote $\Lambda_{\Gamma_1, \Gamma_2}^\infty := \Lambda_{\Gamma_1, \Gamma_2}^\infty$.

It is well known, that the map Λ determine the manifold (M, g) up to an isometry [5]. This is also true for the restriction $\Lambda_{\Gamma, \Gamma}^T$ when Γ is nonempty and T is sufficiently large [24].

In many applications observations of physical fields can not be done on the same locations where the sources of the fields are. For instance, in imaging in Earth Sciences, elastic or acoustic fields are often implemented using explosions [40, 38]. In such a case observation devices need to be far away from the sources.

Similarly, in electromagnetic imaging, it is technically difficult to use electrodes at the same time as sources and for making observations. These are typical examples

of cases where the observation devices and the sources of the fields are supported on disjoint sets.

In this paper we show, that for certain collections of pairs (Γ_1, Γ_2) of open and disjoint subsets of ∂M , the operators $\Lambda_{\Gamma_1, \Gamma_2}^T$ determine the manifold (M, g) up to an isometry.

Theorem 1. *Let $\Gamma_1, \Gamma_2, \Sigma \subset \partial M$ be open sets such that $\bar{\Gamma}_1, \bar{\Gamma}_2 \subset \Sigma$ and $\bar{\Gamma}_1 \cap \bar{\Gamma}_2 \neq \emptyset$. Then Σ , given as a smooth manifold, and the operator $\Lambda_{\Gamma_1, \Gamma_2}$ determine the manifold (M, g) up to an isometry.*

Theorem 2. *Let $\Gamma_1, \Gamma_2, \Gamma_3 \subset \partial M$ be open and nonempty. Then the smooth manifolds Γ_p , $p = 1, 2, 3$, and the operators*

$$\Lambda_{\Gamma_1, \Gamma_2}, \Lambda_{\Gamma_1, \Gamma_3}, \Lambda_{\Gamma_2, \Gamma_3}$$

determine the manifold (M, g) up to an isometry.

For measurements on a finite time interval, we prove a theorem similar to Theorem 2 under an additional controllability assumption:

(A) For any $w \in L^2(M)$ there is a boundary value $f \in L^2(\partial M \times (0, \infty))$ satisfying

$$v^f(T/2) = w, \quad \text{supp}(f) \subset \Gamma_3 \times (0, T),$$

where v^f is the solution of the equation (3) and $\Gamma_3 \subset \partial M$.

Let us comment the controllability assumption (A) when M is embedded in \mathbb{R}^n . Property (A) follows from the geometric control condition of Bardos, Lebeau and Rauch [2], which yields exact controllability of the wave equation. Property (A) follows also from existence of a strictly convex function h on \bar{M} with respect to the Riemannian metric g .

Suppose that $h \in C^2(\bar{M})$ is strictly convex and that $\rho > 0$ is a lower bound for the Hessian of h in the Riemannian metric g , that is

$$D^2h(X, X) \geq \rho|X|_g, \quad X \in T_x M, \quad x \in M.$$

By [31], (A) holds if

$$T > \frac{4}{\rho} \max_{x \in \bar{M}} |\nabla_g h|_g, \quad \sup_{x \in \Gamma_3} \nabla_g h(x) \cdot \bar{\nu}(x) \leq 0,$$

where ∇_g and $|\cdot|_g$ are the gradient and length with respect to the Riemannian metric g , $\bar{\nu}$ is the Euclidean unit outward normal to $\partial M \subset \mathbb{R}^n$ and $\nabla_g h \cdot \bar{\nu}$ is the Euclidean inner product. We refer to [31] for examples of Riemannian manifolds (M, g) having a strictly convex function h .

Theorem 3. *Let $\Gamma_1, \Gamma_2, \Gamma_3 \subset \partial M$ be open and nonempty. If the controllability assumption (A) holds for Γ_3 and $T > 0$, and*

$$4d(x, y) < T, \quad x \in \Gamma_p, \quad y \in M, \quad p = 1, 2, 3,$$

then the Riemannian manifolds $(\Gamma_p, g|_{\Gamma_p})$, $p = 1, 2, 3$, and the operators

$$\Lambda_{\Gamma_1, \Gamma_2}^T, \Lambda_{\Gamma_1, \Gamma_3}^T, \Lambda_{\Gamma_2, \Gamma_3}^T$$

determine the manifold (M, g) up to an isometry.




Figure 1. On left, the measurements $\Lambda_{\Gamma_1, \Gamma_2}$, $\Lambda_{\Gamma_1, \Gamma_3}$, $\Lambda_{\Gamma_2, \Gamma_3}$ are shown as arrows pointing from the support of sources to the support of observations. Using Lemma 1 we can change the direction of any arrow in the picture on left. Hence also the measurements shown on right are covered by Theorems 2 and 3.

The proofs of these theorems consist of showing that the data determine, up to a gauge transformation, the boundary spectral data of the operator A on a part of the boundary. The manifold is then determined up to an isometry, as can be seen using the boundary control method [3, 5, 24, 25].

Notice also, that the operator $\Lambda_{\Gamma_1, \Gamma_2}^T$ determines the operator $\Lambda_{\Gamma_2, \Gamma_1}^T$ by a time reversal argument.

Lemma 1. *Let $\Gamma_1, \Gamma_2 \subset \partial M$ be open and nonempty. Let $T > 0$, and define the time reversal operator $Rf(x, t) := f(x, T-t)$. If $f \in C_0^\infty(\Gamma_1 \times (0, T))$ and $h \in C_0^\infty(\Gamma_2 \times (0, T))$, then*

$$(f, \Lambda_{\Gamma_2, \Gamma_1}^T h)_{L^2(\partial M \times (0, T))} = (R\Lambda_{\Gamma_1, \Gamma_2}^T Rf, h)_{L^2(\partial M \times (0, T))}.$$

Hence $(\Gamma_j, g|_{\Gamma_j})$, $j = 1, 2$, given as Riemannian manifolds, and the operator $\Lambda_{\Gamma_1, \Gamma_2}^T$ determine the operator $\Lambda_{\Gamma_2, \Gamma_1}^T = (R\Lambda_{\Gamma_1, \Gamma_2}^T R)^t$.

This result is relatively well known, see e.g. [6, 12], but for the sake of completeness, we will give a proof in the appendix.

Let us review previous results on the topic. The inverse problem for isotropic wave equation on a compact manifold with measurements on the whole boundary was solved by Belishev and Kurylev [5]. This was based on the boundary control method originally developed in [3] for wave equation on a bounded domain of \mathbb{R}^n . The inverse problems for more general hyperbolic equations on a compact Riemannian manifold with sources and observations on the same open subset Γ of the boundary has been studied by Katchalov and Kurylev [24], see also [29]. Similar problem has recently been studied for non-compact manifolds in [23, 26].

Inverse problems for elliptic equations with data on a part of the boundary have been studied intensively as they are the natural generalization of the Calderón's inverse problem for the conductivity equation [10].

When measurements are given on the whole boundary, the inverse problem for Schrödinger equation on a bounded domain of \mathbb{R}^n , $n \geq 3$, and hence for isotropic conductivity equation, was solved by Sylvester and Uhlmann in [39]. The corresponding two dimensional problem for isotropic conductivity equation was solved first by Nachman in [36] for C^2 conductivities, and for L^∞ conductivities, for which Calderón's inverse problem was originally posed, by Astala and Päivärinta in [1]. Recently, also the inverse problem for Schrödinger equation on a bounded domain of dimension two with measurements on the whole boundary was solved by Bukgheim in [8]. The corresponding problem on a compact Riemannian surface was later solved in [17].

The inverse problem for Schrödinger equation on a bounded domain of \mathbb{R}^n , $n \geq 3$, with observations on an open subset Γ of the boundary was solved in [28]. The inverse problem for Schrödinger equation on a bounded domain of \mathbb{R}^n , $n = 2$, with sources and observations on the same open subset Γ of the boundary was solved by Imanuvilov, Uhlmann and Yamamoto in [21]. The corresponding problem on a compact Riemannian surface was later solved in [16]. For related results with measurements on a part of the boundary, see [9, 15, 22].

The inverse problem for the Laplace-Beltrami operator Δ_g on a compact Riemannian manifold with sources and observations on the same open subset Γ of the boundary has been studied on analytic Riemannian manifolds of dimension $n \geq 3$ in [34, 32], and on Riemannian surfaces in [33], see also [18, 19]. The inverse problem for the Laplace-Beltrami operator in dimensions $n \geq 3$ is open in general, even when measurements are given on the whole boundary. For positive results under certain geometrical conditions see [13].

2. Spectral analysis of the data

Denote by $(\lambda_j)_{j \in \mathbb{N}}$ the increasing sequence of distinct eigenvalues of the operator A and let $(\phi_k)_{k \in \mathbb{N}}$ be an orthonormal basis of real-valued C^∞ -smooth eigenfunctions. Moreover, let $(I_j)_{j \in \mathbb{N}}$ be a partition of \mathbb{N} such that $(\phi_k)_{k \in I_j}$ is a basis for the space of eigenfunctions corresponding λ_j .

Let $f \in C_c^\infty(\partial M \times (0, \infty))$, and consider Λf also as a function in $C^\infty(\partial M \times \mathbb{R})$ by defining $\Lambda f(\cdot, t) = 0$ for $t \leq 0$. There is a constant $C > 0$ such that for $x \in \partial M$, the Fourier transform $\mathcal{F}_{t \rightarrow \tau} \Lambda f(x)$ is an analytic function of τ when $\text{Im } \tau < -C$. It is known (see e.g. [25]), that $\mathcal{F}_{t \rightarrow \tau} \Lambda f(x)$ extends to a meromorphic function of $\tau \in \mathbb{C}$, and that it may have poles only at points $\sqrt{\lambda_j}$. Moreover, the residues at these points are

$$\text{res}_{\tau=\sqrt{\lambda_j}} \mathcal{F}_{t \rightarrow \tau} \Lambda f(x) = \sum_{k \in I_j} \left(\hat{f}(\cdot, \sqrt{\lambda_j}), \partial_\nu \phi_k \right)_{L^2(\partial M, dS_g)} \partial_\nu \phi_k(x),$$

where $\hat{f}(x, \tau) = (\mathcal{F}_{t \rightarrow \tau} f)(x, \tau)$ and dS_g is the Riemannian surface measure.

If $j \in \mathbb{N}$, a_k are constants for $k \in I_j$, and the linear combination

$$\sum_{k \in I_j} a_k \partial_\nu \phi_k = \partial_\nu \left(\sum_{k \in I_j} a_k \phi_k \right)$$

vanish on a nonempty open subset of ∂M , then $a_k = 0$ for all $k \in I_j$ by unique continuation, see e.g. [35]. Hence for an open nonempty set $\Gamma \subset \partial M$ and $j \in \mathbb{N}$, the functions $(\partial_\nu \phi_k|_\Gamma)_{k \in I_j}$ are linearly independent.

Let $\Gamma_1, \Gamma_2 \subset \partial M$ be open and nonempty. By linear independence and smoothness of $\partial_\nu \phi_k$, there are $f \in C_0^\infty(\Gamma_1)$ and $x \in \Gamma_2$ such that

$$\sum_{k \in I_j} (f, \partial_\nu \varphi_k)_{L^2(\partial M, dS_g)} \partial_\nu \varphi_k(x) \neq 0.$$

Moreover, for fixed $\tau \in \mathbb{C}$, the map $f \mapsto \widehat{f}(\cdot, \tau)$ from $C_0^\infty(\Gamma_1 \times (0, \infty))$ to $C_0^\infty(\Gamma_1)$ is surjective.

Hence the operator $\Lambda_{\Gamma_1, \Gamma_2}$ determines the eigenvalues λ_j and the operators

$$\begin{aligned} L_{\Gamma_1, \Gamma_2; j} &: C_c^\infty(\Gamma_1 \times (0, \infty)) \rightarrow C^\infty(\Gamma_2 \times (0, \infty)), \\ L_{\Gamma_1, \Gamma_2; j} f &:= \sum_{k \in I_j} (f, \partial_\nu \varphi_k)_{L^2(\partial M, dS_g)} \partial_\nu \varphi_k|_{\Gamma_2}. \end{aligned} \quad (4)$$

3. Inverse problem with disjoint sources and observations

In this section we prove Theorem 1.

Lemma 2. *Suppose that $f, h \in C^\infty(\mathbb{R})$ are such that*

$$\partial_x^j \partial_y^k (h(x)f(x)f(y))|_{x=0, y=0} = \partial_x^k \partial_y^j (h(x)f(x)f(y))|_{x=0, y=0}$$

for all $j, k \in \mathbb{N}$. Then $\partial^j f(0) = 0$ for all $j \in \mathbb{N} := \{0, 1, 2, \dots\}$ or $\partial^k h(0) = 0$ for all positive $k \in \mathbb{N}$.

Proof. Assume that the claim is not valid. Then there exist $j \in \mathbb{N}$ and $k \in \mathbb{N} \setminus \{0\}$ such that $\partial^j f(0) \neq 0$ and $\partial^k h(0) \neq 0$. Let us next consider the smallest such integers j and k .

By Leibniz's formula

$$\begin{aligned} 0 &= \partial_x^{j+k} \partial_y^j (h(x)f(x)f(y))|_{x=0, y=0} - \partial_x^j \partial_y^{j+k} (h(x)f(x)f(y))|_{x=0, y=0} \\ &= \sum_{l=0}^{j+k} \binom{j+k}{l} \partial^l h(0) \partial^{j+k-l} f(0) \partial^j f(0) \\ &\quad - \sum_{m=0}^j \binom{j}{m} \partial^m h(0) \partial^{j-m} f(0) \partial^{j+k} f(0) \\ &= S_1 + \binom{j+k}{k} \partial^k h(0) \partial^j f(0) \partial^j f(0) + S_2 - S_3, \end{aligned}$$

where

$$\begin{aligned} S_1 &:= \sum_{l=1}^{k-1} \binom{j+k}{l} \partial^l h(0) \partial^{j+k-l} f(0) \partial^j f(0), \\ S_2 &:= \sum_{l=k+1}^{j+k} \binom{j+k}{l} \partial^l h(0) \partial^{j+k-l} f(0) \partial^j f(0), \end{aligned}$$

$$S_3 := \sum_{m=1}^j \binom{j}{m} \partial^m h(0) \partial^{j-m} f(0) \partial^{j+k} f(0),$$

and the terms with indices $l = 0$ and $m = 0$ have cancelled each other out.

As k is the smallest positive integer such that $\partial^k h(0) \neq 0$, we have $\partial^l h(0) = 0$ in the sum S_1 , and so $S_1 = 0$. As j is the smallest integer such that $\partial^j f(0) \neq 0$, we have $\partial^{j+k-l} f(0) = 0$ in the sum S_2 and $\partial^{j-m} f(0) = 0$ in the sum S_3 , thus $S_2 = S_3 = 0$.

Hence $\partial^k h(0)(\partial^j f(0))^2 = 0$, which is a contradiction with the assumption that $\partial^j f(0) \neq 0$ and $\partial^k h(0) \neq 0$. This proves the claim. \square

In the proof of the next lemma we use the equation

$$\partial_t^j f(tv) = \sum_{|\alpha|=j} \frac{j!}{\alpha!} \partial^\alpha f(tv) v^\alpha, \quad (5)$$

where $f \in C^\infty(\mathbb{R}^n)$, $t \in \mathbb{R}$, $v \in \mathbb{R}^n$ and $j \in \mathbb{N}$,

Lemma 3. *Suppose that $f, h \in C^\infty(\mathbb{R}^n)$ are such that*

$$\partial_x^\alpha \partial_y^\beta (h(x)f(x)f(y))|_{x=0,y=0} = \partial_x^\beta \partial_y^\alpha (h(x)f(x)f(y))|_{x=0,y=0}$$

for all multi-indices $\alpha, \beta \in \mathbb{N}^n$. Then $\partial^\alpha f(0) = 0$ for all multi-indices $\alpha \in \mathbb{N}^n$ or $\partial^\beta h(0) = 0$ for all nonzero multi-indices $\beta \in \mathbb{N}^n$.

Proof. Let $j, k \in \mathbb{N}$ and $v \in \mathbb{R}^n$. By (5)

$$\begin{aligned} & \partial_t^j \partial_s^k (h(tv)f(tv)f(sv))|_{t=0,s=0} - \partial_t^k \partial_s^j (h(tv)f(tv)f(sv))|_{t=0,s=0} \\ &= \sum_{|\alpha|=j} \sum_{|\beta|=k} \frac{j!}{\alpha!} \frac{k!}{\beta!} v^\alpha v^\beta (\partial^\alpha (hf)(0) \partial^\beta f(0) - \partial^\beta (hf)(0) \partial^\alpha f(0)) = 0. \end{aligned}$$

Hence

$$\partial_t^j \partial_s^k (h(tv)f(tv)f(sv))|_{t=0,s=0} = \partial_t^k \partial_s^j (h(tv)f(tv)f(sv))|_{t=0,s=0}$$

for all $j, k \in \mathbb{N}$ and $v \in \mathbb{R}^n$.

Define the sets

$$F := \{v \in \mathbb{R}^n : \partial_t^j f(tv)|_{t=0} = 0 \text{ for all } j \in \mathbb{N}\},$$

$$H := \{v \in \mathbb{R}^n : \partial_t^k h(tv)|_{t=0} = 0 \text{ for all positive } k \in \mathbb{N}\}.$$

The sets F and H are closed by smoothness of f and h , respectively. Lemma 2 gives that $F \cup H = \mathbb{R}^n$. If $F \neq \mathbb{R}^n$, then $\mathbb{R}^n \setminus F$ is open, nonempty and contained in H . Thus F or H contains an open nonempty subset.

Suppose that $U \subset F$ is open and nonempty. Let $j \in \mathbb{N}$, and define the polynomial

$$p(v) := \sum_{|\alpha|=j} \frac{j!}{\alpha!} \partial^\alpha f(0) v^\alpha.$$

By (5), $p(v) = \partial_t^j f(tv)|_{t=0}$, and p vanish in U . Using unique continuation for real analytic functions we see that $p = 0$ in \mathbb{R}^n , and so the coefficients of p vanish. As j can be chosen freely, $\partial^\alpha f(0) = 0$ for all multi-indices α .

Similarly, if there exists an open and nonempty $V \subset H$, then $\partial^\alpha h(0) = 0$ for all nonzero multi-indices α . \square

Remark 1. Let U be a C^∞ -smooth manifold of dimension n , $f \in C^\infty(U)$ and $p \in U$. If $\partial^\alpha f(0) = 0$ for all multi-indices $\alpha \in \mathbb{N}^n$ in some local coordinates taking p to 0, then $\partial^\alpha f(0) = 0$ for all multi-indices $\alpha \in \mathbb{N}^n$ in all local coordinates taking p to 0.

Lemma 4. Let ϕ be an eigenfunction of the operator A corresponding to an eigenvalue λ , and let $p_0 \in \partial M$. Then in any local coordinates of ∂M taking p_0 to 0, there is a multi-index $\alpha \in \mathbb{N}^{n-1}$ such that $\partial^\alpha \partial_\nu \phi(0) \neq 0$.

Proof. Assume that the claim is not valid. Then $\partial^\alpha \partial_\nu \phi(0) = 0$ for all $\alpha \in \mathbb{N}^{n-1}$ in some local coordinates of ∂M taking p_0 to 0.

Consider boundary normal coordinates of \overline{M} taking p_0 to 0. We may suppose that the coordinates map a small neighborhood V of p_0 onto $B(0, \epsilon) \times [0, \epsilon]$, where $B(0, \epsilon) \subset \mathbb{R}^{n-1}$ is a ball of radius $\epsilon > 0$ centered at the origin. Then these coordinates take a boundary point $p' \in \partial M \cap V$ to a point $(x', 0) \in B(0, \epsilon) \times \{0\}$, where $x' = (x^1, \dots, x^{n-1})$.

The special property of the boundary normal coordinates is, that a point $p \in \overline{M} \cap V$ has coordinates $(x', x^n) \in B(0, \epsilon) \times [0, \epsilon]$, where $x^n = d(p, \partial M)$ and x' are the coordinates of the unique boundary point $p' \in \partial M$ such that $d(p, p') = d(p, \partial M)$.

Moreover, in the coordinates (x', x^n) the equation

$$(-\Delta_g + q)\phi = \lambda\phi,$$

has the form

$$-\partial_{x^n}^2 \phi - \sum_{j,k=1}^{n-1} g^{jk} \partial_{x^j} \partial_{x^k} \phi + \sum_{j=1}^n a^j \partial_{x^j} \phi + a^0 \phi = \lambda\phi, \quad (6)$$

for some $a^j \in C^\infty(B(0, \epsilon) \times [0, \epsilon])$, $j = 0, \dots, n$, see e.g. [11].

Let us show, that $\phi = 0$. Let $b \in \mathbb{N}$ and $\alpha \in \mathbb{N}^{n-1}$. By applying the operator $\partial_{x'}^\alpha \partial_{x^n}^b$ on the both sides of (6), we get

$$\partial_{x'}^\alpha \partial_{x^n}^{b+2} \phi(0) = \partial_{x'}^\alpha \partial_{x^n}^b \left(- \sum_{j,k=1}^{n-1} g^{jk} \partial_{x^j} \partial_{x^k} \phi + \sum_{j=1}^n a^j \partial_{x^j} \phi + a^0 \phi - \lambda\phi \right) |_{x=0}.$$

The right hand side of this equation is a linear combination of functions

$$\partial_{x'}^{\alpha'} \partial_{x^n}^{b'} \phi, \quad \alpha' \in \mathbb{N}, b' \leq b+1$$

at the point $x = 0$.

The equations $\partial_{x'}^\alpha \phi(0) = 0$ hold for all $\alpha \in \mathbb{N}^{n-1}$ by the boundary condition $\phi|_{\partial M} = 0$. Furthermore, we have by Remark 1, that $\partial_{x'}^\alpha \partial_{x^n}^b \phi(0) = 0$ for all $\alpha \in \mathbb{N}^{n-1}$. Using induction we see that $\partial_{x'}^\alpha \partial_{x^n}^b \phi(0) = 0$ for all $\alpha \in \mathbb{N}^{n-1}$ and $b \in \mathbb{N}$.

Define the odd and even reflection operators

$$R_o f(x', x^n) := (\text{sign } x^n) f(x', |x^n|), \quad R_e f(x', x^n) := f(x', |x^n|),$$

where $f \in C(B(0, \varepsilon) \times [0, \varepsilon))$, $x' \in B(0, \varepsilon)$ and $x^n \in (-\varepsilon, \varepsilon)$.

Also, define $\tilde{\phi} := R_o \phi$, $\tilde{g}^{jk} := R_e g^{jk}$, $\tilde{a}^n := R_o a^n$ and $\tilde{a}^j := R_e a^j$ for $j = 0, \dots, n-1$. Denote $U := B(0, \varepsilon) \times (-\varepsilon, \varepsilon)$.

As $\phi|_{x^n=0} = 0$, we see that $\tilde{\phi} \in H^2(U)$ and

$$\partial_x^\alpha \partial_{x^n}^b \tilde{\phi}(x', x^n) = (\text{sign } x^n)^{b+1} \partial_x^\alpha \partial_{x^n}^b \phi(x', |x^n|), \quad |\alpha| + b \leq 2.$$

Moreover, \tilde{g}^{jk} is Lipschitz continuous in U , $a^j \in L^\infty(U)$ for $j = 0, \dots, n$, and

$$-\partial_{x^n}^2 \tilde{\phi} - \sum_{j,k=1}^{n-1} \tilde{g}^{jk} \partial_{x^j} \partial_{x^k} \tilde{\phi} + \sum_{j=1}^n \tilde{a}^j \partial_{x^j} \tilde{\phi} + \tilde{a}^0 \tilde{\phi} = \lambda \tilde{\phi},$$

where the both sides are considered as functions in $L^2(U)$. Hence for some constant $C > 0$

$$|\partial_{x^n}^2 \tilde{\phi} + \sum_{j,k=1}^{n-1} \tilde{g}^{jk} \partial_{x^j} \partial_{x^k} \tilde{\phi}| \leq C \sum_{|\alpha|+b \leq 1} |\partial_x^\alpha \partial_{x^n}^b \tilde{\phi}|, \quad \text{in } U.$$

Since $\phi \in C^\infty(B(0, \varepsilon) \times [0, \varepsilon))$ vanishes up to arbitrary degree in origin, Taylor's formula gives for any $m \in \mathbb{N}$ a constant $C_m > 0$ such that

$$\int_{B(0,r)} \int_0^r |\phi(x)|^2 dx^n dx' \leq \int_{B(0,r)} \int_0^r C_m r^m dx^n dx', \quad \text{as } r \rightarrow 0.$$

Hence for any $m \in \mathbb{N}$, there is a constant $C'_m > 0$ such that

$$\int_{B(0,r)} \int_{-r}^r |\tilde{\phi}(x)|^2 dx^n dx' \leq C'_m r^m, \quad \text{as } r \rightarrow 0.$$

By Hörmander's strong unique continuation result [20] this yields, that $\tilde{\phi} = 0$ in U . In particular, $\phi = 0$ around some point $q \in M$. As M is connected, unique continuation gives that $\phi = 0$ in M . This is a contradiction with the assumption that ϕ is an eigenfunction, and the claim is proved. \square

Remark 2. Let X and Y be Hilbert spaces, and let $u_1, \dots, u_N \in X$ and $v_1, \dots, v_N \in Y$ be linearly independent. Suppose that $D \subset X$ is a dense subspace, and define

$$L : D \rightarrow Y, \quad Lf := \sum_{k=1}^N (f, u_k)_X v_k.$$

Then L determines the unique bounded extension $\tilde{L} : X \rightarrow Y$, and its adjoint $\tilde{L}^* : Y \rightarrow X$. Hence, L determines the spaces

$$\text{span}(v_1, \dots, v_N) = \tilde{L}(X), \quad \text{span}(u_1, \dots, u_N) = \tilde{L}^*(Y).$$

Theorem 4. Let $\Gamma_1, \Gamma_2, \Sigma \subset \partial M$ be open, $\bar{\Gamma}_1, \bar{\Gamma}_2 \subset \Sigma$, and $\bar{\Gamma}_1 \cap \bar{\Gamma}_2 \neq \emptyset$. Then the smooth manifold Σ and the collection

$$\{(\lambda_j, L_{\Gamma_1, \Gamma_2; j}) \mid j \in \mathbb{N}\} \tag{7}$$

determine boundary spectral data up to a constant gauge transformation on Γ_2 . That is, one can find a collection

$$\{(\lambda_j, (\partial_\nu \psi_k|_{\Gamma_2})_{k \in I_j}) \mid j \in \mathbb{N}\}, \quad (8)$$

where for an unknown constant $C > 0$ not depending on j or k , $(C\psi_k)_{k \in I_j}$ is an orthonormal basis of eigenfunctions in $L^2(M)$ corresponding the eigenvalue λ_j .

Proof. Choose a smooth positive measure $d\mu$ on Σ . Then there is a positive function $\eta \in C^\infty(\Sigma)$ such that $\eta d\mu = dS_g|_\Sigma$. As $\eta > 0$, the functions $(\eta \partial_\nu \phi_k|_{\Gamma_1})_{k \in I_j}$ are linearly independent for all $j \in \mathbb{N}$.

For all $j \in \mathbb{N}$, denote $L_j := L_{\Gamma_1, \Gamma_2; j}$, and define

$$l_j(x, y) := \sum_{k \in I_j} \eta(x) \partial_\nu \phi_k(x) \partial_\nu \phi_k(y), \quad x, y \in \partial M,$$

$$\tilde{E}_j^1 := \text{span}(\eta \partial_\nu \phi_k|_{\Gamma_1})_{k \in I_j}, \quad E_j^2 := \text{span}(\partial_\nu \phi_k|_{\Gamma_2})_{k \in I_j}.$$

Note, that for the smallest eigenvalue λ_0 , the space of eigenfunctions is one dimensional (see e.g. [14, Thm 6.5.2]), and so

$$l_0(x, y) = \eta(x) \partial_\nu \phi_0(x) \partial_\nu \phi_0(y).$$

Consider a positive function $\tilde{\eta} \in C^\infty(\Sigma)$ and real-valued functions $e_k \in C^\infty(\Sigma)$, $k \in \mathbb{N}$, such that the following three conditions hold:

(A1) If $x_0 \in \bar{\Gamma}_1 \cap \bar{\Gamma}_2$, then in local coordinates of Σ taking x_0 to 0

$$\partial_x^\alpha \partial_y^\beta \left(\frac{l_0(x, y)}{\tilde{\eta}(x)} \right) \Big|_{x=0, y=0} = \partial_x^\beta \partial_y^\alpha \left(\frac{l_0(x, y)}{\tilde{\eta}(x)} \right) \Big|_{x=0, y=0}$$

for all multi-indices $\alpha, \beta \in \mathbb{N}^{n-1}$.

(A2) $\text{span}(\tilde{\eta} e_k|_{\Gamma_1})_{k \in I_j} = \tilde{E}_j^1$ and $\text{span}(e_k|_{\Gamma_2})_{k \in I_j} = E_j^2$ for all $j \in \mathbb{N}$.

(A3) $\tilde{L}_j = L_j$ for all $j \in \mathbb{N}$, where

$$\tilde{L}_j f(y) := \sum_{k \in I_j} (f, \tilde{\eta} e_k)_{L^2(\partial M, d\mu)} e_k(y), \quad f \in C_0^\infty(\Gamma_1), \quad y \in \Gamma_2.$$

Such functions $\tilde{\eta}$ and e_k exist. For example, $\tilde{\eta} = \eta$ and $e_k = \partial_\nu \phi_k|_\Sigma$ satisfy the conditions.

Next we show the following two statements.

- (i) We can verify using the data (7), whether any given functions $\tilde{\eta} \in C^\infty(\Sigma)$ and $e_k \in C^\infty(\Sigma)$, $k \in \mathbb{N}$, satisfy the conditions (A1), (A2) and (A3).
- (ii) There is an orthonormal basis of eigenfunctions $(\psi_k)_{k \in \mathbb{N}}$ of operator A and a constant $C > 0$, not depending on k , such that

$$e_k|_{\Gamma_2} = C \partial_\nu \psi_k|_{\Gamma_2}, \quad k \in \mathbb{N}.$$

If (i) holds, then the data (7) determine the nonempty collection

$$\{(e_k|_{\Gamma_2})_{k \in \mathbb{N}} \mid (A1)-(A3) \text{ hold with a positive } \tilde{\eta} \in C^\infty(\Sigma)\},$$

and if (ii) holds, then any element from this collection determines a collection of type (8). So the claim of the theorem is proved after proving (i) and (ii).

Let us show the claim (i). Clearly, the condition (A3) can be verified using the data (7). As

$$L_j f(y) = (f, \sum_{k \in I_j} (\partial_\nu \phi_k(y)) \eta \partial_\nu \phi_k)_{L^2(\partial M, d\mu)}, \quad f \in C_0^\infty(\Gamma_1), y \in \Gamma_2,$$

by varying $f \in C_0^\infty(\Gamma_1)$, we see that the map L_j determines the function $l_j|_{\Gamma_1 \times \Gamma_2}$.

Let $x_0 \in \bar{\Gamma}_1 \cap \bar{\Gamma}_2$. Given $l_0|_{\Gamma_1 \times \Gamma_2}$ and $\tilde{\eta}$, it is possible to compute

$$\partial_x^\alpha \partial_y^\beta \left(\frac{l_0(x, y)}{\tilde{\eta}(x)} \right), \quad (x, y) \in \Gamma_1 \times \Gamma_2, \quad \alpha, \beta \in \mathbb{N}^{n-1},$$

in any local coordinates Σ taking x_0 to 0. By smoothness of η , $\tilde{\eta}$ and $\partial_\nu \phi_0$, these functions are known also at $(x, y) = (0, 0)$. Hence the condition (A1) can be verified using the data (7).

Taking $X = L^2(\Gamma_1, d\mu)$, $Y = L^2(\Gamma_2, d\mu)$ and $D = C_0^\infty(\Gamma_1)$ in the formulation of Remark 2, we see that the map L_j determines the spaces \tilde{E}_j^1 and E_j^2 . Hence the condition (A2) can be verified using the data (7), and the claim (i) is proved.

Let us show the claim (ii). Let $x_0 \in \bar{\Gamma}_1 \cap \bar{\Gamma}_2$. Lemma 4 gives that, in local coordinates of ∂M taking x_0 to 0, there is a multi-index $\alpha \in \mathbb{N}^{n-1}$ such that $\partial^\alpha \partial_\nu \phi_0(0) \neq 0$. Hence the condition (A1) and Lemma 3 imply, that $\partial^\beta (\eta \tilde{\eta}^{-1})(0) = 0$ for all nonzero multi-indices $\beta \in \mathbb{N}^{n-1}$.

Fix $j \in \mathbb{N}$ and, to simplify the notation, drop the subindices j from now on. By the condition (A2)

$$\tilde{\eta} e_l|_{\Gamma_1} = \sum_{k \in I} a_{lk} \eta \partial_\nu \phi_k|_{\Gamma_1}, \quad e_l|_{\Gamma_2} = \sum_{k \in I} b_{lk} \partial_\nu \phi_k|_{\Gamma_2} \quad l \in I,$$

for some constant matrices $A := (a_{lk})_{l, k \in I}$ and $B := (b_{lk})_{l, k \in I}$.

Fix $x_0 \in \bar{\Gamma}_1 \cap \bar{\Gamma}_2$, let $l \in I$ and define the function

$$\phi(p) := \sum_{k \in I} \left(a_{lk} \frac{\eta(x_0)}{\tilde{\eta}(x_0)} - b_{lk} \right) \phi_k(p), \quad p \in M.$$

We have seen that, in local coordinates of ∂M taking x_0 to 0, the equation $\partial^\beta (\eta \tilde{\eta}^{-1})(0) = 0$ holds for all nonzero multi-indices $\beta \in \mathbb{N}^{n-1}$. Hence for any multi-index $\alpha \in \mathbb{N}^{n-1}$

$$\begin{aligned} 0 &= \partial^\alpha e_l(0) - \partial^\alpha e_l(0) \\ &= \partial_x^\alpha \left(\sum_{k \in I} a_{lk} \frac{\eta(x)}{\tilde{\eta}(x)} \partial_\nu \phi_k(x) \right) \Big|_{x=0} - \partial_y^\alpha \left(\sum_{k \in I} b_{lk} \partial_\nu \phi_k(y) \right) \Big|_{y=0} \\ &= \sum_{k \in I} \left(a_{lk} \frac{\eta(0)}{\tilde{\eta}(0)} - b_{lk} \right) \partial^\alpha \partial_\nu \phi_k(0) = \partial^\alpha \partial_\nu \phi(0). \end{aligned}$$

By Lemma 4, the coefficients

$$a_{lk} \frac{\eta(0)}{\tilde{\eta}(0)} - b_{lk}, \quad k, l \in I$$

vanish, and so $\eta(0)\tilde{\eta}(0)^{-1}A = B$.

Moreover

$$\begin{aligned} & (h, \tilde{L}f)_{L^2(\partial M, dS_g)} \\ &= \sum_{k \in I} (f, \tilde{\eta}e_k)_{L^2(\partial M, d\mu)} (h, e_k)_{L^2(\partial M, dS_g)} \\ &= \sum_{k \in I} (f, \sum_{l \in I} a_{kl} \eta \partial_\nu \phi_l)_{L^2(\partial M, d\mu)} (h, \sum_{m \in I} b_{km} \partial_\nu \phi_m)_{L^2(\partial M, dS_g)} \\ &= \sum_{l, m \in I} \left(\sum_{k \in I} a_{kl} b_{km} \right) (f, \partial_\nu \phi_l)_{L^2(\partial M, dS_g)} (h, \partial_\nu \phi_m)_{L^2(\partial M, dS_g)}, \end{aligned}$$

for all $f \in C_0^\infty(\Gamma_1)$ and $h \in C_0^\infty(\Gamma_2)$. On the other hand, the condition (A3) gives

$$\begin{aligned} (h, \tilde{L}f)_{L^2(\partial M, dS_g)} &= (h, Lf)_{L^2(\partial M, dS_g)} \\ &= \sum_{k \in I} (f, \partial_\nu \phi_k)_{L^2(\partial M, dS_g)} (h, \partial_\nu \phi_k)_{L^2(\partial M, dS_g)}, \end{aligned}$$

for all $f \in C_0^\infty(\Gamma_1)$ and $h \in C_0^\infty(\Gamma_2)$.

Denote $(\cdot, \cdot) := (\cdot, \cdot)_{L^2(\partial M, dS_g)}$. By density of $C_0^\infty(\Gamma_p)$ in $L^2(\Gamma_p, dS_g)$, $p = 1, 2$,

$$\sum_{l, m \in I} \left(\sum_{k \in I} a_{kl} b_{km} \right) (f, \partial_\nu \phi_l) (h, \partial_\nu \phi_m) = \sum_{k \in I} (f, \partial_\nu \phi_k) (h, \partial_\nu \phi_k),$$

for all $f \in L^2(\Gamma_1, dS_g)$ and $h \in L^2(\Gamma_2, dS_g)$.

Let $(f_l)_{l \in I}$ be biorthogonal with $(\partial_\nu \phi_l|_{\Gamma_1})_{l \in I}$ in $L^2(\Gamma_1, dS_g)$, and let $(h_m)_{m \in I}$ be biorthogonal with $(\partial_\nu \phi_m|_{\Gamma_2})_{m \in I}$ in $L^2(\Gamma_2, dS_g)$, that is,

$$(f_{l'}, \partial_\nu \phi_l) = \delta_{l'l}, \quad (h_{m'}, \partial_\nu \phi_m) = \delta_{m'm}, \quad l', l, m, m' \in I.$$

Then

$$\begin{aligned} \sum_{k \in I} a_{kl'} b_{km'} &= \sum_{l, m \in I} \left(\sum_{k \in I} a_{kl} b_{km} \right) (f_{l'}, \partial_\nu \phi_l) (h_{m'}, \partial_\nu \phi_m) \\ &= \sum_{k \in I} (f_{l'}, \partial_\nu \phi_k) (h_{m'}, \partial_\nu \phi_k) = \delta_{l'm'}, \quad l', m' \in I. \end{aligned}$$

Denote $c := \eta(0)^{-1}\tilde{\eta}(0) > 0$. We have shown, that $I = A^T B = cB^T B$. Hence the matrix $\sqrt{c}B$ is orthogonal. To conclude, we observe that

$$e_l|_{\Gamma_2} = \frac{1}{\sqrt{c}} \partial_\nu \sum_{k \in I} \sqrt{c} b_{lk} \phi_k|_{\Gamma_2}, \quad l \in I,$$

where $(\sum_{k \in I} \sqrt{c} b_{lk} \phi_k)_{l \in I}$ is an orthonormal basis of eigenfunctions corresponding the eigenvalue λ_j . \square

As discussed in the previous section, the operator $\Lambda_{\Gamma_1, \Gamma_2}$ determines the collection (7). So by previous theorem, if $\bar{\Gamma}_1 \cap \bar{\Gamma}_2 \neq \emptyset$, then the operator $\Lambda_{\Gamma_1, \Gamma_2}$ determines the collection (8). The collection (8) determines the manifold up to an isometry by [25, Chapter 4.4]. This proves Theorem 1.

4. Inverse problem with observations far away from sources

In this section we prove Theorems 2 and 3.

Proof of Theorem 2. Denote $L_j^{p \rightarrow q} = L_{\Gamma_p, \Gamma_q; j}$. It is enough to show, that the collection

$$\{(\lambda_j, L_j^{1 \rightarrow 2}, L_j^{1 \rightarrow 3}, L_j^{2 \rightarrow 3}) \mid j \in \mathbb{N}\} \quad (9)$$

determines a collection of type (8).

Denote $\Sigma = \Gamma_1 \cup \Gamma_2 \cup \Gamma_3$, and choose a smooth positive measure $d\mu$ on Σ . There is a positive function $\eta \in C^\infty(\Sigma)$ such that $\eta d\mu = dS_g|_\Sigma$. Define for all $j \in \mathbb{N}$

$$\begin{aligned} \tilde{E}_j^p &:= \text{span}(\eta \partial_\nu \phi_k|_{\Gamma_p})_{k \in I_j}, \quad p = 1, 2, \\ E_j^q &:= \text{span}(\partial_\nu \phi_k|_{\Gamma_q})_{k \in I_j}, \quad q = 2, 3. \end{aligned}$$

Choose a positive function $\tilde{\eta} \in C^\infty(\Sigma)$ and real-valued functions $e_k \in C^\infty(\Sigma)$, $k \in \mathbb{N}$, such that the following two conditions hold:

(B1) for all $j \in \mathbb{N}$

$$\begin{aligned} \text{span}(\tilde{\eta} e_k|_{\Gamma_p})_{k \in I_j} &= \tilde{E}_j^p, \quad p = 1, 2, \\ \text{span}(e_k|_{\Gamma_q})_{k \in I_j} &= E_j^q, \quad q = 2, 3. \end{aligned}$$

(B2) $\tilde{L}_j^{p \rightarrow q} = L_j^{p \rightarrow q}$ for all $j \in \mathbb{N}$ and $(p, q) = (1, 2), (1, 3), (2, 3)$, where

$$\tilde{L}_j^{p \rightarrow q} f(y) := \sum_{k \in I_j} (f, \tilde{\eta} e_k)_{L^2(\partial M, d\mu)} e_k(y) \quad f \in C_0^\infty(\Gamma_p), y \in \Gamma_q.$$

Again, such functions $\tilde{\eta}$ and e_k exist, as $\tilde{\eta} = \eta$ and $e_k = \partial_\nu \phi_k|_\Sigma$ satisfy the conditions (B1) and (B2). It is enough to show the following two statements:

- (i) We can verify using the data (9), whether any given functions $\tilde{\eta} \in C^\infty(\Sigma)$ and $e_k \in C^\infty(\Sigma)$, $k \in \mathbb{N}$, satisfy the conditions (B1) and (B2).
- (ii) There is an orthonormal basis of eigenfunctions $(\psi_k)_{k \in \mathbb{N}}$ of operator A and a constant $C > 0$, not depending on k , such that

$$e_k|_{\Gamma_2} = C \partial_\nu \psi_k|_{\Gamma_2}, \quad k \in \mathbb{N}.$$

Analogously with the proof of Theorem 4, the maps $L_j^{1 \rightarrow 2}$, $L_j^{1 \rightarrow 3}$ and $L_j^{2 \rightarrow 3}$ determine the spaces \tilde{E}_j^1 , E_j^2 , E_j^3 and \tilde{E}_j^2 . Hence the claim (i) is proved.

Let us show the claim (ii). Fix $j \in \mathbb{N}$ and, to simplify the notation, drop the subindices j from now on. The condition (B1) gives, that for all $l \in I$

$$\begin{aligned} \tilde{\eta} e_l|_{\Gamma_1} &= \sum_{k \in I} a_{lk} \eta \partial_\nu \phi_k|_{\Gamma_1}, \quad \tilde{\eta} e_l|_{\Gamma_2} = \sum_{k \in I} \tilde{b}_{lk} \eta \partial_\nu \phi_k|_{\Gamma_2}, \\ e_l|_{\Gamma_2} &= \sum_{k \in I} b_{lk} \partial_\nu \phi_k|_{\Gamma_2}, \quad e_l|_{\Gamma_3} = \sum_{k \in I} c_{lk} \partial_\nu \phi_k|_{\Gamma_3}, \end{aligned}$$

for some constant matrices

$$A := (a_{lk})_{l,k \in I}, \quad \tilde{B} := (\tilde{b}_{lk})_{l,k \in I}, \quad B := (b_{lk})_{l,k \in I}, \quad C := (c_{lk})_{l,k \in I}.$$

For the smallest eigenvalue λ_0 , the space of eigenfunctions is one dimensional, and so

$$0 = e_0(y) - e_0(y) = \left(\frac{\eta(y)}{\tilde{\eta}(y)} \tilde{b}_{00} - b_{00} \right) \partial_\nu \phi_0(y)$$

for all $y \in \Gamma_2$. By Lemma 4, the set

$$N := \{y \in \Gamma_2 : \partial_\nu \phi_0(y) = 0\}$$

does not contain a nonempty open set of ∂M . Hence $\overline{\Gamma_2 \setminus N} = \overline{\Gamma_2}$. Moreover, $\eta \tilde{\eta}^{-1} \tilde{b}_{00} - b_{00} = 0$ in $\Gamma_2 \setminus N$ and by continuity also in the whole set Γ_2 .

Denote by c the constant $\eta^{-1} \tilde{\eta}|_{\Gamma_2} > 0$. As $(\partial_\nu \phi_k|_{\Gamma_2})_{k \in I}$ are linearly independent, $\tilde{B} = cB$ for all $j \in \mathbb{N}$. Moreover, we may use the condition (B2) as we used the corresponding condition in the proof of Theorem 4, and get

$$A^T B = I, \quad A^T C = I, \quad \tilde{B}^T C = I.$$

Hence $B = C$ and $cB^T B = I$.

To conclude, we observe that

$$e_l|_{\Gamma_2} = \frac{1}{\sqrt{c}} \partial_\nu \sum_{k \in I} \sqrt{c} b_{lk} \phi_k|_{\Gamma_2}, \quad l \in I,$$

where $(\sum_{k \in I} \sqrt{c} b_{lk} \phi_k)_{l \in I}$ is an orthonormal basis of eigenfunctions corresponding the eigenvalue λ_j . \square

We prove Theorem 3 by reduction to Theorem 2 using a time continuation argument similar to [30].

Lemma 5. *Suppose that $\Gamma_1, \Gamma_2 \subset \partial M$ are open and nonempty. Denote*

$$T^* := 2 \max\{d(x, y) : x \in \Gamma_1, y \in M\}.$$

If $T^ < t_0 < T$, then the smooth manifolds Γ_1, Γ_2 , the operator $\Lambda_{\Gamma_1, \Gamma_2}^T$ and the inner products*

$$(u^f(t_0), u^h(t_0))_{L^2(M)}, \quad f, h \in C_0^\infty(\Gamma_1 \times (0, T)) \quad (10)$$

determine the operator $\Lambda_{\Gamma_1, \Gamma_2}^{T+\delta}$ for $\delta < t_0 - T^$.*

Proof. Denote by Y_s the time delay operator

$$Y_s f(\cdot, t) := f(\cdot, t - s), \quad t, s \in \mathbb{R}.$$

As the coefficients of the wave equation (3) are time-independent, $v^{Y_s f}(x, t) = (Y_s v^f)(x, t)$ and $(\Lambda Y_s f)(x, t) = (Y_s \Lambda f)(x, t)$.

Let $f \in C_0^\infty(\Gamma_1 \times (0, T + \delta))$ and choose $h \in C_0^\infty(\Gamma_1 \times (0, t_0))$ and $h' \in C_0^\infty(\Gamma_1 \times (\delta, T + \delta))$ such that $f = h + h'$.

Let $\epsilon \in (0, \delta)$. As $\text{supp}(Y_{-\delta}h') \subset \Gamma_1 \times (0, T)$, the operator $\Lambda_{\Gamma_1, \Gamma_2}^T$ determine the function

$$\Lambda h'(\cdot, T + \epsilon) = (Y_{-\delta}\Lambda h')(\cdot, T - (\delta - \epsilon)) = (\Lambda_{\Gamma_1, \Gamma_2}^T Y_{-\delta}h')(\cdot, T - (\delta - \epsilon)),$$

in Γ_2 . Therefore, it is enough to show that the given data determine also

$$\Lambda h(x, T + \epsilon), \quad x \in \Gamma_2.$$

Consider a sequence $(h_j)_{j=1}^\infty \subset C_0^\infty(\Gamma_1 \times (0, t_0 - \delta))$ satisfying the following two conditions.

- (C1) $\lim_{j \rightarrow \infty} v^{Y_\delta h_j}(t_0) = v^h(t_0)$ in $H^1(M)$,
 (C2) $\lim_{j \rightarrow \infty} \partial_t v^{Y_\delta h_j}(t_0) = \partial_t v^h(t_0)$ in $L^2(M)$.

Such a sequence exists, since $t_0 > T^*$, and thus we see exactly as in [25, Thm. 4.28] that the set

$$\{(v^f(t_0 - \delta), \partial_t v^f(t_0 - \delta)) : f \in C_0^\infty(\Gamma_1 \times (0, t_0 - \delta))\}$$

is dense in $H_0^1(M) \times L^2(M)$.

Let us prove, that (C1) is equivalent with

(C1') For all $c > 0$

$$\lim_{j \rightarrow \infty} ((-\Delta_g w_j + q w_j, w_j)_{L^2(M)} + c(w_j, w_j)_{L^2(M)}) = 0,$$

where $w_j := v^{Y_\delta h_j}(t_0) - v^h(t_0)$.

As $\text{supp}(Y_\delta h_j) \subset \Gamma_1 \times (0, t_0)$ and $\text{supp}(h) \subset \Gamma_1 \times (0, t_0)$, we have that $w_j|_{\partial M} = 0$. Hence

$$-(\Delta_g w_j, w_j)_{L^2(M)} = (dw_j, dw_j)_{L^2(M)},$$

where d is the exterior derivative on M . If (C1) holds, then

$$\begin{aligned} & |(-\Delta_g w_j + q w_j, w_j)_{L^2(M)} + c(w_j, w_j)_{L^2(M)}| \\ & \leq \|dw_j\|_{L^2(M)}^2 + \|q + c\|_{L^\infty(M)} \|w_j\|_{L^2(M)}^2 \\ & \rightarrow 0, \quad \text{as } j \rightarrow \infty. \end{aligned}$$

For large enough $c > 0$ there is a constant $c_0 > 0$ such that $q + c \geq c_0$. Hence if (C1') holds, then

$$\|dw_j\|_{L^2(M)}^2 + c_0 \|w_j\|_{L^2(M)}^2 \leq (-\Delta_g w_j + q w_j, w_j)_{L^2(M)} + c(w_j, w_j)_{L^2(M)} \rightarrow 0,$$

as $j \rightarrow \infty$, and (C1) holds. Therefore (C1) and (C1') are equivalent.

Next we observe that

$$\begin{aligned} & -\partial_s^2(v^{Y_\delta(Y_\delta h_j - h)}(t_0), v^{Y_\delta h_j - h}(t_0))_{L^2(M)}|_{s=0} \\ & = (-\partial_t^2 w_j, w_j)_{L^2(M)} = ((-\Delta_g + q)w_j, w_j)_{L^2(M)}. \end{aligned}$$

Hence the condition (C1') can be verified for given functions $(h_j)_{j=1}^\infty$ and h using the inner products (10).

Similarly,

$$\partial_{s_1} \partial_{s_2} (v^{Y_{s_1}(Y_\delta h_j - h)}(t_0), v^{Y_{s_2}(Y_\delta h_j - h)}(t_0))_{L^2(M)}|_{s_1=0, s_2=0} = (\partial_t w_j, \partial_t w_j)_{L^2(M)},$$

and condition (C2) can be verified for given functions $(h_j)_{j=1}^\infty$ and h using the inner products (10).

As $v^{Y_\delta h_j} - v^h = 0$ on $\partial M \times [t_0, \infty)$, conditions (C1) and (C2) together with the continuous dependency of the solution of the wave equation on the initial data, see e.g. [25, Thm. 2.30], give

$$\lim_{j \rightarrow \infty} \partial_\nu (v^{Y_\delta h_j} - v^h) = 0, \quad \text{in } L^2(\partial M \times (t_0, \infty)).$$

We have seen that, the inner products (10) determine for any $h \in C_0^\infty(\Gamma_1 \times (0, t_0))$ the nonempty set

$$\{(h_j)_{j=1}^\infty \subset C_0^\infty(\Gamma_1 \times (0, t_0 - \delta)) \mid (C1) \text{ and } (C2) \text{ hold}\},$$

and that any sequence in this set satisfies

$$\Lambda h(x, T + t) = \lim_{j \rightarrow \infty} (\Lambda Y_\delta h_j)(x, T + t) = \lim_{j \rightarrow \infty} \Lambda_{\Gamma_1, \Gamma_2}^T h_j(x, T - (\delta - t))$$

in $L^2(\Gamma_2 \times (T, T + \delta))$. As $\Lambda h \in C^\infty(\Gamma_2 \times (0, T + \delta))$, the inner products (10) and the operator $\Lambda_{\Gamma_1, \Gamma_2}^T$ determine $\Lambda h(x, T + \epsilon)$ pointwise for $x \in \Gamma_2$ and $\epsilon \in (0, \delta)$. \square

Next we prove the last of the three main theorems formulated in the introduction.

Proof of Theorem 3. By Lemma 1 the operators $\Lambda_{\Gamma_1, \Gamma_2}^T, \Lambda_{\Gamma_1, \Gamma_3}^T, \Lambda_{\Gamma_2, \Gamma_3}^T$ determine the operators

$$\Lambda_{\Gamma_p, \Gamma_q}^T, \quad p, q = 1, 2, 3, \quad p \neq q, \quad (11)$$

We use the time delay operator Y_s defined in the proof of Lemma 5. Define

$$B[f, h] := \int_0^T \int_{\partial M} \left(\partial_\nu v^f \overline{v^h} - v^f \overline{\partial_\nu v^h} \right) dS_g dt, \quad f, h \in C_0^\infty(\partial M \times (0, \infty)),$$

and let $t_0 := T/2$. We recall Blagovestchenskii identity [25, Lem. 4.16], originating from [7],

$$(v^f(t_0), v^h(t_0))_{L^2(M)} = \frac{1}{2} \int_{-t_0}^{t_0} (\text{sign } s) B[Y_{t_0+s} f, Y_{t_0-s} h] ds,$$

where $f, h \in C_0^\infty(\partial M \times (0, T))$. By this identity, the operators (11) determine the inner products

$$(v^f(t_0), v^h(t_0))_{L^2(M)}, \quad f \in C_0^\infty(\Gamma_p \times (0, T)), \quad h \in C_0^\infty(\Gamma_q \times (0, T)), \quad (12)$$

for $p, q = 1, 2, 3, p \neq q$.

Next we will show that the operators (11) determine the inner products (12) also for $p = q, p = 1, 2, 3$.

Let $f \in C_0^\infty(\Gamma_2 \times (0, T))$ and consider a sequence $(\tilde{f}_j)_{j=1}^\infty \subset C_0^\infty(\Gamma_3 \times (0, T))$ satisfying the following two conditions.

(D1) For all $h \in C_0^\infty(\Gamma_1 \times (0, T))$

$$\lim_{j \rightarrow \infty} (v^f(t_0) - v^{\tilde{f}_j}(t_0), v^h(t_0))_{L^2(M)} = 0.$$

(D2) The sequence $(\tilde{f}_j)_{j=1}^\infty$ is bounded in $L^2(\Gamma_3 \times (0, T))$.

By assumption (A), there is $\tilde{f} \in L^2(\Gamma_3 \times (0, T))$ such that $v^f(t_0) = v^{\tilde{f}}(t_0)$. Thus, there is a sequence $(\tilde{f}_j)_{j=1}^\infty \subset C_0^\infty(\Gamma_3 \times (0, T))$ such that

$$\lim_{j \rightarrow \infty} \tilde{f}_j = \tilde{f}, \quad \text{in } L^2(\Gamma_3 \times (0, T)).$$

By [25, Lem. 2.42], $v^{\tilde{f}_j}(t_0) \rightarrow v^{\tilde{f}}(t_0)$ in $L^2(M)$ as $j \rightarrow \infty$. Hence a sequence satisfying the conditions (D1) and (D2) exists.

Let us next show that

$$\|v^f(t_0)\|_{L^2(M)}^2 = \lim_{j \rightarrow \infty} (v^{\tilde{f}_j}(t_0), v^f(t_0))_{L^2(M)}. \quad (13)$$

As $t_0 > 2d(x, y)$ for all $x \in \Gamma_1$ and $y \in M$, [25, Thm. 3.10] gives that the set

$$\{v^h(t_0) \mid h \in C_0^\infty(\Gamma_1 \times (0, T))\}$$

is dense in $L^2(M)$. Let $\epsilon > 0$ and choose $h \in C_0^\infty(\Gamma_1 \times (0, T))$ such that

$$\|v^f(t_0) - v^h(t_0)\|_{L^2(M)} < \epsilon.$$

By [25, Lem. 2.42] there is $C > 0$, and by the condition (D1) there is $J \in \mathbb{N}$ such that for $j \geq J$

$$\begin{aligned} & \left| (v^f(t_0) - v^{\tilde{f}_j}(t_0), v^f(t_0))_{L^2(M)} \right| \\ & \leq \left| (v^f(t_0) - v^{\tilde{f}_j}(t_0), v^f(t_0) - v^h(t_0))_{L^2(M)} \right| \\ & \quad + \left| (v^f(t_0) - v^{\tilde{f}_j}(t_0), v^h(t_0))_{L^2(M)} \right| \\ & \leq C \|f - \tilde{f}_j\|_{L^2(\partial M \times (0, T))} \epsilon + \epsilon. \end{aligned}$$

By the condition (D2)

$$\sup_{j \in \mathbb{N}} \|f - \tilde{f}_j\|_{L^2(\partial M \times (0, T))} < \infty.$$

Hence the equation (13) is valid.

By [25, Thm. 3.10] the functions $v^f(t_0)$, $f \in C_0^\infty(\Gamma_2 \times (0, T))$, are dense in $L^2(M)$. Hence

$$\|v^h(t_0)\|_{L^2(M)} = \sup (v^h(t_0), v^f(t_0))_{L^2(M)} \quad (14)$$

where $h \in C_0^\infty(\Gamma_p \times (0, T))$, $p = 1, 3$, and the supremum is taken over all $f \in C_0^\infty(\Gamma_2 \times (0, T))$ such that $\|v^f(t_0)\|_{L^2(M)} = 1$.

The condition (D1) can be verified for any f and $(f_j)_{j=1}^\infty$ using the inner products (12) for $p = 2, 3$, $q = 1$. Therefore, these inner products determine for any $f \in C_0^\infty(\Gamma_2 \times (0, T))$ the nonempty set

$$\{(\tilde{f}_j)_{j=1}^\infty \subset C_0^\infty(\Gamma_3 \times (0, T)) \mid (D1), (D2) \text{ hold}\}.$$

By equation (13) any sequence in this set together with inner products (12) for $p = 3$ and $q = 2$ determine $\|v^f(t_0)\|_{L^2(M)}$.

As $f \in C_0^\infty(\Gamma_2 \times (0, T))$ can be chosen freely, the inner products (12) for $p = 2, 3$, $q = 1$ and for $p = 3$, $q = 2$ together with polarization identity determine the inner products (12) for $p = q = 2$.

The equation (14), polarization identity and the inner products (12) for $p = 1, 2, 3$, $q = 2$ determine the inner products (12) for $p = q = 1, 3$.

Therefore, the operators (11) determine the inner products

$$(v^f(t_0), v^h(t_0))_{L^2(M)}, \quad f, h \in C_0^\infty(\Gamma_p \times (0, T)), \quad p = 1, 2, 3. \quad (15)$$

Choose $\delta \in (0, t_0 - T^*)$, where T^* is defined as in Lemma 5. By Lemma 5 the operators (11) and the inner products (15) determine the operators

$$\Lambda_{\Gamma_p, \Gamma_q}^{T+\delta}, \quad p, q = 1, 2, 3, \quad p \neq q. \quad (16)$$

Repeating this construction, we see that the operators

$$\Lambda_{\Gamma_p, \Gamma_q}^{T+m\delta}, \quad p, q = 1, 2, 3, \quad p \neq q,$$

are determined for all $m \in \mathbb{N}$. The claim follows from Theorem 2. \square

Appendix

Next we prove the Lemma 1 stating that the operator $\Lambda_{\Gamma_1, \Gamma_2}^T$ determines the operator $\Lambda_{\Gamma_2, \Gamma_1}^T$.

Proof of Lemma 1. Define $u := Rv^{Rf}$, where v^{Rf} is the solution of the equation (3) with the boundary data $Rf \in C_0^\infty(\Gamma_1 \times (0, T))$. Then $u(x, t) = v^{Rf}(x, T - t)$ satisfies the equation

$$\partial_t^2 u + a(x, D)u = 0, \quad \text{in } M \times (0, T),$$

$$u|_{\partial M \times (0, T)} = f,$$

$$u|_{t=T} = \partial_t u|_{t=T} = 0.$$

Integration by parts gives

$$\begin{aligned} & (f, \Lambda_{\Gamma_2, \Gamma_1}^T h)_{L^2(\partial M \times (0, T))} - (R\Lambda_{\Gamma_1, \Gamma_2}^T Rf, h)_{L^2(\partial M \times (0, T))} \\ &= \int_0^T \int_{\partial M} (u(x, t) \partial_\nu v^h(x, t) - (\partial_\nu v^{Rf})(x, T - t) v^h(x, t)) dS_g(x) dt \\ &= \int_0^T \int_M (u(x, t) \Delta_g v^h(x, t) - (\Delta_g u)(x, t) v^h(x, t)) dV_g(x) dt \end{aligned}$$

$$\begin{aligned}
&= \int_0^T \int_M (u(x, t) \partial_t^2 v^h(x, t) - (\partial_t^2 u)(x, t) v^h(x, t)) dV_g(x) dt \\
&= \left[\int_M (u(x, t) \partial_t v^h(x, t) - (\partial_t u)(x, t) v^h(x, t)) dV_g(x) \right]_{t=0}^{t=T} = 0,
\end{aligned}$$

since $u|_{t=T} = \partial_t u|_{t=T} = 0$ and $v^f|_{t=0} = \partial_t v^f|_{t=0} = 0$. \square

Acknowledgements. The authors were partly supported by Finnish Centre of Excellence in Inverse Problems Research, Academy of Finland COE 213476. LO was partly supported also by Finnish Graduate School in Computational Sciences.

References

- [1] K. Astala, L. Päivärinta, Calderón's inverse conductivity problem in the plane. *Ann. of Math.* (2) 163 (2006), no. 1, 265–299.
- [2] C. Bardos, G. Lebeau, J. Rauch, Sharp sufficient conditions for the observation, control and stability of waves from the boundary, *SIAM J. Control Optim.* 30, 1992, 1024–1065.
- [3] M. Belishev, An approach to multidimensional inverse problems for the wave equation. (Russian) *Dokl. Akad. Nauk SSSR* 297 (1987), no. 3, 524–527; translation in *Soviet Math. Dokl.* 36 (1988), no. 3, 481–484.
- [4] M. Belishev, The Calderón problem for two-dimensional manifolds by the BC-method. *SIAM J. Math. Anal.* 35 (2003), no. 1, 172–182.
- [5] M. Belishev, Y. Kurylev, To the reconstruction of a Riemannian manifold via its spectral data (BC-method), *Comm. Partial Differential Equations* 17, 1992, no. 5-6, 767–804.
- [6] K. Bingham, Y. Kurylev, M. Lassas, S. Siltanen, Iterative time reversal control for inverse problems. *Inverse Problems and Imaging* 2 (2008), 63–81.
- [7] A.S. Blagovestchenskii, The inverse problem of the theory of seismic wave propagation. (Russian) *Probl. of Math. Phys.*, No. 1, pp. 6881. Izdat. Leningrad. Univ., Leningrad, 1966.
- [8] A. L. Bukhgeim, Recovering a potential from Cauchy data in the two-dimensional case. *J. Inverse Ill-Posed Probl.* 16 (2008), no. 1, 19–33.
- [9] A. L. Bukhgeim, G. Uhlmann, Recovering a potential from partial Cauchy data. *Comm. Partial Differential Equations* 27 (2002), no. 3-4, 653–668.
- [10] A. P. Calderón, On an inverse boundary value problem. *Seminar on Numerical Analysis and its Applications to Continuum Physics* (Rio de Janeiro, 1980), pp. 65–73, Soc. Brasil. Mat., Rio de Janeiro, 1980.
- [11] I. Chavel, *Riemannian geometry: A modern introduction*, Cambridge Studies in Advanced Mathematics 98, Cambridge University Press, 2006, xvi+471 pp.
- [12] M. Dahl, A. Kirpichnikova, M. Lassas, Focusing waves in unknown media by modified time reversal iteration. *SIAM J. Control Optim.* 48 (2009), no. 2, 839–858.
- [13] D. Dos Santos Ferreira, C. Kenig, M. Salo, G. Uhlmann, Limiting Carleman weights and anisotropic inverse problems. *Invent. Math.* 178 (2009), no. 1, 119–171.
- [14] L. Evans, *Partial differential equations*. Graduate Studies in Mathematics 19, American Mathematical Society, 1998, xviii+662 pp.
- [15] A. Greenleaf, G. Uhlmann Local uniqueness for the Dirichlet-to-Neumann map via the two-plane transform *Duke Math. J.* 108 (2001), 599–617.
- [16] C. Guillarmou, L. Tzou, Calderón inverse Problem with partial data on Riemann Surfaces. [arXiv:0908.1417v2](https://arxiv.org/abs/0908.1417v2).
- [17] C. Guillarmou, L. Tzou, Calderón inverse Problem for the Schrödinger Operator on Riemann Surfaces. [arXiv:0904.3804](https://arxiv.org/abs/0904.3804).

- [18] G. Henkin, V. Michel, On the explicit reconstruction of a Riemann surface from its Dirichlet-Neumann operator. *Geom. Funct. Anal.* 17 (2007), no. 1, 116–155
- [19] G. Henkin, V. Michel, Inverse conductivity problem on Riemann surfaces. *J. Geom. Anal.* 18 (2008), no. 4, 1033–1052.
- [20] L. Hörmander, Uniqueness Theorems for Second Order Elliptic Differential Equations, *Comm. in Partial Differential Equations*, 8(1), 21-64, 1983.
- [21] O. Imanuvilov, G. Uhlmann, M. Yamamoto, Global uniqueness from partial Cauchy data in two dimensions. *arXiv:0810.2286v1*.
- [22] V. Isakov, On the Uniqueness in the Inverse Conductivity Problem with Local Data. *Inverse Problems and Imaging*, Vol. 1 (2007), no. 1, 95-105.
- [23] H. Isozaki, Y. Kurylev, M. Lassas, Forward and Inverse scattering on manifolds with asymptotically cylindrical ends. To appear in *Journal of Functional Analysis*. doi:10.1016/j.jfa.2009.11.009
- [24] A. Katchalov, Ya. Kurylev, Multidimensional inverse problem with incomplete boundary spectral data, *Comm. Partial Differential Equations* 23, 1998, no. 1-2, 55-95.
- [25] A. Katchalov, Y. Kurylev, M. Lassas, *Inverse Boundary Spectral Problems*, *Monographs and Surveys in Pure and Applied Mathematics* 123, Chapman Hall/CRC-press, 2001, xi+290 pp.
- [26] A. Katchalov, Y. Kurylev, M. Lassas, Energy measurements and equivalence of boundary data for inverse problems on non-compact manifolds. *IMA volumes in Mathematics and Applications* (Springer Verlag) "Geometric Methods in Inverse Problems and PDE Control" Ed. C. Croke, I. Lasiecka, G. Uhlmann, M. Vogelius, 2004, pp. 183-214.
- [27] A. Katchalov, Y. Kurylev, M. Lassas, N. Mandache, Equivalence of time-domain inverse problems and boundary spectral problems. *Inverse Problems* 20 (2004), no. 2, 419–436.
- [28] C. Kenig, J. Sjöstrand, G. Uhlmann, The Calderón problem with partial data. *Ann. of Math. (2)* 165 (2007), no. 2, 567–591.
- [29] Kurylev, M. Lassas, Hyperbolic inverse problem with data on a part of the boundary. *Differential equations and mathematical physics* (1999), 259–272, *AMS/IP Stud. Adv. Math.*, 16, Amer. Math. Soc., 2000.
- [30] Kurylev, M. Lassas, Hyperbolic inverse boundary-value problem and time-continuation of the non-stationary Dirichlet-to-Neumann map. *Proc. Roy. Soc. Edinburgh Sect. A* 132 (2002), no. 4, 931–949.
- [31] I. Lasiecka, R. Triggiani, P. Yao, Exact controllability for second-order hyperbolic equations with variable coefficient-principal part and first-order terms. *Proceedings of the Second World Congress of Nonlinear Analysts, Part 1* (Athens, 1996). *Nonlinear Anal.* 30 (1997), no. 1, 111–122.
- [32] M. Lassas, M. Taylor, G. Uhlmann, The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary. *Comm. Anal. Geom.* 11 (2003), no. 2, 207–221.
- [33] M. Lassas, G. Uhlmann, On determining a Riemannian manifold from the Dirichlet-to-Neumann map. *Ann. Sci. cole Norm. Sup. (4)* 34 (2001), no. 5, 771–787.
- [34] J. Lee, G. Uhlmann, Determining anisotropic real-analytic conductivities by boundary measurements. *Comm. Pure Appl. Math.* 42 (1989), no. 8, 1097–1112.
- [35] R. Leis, *Initial-boundary value problems in mathematical physics*. B. G. Teubner, Stuttgart; John Wiley & Sons, Ltd., Chichester, 1986. viii+266 pp.
- [36] A. Nachman, Reconstructions from boundary measurements. *Ann. of Math. (2)* 128 (1988), no. 3, 531–576.
- [37] A. Nachman, J. Sylvester, G. Uhlmann, An n -dimensional Borg-Levinson theorem. *Comm. Math. Phys.* 115 (1988), no. 4, 595–605.
- [38] Rakesh, A linearised inverse problem for the wave equation. *Comm. Partial Differential Equations* 13 (1988), no. 5, 573–601.
- [39] J. Sylvester, G. Uhlmann, A global uniqueness theorem for an inverse boundary value problem. *Ann. of Math. (2)* 125 (1987), no. 1, 153–169.
- [40] W. Symes, The seismic reflection inverse problem. *Inverse Problems* 25 (2009), no. 12.