

CANNON-THURSTON MAPS FOR KLEINIAN GROUPS

MAHAN MJ

ABSTRACT. We show that Cannon-Thurston maps exist for degenerate free groups without parabolics, i.e. for handlebody groups. Combining these techniques with earlier work proving the existence of Cannon-Thurston maps for surface groups, we show that Cannon-Thurston maps exist for arbitrary finitely generated Kleinian groups without parabolics, proving conjectures of Thurston and McMullen. We also show that point pre-images under Cannon-Thurston maps for degenerate free groups without parabolics correspond to end-points of leaves of an ending lamination in the Masur domain, whenever a point has more than one pre-image. This proves a conjecture of Otal. We also prove a similar result for point pre-images under Cannon-Thurston maps for arbitrary finitely generated Kleinian groups without parabolics.

To Bill Thurston for lasting inspiration.

CONTENTS

1. Introduction	2
1.1. Outline of Paper and Scheme of Proof	4
2. Preliminaries	5
2.1. Relative Hyperbolicity	5
2.2. Cannon-Thurston Maps	8
3. Split Geometry	9
4. Free Groups and Finitely Generated Kleinian Groups	15
4.1. The Masur Domain	15
4.2. Incompressibility of Split Components	17
4.3. Constructing Quasidisks	18
4.4. Reduction Lemma	19
4.5. Admissible Quasigeodesics	21
4.6. Cannon-Thurston Maps for Free Groups	23
4.7. Finitely Generated Kleinian Groups	25
5. Point pre-images of the Cannon-Thurston Map	27
5.1. EL leaves are CT leaves	28
5.2. CT leaves are EL leaves	31
6. Applications	33
6.1. Primitive Stable Representations	33
6.2. Discreteness of Commensurators	34
6.3. Extending the Sullivan-McMullen Dictionary	34
7. Appendix: Model Manifold for compressible boundary	35
References	37

1. INTRODUCTION

Let G be a Kleinian group, L_G its limit set in the boundary sphere S^2 , and $D_G(= S^2 \setminus L_G)$ its domain of discontinuity. We paraphrase Question 14 of Thurston's problem-list [Thu82] below:

Question 1.1. *Suppose that Γ is a geometrically finite Kleinian group and G an arbitrary Kleinian group abstractly isomorphic to Γ via a weakly type-preserving isomorphism, i.e. an isomorphism taking parabolics of Γ to parabolics of G . Then is it true that there is a continuous map g from the limit set L_Γ (of Γ) onto the limit set L_G (of G) taking the fixed point(s) of an element γ to the fixed point(s) of the corresponding element γ' ?*

A continuous map g as in Question 1.1 is called a **Cannon-Thurston map** because of Cannon and Thurston's seminal paper [CT85, CT07]. We refer to the Introduction of [Mj14a] for a detailed history of the problem and mention only that in [Mj14a] we showed that simply or doubly degenerate surface Kleinian groups without accidental parabolics admit Cannon-Thurston maps. In [Mj14b] we had shown that point pre-images of the Cannon-Thurston map for simply or doubly degenerate groups without accidental parabolics correspond to endpoints of leaves of ending laminations whenever a point has more than one pre-image. The main aim of this paper is to apply the techniques developed in [Mj14a] and [Mj14b] to extend these results to arbitrary finitely generated Kleinian groups without parabolics (and more generally to manifolds whose ends admit a Minsky model), thus answering affirmatively Question 1.1 as also questions of McMullen [McM01] and Otal [Ota88]. This completes the project starting with [Mj14a] and proceeding through [Mj14b, DM16].¹

One more ingredient is necessary before we proceed with statements of the main results. A geodesic lamination on a hyperbolic surface is a foliation of a closed subset by geodesics. Let E be a degenerate end of a hyperbolic 3-manifold with boundary surface $S = \partial E$. Then there exists a sequence of simple closed curves $\{\sigma_n\}$ on S whose geodesic realizations exit the end E (this innocent sounding statement is a consequence of deep work of several authors including Thurston [Thu80], Bonahon [Bon86], Canary [Can93], Agol [Ago04] and Calegari-Gabai [GC06]). Then the limit of such a sequence (in the space of projectivized measured laminations $\mathcal{PML}(S)$; for the time being, the reader will not be too far off if (s)he thinks of the Hausdorff limit on the bounding surface S of E) is a lamination λ . It turns out that λ is independent of the sequence $\{\sigma_n\}$ and is called the **ending lamination** for the end E . The following provides one of the main theorems of this paper.

Theorem 5.5 *Let G be a finitely generated free degenerate Kleinian group without parabolics. Let $i : \Gamma_G \rightarrow \mathbb{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbb{H}^3$. Then i extends continuously to a map $\hat{i} : \widehat{\Gamma}_G \rightarrow \mathbb{D}^3$ between the compactifications $\widehat{\Gamma}_G, \mathbb{D}^3$ of $\Gamma_G, \mathbb{H}^3$ respectively. Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial \Gamma_G$ of Γ_G . Then $\partial i(a) = \partial i(b)$ for $a \neq b \in \partial \Gamma_G$ if and only if a, b are either ideal end-points of a leaf of an ending lamination of G , or ideal boundary points of a complementary ideal polygon.*

¹An earlier version of some parts of this paper existed in draft form in an earlier version of [Mj14a]. The division of material between [Mj14a] and the present paper is in the interests of readability.

The hyperbolic boundary $\partial\Gamma_G$ is (G -equivariantly) homeomorphic to the limit set of some (any) geometrically finite group (without parabolics) isomorphic to G . Hence Theorem 5.5 above provides a positive answer to Question 1.1 for free degenerate Kleinian group without parabolics. There are three main new ingredients of the proof over and above [Mj14a]:

- a) We need to show that split components sufficiently deep inside an end are incompressible (Proposition 4.5). This was automatic in [Mj14a].
- b) A crucial idea in the proof of the existence of the Cannon-Thurston map $\hat{i}$ in Theorem 5.5 goes back to work of Miyachi [Miy02] (see also [Sou06]) in the case of bounded geometry. There a collection of disjoint embedded disks are constructed in the (compact) core handlebody cutting it up into a ball. The boundary circles of these disks are flowed out the end giving rise to a collection of quasiconvex disks.
- c) The broad idea here is similar to (b). The basic difference between Miyachi's approach and ours is that we are forced to use a coarse model rather than a continuous one, forcing the methods of this paper to be technically quite a bit more involved. To tackle this issue we need to use the split geometry model [Mj14a], recalled in Section 3, and introduce certain "admissible paths" in Section 4.5.

The proof of Theorem 5.5 generalizes with some modifications to arbitrary finitely generated Kleinian groups. The relative hyperbolic boundary $\partial\Gamma_G$ [Bow12] of a Kleinian group G is (G -equivariantly) homeomorphic to the limit set of some (any) geometrically finite Kleinian group isomorphic to G , provided the isomorphism is strictly type-preserving (i.e. it maps parabolics to parabolics and pulls back parabolics to parabolics).

Theorems 4.20 and 5.6 *Let G be a finitely generated Kleinian group. Let $i : \Gamma_G \rightarrow \mathbb{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbb{H}^3$. Let $M = \mathbb{H}^3/G$ and assume that each degenerate end E of M admits a bi-Lipschitz Minsky model (for instance if M has no parabolics, see Remark 1.2 below). Then i extends continuously to a map $\hat{i} : \widehat{\Gamma}_G \rightarrow \mathbb{D}^3$, where $\widehat{\Gamma}_G$ denotes the (relative) hyperbolic compactification of Γ_G . Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial\Gamma_G$ of Γ_G .*

Let E be a degenerate end of $N^h = \mathbb{H}^3/G$ and $\tilde{E}$ a lift of E to $\tilde{N}^h$ and let M_{gf} be an augmented Scott core of N^h . Then the ending lamination $\mathcal{L}_E$ for the end E lifts to a lamination on $\widetilde{M_{gf}} \cap \tilde{E}$. Each such lift $\mathcal{L}$ of the ending lamination of a degenerate end defines a relation $\mathcal{R}_{\mathcal{L}}$ on the (Gromov) hyperbolic boundary $\partial\widetilde{M_{gf}}$ (equal to the relative hyperbolic boundary $\partial\Gamma_G$ of Γ_G), given by $a\mathcal{R}_{\mathcal{L}}b$ if and only if a, b are end-points of a leaf of $\mathcal{L}$. Let $\{\mathcal{R}_i\}_i$ be the entire collection of relations on $\partial\widetilde{M_{gf}}$ obtained this way. Let $\mathcal{R}$ be the transitive closure of the union $\bigcup_i \mathcal{R}_i$. Then $\partial i(a) = \partial i(b)$ if and only if $a\mathcal{R}b$.

Remark 1.2. *Theorem 4.20 gives an affirmative answer to a conjecture of McMullen [McM01] and Theorem 5.5 gives an affirmative answer to a conjecture of Otal [Ota88] under the assumption of the existence of a Minsky model. The existence of such a model was established in [Min10, BCM12] for manifolds with incompressible boundary and announced for the general (compressible boundary) case in [BCM14]. Since [BCM14] has not yet appeared we give a sketch of a proof of the*

existence of a Minsky model in the special case that M has no parabolics following ideas of Brock, Bromberg and Souto in the Appendix 7.

For ease of exposition, throughout this paper, we shall often first work out the problem for free groups and then indicate the generalization to arbitrary finitely generated Kleinian groups.

Acknowledgments: I am grateful to Jean-Pierre Otal for suggesting the problem of finding point pre-images of the Cannon-Thurston map for handlebodies; and for giving me a copy of his thesis [Ota88], where the structure of Cannon-Thurston maps for handlebody groups is conjectured. I thank Ken Bromberg for explaining the grafting construction in [Bro07] and the proof in the Appendix to me. I would also like to thank the referee for detailed suggestions and corrections. Research of the author is supported in part by CEFIPRA Indo-French Research grant 4301-1 and in part by a DST JC Bose Fellowship.

1.1. Outline of Paper and Scheme of Proof. After discussing some preliminary material on Relative Hyperbolicity and Cannon-Thurston maps in Section 2, we recall the essential technical tools from [Mj14a] in Section 3. Section 3 has to do with the fact that any end admits a *split geometry* structure. Consider a geometrically infinite end E homeomorphic to $S \times [0, \infty)$ where S is a compact hyperbolic surface (possibly with boundary). Split geometry roughly gives a sequence of embedded surfaces $\{\Sigma_i\}$ exiting E , such that successive surfaces Σ_i, Σ_{i+1} bound between them a block B_i , which either has bounded geometry, (i.e. is uniformly bi-Lipschitz homeomorphic to $S \times [0, 1]$ where the latter has the product metric) or contains a thin Margulis tube running vertically from Σ_i to Σ_{i+1} and ‘splitting’ B_i into split components. Blocks of the latter kind are called *split blocks*. Electrocuting the lifts of split components in the universal cover gives rise to a combinatorial metric d_G called a *graph metric* on $\tilde{E}$.

Section 4 is the core of the paper and proves the existence of Cannon-Thurston maps for arbitrary finitely generated Kleinian groups G under the additional assumption that each degenerate end of M admits a Minsky model. Modulo [Mj14a, DM16] the proof reduces to proving this for manifolds $M = \mathbb{H}^3/G$ with compressible core. The prototypical case is that of free degenerate Kleinian groups without parabolics, which is what we elaborate on here in the Introduction.

We briefly recall the proof of Miyachi [Miy02] (or Souto [Sou06]) in the case of bounded geometry free degenerate Kleinian groups without parabolics. Let H denote the handlebody compact core of M . The end $E(= M \setminus H)$ is equipped with a Sol-type metric as in [CT07]. A finite collection of disjoint disks $D_1, \dots, D_k$ on H are chosen cutting H up into a ball. Let σ_i denote the boundary curve of D_i . Then σ_i can be canonically ‘flowed’ out the end E using the metric on E . In fact E may be thought of as (bi-Lipschitz homeomorphic to) the universal curve over a Teichmüller geodesic ray η and each σ_i gives rise to an annulus C_i – the ‘flow image’ of σ_i by the flow given by the one-parameter family of Teichmüller maps along η . The collection $\{A_i(= D_i \cup C_i)\}$ cut M up into a (non-compact) ball. The crucial point in [Miy02] (or [Sou06]) that makes their proof work is the quasiconvexity of each A_i . This is proved there using techniques similar to [CT07]. Once this is done, it follows more or less automatically that if a geodesic segment in $\tilde{H}$ joins a pair of points a, b in a complementary component of a lift D (of one of the D_i ’s) then

the geodesic in $\widetilde{M}$ joining a, b lies (coarsely) in the corresponding component of $(\widetilde{M} \setminus A)$, where A is the lift of A_i containing D .

In this paper, we do not have the luxury of flowing along a Teichmüller geodesic. Instead we use the split geometry model of the end E to

- (1) discretize the problem by using the level surfaces Σ_i given by split geometry.
- (2) construct a discretized flow image, by considering a ‘coarse annulus’ replacement of A_i by taking the union (over i) of closed geodesics on Σ_i in the same homotopy class as σ_i .

We need to work in $\widetilde{M}$ rather than $\widetilde{E}$. Hence, in order to carry over the split geometry machinery (especially the crucially important graph metric d_G) in the context of compressible cores, we need to ensure that split components are actually incompressible sufficiently deep into the end E . This is proved in Section 4.2. With this in place, we prove in Section 4.3 that each A_i is quasiconvex in $\widetilde{M}$ equipped with the graph metric d_G (see Lemma 4.9, Corollary 4.10 and Lemma 4.11 in particular).

Next, while in [Miy02, Sou06], it is clear for trivial topological reasons that each A_i separates $\widetilde{M}$, this is no longer true in our case. We do have quasiconvexity of A_i in the d_G -metric however. It is also true that $A_i \cap \widetilde{\Sigma}_i$ separates the lift $\widetilde{\Sigma}_i$ of the split surface Σ_i . In order to use this weaker separation property, we need to ensure that if a geodesic segment in $\widetilde{H}$ joins a pair of points a, b , then the geodesic in $\widetilde{M}$ joining a, b has approximants built up of vertical pieces in blocks $\widetilde{B}_i$ and horizontal pieces (lying on $\widetilde{\Sigma}_i$). We call such approximants *admissible quasigeodesics* and deal with their construction in Section 4.5.

Once all these ingredients are in place, we prove the existence of Cannon-Thurston maps for free degenerate Kleinian groups without parabolics in Theorem 4.19.

Section 5 then generalizes the work of [Mj14b] to show that point pre-images of multiple points are given by ending laminations. We end by mentioning some applications, especially work of Jeon, Kim, Lecuire and Ohshika [JKLO14] on primitive stable representations and that of the author [Mj11] on discreteness of commensurators of Kleinian groups. In Appendix 7, we sketch a proof of the existence of a Minsky model when M has no parabolics.

2. PRELIMINARIES

2.1. Relative Hyperbolicity. We refer the reader to [Far98] for terminology and details on relative hyperbolicity and electric geometry.

Definition 2.1. Given a metric space (X, d_X) and a collection $\mathcal{H}$ of subsets, let $\mathcal{E}(X, \mathcal{H}) = X \sqcup_{H \in \mathcal{H}} (H \times [0, \frac{1}{2}])$ be the identification space obtained by identifying $(h, 0) \in H \times [0, \frac{1}{2}]$ with $h \in X$. Each $\{h\} \times [0, \frac{1}{2}]$ is declared to be isometric to the interval $[0, \frac{1}{2}]$ and $H \times \{\frac{1}{2}\}$ is equipped with the zero metric. $\mathcal{E}(X, \mathcal{H})$ is given a path pseudo-metric as follows.

Only such paths in $\mathcal{E}(X, \mathcal{H})$ are allowed whose intersection with any $\{h\} \times (0, \frac{1}{2})$ is either all of $\{h\} \times (0, \frac{1}{2})$ or is empty. The distance between two points in $\mathcal{E}(X, \mathcal{H})$ is the infimum of lengths of such allowable paths.

The resulting pseudo-metric space $\mathcal{E}(X, \mathcal{H})$ is the **electric space** associated to X and the collection $\mathcal{H}$.

We shall say that $\mathcal{E}(X, \mathcal{H})$ is constructed from X by **electrocuting** the collection

$\mathcal{H}$ and the induced pseudo-metric d_e will be called the **electric metric**. (Quasi) geodesics in the electric metric will be referred to as electric (quasi) geodesics. If $\mathcal{E}(X, \mathcal{H})$ is (Gromov) hyperbolic, we say that X is **weakly hyperbolic** relative to $\mathcal{H}$.

Note that since $\mathcal{E}(X, \mathcal{H}) = X \sqcup_{H \in \mathcal{H}} (H \times [0, \frac{1}{2}])$, X can be naturally identified with a subspace of $\mathcal{E}(X, \mathcal{H})$. Paths in (X, d_X) can therefore be regarded as paths in $\mathcal{E}(X, \mathcal{H})$, but are very far from being quasi-isometrically embedded in general.

A collection $\mathcal{H}$ of subsets of (X, d_X) is said to be D -separated if $d_X(H_1, H_2) \geq D$ for all $H_1, H_2 \in \mathcal{H}; H_1 \neq H_2$. D -separatedness is only a technical restriction as the collection $\{H \times \{\frac{1}{2}\} : H \in \mathcal{H}\}$ is 1-separated in $\mathcal{E}(X, \mathcal{H})$.

Definition 2.2. Given a collection $\mathcal{H}$ of C -quasiconvex, D -separated sets in a (Gromov) hyperbolic metric space (X, d_X) we shall say that a geodesic (resp. quasi-geodesic) γ is a geodesic (resp. quasigeodesic) **without backtracking** if γ does not return to H after leaving it, for any $H \in \mathcal{H}$.

There is a distinguished collection of 1-separated subsets of $\mathcal{E}(X, \mathcal{H})$ given by $\{H \times \{\frac{1}{2}\} : H \in \mathcal{H}\}$. An electric quasigeodesic **without backtracking** in $\mathcal{E}(X, \mathcal{H})$ is an electric quasigeodesic that does not return to $H \times \{\frac{1}{2}\}$ after leaving it, for any $H \in \mathcal{H}$.

Notation: For any pseudo metric space (Z, ρ) and $A \subset Z$, we shall use the notation $N_R(A, \rho) = \{x \in Z : \rho(x, A) \leq R\}$ as for metric spaces.

Lemma 2.3. (Lemma 4.5 and Proposition 4.6 of [Far98]; Theorem 5.3 of [Kla99]; [Bow12])

Given δ, C there exists Δ such that if (X, d_X) is a δ -hyperbolic metric space with a collection $\mathcal{H}$ of C -quasiconvex sets. Then,

- 1) Electric quasi-geodesics electrically track (Gromov) hyperbolic geodesics, i.e. for all $P > 0$, there exists $K > 0$ such that if β is any electric P -quasigeodesic from x to y , and γ is a geodesic in (X, d_X) from x to y , then $\beta \subset N_K(\gamma, d_e)$.
- 2) $\gamma \subset N_K((N_0(\beta, d_e)), d_X)$.
- 3) Relative Hyperbolicity: X is weakly hyperbolic relative to $\mathcal{H}$. $\mathcal{E}(X, \mathcal{H})$ is Δ -hyperbolic.

Note that we do not need D -separatedness in the hypothesis of Lemma 2.3 as the definition of electrocution takes care of this..

Let (X, d_X) be a δ -hyperbolic metric space, and $\mathcal{H}$ a family of C -quasiconvex, collection of subsets. Let $\alpha = [a, b]$ be a geodesic in (X, d_X) and β an electric P -quasigeodesic without backtracking in $\mathcal{E}(X, \mathcal{H})$ joining a, b . Order from the left the collection of maximal subsegments of β contained entirely in some $H \times \{\frac{1}{2}\} : H \in \mathcal{H}$. Let $\{[p_i, q_i] \times \{\frac{1}{2}\} \subset H_i \times \{\frac{1}{2}\}\}_i$ be the collection of maximal subsegments. Replace, as per this order, each path of the form $\{p_i\} \times [0, \frac{1}{2}] \cup [p_i, q_i] \times \{\frac{1}{2}\} \cup \{q_i\} \times [0, \frac{1}{2}] \subset H_i \times [0, \frac{1}{2}]$ by a geodesic $[p_i, q_i]$ in X . The resulting **connected** path β_q in X is called an *electro-ambient representative* of β in X , or simply as the electro-ambient quasigeodesic joining the end-points of β .

Lemma 2.4. (See Proposition 4.3 of [Kla99], Lemma 2.5 of [Mj14a]) Given δ, C, P there exists C_3 such that the following holds:

Let (X, d_X) be a δ -hyperbolic metric space and $\mathcal{H}$ a family of C -quasiconvex subsets.

Let (X, d_e) denote the electric space obtained by electrocuting elements of $\mathcal{H}$. Then, if α, β_q denote respectively a (Gromov) hyperbolic geodesic and an electro-ambient P -quasigeodesic with the same end-points in X , then α lies in a (Gromov hyperbolic d_X -) C_3 neighborhood of β_q .

Two paths β, γ in (X, d_X) with the same endpoints are said to have *similar intersection patterns* with $\mathcal{H}$ if there exists $\epsilon > 0$, depending only on $(X, \mathcal{H})$, such that:

- **Similar Intersection Patterns 1:** If precisely one of $\{\beta, \gamma\}$ meets some $H \in \mathcal{H}$, then the d_X -distance from the entry point to the exit point is at most D .
- **Similar Intersection Patterns 2:** If both $\{\beta, \gamma\}$ meet some $H \in \mathcal{H}$, then the distance from the entry point of β to that of γ is at most D , and similarly for the exit points.

Definition 2.5. [Far98] Suppose that X is weakly hyperbolic relative to $\mathcal{H}$. Suppose that any two electric quasigeodesics without backtracking and with the same endpoints have similar intersection patterns with respect to the collection $\{H \times \frac{1}{2} : H \in \mathcal{H}\}$. Then $(X, \mathcal{H})$ is said to satisfy **bounded penetration** and X is said to be **strongly hyperbolic** relative to $\mathcal{H}$.

The next condition ensures that $(X, \mathcal{H})$ is **strongly hyperbolic** relative to $\mathcal{H}$.

Definition 2.6. A collection $\mathcal{H}$ of uniformly C -quasiconvex sets in a δ -hyperbolic metric space X is said to be **mutually D-cobounded** if for all $H_i, H_j \in \mathcal{H}$, $\pi_i(H_j)$ has diameter less than D , where π_i denotes a nearest point projection of X onto H_i . A collection is **mutually cobounded** if it is mutually D-cobounded for some D .

Lemma 2.7. [Far98, Proposition 4.6], [Bow12] *Given $C, \delta \geq 0$, there exists P such that the following holds:*

Let X be a δ -hyperbolic metric space and $\mathcal{H}$ a collection of ϵ neighborhoods of mutually cobounded C -quasiconvex sets; then any electro-ambient quasigeodesic is a (P, P) quasigeodesic in X .

Partial Electrocution

Let M be a (not necessarily simply connected) convex hyperbolic 3-manifold with a neighborhood of the cusps excised. Then we can ensure that each boundary component of M is isometric to $\sigma \times P$, where P is either an interval or a circle, and σ is a horocycle of some fixed length e_0 . In the universal cover $\widetilde{M}$, if we excise (open) horoballs, we are left with a manifold whose boundaries are flat horospheres of the form $\tilde{\sigma} \times \tilde{P}$. Note that $\tilde{P} = P$ if P is an interval, and $\mathbb{R}$ if P is a circle (the case for a $(Z + Z)$ -cusp).

Let Y be a convex simply connected hyperbolic 3-manifold. Let $\mathcal{B}$ denote a collection of horoballs. Let X denote Y minus the interior of the horoballs in $\mathcal{B}$. Let $\mathcal{H}$ denote the collection of boundary horospheres. Then each $H \in \mathcal{H}$ with the induced metric is isometric to a Euclidean product $E^1 \times L$ for an interval $L \subset \mathbb{R}$. Here E^1 denotes Euclidean 1-space. **Partially electrocute** each H by giving it the product of the zero metric with the Euclidean metric, i.e. on E^1 put the zero metric and on L put the Euclidean metric. The resulting space is essentially what one would get (in the spirit of [Far98]) by gluing to each H the mapping cylinder

of the projection of H onto the L -factor. Let d_{pel} denote the partially electrocuted pseudometric on X .

The above construction can be done in the base manifold M itself by equipping the boundary component $\sigma \times P$ with the product of a zero metric in the σ direction and the Euclidean metric in the P -direction.

Lemma 2.8. [MP11, Lemma 1.20] *(X, d_{pel}) is a (Gromov) hyperbolic metric space.*

2.2. Cannon-Thurston Maps. Let (X, d_X) and (Y, d_Y) be hyperbolic metric spaces. By adjoining the Gromov boundaries ∂X and ∂Y to X and Y , one obtains their compactifications $\hat{X}$ and $\hat{Y}$ respectively.

Let $i : Y \rightarrow X$ denote a proper map.

Definition 2.9. *Let X and Y be hyperbolic metric spaces and $i : Y \rightarrow X$ be a proper map. A **Cannon-Thurston map** $\hat{i}$ from $\hat{Y}$ to $\hat{X}$ is a continuous extension of i .*

Lemma 2.1 of [Mit98] below gives a necessary and sufficient condition for the existence of Cannon-Thurston maps.

Lemma 2.10. [Mit98] *A Cannon-Thurston map from $\hat{Y}$ to $\hat{X}$ exists iff the following condition is satisfied:*

Given $y_0 \in Y$, there exists a non-negative function $f(n)$, such that $f(n) \rightarrow \infty$ as $n \rightarrow \infty$ and for all geodesic segments λ lying outside an n -ball around $y_0 \in Y$ any geodesic segment in X joining the end-points of $i(\lambda)$ lies outside the $f(n)$ -ball around $i(y_0) \in X$.

We shall now give a criterion for the existence of Cannon-Thurston maps between relatively hyperbolic spaces. Let X and Y be strongly hyperbolic relative to the collections $\mathcal{H}_X$ and $\mathcal{H}_Y$ respectively. Let $i : Y \rightarrow X$ be a weakly type-preserving proper embedding, i.e. for $H_Y \in \mathcal{H}_Y$ there exists $H_X \in \mathcal{H}_X$ such that $i(H_Y) \subset H_X$ and images of distinct elements of $\mathcal{H}_Y$ lie in distinct elements of $\mathcal{H}_X$.

In the Lemma below, we specialize to the case where X, Y are convex simply connected complete hyperbolic manifolds with some disjoint (open) horoballs removed. $\mathcal{H}_X$ and $\mathcal{H}_Y$ will denote the resulting collections of horospheres.

Lemma 2.11. [MP11, Lemma 1.28] *A Cannon-Thurston map for a weakly type-preserving proper embedding $i : Y \rightarrow X$ exists if and only if there exists a non-negative function $f(n)$ with $f(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that the following holds:*

Suppose $y_0 \in Y$, and $\hat{\lambda}$ in $\hat{Y} = \mathcal{E}(Y, \mathcal{H}_Y)$ is an electric quasigeodesic segment starting and ending outside horospheres. If $\lambda^b = \hat{\lambda} \setminus \bigcup_{K \in \mathcal{H}_Y} K$ lies outside $B_n(y_0) \subset Y$, then for any electric quasigeodesic $\hat{\beta}$ joining the end points of $\hat{i}(\hat{\lambda})$ in $\hat{X} = \mathcal{E}(X, \mathcal{H}_X)$, $\beta^b = \hat{\beta} \setminus \bigcup_{H \in \mathcal{H}_X} H$ lies outside $B_{f(n)}(i(y_0)) \subset X$.

We shall describe this informally as follows:

*If λ lies outside a large ball **modulo horoballs** in Y then so does any geodesic in X joining its endpoints.*

In [DM16] we proved the existence of Cannon-Thurston maps for Kleinian groups corresponding to pared manifolds whose boundary is incompressible away from cusps.

Definition 2.12. A **pared manifold** is a pair (M, P) , where P , contained in the boundary ∂M of M , is a (possibly empty) 2-dimensional submanifold with boundary such that

- (1) the fundamental group of each component of P injects into the fundamental group of M
- (2) P is a union of annuli and tori.
- (3) any cylinder $C : (S^1 \times I, \partial(S^1 \times I)) \rightarrow (M, P)$ such that $\pi_1(C)$ is injective is homotopic rel boundary into P .
- (4) P contains every torus component of ∂M .

A hyperbolic structure adapted to (M, P) is a hyperbolic structure on M such that the parabolics are precisely the elements of P . The collection of such structures is denoted as $H(M, P)$.

A pared manifold (M, P) is said to have **incompressible boundary** if each component of $\partial_0 M = \partial M \setminus P$ is incompressible in M .

The following Theorem summarizes the main results of [Mj14a, Mj14b, DM16]. It proves the existence of Cannon-Thurston maps for Kleinian groups corresponding to pared manifolds whose boundary is incompressible away from cusps. It also describes the structure of these maps in terms of ending laminations.

Theorem 2.13. [DM16] Suppose that $N^h \in H(M, P)$ is a hyperbolic structure on a pared manifold (M, P) with incompressible boundary. Let M_{gf} denote a geometrically finite hyperbolic structure adapted to (M, P) . Then the map $i : \widetilde{M}_{gf} \rightarrow \widetilde{N}^h$ extends continuously to the boundary $\partial i : \partial \widetilde{M}_{gf} \rightarrow \partial \widetilde{N}^h$.

Let E be a degenerate end of N^h and $\tilde{E}$ a lift of E to $\widetilde{N}^h$. Then the ending lamination $\mathcal{L}_E$ for the end E lifts to a lamination on $\widetilde{M}_{gf} \cap \tilde{E}$. Each such lift $\mathcal{L}$ of the ending lamination of a degenerate end defines a relation $\mathcal{R}_{\mathcal{L}}$ on the (Gromov) hyperbolic boundary $\partial \widetilde{M}_{gf}$ given by $a \mathcal{R}_{\mathcal{L}} b$ iff a, b are end-points of a leaf of $\mathcal{L}$. Let $\{\mathcal{R}_i\}_i$ be the entire collection of relations on $\partial \widetilde{M}_{gf}$ obtained this way. Let $\mathcal{R}$ be the transitive closure of the union $\bigcup_i \mathcal{R}_i$. Then $\partial i(a) = \partial i(b)$ iff $a \mathcal{R} b$.

3. SPLIT GEOMETRY

We recapitulate the essential aspects of split geometry from [Min10, Mj14a].

Split level Surfaces

A **pants decomposition** of a compact surface S , possibly with boundary, is a disjoint collection of 3-holed spheres $P_1, \dots, P_n$ embedded in S such that $S \setminus \bigcup_i P_i$ is a disjoint collection of non-peripheral annuli in S , no two of which are homotopic.

Let N be the convex core of a hyperbolic 3-manifold minus an open neighborhood of the cusp(s). Then any end E of N is simply degenerate [Ago04, GC06, Can93] and homeomorphic to $S \times [0, \infty)$, where S is a compact surface, possibly with boundary. A closed geodesic in an end E homeomorphic to $S \times [0, \infty)$ is **unknotted** if it is isotopic in E to a simple closed curve in $S \times \{0\}$ via the homeomorphism. A **tube** in an end $E \subset N$ is a regular R -neighborhood $N(\gamma, R)$ of an unknotted geodesic γ in E .

Let $\mathcal{T}$ denote a collection of disjoint, uniformly separated tubes in ends of N such that

- a) Otal [Ota95] showed the existence of an $\epsilon_{ot} > 0$ (depending only on the topology of the end E) such that any primitive closed geodesic of length at most $\epsilon_{ot} > 0$ (referred to below as the Margulis-Otal constant) is unknotted. We shall refer to tubes around such geodesics as Margulis tubes (strictly speaking, ϵ_{ot} is smaller than the Margulis constant; so this is a slight abuse of terminology). All Margulis tubes in E belong to $\mathcal{T}$ for all ends E of N .
- b) there exists $\epsilon_0 > 0$ such that the injectivity radius $injr_{ad_x}(E) > \epsilon_0$ for all $x \in E \setminus \bigcup_{T \in \mathcal{T}} Int(T)$ and all ends E of N .

In [Min10], Minsky constructs a model manifold M biLipschitz homeomorphic to N and equipped with a piecewise Riemannian structure. We shall refer to this model as the **Minsky model**. The features we shall use for M will be given below. For the time being, we note that M has a collection of tubes, each with a Euclidean structure on its boundary. The complement of these tubes may be decomposed as a union of blocks of some standard types. A Lipschitz map F was constructed in [Min10] from N to M and it was shown by Brock-Canary-Minsky [BCM12] (see [Bow11] for the general Kleinian groups case) that F was, in fact bi-Lipschitz.² Thus, let $F : N \rightarrow M$ be a bi-Lipschitz homeomorphism to the model manifold M and let $M(0)$ be the image of $N \setminus \bigcup_{T \in \mathcal{T}} Int(T)$ in M under F . Let $\partial M(0)$ (resp. ∂M) denote the boundary of $M(0)$ (resp. M). The metrics on M and $\widetilde{M}$ will be denoted by d_M .

Let $(Q, \partial Q)$ be the unique hyperbolic pair of pants such that each component of ∂Q has length one. Q will be called the *standard* pair of pants. An isometrically embedded copy of $(Q, \partial Q)$ in $(M(0), \partial M(0))$ will be said to be *flat*.

Definition 3.1. A **split level surface** associated to a pants decomposition $\{Q_1, \dots, Q_n\}$ of a compact surface S (possibly with boundary) in $M(0) \subset M$ is an embedding $f : \bigcup_i (Q_i, \partial Q_i) \rightarrow (M(0), \partial M(0))$ such that

- 1) Each $f(Q_i, \partial Q_i)$ is flat
- 2) f extends to an embedding (also denoted f) of S into M such that the interior of each annulus component of $f(S \setminus \bigcup_i Q_i)$ lies entirely in $F(\bigcup_{T \in \mathcal{T}} Int(T))$.

Let S_i^s denote the union of the collection of flat pairs of pants in the image of the embedding S_i . Note that $S_i \setminus S_i^s$ consists of annuli properly embedded in Margulis tubes.

The class of *all* topological embeddings from S to M that agree with a split level surface f associated to a pants decomposition $\{Q_1, \dots, Q_n\}$ on $Q_1 \cup \dots \cup Q_n$ will be denoted by $[f]$.

We define a partial order $\leq_E$ on the collection of split level surfaces in an end E of M as follows:

$f_1 \leq_E f_2$ if there exist $g_i \in [f_i]$, $i = 1, 2$, such that $g_2(S)$ lies in the unbounded component of $E \setminus g_1(S)$.

² The bi-Lipschitz homeomorphism between the model manifold M and the hyperbolic manifold N is established by Brock-Canary-Minsky in [BCM12] for (pared) manifolds with incompressible boundary. The extension to the general case was sketched briefly in [BCM12], but details have not appeared in print. The paper [Bow11] is also in preprint form as of date. To get around this, we provide a brief sketch in the Appendix 7 using published work of several authors, reducing the general case to the case with incompressible boundary.

A sequence S_i of split level surfaces is said to exit an end E if $i < j$ implies $S_i \leq_E S_j$ and further for all compact subsets $B \subset E$, there exists $L > 0$ such that $S_i \cap B = \emptyset$ for all $i \geq L$.

Definition 3.2. A curve v in $S \subset E$ is **l -thin** if the core curve of the Margulis tube $T_v(\subset E \subset N)$ has length less than or equal to l . A tube $T \in \mathcal{T}$ is l -thin if its core curve is l -thin. A tube $T \in \mathcal{T}$ is l -thick if it is not l -thin.

A curve v is said to split a pair of split level surfaces S_i and S_j ($i < j$) if v occurs as a boundary curve of both S_i and S_j . A pair of split level surfaces S_i and S_j ($i < j$) is said to be an **l -thin pair** if there exists an l -thin curve v splitting both S_i and S_j .

The collection of all l -thin tubes is denoted as $\mathcal{T}_l$. The union of all l -thick tubes along with $M(0)$ is denoted as $M(l)$. Unless otherwise indicated, we shall choose l to be the Margulis-Otal constant, though the discussion below will go through for any $l < \epsilon_{ot}$.

Definition 3.3. A pair of split level surfaces S_i and S_j ($i < j$) is said to be **k -separated** if

- a) for all $x \in S_i^s$, $d_M(x, S_j^s) \geq k$
- b) Similarly, for all $x \in S_j^s$, $d_M(x, S_i^s) \geq k$.

Definition 3.4. An L -bi-Lipschitz **split surface** in $M(l)$ associated to a pants decomposition $\{Q_1, \dots, Q_n\}$ of S and a collection $\{A_1, \dots, A_m\}$ of complementary annuli (not necessarily all of them) in S is an embedding $f : \cup_i Q_i \cup \cup_i A_i \rightarrow M(l)$ such that

- 1) the restriction $f : \cup_i (Q_i, \partial Q_i) \rightarrow (M(0), \partial M(0))$ is a split level surface
- 2) the restriction $f : A_i \rightarrow M(l)$ is an L -bi-Lipschitz embedding.
- 3) f extends to an embedding (also denoted f) of S into M such that the interior of each annulus component of $f(S \setminus (\cup_i Q_i \cup \cup_i A_i))$ lies entirely in $F(\bigcup_{T \in \mathcal{T}_l} \text{Int}(T))$.

Note: The difference between a split level surface and a split surface is that the latter may contain bi-Lipschitz annuli in addition to flat pairs of pants.

We denote split surfaces by Σ_i to distinguish them from split level surfaces S_i . Let Σ_i^s denote the union of the collection of flat pairs of pants and bi-Lipschitz annuli in the image of the split surface (embedding) Σ_i . The next Theorem is one of the technical tools from [Mj14a]. For the convenience of the reader, we recall the following representative situation from the Introduction to [Mj14a].

- (1) there exists a sequence $\{S_i\}$ of disjoint, embedded, bounded geometry surfaces exiting E . These are ordered in a natural way along E , i.e. $i < j$ implies that S_j is contained in the unbounded component of $E \setminus S_i$. The topological product region between S_i and S_{i+1} is denoted B_i .
- (2) Each product region B_i is of two types:
 Either the product region B_i is **thick**, i.e. there exists a uniform (independent of i) constant $K' \geq 1$ such that B_i is K' -biLipschitz homeomorphic to $S_i \times [0, 1]$. Such product regions are called **thick blocks**;
 Or, corresponding to the product region B_i , there exists a Margulis tube T_i such that $T_i \subset B_i$. Further, $T_i \cap S_i$ and $T_i \cap S_{i+1}$ are annuli on S_i and S_{i+1} respectively, with core curves homotopic to the core curve of T_i . These are examples of **split blocks**.

Thus, for split blocks, the T_i split both S_i and S_{i+1} . The complementary pieces (and their lifts to the universal cover) are examples of *split components*. Note that we have little control, however, on the geometry of the split components. This situation generalizes to give a sequence of split surfaces (rather than actual surfaces as in the representative situation above) exiting the end, such that successive pairs are split by some Margulis tubes:

Theorem 3.5. [Mj14a, Theorem 4.8] *Let $N, M, M(0), S, F$ be as above and E an end of M . Fix l less than the Margulis-Otal constant, and let $M(l) = \{F(x) : \text{inrad}_x(N) \geq l\}$. Fix a hyperbolic metric on S such that each component of ∂S is totally geodesic of length one (this is a normalization condition). There exist $L_1 \geq 1$, $\epsilon_1 > 0$, $n \in \mathbb{N}$, and a sequence Σ_i of L_1 -bi-Lipschitz, ϵ_1 -separated split surfaces exiting the end E of M such that for all i , one of the following occurs:*

- (1) *An l -thin curve v splits the pair (Σ_i, Σ_{i+1}) , i.e. v splits the associated split level surfaces (S_i, S_{i+1}) , which in turn form an l -thin pair.*
- (2) *there exists an L_1 -bi-Lipschitz embedding*

$$G_i : (S \times [0, 1], (\partial S) \times [0, 1]) \rightarrow (M, \partial M),$$

(equipping $S \times [0, 1]$ with the product metric) such that $\Sigma_i^s = G_i(S \times \{0\})$ and $\Sigma_{i+1}^s = G_i(S \times \{1\})$

Finally, each l -thin curve in S splits at most n split level surfaces in the sequence $\{\Sigma_i\}$.

In Theorem 3.5 above, n depends on the genus of the surface S . The exact nature of this dependence is in terms of hierarchies and is explicated in [Mj14a]; it is not important for this paper.

A model manifold M all of whose ends are equipped with a collection of exiting split surfaces satisfying the conclusions of Theorem 3.5 is said to be equipped with a **weak split geometry** structure.

As mentioned in Definition 3.2, pairs of split surfaces satisfying Alternative (1) of Theorem 3.5 will be called an l -thin pair of split surfaces (or simply a thin pair if l is understood). Similarly, pairs of split surfaces satisfying Alternative (2) of Theorem 3.5 will be called an l -thick pair (or simply a thick pair) of split surfaces.

Definition 3.6. *Let $(\Sigma_i^s, \Sigma_{i+1}^s)$ be a thick pair of split surfaces in M . The closure of the bounded component of $M \setminus (\Sigma_i^s \cup \Sigma_{i+1}^s)$ between $\Sigma_i^s, \Sigma_{i+1}^s$ will be called a thick block.*

Note that a thick block is uniformly bi-Lipschitz to the product $S \times [0, 1]$ and that its boundary components are $\Sigma_i^s, \Sigma_{i+1}^s$.

Definition 3.7. *Let $(\Sigma_i^s, \Sigma_{i+1}^s)$ be an l -thin pair of split surfaces in M and $F(\mathcal{T}_i)$ be the collection of l -thin Margulis tubes that split both $\Sigma_i^s, \Sigma_{i+1}^s$. The closure of the union of the bounded components of $M \setminus ((\Sigma_i^s \cup \Sigma_{i+1}^s) \bigcup_{F(T) \in F(\mathcal{T}_i)} F(T))$ between $\Sigma_i^s, \Sigma_{i+1}^s$ will be called a **split block**. The closure of any bounded component is called a **split component**.*

Note that each split component may contain Margulis tubes that *do not* split both $\Sigma_i^s, \Sigma_{i+1}^s$.

Remark 3.8. For each lift $\tilde{K} \subset \tilde{M}$ of a split component K of a split block of $M(l) \subset M$, there are lifts of l -thin Margulis tubes that share the boundary of $\tilde{K}$ in $\tilde{M}$. Adjoining these lifts to $\tilde{K}$ we obtain **extended split components**. Let $\mathcal{K}'$ denote the collection of extended split components in $\tilde{M}$. Denote the collection of split components in $\tilde{M}(l) \subset \tilde{M}$ by $\mathcal{K}$. Let $\tilde{M}(l)$ denote the lift of $M(l)$ to $\tilde{M}$. Then the inclusion of $\tilde{M}(l)$ into $\tilde{M}$ gives a quasi-isometry between $\mathcal{E}(\tilde{M}(l), \mathcal{K})$ and $\mathcal{E}(\tilde{M}, \mathcal{K}')$ equipped with the respective electric metrics. This follows from the last assertion of Theorem 3.5.

Note here that two split components may intersect along a flat subsurface along a common horizontal boundary component, and that two extended split components may intersect along Margulis tubes in addition. However, for the electrocution operation, this does not pose any problems. This is because while electrocuting, products with the unit interval of the form $\tilde{K} \times [0, \frac{1}{2}]$ are attached to $\tilde{M}$ by identifying $\tilde{K} \times \{0\}$ with $\tilde{K} \subset \tilde{M}$ and then each element of the collection $\tilde{K} \times \{\frac{1}{2}\}$ is given the zero metric.

The electric metric on $\mathcal{E}(\tilde{M}, \mathcal{K}')$ is called the **graph-metric** and is denoted by d_G . The electric space will be denoted as $(\tilde{M}, d_G)$. If there are no thick blocks, the graph metric between two points x, y roughly measures the minimal number of split components one has to pass through to go from x to y .

The electric metric on $\mathcal{E}(\tilde{M}, \mathcal{K} \cup \mathcal{T}_l)$ is quasi-isometric to the electric metric on $\mathcal{E}(\tilde{M}, \mathcal{K}')$, again by the last assertion of Theorem 3.5. The electric space will be denoted as $(\tilde{M}, d_G^1)$.

Note also that in $\mathcal{E}(\tilde{M}, \mathcal{K} \cup \mathcal{T}_l)$, elements of $\mathcal{K}$ and $\mathcal{T}$ are electrocuted separately. On the other hand each element in $\mathcal{K}'$ is a union of an element of $\mathcal{K}$ and abutting elements of $\mathcal{T}$. Thus we cannot say that $\mathcal{E}(\tilde{M}, \mathcal{K} \cup \mathcal{T}_l)$ and $\mathcal{E}(\tilde{M}, \mathcal{K}')$ are isometric.

Definition 3.9. Let $Y \subset \tilde{N}$ and $X = F(Y)$. $X \subset \tilde{M}$ is said to be Δ -graph quasiconvex if for any hyperbolic geodesic μ joining $a, b \in Y$, $F(\mu)$ lies inside $N_\Delta(X, d_G) \subset \mathcal{E}(\tilde{M}, \mathcal{K}')$.

For $X (= F(Y))$ a split component in a manifold, define $CH(X) = F(CH(Y))$, where $CH(Y)$ is the convex hull of Y in $\tilde{N}$, provided the ends of N have no cusps, i.e. $N = N^h$. Else define $CH(X)$ to be the image under F of $CH(Y)$ minus cusps. Further, in order to ensure hyperbolicity of the universal cover, we partially electrocute the cusps of M (cf. Lemma 2.8).

Then Δ -graph quasiconvexity of X is equivalent to the condition that $dia_G(CH(X))$ is bounded by $\Delta' = \Delta'(\Delta)$ as any split component has diameter one in $(\tilde{M}, d_G)$.

A split component $K (\subset E) \subset N$ is incompressible if the map $i_* : \pi_1(K) \rightarrow \pi_1(N)$ induced by the inclusion is injective. Lemma 3.10, Proposition 3.11 and Proposition 3.12 below were proved in [Mj14a] for M homotopy equivalent to a surface, where all split components are automatically incompressible. However the proofs in [Mj14a] require only that the split components be incompressible in M .

Lemma 3.10. [Mj14a, Lemma 4.16] *Let E be a simply degenerate end of a hyperbolic 3-manifold N equipped with a weak split geometry model M . For K an incompressible split component contained in E , let $\tilde{K}$ be a lift to $\tilde{N}$. Then there exists*

$C_0 = C_0(K)$ such that the convex hull of $\tilde{K}$ minus cusps lies in a C_0 -neighborhood of $\tilde{K}$ in $\tilde{N}$.

Proposition 3.11. [Mj14a, Proposition 4.23] *If K is an incompressible split component, then $\tilde{K}$ is uniformly graph-quasiconvex in $\tilde{M}$, i.e. there exists Δ' such that $\text{dia}_G(CH(\tilde{K})) \leq \Delta'$ for all incompressible split components $\tilde{K}$.*

Proposition 3.12. [Mj14a, Corollary 4.30] *Suppose that all split components of $\tilde{M}$ are incompressible. Then $(\tilde{M}, d_G)$ and hence $(\tilde{M}, d_G^1)$ are Gromov-hyperbolic.*

In fact electro-ambient quasigeodesics in $(\tilde{M}, d_G)$ and $(\tilde{M}, d_G^1)$ have the following relation.

Lemma 3.13. *Let M be a model of split geometry such that all split components are incompressible. Let $(\tilde{M}, d_G) (= \mathcal{E}(\tilde{M}, \mathcal{K}'))$ and $(\tilde{M}, d_G^1) (= \mathcal{E}(\tilde{M}, \mathcal{K} \cup \mathcal{T}_l))$ be as above. Given $o \in \tilde{M}$ and $C_0 > 0$, there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that the following holds.*

For any $a, b \in \tilde{M}$, let β_{ea}^h be an electro-ambient C_0 -quasigeodesic without backtracking in $(\tilde{M}, d_G)$ joining a, b . Let $\beta_{ea} = \beta_{ea}^h \setminus \partial\tilde{M}$ be the part of β_{ea}^h lying away from the (bi-Lipschitz) horospherical boundary of $\tilde{M}$. Again, let β_{ea1}^h be an electro-ambient C_0 -quasigeodesic without backtracking in $(\tilde{M}, d_G^1)$ joining a, b . Let $\beta_{ea1} = \beta_{ea1}^h \setminus \partial\tilde{M}$ be the part of β_{ea1}^h lying away from the (bi-Lipschitz) horospherical boundary of $\tilde{M}$, where we refer to the bi-Lipschitz image under F of the horospherical boundary of $\tilde{N}$ as the (bi-Lipschitz) horospherical boundary of $\tilde{M}$.

Then $d_M(\beta_{ea}, o) \geq n$ implies that $d_M(\beta_{ea1}, o) \geq \Theta(n)$. Conversely, $d_M(\beta_{ea1}, o) \geq n$ implies that $d_M(\beta_{ea}, o) \geq \Theta(n)$.

Proof. Let K' be an extended split component in $\mathcal{K}'$ and $\tilde{K}'$ denote its universal cover. Let $\tilde{K}' = \tilde{K} \cup \tilde{T}^i$ where $\tilde{T}^i$ are the universal covers of l -thin Margulis tubes abutting the split component $\tilde{K}$ in $\tilde{K}'$. Suppose K is contained in B_i , the i -th block in an end E .

Then $\tilde{K}'$ is hyperbolic and is contained in a $C (= C(K'))$ -neighborhood of $\tilde{K}$. The argument is now a reprise of similar arguments in Section 6 (e.g. Lemma 6.8 and 6.10) of [Mj14a]:

For all i , there exists $C(i)$, such that $\beta_{ea1} \cap \tilde{B}_i$ lies in a $C(i)$ -neighborhood of $\beta_{ea} \cap \tilde{B}_i$ in $\tilde{M}$. Suppose $d_M(\beta_{ea}, o) \geq n$. Hence, by uniform k_0 -separatedness of split surfaces (Theorem 3.5), $d_M(\beta_{ea1} \cap \tilde{B}_i, o) \geq \max(n - C(i), ik_0)$.

Let $D(i) = \max_{1 \leq j \leq i} C(j)$. Then $d_M(\beta_{ea1}, o) \geq \max(n - D(i), ik_0)$ for all i . One direction of the Lemma follows.

The converse direction is similar. $\square$

We summarize the conclusions of the above propositions below.

Definition 3.14. *A model manifold of weak split geometry is said to be of **split geometry** if*

- (1) *Each split component $\tilde{K}$ is quasiconvex (not necessarily uniformly) in the hyperbolic metric on $\tilde{N}$.*

- (2) Equip $\widetilde{M}$ with the graph-metric d_G obtained by electrocuting (extended) split components $\widetilde{K}$. Then the convex hull $CH(\widetilde{K})$ of any split component $\widetilde{K}$ has uniformly bounded diameter in the metric d_G .

Hence by Lemma 3.10 and Proposition 3.11 we have the following (where we refer the reader to the Appendix, Section 7, for a sketch of a proof of the existence of a Minsky model for a general finitely generated Kleinian group without parabolics).

Theorem 3.15. *Any degenerate end E of a hyperbolic 3-manifold M admitting a Minsky model also has a model of split geometry. In particular, if $j : E \rightarrow M$ denotes the inclusion and if no element of $j_*(\pi_1(E))$ is a parabolic, then E admits a model of split geometry.*

4. FREE GROUPS AND FINITELY GENERATED KLEINIAN GROUPS

Let G be a free geometrically infinite Kleinian group. Agol [Ago04], and independently, Gabai and Calegari [GC06] have shown that $N = \mathbf{H}^3/G$ is topologically tame, and hence, by work of Canary [Can93], geometrically tame. Then any manifold M bi-Lipschitz to N is homeomorphic to the interior of a handlebody with boundary S .

More generally, let G be a finitely generated Kleinian group and G_f be a geometrically finite Kleinian group, abstractly isomorphic to G via a strictly type-preserving isomorphism. Let H denote the convex core of $\mathbf{H}^3/G_f$ and let M be a model manifold bi-Lipschitz homeomorphic to $N = \mathbf{H}^3/G$. Then there is a natural identification $i : H \rightarrow M$ of H with the augmented Scott core (i.e. Scott core plus parabolics) of M . Let $\widetilde{i}$ indicate the lift of i to the universal cover. For most of the discussion below, it might be helpful at first reading to have in mind a free geometrically infinite Kleinian group without parabolics. We fix this notation for H, M, S throughout this section.

Standing Assumption: For the purposes of this Section, we assume that each degenerate end E of M admits a Minsky model. As mentioned in Remark 1.2 this is expected to be satisfied always and a proof in the special case that M has no parabolics is sketched in Section 7.

4.1. The Masur Domain.

Definition 4.1. *Let E be an end of a hyperbolic manifold such that E is homeomorphic to $S \times [0, \infty)$ for S a finite area hyperbolic surface. A map $h : E \rightarrow S \times [0, \infty)$ is said to be type-preserving, if all and only the cusps of E are mapped to cusps of $S \times [0, \infty)$.*

Theorem 4.2. [Ago04, GC06, Bow11, BCM14] *Let G be a finitely generated Kleinian group and $M = \mathbf{H}^3/G$. Let H denote an augmented Scott core of M . Let E_1 be a geometrically infinite end of $M \setminus H$. Then E_1 is homeomorphic (via a type-preserving homeomorphism) to a topological product $S \times [0, \infty)$ for a hyperbolic surface S of finite area. Further, there exists a neighborhood E of the end corresponding to E_1 such that E is bi-Lipschitz homeomorphic to a Minsky model for $S \times [0, \infty)$ and hence to a model of split geometry.*

The last part of the last statement follows from Theorem 3.15.

Some ambiguity remains in the statement of Theorem 4.2 above. This lies in the choice of the ending lamination for E used to build the Minsky model. Since

$i : S \subset E$ is type-preserving, no parabolic element of S bounds a compressing disk. Let $\mathcal{ML}(S)$ be the space of measured laminations on S and let $\mathcal{PML}(S)$ denote the space of projectivized measured laminations. Let $\mathcal{D}(S)$ be the subset of $\mathcal{PML}(S)$ consisting of weighted unions of disjoint meridians (boundaries of compression disks lying on S). Let $cl(\mathcal{D}(S))$ denote the closure of $\mathcal{D}(S)$. Define the **Masur domain** of S by

$\mathcal{PMD}(S) = \{\lambda \in \mathcal{PML}(S) : i(\lambda, \mu) > 0\}$ for all $\mu \in cl(\mathcal{D}(S))$, provided S has at least two disjoint isotopy classes of compressing disks.

Else, we define

$\mathcal{PMD}(S) = \{\lambda \in \mathcal{PML}(S) : i(\lambda, \mu) > 0 \text{ for any } \mu \text{ that is disjoint from a compressing disk}\}$.

Here $i(\lambda, \mu)$ denotes the intersection number of λ, μ . Let $\mathcal{MD}(S) (\subset \mathcal{ML}(S))$ be the set of measured geodesic laminations whose projective class lies in $\mathcal{PMD}(S)$.

Now, let $M = H \cup_i E_i$, where H is an augmented Scott core. We consider the ends E_j which have compressible boundary, i.e. $\partial E_j (= E_j \cap H)$ is compressible in H . Let E be such an E_j and $S = H \cap E$ be the corresponding boundary component of H . Then any compressible simple closed curve on ∂H lies on such an S . We fix an S for now and proceed. Let $Mod_0(S)$ denote the subgroup of the mapping class group of S generated by Dehn twists along essential simple closed curves that bound embedded disks in H . $Mod_0(S)$ acts on $\mathcal{PML}(S)$. It was shown by Otal [Ota88] (see also McCarthy and Papadopoulos [MP89]) that under this action, $Mod_0(S)$ acts properly discontinuously on $\mathcal{PMD}(S) (\subset \mathcal{PML}(S))$ with limit set $cl(\mathcal{D}(S)) (\subset \mathcal{PML}(S))$.

Remark 4.3. Classically [Luf78, Suz77, MM86], $Mod_0(S)$ is defined as the subgroup of the mapping class group $Mod(S)$ of S which extend to the trivial outer automorphism of $\pi_1(H)$. Luft [Luf78], Suzuki [Suz77] and McCullough-Miller [MM86] prove that this group is exactly the subgroup of the mapping class group of S generated by Dehn twists along essential simple closed curves that bound embedded disks in H .

As mentioned in the introduction to the paper, for any degenerate end E with boundary surface S , there exists a sequence of simple closed curves $\{\sigma_n\}$ on S , whose geodesic realizations exit the end E . The limit of these curves in $\mathcal{PML}(S)$ defines an ending lamination for E . However, since S is compressible (in H), we may obtain a different sequence of simple closed curves $\{\sigma'_n\}$ on S by acting on σ_n by different elements of $Mod_0(S)$. Note that σ'_n and σ_n are homotopic in $H \cup E$ and hence have the same geodesic realization in E . Otal [Ota88] (see also [KS03] for the case with parabolics) shows that any two essential simple closed curves on S in the Masur domain that are freely homotopic in M lie in the same $Mod_0(S)$ -orbit. We refer the reader to [Can93, Proposition 3.3] for a published proof. Otal [Ota88] (see also [KS03]) further shows that the subset of weighted multicurves in $\mathcal{PMD}(S)/Mod_0(S)$ injects homeomorphically into the space of currents on M . It follows that the ending lamination for E is well-defined up to the action of $Mod_0(S)$:

Theorem 4.4. *For any finitely generated Kleinian group, the ending lamination λ facing a surface S with a compressing disk lies in the Masur Domain and is a well-defined element of $\mathcal{PMD}(S)/Mod_0(S)$.*

For our purposes we shall mostly be satisfied with the fact that E is bi-Lipschitz homeomorphic to *some* Minsky model, and hence, by Theorem 4.2 to a model of split geometry.

4.2. Incompressibility of Split Components. Recall that we are working in the setup where M is a hyperbolic 3-manifold and H its Scott core (augmented Scott core, when M has parabolics). Let S be a boundary component of H that is compressible (rel. cusps when M , and hence H , has parabolics). Let E be the end with S as its boundary. We would like to show that sufficiently deep within E , all split components are incompressible in M . Recall that splitting tubes correspond to *thin Margulis tubes* in the split geometry model built from the Minsky model.

Proposition 4.5. *Let M, H, E be as above. Equip E with a split geometry structure. Then there exists (a "sub-end") $E_2 \subset E$ such that*

- (1) E_2 is homeomorphic to $S \times [0, \infty)$ by a type-preserving homeomorphism and consists of a union of blocks and tubes from the split geometry model for E .
- (2) All split components of E_2 are incompressible, i.e. if K is a split component of E_2 , then the inclusion $i : K \rightarrow M$ induces an injective map $i_* : \pi_1(K) \rightarrow \pi_1(M)$.

Proof. Suppose not. Then there exists a sequence of split components K_i exiting the end E such that $i_* : \pi_1(K) \rightarrow \pi_1(M)$ is not injective. Since K_i are split components, $K_i = S_i \times I$ for some subsurface S_i of S . By the Loop Theorem (see for instance, Hempel [Hem76]), there exist simple closed curves $\sigma_i \subset S_i$ such that σ_i bound embedded topological disks in H . Hence $\sigma_i \in \mathcal{D}(S)$. Let T_i be a splitting tube bounding K_i . Then T_i exit E and has a core curve α_i which in turn corresponds to a simple closed curve on S . It follows that α_i is disjoint from an essential disk and hence some $\sigma'_i \in \mathcal{D}(S)$. Since α_i are simple closed curves whose geodesic realizations exit E , it follows that any such sequence of curves α_i converges (in $\mathcal{PML}(S)$) to the ending lamination λ corresponding to the end E . Hence $\lambda \in \mathcal{cl}(\mathcal{D}(S))$ and cannot lie in the Masur domain. This contradicts Theorem 4.4. The proposition follows. $\square$

Henceforth we shall choose l (while fixing l -thin Margulis tubes in the construction of the split geometry model) to be small enough, so that $(E \setminus E_2)$ is contained in a thick block and hence all split components are incompressible. As an immediate consequence we have the following.

Lemma 4.6. *If K is a split component, then $\pi_1(K) \subset \pi_1(M)$ is geometrically finite (Schottky, in the absence of parabolics).*

Proof. Follows from Lemma 3.10. $\square$

Proposition 4.7. *If K is a split component, then $\tilde{K}$ is uniformly graph-quasiconvex in $\tilde{M}$.*

Proof. Follows from Proposition 3.11. $\square$

Proposition 4.8. *$(\tilde{M}, d_G)$ is a hyperbolic metric space.*

Proof. Follows from Proposition 3.12. $\square$

4.3. Constructing Quasidisks. The construction in this subsection may be regarded as a *graph-metrized coarse* analogue of an unpublished construction due to Miyachi [Miy02] (see also Souto [Sou06]). The main technical difference between Miyachi's construction and ours is that Miyachi constructs continuous images of disks that actually separate the universal cover $\widetilde{M}$, whereas we only construct quasidisks. As a consequence it becomes technically more difficult for us to prove that quasidisks coarsely separate. This is why we need a special family of paths which we shall call 'admissible paths' in the next subsection which either intersect or come close to the quasidisks we construct below.

Recall that M is a hyperbolic 3-manifold and H its Scott core (augmented Scott core, when M has parabolics). Also, E is an end such that $S = H \cap E$ is compressible. We choose a collection of essential simple closed curves $\sigma_1 \cdots \sigma_g$ on S bounding disks $D_1 \cdots D_g$ with neighborhoods $D_i \times (-\epsilon, \epsilon)$ such that each component of $H \setminus \bigcup_i D_i \times (-\epsilon, \epsilon)$ is either a ball or has incompressible boundary (rel. cusps). Also assume that σ_i are geodesics in the intrinsic metric on S . To avoid multiple indices we fix an end E of M and describe the construction of quasidisks in E . Next, fix a split geometry structure on E as a union of contiguous blocks B_k , where each B_k is either a split block, or a thick block.

Further, let $\partial B_k = S_{k-1} \cup S_k$ with S_k the *upper boundary* and S_{k-1} the *lower boundary*. Also let $S = S_0$ and $\sigma_i = \sigma_{i0}$. Let σ_{ik} be the shortest closed curve in the split metric on S_k (i.e. in the pseudometric obtained by electrocuting annular intersections of splitting Margulis tubes with the split level surface S_k) homotopic in E to $\sigma_i = \sigma_{i0}$. Let $\overline{A_i} = D_i \bigcup_k \sigma_{ik} \subset M$ be the union of the disk D_i and the quasi-annulus $\bigcup_k \sigma_{ik}$. Then any lift A_i of $\overline{A_i}$ to $\widetilde{M}$ is isometric to $\overline{A_i}$ as D_i is homotopically trivial and σ_{ik} are all freely homotopic to $\sigma_i = \sigma_{i0} = \partial D_i$.

We want to show that A_i are quasiconvex in $(\widetilde{M}, d_G)$ which in turn is hyperbolic by Proposition 4.8.

qi Rays

Fix a σ and the disk D it bounds. Let $A = D \bigcup_k \sigma_k \subset M$, where $\sigma_k \subset S_k$. Lift $\bigcup_k \sigma_k$ to the universal cover $\widetilde{E}$ of E such that any lift $\widetilde{\sigma}_k$ lies in the universal cover $\widetilde{S}_k$ of S_k (We are using this notation to distinguish from lifts to $\widetilde{M}$). Let λ_k be any such lift $\widetilde{\sigma}_k$. We then have the following from [Mj14a].

Lemma 4.9. [Mj14a, Lemma 5.9] *There exists $C \geq 0$ and for all $k \in \mathbb{N}$, there exists $B(k) > 0$ such that the following hold:*

- (1) *For $x_k \in \lambda_k$ there exists $x_{k-1} \in \lambda_{k-1}$ with $d_G(x_k, x_{k-1}) \leq C$ and $d_M(x_k, x_{k-1}) \leq B(k)$.*
- (2) *Similarly there exists $x_{k+1} \in \lambda_{k+1}$ with $d_G(x_k, x_{k+1}) \leq C$ and $d_M(x_k, x_{k+1}) \leq B(k)$.*

Hence, for all n and $x \in \lambda_n$, there exists a C -quasigeodesic ray r (in the d_G -metric) such that $r(k) \in \lambda_k$ for all k and $r(n) = x$.

By construction of split blocks, $d_G(x_i, S_{i-1}) = 1$. Therefore inductively, $d_G(x_i, S_j) = |i - j|$. Hence $d_G(x_i, x_j) \geq |i - j|$. By construction, $d_G(x_i, x_j) \leq C|i - j|$.

Hence, given $p \in \lambda_i$ the sequence of points $x_n, n \in \mathbb{N} \cup \{0\}$ with $x_i = p$ gives by Lemma 4.9 above, a quasigeodesic in the d_G -metric. Such quasigeodesics shall be referred to as *d_G -quasigeodesic rays*.

After projecting $\widetilde{E}$ to $\widetilde{M} \setminus \widetilde{H}$ we have the following conclusion.

Lemma 4.10. *There exists $C \geq 0$ and for all k there exists B_k satisfying the following:*

For all $x_{ik} \in \sigma_{ik}$ there exists $x_{i,k-1} \in \sigma_{i,k-1}$ with $d_G(x_{ik}, x_{i,k-1}) \leq C$ and $d(x_{ik}, x_{i,k-1}) \leq B_k$.

The following Corollary will turn out to be quite useful. Here we replace A by the collection A_i of annuli constructed in the third paragraph of this section.

Corollary 4.11. *There exists $C \geq 0$ such that for all k and all $x_{ik} \in \sigma_{ik}$ there exists $q \in \sigma_{i0}$ and a sequence of points $p = x_{ik}, \dots, x_{i0} = q$ which is a quasigeodesic in $(\widetilde{M}, d_G)$.*

Proof. By construction of split blocks, $d_G(x_{ik}, S_{k-1}) = 1$.

Hence, given $p \in \sigma_{ik}$ the sequence of points $p = x_{ik}, \dots, x_{i0}$ gives by Lemma 4.10 above, a quasigeodesic in the d_G -metric lying entirely on A_i joining p to a point $q \in D_i$. $\square$

We can choose a point $z_i \in D_i$ (quite arbitrarily) and extend any quasigeodesic constructed as above by adding on a path from q to z_i lying entirely in D_i and having uniformly bounded length since D_i has bounded diameter.

Proposition 4.12. *There exists $C_0 \geq 0$ such that each A_j is C_0 -quasiconvex in $(\widetilde{M}, d_G)$.*

Proof: By Lemma 4.10 and Corollary 4.11 above, it follows that there exist $K \geq 1$ such that for any two points p_1, p_2 in A_j there exist K -quasi-geodesics γ_1, γ_2 to $z_j \in D_j$. By Proposition 4.8 we also have that $(\widetilde{M}, d_G)$ is hyperbolic. Hence any geodesic α_i ($i = 1, 2$) joining p_i to z_j lies in some K_1 neighborhood of γ_i . Further, by hyperbolicity of $(\widetilde{M}, d_G)$, we conclude that a geodesic β joining p_1, p_2 lies in a K_2 -neighborhood of $\alpha_1 \cup \alpha_2$. Hence, finally, β lies in a $(K_1 + K_2)$ neighborhood of $\gamma_1 \cup \gamma_2 \subset A_j$. Choosing $C_0 = K_1 + K_2$, we are through. $\square$

The quasidisks constructed above have the following property.

Lemma 4.13. *Let M be a model manifold of split geometry. Let H be a Scott core of M (augmented Scott core if M has parabolics) and $\{D_i\}$ a maximal collection of disjoint compressing disks in H . Then there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that for all $o \in \widetilde{H}$ the following holds.*

Let D be a lift of one of the D_i 's to $\widetilde{M}$ and let A be the quasidisk in $\widetilde{M}$ constructed from D as above.

Then $d_M(D, o) \geq n$ implies that $d_M(A, o) \geq \Theta(n)$.

Proof. By Lemma 4.10 and Corollary 4.11, there exists $b_1, \dots, b_k, \dots$ and $z \in D$ such that for all $x_k \in \sigma_k \subset A$, $d_M(x_k, z) \leq (b_1 + \dots + b_k) = c_k$ (say). Hence $d_M(x_k, 0) \geq (n - c_k)$.

By uniform ϵ_0 -separatedness of split surfaces (Theorem 3.5), $d_M(x_k, 0) \geq k\epsilon_0$. Hence $d_M(x_k, 0) \geq \max((n - c_k), k\epsilon_0)$. Choosing $\Theta(n)$ to be the largest value of $k\epsilon_0$ such that $k\epsilon_0 \leq n - c_k$ we are done. $\square$

4.4. Reduction Lemma. Before we get into the proof of the existence of Cannon-Thurston maps, we recall some material from Section 6 of [Mj14a] that will help streamline the proof.

The next Lemma allows us to apply the criterion for existence of Cannon-Thurston maps in Lemmas 2.10 and 2.11 to electro-ambient quasigeodesics in $\widetilde{M}$

rather than hyperbolic geodesics in $\tilde{N}$. Lemma 4.14 below is a paraphrasing of what Lemmas 6.8 and 6.10 of [Mj14a] prove in the context of this paper. (Note that though [Mj14a, Lemmas 6.8 and 6.10] are stated for simply degenerate surface groups, the relevant parts of the proofs only use incompressibility of split components. This is pointed out in [Mj14a, Lemma 8.7].)

Lemma 4.14. [Mj14a, Lemma 8.7] *Let N be the convex core of a complete hyperbolic 3-manifold N^h minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let M be the resulting model of split geometry and $F : N \rightarrow M$ be the bi-Lipschitz homeomorphism between the two. Let $\tilde{F}$ be a lift of F to the universal covers. Then for all $C_0 > 0$, and $o \in \tilde{N}$ there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that the following holds.*

For any $a, b \in \tilde{N} \subset \tilde{N}^h$, let λ^h be the hyperbolic geodesic in $\tilde{N}^h$ joining them and let $\lambda_{thick}^h = \lambda^h \cap \tilde{N}$. Similarly let β_{ea}^h be an electro-ambient C_0 -quasigeodesic without backtracking in $\tilde{M} \subset \mathcal{E}(\tilde{M}, \mathcal{K}')$ joining $\tilde{F}(a)$, $\tilde{F}(b)$. Let $\beta_{ea} = \beta_{ea}^h \setminus \partial\tilde{M}$ be the part of β_{ea}^h lying away from the (bi-Lipschitz) horospherical boundary of $\tilde{M}$.

Then $d_M(\beta_{ea}, \tilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}(\lambda_{thick}^h, o) \geq \Theta(n)$.

Combining Lemma 4.14 with Lemma 3.13, we have the following.

Corollary 4.15. *Let N be the convex core of a complete hyperbolic 3-manifold N^h minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let M be the resulting model of split geometry and $F : N \rightarrow M$ be the bi-Lipschitz homeomorphism between the two. Let $\tilde{F}$ be a lift of F to the universal covers. Then for all $C_0 > 0$, and $o \in \tilde{N}$ there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that the following holds.*

For any $a, b \in \tilde{N} \subset \tilde{N}^h$, let λ^h be the hyperbolic geodesic in $\tilde{N}^h$ joining them and let $\lambda_{thick}^h = \lambda^h \cap \tilde{N}$. Similarly let β_{ea}^h be an electro-ambient C_0 -quasigeodesic without backtracking in $\tilde{M} \subset \mathcal{E}(\tilde{M}, \mathcal{K} \cup \mathcal{T}_1)$ joining $\tilde{F}(a)$, $\tilde{F}(b)$. Let $\beta_{ea} = \beta_{ea}^h \setminus \partial\tilde{M}$ be the part of β_{ea}^h lying away from the (bi-Lipschitz) horospherical boundary of $\tilde{M}$.

Then $d_M(\beta_{ea}, \tilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}(\lambda_{thick}^h, o) \geq \Theta(n)$.

Again, combining Lemma 4.14 with Lemma 4.13, we have the following.

Corollary 4.16. *Let N be the convex core of a complete hyperbolic 3-manifold N^h minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let M be the resulting model of split geometry and $F : N \rightarrow M$ be the bi-Lipschitz homeomorphism between the two. Let $\tilde{F}$ be a lift of F to the universal covers. Let H be a Scott core of N and $\{D_i\}$ a maximal collection of compressing disks in H . Then there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that for all $o \in \tilde{H}$ the following holds.*

Let D be a lift of one of the D_i 's to $\tilde{M}$ and let A be the quasidisk in $\tilde{M}$ constructed from D as above. For any $a, b \in A$, let $[a, b]_h$ be the hyperbolic geodesic in $\tilde{N}^h$ joining $\tilde{F}^{-1}(a)$, $\tilde{F}^{-1}(b)$ and let $[a, b] = [a, b]_h \cap \tilde{N}$.

Then $d_M(D, \tilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}([a, b], o) \geq \Theta(n)$.

Proof. By Lemma 4.13 there exists $z \in D$, a function $\Theta_0 : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta_0(n) \rightarrow \infty$ as $n \rightarrow \infty$ and electro-ambient quasigeodesics β_a, β_b in $(\widetilde{M}, d_G)$, joining a to z and b to z respectively, such that $d_M(D, \widetilde{F}(o)) \geq n$ implies that $d_M(\beta_a \cup \beta_b, \widetilde{F}(o)) \geq \Theta_0(n)$.

Let $[a, z]_h, [b, z]_h$ be the hyperbolic geodesic in $\widetilde{N}$ joining $\widetilde{F}^{-1}(a), \widetilde{F}^{-1}(b)$ respectively to $\widetilde{F}^{-1}(z)$ and let $[a, z] = [a, z]_h \cap \widetilde{N}$, $[b, z] = [b, z]_h \cap \widetilde{N}$.

Then by Lemma 4.14 there exists a function $\Theta_1 : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta_1(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that $d_M(D, \widetilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}([a, z] \cup [b, z], o) \geq \Theta_1(n)$.

Let $\delta > 0$ be such that all geodesic triangles in $\mathbf{H}^3$ are δ -thin. Taking $\Theta(n) = \Theta_1(n) - \delta$, it follows that $d_M(D, \widetilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}([a, b], o) \geq \Theta(n)$. $\square$

4.5. Admissible Quasigeodesics. We shall need a collection of paths consisting of horizontal and vertical segments approximating electro-ambient quasigeodesics. We shall call these admissible quasigeodesics. Let M be a model manifold each of whose ends is equipped with a split geometry structure such that all split components are incompressible. Recall that each thick block and each split block in M is homeomorphic to a product $\Sigma_i^s \times I$. We fix such a product structure for each block. Let $t_i = \sup\{\text{length}(\{x\} \times I) : x \in \Sigma_i^s\}$ be the *thickness* of the i -th block.

Recall that $F : N \rightarrow M$ is a bi-Lipschitz homeomorphism from a hyperbolic manifold N (minus cusps) to M and let $\widetilde{F}$ denote its lift to the universal cover.

An **elementary admissible path** in $\widetilde{M}$ is one of the following:

Type 1: A ‘horizontal’ geodesic in the intrinsic path metric on some lift $\widetilde{\Sigma}_i^s$ of a split surface to $\widetilde{M}$.

Type 2: A ‘vertical’ path of the form $x \times I$ (with respect to the fixed product structure above) in the lift to $\widetilde{M}$ of either a thick block or a split block.

Let B_i be a thick block of M and let S_i^s, S_{i+1}^s be its horizontal boundary components. Recall that a product structure $B_i = S \times I$ has been fixed. Let $\mu = [a, b]$ be a geodesic in $\widetilde{B}_i \subset \widetilde{M}$ such that its end-points a, b lie on the horizontal boundary components. Let $P(\mu)$ denote the projection of μ onto the horizontal boundary component (S_i^s or S_{i+1}^s) containing a . If b belongs to the same horizontal boundary component as a , define $\mu_{adm} = P(\mu)$. Else define $\mu_{adm} = P(\mu) \cup \{b\} \times I$, where $\{b\} \times I$ is the elementary vertical path through b . μ_{adm} will be called the **admissible quasigeodesic** corresponding to μ in the thick block $\widetilde{B}_i$.

Let $\mathcal{B}$ denote the collection of thick blocks.

Definition 4.17. Let β_{ea} be an electro-ambient C_0 -quasigeodesic (without backtracking), such that it enters or leaves split components at split level surfaces. Then an **admissible quasigeodesic** β_{adm} in $\widetilde{M}$ corresponding to β_{ea} in $\widetilde{M} \subset \mathcal{E}(\widetilde{M}, \mathcal{K} \cup \mathcal{T}_l)$ is a path such that $\beta_{adm} \cap (\widetilde{M} \setminus \bigcup_{K \in \mathcal{K}} \widetilde{K} \cup \bigcup_{B \in \mathcal{B}} \widetilde{B}) = \beta_{ea} \cap (\widetilde{M} \setminus \bigcup_{K \in \mathcal{K}} \widetilde{K} \cup \bigcup_{B \in \mathcal{B}} \widetilde{B})$.

Further, for each $\widetilde{K}$, $\beta_{adm} \cap \widetilde{K}$ is a union of elementary admissible paths with disjoint interiors such that

- (1) $\beta_{adm} \cap \widetilde{K}$ has at most one vertical path of type (2) above.
- (2) for any connected horizontal boundary component $\widetilde{\Sigma}_0^s$ of $\widetilde{K}$, $\beta \cap \widetilde{\Sigma}_0^s$ has at most one ‘horizontal’ geodesic of type (1) above.

Finally for each $B \in \mathcal{B}$, $\beta_{adm} \cap \tilde{B}$ is the admissible quasigeodesic corresponding to $\beta_{ea} \cap \tilde{B}$ in $\tilde{B}$.

Definition 4.17 allows us to replace an electro-ambient quasigeodesic by a path with greater control. We shall be concerned with electro-ambient quasigeodesics β_{ea} in $\tilde{M}$ starting and ending at points in $\tilde{H}$. Any such electro-ambient quasigeodesic has a representative entering and leaving split components at split level surfaces. This follows from the observation that any electro-ambient quasigeodesic necessarily enters and leaves split blocks as well as thick blocks along (horizontal) split level surfaces except at most for maximal pieces that have end-points in one of the thin tubes in $\mathcal{T}_l$. Hence the restriction on electro-ambient quasigeodesics β_{ea} given by (the first sentence of) Definition 4.17 may be regarded just as a choice of a representative of β_{ea} or as a mild normalization condition. The construction of the admissible quasigeodesic β_{adm} corresponding to β_{ea} changes this representative of β_{ea} by replacing the intersection of β_{ea} with each thick block $\tilde{B}$ or split component $\tilde{K}$ by one horizontal and at most one vertical piece. The reader should thus think of an admissible quasigeodesic β_{adm} corresponding to β_{ea} as an electro-ambient quasigeodesic

- (1) with the same entry and exit points as β_{ea} into either thick blocks (elements of $\tilde{B}$) or split components (elements of $\tilde{K}$).
- (2) if the entry and exit points p, q of β_{ea} into some $\tilde{B}$ (or $\tilde{K}$) lie on the same horizontal level, join p, q by a horizontal geodesic in the corresponding level surface to obtain the corresponding piece of β_{adm} .
- (3) if the entry and exit points p, q of β_{ea} into some $\tilde{B}$ (or $\tilde{K}$) lie on levels i and $i + 1$, say, then project q (using the product structure of the block B or K) to $P(q)$ on level i . Join $p, P(q)$ by a horizontal geodesic in the i -th level surface and then follow it by the vertical segment joining $P(q)$ to q .

The choice of an admissible quasigeodesic corresponding to an electro-ambient quasigeodesic is not unique. In each piece of the third type above, we can also choose the vertical segment at p and then follow by a horizontal geodesic in the $(i - 1)$ -th level surface. The ambiguity is bounded however in a sense made precise in Lemma 4.18 below.

The next Lemma allows us to apply the criterion for existence of Cannon-Thurston maps in Lemma 2.10 to admissible quasigeodesics in $\tilde{M}$ rather than electro-ambient quasigeodesics in $\tilde{M}$. The proof of Lemma 4.18 is exactly like Lemma 6.5 of [Mj14a] and we omit it here (see also the proof of Lemma 3.13 above).

Lemma 4.18. *Let N be the convex core of a complete hyperbolic 3-manifold N^h minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let M be the resulting model of split geometry. Then for all $C_0 > 0$, and $o \in \tilde{M}$ there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that the following holds.*

For any $a, b \in \tilde{M}$, let β_{ea}^h be an electro-ambient C_0 -quasigeodesic without backtracking in $\tilde{M}$ joining a, b . Let $\beta_{ea} = \beta_{ea}^h \setminus \partial\tilde{M}$ be the part of β_{ea}^h lying away from the (bi-Lipschitz) horospherical boundary of $\tilde{M}$. Again, let β_{adm}^h be the admissible quasigeodesic corresponding to β_{ea}^h and let $\beta_{adm} = \beta_{adm}^h \setminus \partial\tilde{M}$ be the part of β_{adm}^h lying away from the (bi-Lipschitz) horospherical boundary of $\tilde{M}$.

Then $d_M(\beta_{ea}, o) \geq n$ implies that $d_M(\beta_{adm}, o) \geq \Theta(n)$. Conversely, $d_M(\beta_{adm}, o) \geq n$ implies that $d_M(\beta_{ea}, o) \geq \Theta(n)$.

4.6. Cannon-Thurston Maps for Free Groups. We are now in a position to prove the existence of Cannon-Thurston maps for arbitrary finitely generated Kleinian groups. In this subsection we shall describe the proof for handlebody groups where the book-keeping is minimal. In the next subsection we shall indicate the modifications necessary for arbitrary finitely generated Kleinian groups. We identify H with its bi-Lipschitz image $F(H)$ contained in M under the bi-Lipschitz homeomorphism $F : N \rightarrow M$ from the hyperbolic manifold N to the model manifold M .

We now want to show that if $\lambda = [a, b]$ is a geodesic in the intrinsic metric on $\tilde{H}$ joining $a, b \in \tilde{H}$, and lying outside a large ball about a fixed reference point $p \in \tilde{H} \subset \tilde{M}$, then the (bi-Lipschitz) hyperbolic geodesic λ^h joining $a, b \in \tilde{M}$ also lies outside a large ball about p in $\tilde{M}$. This would guarantee the existence of a Cannon-Thurston Map by Lemma 2.10.

To fix notation, let Q be a free Kleinian group without parabolics. Let $N = \mathbb{H}^3/Q$, $M = F(N)$ and H a compact (Scott) core of M . $\tilde{H}$ with its intrinsic metric is quasi-isometric to the Cayley graph Γ_Q and so its intrinsic boundary may be identified with the Cantor set ∂Q thought of as the Gromov boundary of Γ_Q . Let $\hat{H}$ and $\hat{M}$ denote the compactifications by adjoining ∂Q and the limit set Λ_Q to $\tilde{H}$ and $\tilde{M}$ respectively.

Theorem 4.19. Cannon-Thurston for Free Groups *The inclusion $i : \tilde{H} \rightarrow \tilde{M}$ extends continuously to a map $\hat{i} : \hat{H} \rightarrow \hat{M}$.*

Proof. Let $p \in \tilde{H}$ be a base-point, and $\lambda = [a, b]$ be a geodesic in the intrinsic metric on $\tilde{H}$ and λ^h be the (bi-Lipschitz) hyperbolic geodesic joining its end-points in $\tilde{M}$. By Lemma 2.10 it suffices to show that if λ lies outside a large ball about p in $\tilde{H}$, then λ^h lies outside a large ball about p in $\tilde{M}$.

Suppose that λ lies outside an n -ball about p in $\tilde{H}$, i.e. $d_{\tilde{H}}(\lambda, p) \geq n$. Let $\{D_i\}$ be a finite collection of compressing disks in H such that each component of $\partial H \setminus \bigcup_i \partial D_i$ is a pair of pants.

Since each (lift of) D_i separates $\tilde{H}$ and since λ lies outside a large n -ball about p in $\tilde{H}$, we conclude that there exists such a lift D lying outside an $m = m(n)$ -ball about p in $\tilde{M}$, (i.e. $d_{\tilde{M}}(D, p) \geq m(n)$) and that λ lies in the component of $\tilde{H} \setminus D$ not containing p , where $m(n) \rightarrow \infty$ as $n \rightarrow \infty$. Let $A = D \cup \bigcup_i \sigma_i$ be the quasidisk containing D constructed in Section 4.3, where σ_i is a closed curve on the lift $\tilde{\Sigma}_i^s$ of the i -th split surface Σ_i^s to $\tilde{M}$. Also, let $\tilde{\Sigma}_i^s \setminus \sigma_i = \tilde{\Sigma}_{i+}^s \cup \tilde{\Sigma}_{i-}^s$ where $\tilde{\Sigma}_{i+}^s, \tilde{\Sigma}_{i-}^s$ are the two components of $\tilde{\Sigma}_i^s \setminus \sigma_i$. Similarly, let $\tilde{H} \setminus D = \tilde{H}_+ \cup \tilde{H}_-$, where $\tilde{\Sigma}_{0+}^s \subset \partial \tilde{H}_+$ and $\tilde{\Sigma}_{0-}^s \subset \partial \tilde{H}_-$. Assume without loss of generality that $\lambda \subset \tilde{H}_+$ and $p \in \tilde{H}_-$. Let $\tilde{M}_+ = \tilde{H}_+ \cup \bigcup_i \tilde{\Sigma}_{i+}^s$ and $\tilde{M}_- = \tilde{H}_- \cup \bigcup_i \tilde{\Sigma}_{i-}^s$. Also let $\tilde{M}_H = \bigcup_i \tilde{\Sigma}_i^s$ be the union of all the horizontal split surfaces lifted to $\tilde{M}$.

Let α be an electro-ambient quasigeodesic joining the end-points of λ in $\tilde{M}$ and β be an admissible quasigeodesic corresponding to α .

Since β is admissible, it consists of horizontal and vertical pieces. Two cases arise:

- a) $\beta \cap \widetilde{M}_H \subset \widetilde{M}_+$
 b) $\beta \cap (\widetilde{M}_+)^- \cap (\widetilde{M}_-)^- \neq \emptyset$.

Roughly speaking Cases (a) and (b) correspond respectively to the cases where β does not or does intersect A coarsely.

Case a: $\beta \cap \widetilde{M}_H \subset \widetilde{M}_+$

By Corollary 4.15 and Lemma 4.18 it suffices to show that there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that the following holds.

$d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\beta, p) \geq \Theta(n)$.

The existence of such a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ follows exactly as in Lemma 4.13. We briefly recount the proof. It suffices to prove that if $d_{\widetilde{M}}(D, p) \geq m(n)$, then $d_{\widetilde{M}}(A, p) \geq \Theta(n)$. As in the proof of Lemma 4.13, there exist constants $b_1, \dots, b_k, \dots$ (depending only on the geometry of E and the split geometry model) and a $z \in D$ such that for any $x_k \in \sigma_k(\subset A)$, we have $d_M(x_k, z) \leq b_1 + \dots + b_k = c_k$ (say). Hence $d_M(x_k, p) \geq m(n) - c_k$. Also, using uniform separatedness, $d_M(x_k, z) \geq k\epsilon_0$. Therefore there exists a proper function $\Theta_0 : \mathbb{N} \rightarrow \mathbb{N}$ such that $d_M(A, p) \geq \Theta_0(n)$. Since β is disjoint from A and is admissible, it follows that every horizontal segment of β is at distance at least $\Theta_0(n)$ from p .

Any vertical segment τ_k of β necessarily joins a pair of points y_k, y_{k+1} , where y_k, y_{k+1} lie on the top and bottom horizontal boundaries of the same block (thick or split) with y_k (resp. y_{k+1}) lying on the same horizontal split level surface as σ_k (resp. σ_{k+1}). Further, the vertical segment has length at most C_k – the thickness of the k -th block. It follows that any point on τ_k lies at distance at least $\Theta_0(n) - C_k$ from p . Again, there exists ϵ_0 (a lower bound on the separation between successive horizontal levels) such that any point on τ_k lies at distance at least $k\epsilon_0$ from p . Choosing $\Theta(n) = \max\{\Theta_0(n) - C_k, k\epsilon_0\}$, we are done.

Case b: $\beta \cap (\widetilde{M}_+)^- \cap (\widetilde{M}_-)^- \neq \emptyset$.

We shall say that β crosses A at x if either $x \in \beta \cap A$, or if there exists a vertical elementary admissible subpath $x \times I \subset \beta$, such that either $(x, 0) \in \widetilde{M}_+$ and $(x, 1) \in \widetilde{M}_-$ or $(x, 1) \in \widetilde{M}_+$ and $(x, 0) \in \widetilde{M}_-$.

Let r, q be the first and last points at which β crosses A . Let β_{ar}, β_{qb} be the subpaths of β joining a, r and b, q respectively. Then again, as in Case (a) above, there exists a function $\Theta : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that $d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\beta_{ar} \cup \beta_{qb}, p) \geq \Theta(n)$. Hence by Corollary 4.15 and Lemma 4.18 it follows that there exists a function $\Theta_2 : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta_2(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that $d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\mu_{ar} \cup \mu_{qb}, p) \geq \Theta_2(n)$, where μ_{ar} (resp. μ_{qb}) are the (bi-Lipschitz) hyperbolic geodesics in $\widetilde{M}$ joining a, r and b, q respectively.

If $r, q \in A$, then, since $d_M(D, p) \geq m(n)$, Corollary 4.16 ensures the existence of a proper function $\Theta_3 : \mathbb{N} \rightarrow \mathbb{N}$ (i.e. $\Theta_3(n) \rightarrow \infty$ as $n \rightarrow \infty$) such that $d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\mu_{rq}, p) \geq \Theta_3(n)$, where μ_{rq} is the (bi-Lipschitz) hyperbolic geodesic in $\widetilde{M}$ joining r and q . Hence by δ -hyperbolicity of $\widetilde{M}$, $d_M(\mu_{ab}, p) \geq \min(\Theta_2(n), \Theta_3(n)) - 2\delta$ and we are done.

Else, let $r \in \widetilde{\Sigma}_i^s$ and $q \in \widetilde{\Sigma}_j^s$. There exists $r' \in \widetilde{\Sigma}_{i+1}^s$, such that r, r' lie on a vertical segment in β . Further they lie on opposite (+ and -) sides of A . Similarly, there exists $s' \in \widetilde{\Sigma}_{j-1}^s$, such that s, s' lie on a vertical segment in β and on opposite (- and +) sides of A . Hence, (depending on thickness of the m -th block) there

exist $C(m), m \in \mathbb{N}$ such that $d_M(r, A) \leq C(i)$ and $d_M(q, A) \leq C(j)$. Choose r_1, q_1 in σ_i, σ_j respectively such that $d_M(r, r_1) \leq C(i)$ and $d_M(q, q_1) \leq C(j)$.

Then, by Corollary 4.16, there exists a function $\Theta_4 : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\Theta_4(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that $d_M(\mu_{r_1 q_1}, p) \geq \Theta_4(n)$, where $\mu_{r_1 q_1}$ is the (bi-Lipschitz) hyperbolic geodesic in $\widetilde{M}$ joining r_1 and q_1 . The existence of $C(m), m \in \mathbb{N}$ now guarantees the existence of a proper function $\Theta_5 : \mathbb{N} \rightarrow \mathbb{N}$ such that $d_M(\mu_{rq}, p) \geq \Theta_5(n)$ (reprising, for instance, the argument in the last paragraph of Case (a) above). Hence by δ -hyperbolicity of $\widetilde{M}$ again, $d_M(\mu_{ab}, p) \geq \min(\Theta_2(n), \Theta_5(n)) - 2\delta$ and we are through. $\square$

4.7. Finitely Generated Kleinian Groups. In this subsection, we indicate the modifications necessary in the proof of Theorem 4.19 to prove the analogous theorem for finitely generated Kleinian groups.

Let N_{gf} denote the augmented Scott core of $N^h = \mathbf{H}^3/G$. Let $i : N_{gf} \rightarrow N^h$ be the natural inclusion map. Thurston showed that there exists a geometrically finite manifold admitting a strictly type-preserving homotopy equivalence with N^h . The convex core of such a manifold admits a proper homeomorphism to N_{gf} . Thus, N_{gf} may be thought of as the convex core of a geometrically finite manifold admitting a strictly type-preserving homotopy equivalence with N^h . Let H be N_{gf} with open neighborhoods of cusps removed. Let N be N^h with open neighborhoods of cusps removed. Then $\widetilde{H}$ is strongly hyperbolic relative to its horospheres and $\widetilde{N}$ is strongly hyperbolic relative to its horospheres. Let $\widehat{H}$ and $\widehat{N}$ denote their relative hyperbolic compactifications. Note that $\widehat{H} = \widehat{N_{gf}}$, where $\widehat{N_{gf}}$ is the Gromov compactification of the hyperbolic space $\widetilde{N_{gf}}$. Similarly, $\widehat{N} = \widehat{N^h}$, where $\widehat{N^h}$ is the Gromov compactification of the hyperbolic space $\widetilde{N^h}$. Let $\widetilde{i} : \widetilde{H} \rightarrow \widetilde{N}$ indicate the lift of i . Let M denote the model manifold for N and let $\widehat{M}$ denote the relative hyperbolic compactification of $\widetilde{M}$. We identify H with its bi-Lipschitz image in M under the bi-Lipschitz homeomorphism $F : N \rightarrow M$. Let d_G be the graph metric on $\widetilde{M}$ equipped with a split geometry structure where all split components are incompressible.

First, suppose that H has incompressible boundary as a pared manifold. Then Theorem 2.13 shows that a Cannon-Thurston map exists for $\widetilde{i} : \widetilde{H} \rightarrow \widetilde{M}$. The point pre-image description is also furnished by Theorem 2.13.

Else H may be decomposed as the disk-connected sum (or boundary-connected sum) $\#_{i=1, \dots, r+s} H_i$ of $H_1, H_2 \dots H_{r+s}$, where

- (1) $H_{r+1}, \dots, H_{r+s}$ are pared manifolds with incompressible boundary.
- (2) $H_1, \dots, H_r$ are handlebodies such that in the boundary connected sum decomposition of H , no two are connected to the same boundary component of $\#_{i=r+1, \dots, r+s} H_i$. (This ensures that the boundary connected sum decomposition is minimal.)

Let $D'_1, \dots, D'_{r+s-1}$ be the compressing disks obtained from the above boundary connected sum decomposition. Next for each handlebody H_i ($i = 1, \dots, r$), we choose a minimal collection of disjoint non-separating compressing disks such that their complement in H_i is a ball. Taking the union of $D'_1, \dots, D'_{r+s-1}$ with all these compressing disks for H_i ($i = 1, \dots, r$), we obtain a collection $D_i, i = 1 \dots m$ of compressing disks. Since $m \geq 1$, there is at least one compressing disk. Note

that the collection ∂D_i , $i = 1 \cdots m$ forms a maximal collection of homotopically distinct compressible simple closed curves on ∂H .

Theorem 4.20. Cannon-Thurston for Kleinian Groups *Let H, M, N, N_{gf}, N^h be as above. Further suppose that each degenerate end of M admits a Minsky model. The inclusion $\tilde{i} : \tilde{H} \rightarrow \tilde{M}$ extends continuously to a map $\hat{i} : \hat{H} \rightarrow \hat{M}$ between the relative hyperbolic compactifications. Equivalently, $\tilde{i} : \tilde{N}_{gf} \rightarrow \tilde{N}^h$ extends continuously to a map $\hat{i} : \hat{N}_{gf} \rightarrow \hat{N}^h$ between the hyperbolic compactifications.*

Proof: From Propositions 4.5, 4.8 and 4.12, we can construct quasidisks A_i corresponding to D_i as before and lift them to $\tilde{M}$ (after partially electrocuting $\mathbb{Z}$ -cusps if any).

Now, let λ be a geodesic segment in $\tilde{H}$ lying outside a large ball $B_N(p)$ for a fixed reference point p . λ may be decomposed into (at most) three pieces λ_- , λ_0 and λ_+ as follows.

- (1) The middle piece λ_0 does not intersect any of the (lifts of the) compressing disks D_i in the interior. (We thus allow for the cases where λ_- and/or λ_+ are empty.)
- (2) the common end-point $\lambda_- \cap \lambda_0$ lies on some D_i . The same is demanded of $\lambda_0 \cap \lambda_+$.
- (3) the point q on λ nearest to p lies on λ_0 .
- (4) If λ intersects exactly one disk D , then λ_0 is defined to be the piece of λ ending on D and containing q ; the other piece being designated λ_- or λ_+ .

We shall consider a sequence of λ 's converging to a point ξ on the boundary of $\hat{H}$ (or $\hat{N}_{gf}$) in the Hausdorff topology. (Equivalently, the sequence of end-points of the λ 's converge to ξ .) Such a sequence must necessarily lie outside larger and larger balls $B_n(p)$ about p . Two cases arise.

Case A: For a sequence of λ 's lying outside larger and larger balls $B_n(p)$ about p , and converging to ξ , the sequence of λ 's eventually lies outside any fixed component of $\tilde{H} \setminus \cup_i D_i$. This is exactly the same case as in the proof of Theorem 4.19. The same proof goes through by Corollary 4.16.

Case B: Either $[p, q]$ does not intersect any D_i or there is (up to subsequencing) a fixed disk D_i that is the last disk that $[p, q]$ intersects. Since D_i is of uniformly bounded diameter, we may shift our base point to a point p' in the component $\tilde{H}_i$ which is the lift of H_i containing q . In this case, there exists a fixed n_0 such that λ lies outside $B_{(n-n_0)}(p')$. By shifting origin, we rewrite p' as p and $(n - n_0)$ as n .

Step 1: Now, $\lambda_0 \subset \tilde{H}_i \subset \tilde{H}$ as it does not meet any disk D_i in its interior. Since H_i is either a handlebody without parabolics, or a pared manifold with incompressible boundary, then by Theorem 4.19 or Theorem 2.13 respectively, a Cannon-Thurston map exists for the inclusion $\tilde{H}_i \subset \tilde{M}$. By (the necessity part of) Lemma 2.10 and Lemma 2.11, it follows that the hyperbolic geodesic λ_0^h joining the end-points of λ_0 in $\tilde{N}^h$ lies outside a large ball about p . Thus, there exists $m_1(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that λ_0^h lies outside a ball of radius $m_1(n)$ about p in $\tilde{N}^h$.

Step 2: If λ_+ (or λ_-) is non-empty, then λ_- (or λ_+) is separated from p by a disk $D_i \subset \tilde{H}$ lying outside $B_n(p)$.

Note: Strictly speaking some uniformly bounded pieces of λ_+ (or λ_-) close to the intersection point with λ could enter $\widetilde{H}_i$. To see the existence of a uniform bound, note first that $\widetilde{H}$ is quasi-isometric to (the Cayley graph of) $\pi_1(M)(= \pi_1(H))$. Then the disks D_i correspond to splittings of the group $\pi_1(H)$. Geodesic words in $\pi_1(H)$ cannot ‘backtrack’, i.e. in the tree of spaces description of $\widetilde{H}$ corresponding to splittings by D_i , such paths cannot re-enter a vertex space after leaving it. Uniform boundedness now follows from the quasi-isometry between $\widetilde{H}$ and (the Cayley graph of) $\pi_1(M)(= \pi_1(H))$.

However they can be replaced first by paths lying entirely on D_i and then pushed out of $\widetilde{H}_i$ altogether. Thus, replacing the geodesic segments λ_- (or λ_+) by quasi-geodesics if necessary, we can ensure that λ_- (or λ_+) is separated from p by a disk $D_i \subset \widetilde{H}$ lying outside $B_n(p)$. We shall ignore this mild technicality.

Recall that M is a bi-Lipschitz model for N , which in turn is N^h with cusps removed. Also recall that d_G is the graph metric on $\widetilde{M}$. Then the quasidisk A_i is quasiconvex in $(\widetilde{M}, d_G)$ and lies outside a large ball of radius $m_2(n)$ about p , where $m_2(n) \rightarrow \infty$ as $n \rightarrow \infty$. Let λ_+^h (resp. λ_-^h) be the hyperbolic geodesics joining the end-points of λ_+ (resp. λ_-). Again, by constructing admissible paths and electro-ambient quasigeodesics as in the proof of Theorem 4.19, we obtain a new function $m_3(n)$ such that $m_3(n) \rightarrow \infty$ as $n \rightarrow \infty$ and so that the hyperbolic geodesics λ_-^h or λ_+^h lie outside a ball of radius $m_3(N)$ about p in $\widetilde{N}^h$.

Step 3: Therefore $\lambda_-^h \cup \lambda_0^h \cup \lambda_+^h$ lies outside a ball of radius $m_4(n) = \min\{m_1(n), m_3(n)\}$.

Finally, since $\widetilde{N}^h$ is δ -hyperbolic, the hyperbolic geodesic λ^h joining the end-points of λ lies outside a ball of radius $m(n) = m_4(n) - 2\delta$ about p . Also, $m(n) \rightarrow \infty$ as $n \rightarrow \infty$. Therefore, by Lemma 2.10 or 2.11, it follows that the inclusion $\tilde{i} : \widetilde{H} \rightarrow \widetilde{M}$ extends continuously to a map $\hat{i} : \widehat{H} \rightarrow \widehat{M}$ or equivalently that $\tilde{i} : \widetilde{N}_{gf} \rightarrow \widetilde{N}^h$ extends continuously to a map $\hat{i} : \widehat{N}_{gf} \rightarrow \widehat{N}^h$. This concludes the proof. $\square$

Remark 4.21. As mentioned in Remark 1.2, the hypothesis that each degenerate end of M admits a Minsky model is expected to be superfluous and is established in Appendix 7 when M has no parabolics.

Let $\widehat{G_F}$ denote the Floyd compactification of a group G (See [Flo80]). McMullen conjectured in [McM01] that there exists a continuous extension of $i : \Gamma_G \rightarrow \widetilde{M}$ to a map from $\widehat{G_F}$ to $\widehat{M}$. It was shown by Floyd in [Flo80] that there is a continuous map from $\widehat{G_F}$ to $\widehat{H}$. Combining this with Theorem 4.20 above for Kleinian groups with parabolics, we get a proof of the following, which proves McMullen’s conjecture.

Theorem 4.22. *Let G be any finitely generated Kleinian group and $M = \mathbf{H}^3/G$. Further suppose that each degenerate end of M admits a Minsky model (for instance if M has no parabolics). Then there is a continuous extension $\hat{i} : \widehat{G_F} \rightarrow \widehat{M}$.*

5. POINT PRE-IMAGES OF THE CANNON-THURSTON MAP

In this section, we first determine the pre-images of multiple points under the Cannon-Thurston map for degenerate free Kleinian groups G . We then indicate the modifications necessary to extend the results to arbitrary finitely generated Kleinian groups. This is done for two reasons:

a) Otal had specifically conjectured [Ota88] the structure of the Cannon-Thurston maps we prove for handlebody groups.

b) The additional work required for arbitrary finitely generated Kleinian groups involves extra book-keeping with respect to a finite family of ends.

The extra generality at this stage would tend to clutter up the exposition.

We shall not have need to distinguish between the hyperbolic manifold and its bi-Lipschitz model any longer and will denote the manifold by M .

We set up some notation for the purposes of this section. Let G be a free degenerate Kleinian group without parabolics. Suppose that G is not geometrically finite. Let $M = \mathbb{H}^3/G$ be the quotient manifold. Note that the limit set of G is all of the sphere at infinity. Hence M is its own convex core. Let H be a compact core of M . H is a handlebody whose inclusion into M induces a homotopy equivalence. In fact, M deformation retracts onto H . Then $\tilde{H}$ is embedded in $\tilde{M} = \mathbb{H}^3$. Let Γ denote the Cayley graph of G with respect to some finite generating set of G . Assume that Γ is equivariantly embedded in $\tilde{H}$ with edges being mapped to geodesic segments. Let S denote the boundary surface of H . We assume that the ending lamination Λ_{EL} is a geodesic lamination on S equipped with some (any) hyperbolic metric. This is well-defined only up to Dehn twists along simple closed curves in S that bound disks in H and gives a well-defined ending lamination in the Masur domain by Theorem 4.4. To make this explicit, we denote the ending lamination in the Masur domain by Λ_{ELH} . $M \setminus \text{Int}(H)$ is homeomorphic to $S \times [0, \infty)$ and is bi-Lipschitz homeomorphic to an end M_S of a simply degenerate hyperbolic manifold without accidental parabolics [Bow05] [BCM14]. Thus $S \times [0, \infty) \subset M$ equipped with its *intrinsic* path metric is bi-lipschitz homeomorphic to M_S . We shall have need to pass interchangeably between these two below.

5.1. EL leaves are CT leaves. Let $i : \tilde{H} \rightarrow \tilde{M}$ denote the inclusion. Let ∂i denote the continuous extension of i to the boundary in Theorem 4.19. Note that the inclusion of Γ into $\tilde{H}$ with its intrinsic metric is a quasi-isometry. So we might as well replace the inclusion of Γ into $\tilde{M}$ by that of $\tilde{H}$ into $\tilde{M}$. We shall show that point pre-images of multiple points under ∂i correspond to end-points of leaves of an ending lamination in the Masur domain.

The inclusion of S into H as its boundary induces a surjection of fundamental groups. Let N denote the kernel. Let $S_N (= \partial \tilde{H})$ denote the cover of S corresponding to N .

To distinguish between the ending lamination Λ_{ELH} (in the Masur domain) and bi-infinite geodesics whose end-points are identified by ∂i , we make the following definition.

Definition 5.1. A **CT leaf** λ_{CT} is a bi-infinite geodesic whose end-points are identified by ∂i .

An **EL leaf** λ_{EL} is a bi-infinite geodesic whose end-points are ideal boundary points of either a leaf of the ending lamination, or a complementary ideal polygon.

We shall show that

- An *EL leaf* is a *CT leaf*.
- A *CT leaf* is an *EL leaf*.

Proposition 5.2. EL is CT Let G be a free degenerate Kleinian group without parabolics. Let u, v be either ideal end-points of a leaf of an ending lamination of

G , or ideal boundary points of a complementary ideal polygon. Then $\partial i(u) = \partial i(v)$.

Proof. This is almost identical to Proposition 3.1 of [Mj14b]. However, since the setup is somewhat different we include a sketch of a proof. Take a sequence of short geodesics s_i exiting the end. Let a_i be geodesics in the intrinsic metric on the boundary S (of H) freely homotopic to s_i . By topological tameness [Ago04] [GC06] and geometric tameness ([Thu80] Ch. 9) we may assume further that a_i 's are simple closed curves on S . Join a_i to s_i by the shortest geodesic t_i in $S \times [0, \infty)$ connecting the two curves. Then the collection a_i may be chosen to converge to the ending lamination on S ([Thu80] Ch. 9). Also, in $S_N \subset \tilde{H} \subset \tilde{M}$, we choose lifts $a_i \subset \tilde{S}$ (of a_i) which are finite segments whose end-points are identified by the covering map $P : S \times [0, \infty) \rightarrow S \times [0, \infty)$. We also assume that P is injective restricted to the interior of a_i 's mapping to a_i . Similarly there exist segments $s_i \subset \tilde{M}$ which are finite segments whose end-points are identified by the covering map $P : \tilde{M} \rightarrow M$. We also assume that P is injective restricted to the interior of s_i 's. The finite segments s_i and a_i are chosen in such a way that there exist lifts t_{1i} , t_{2i} , joining end-points of a_i to corresponding end-points of s_i . The union of these four pieces looks like a trapezium (see below, where we have omitted subscripts for convenience).




Figure: Trapezium

Next, given any leaf λ of the ending lamination, we may choose translates of the finite segments a_i (under the action of $\pi_1(H)$) appropriately, such that they converge to λ in S_N . For each a_i , let

$$b_i = t_{1i} \circ s_i \circ t_{2i}^{-1}$$

where t_{2i}^{-1} denotes t_{2i} with orientation reversed. If the translates of a_i we are considering have end-points lying outside large balls around a fixed reference point $p \in S_N$, it is not hard to see that b_i 's lie outside large balls about p in $\tilde{M}$.

Here is a quick sketch. We think of t_i as vertical and a_i, s_i as horizontal. We justify this heuristic. We choose t_i to be a distance minimizing geodesic between a loop in $S \times \{0\}$ corresponding to a_i and loop in $S \times \{i\}$ corresponding to s_i . Saying that t_i is vertical is meant to convey the idea that t_i is indeed well-behaved with respect to the block structure (coming from split geometry of the end) in terms of the progress it makes out the end. A (coarse) measure of the vertical co-ordinate of $z \in t_i$ is the number of (thick or split) blocks that lie 'below' z . The segments t_{1i}, t_{2i} make *definite progress* out the end $S \times [0, \infty)$ in the sense that the number of

blocks below z (i.e. coarse vertical co-ordinate of z) is a **proper** function (say Φ) of the length of the initial segment of $\underline{t}_i$ up to z . Further, this proper function Φ is independent of the index i . This follows from the fact that a distance minimizing geodesic like $\underline{t}_i$ can spend only a bounded amount of time in each (split or thick) block. The bound depends on the block in general but this is adequate for definite progress.

Next, t_{1i}, t_{2i} are both lifts of $\underline{t}_i$. Similarly s_i is contained in $\tilde{S} \times \{i\}$ and a_i is contained in $\tilde{S} \times \{0\}$ allowing us to legitimize the statement that they are ‘horizontal’. Thus each S_i and paths lying on them are being thought of as horizontal and paths that make definite progress transversely are being thought of as vertical.

Also $\underline{s}_i$ lies at a large vertical height (at least $O(i)$) from H and hence s_i lies at a large distance (at least $O(i)$) from p . Next, the initial point of t_{1i} lies at a large horizontal distance d_{hor} from p . Note that d_{hor} may be chosen to be as large as we like (independent of i). The vertical distance from p increases monotonically (according to the proper function Φ independent of i) as we proceed along t_{1i} . The horizontal component (of a point z on t_{1i}) can go down at each step (as we move from the m -th block to the $(m+1)$ -th block) by a constant depending only on m . Since d_{hor} may be chosen to be arbitrarily large, it follows that for any point on t_{1i} (or t_{2i}) at least one of the vertical or horizontal distances is large.

Note: A parenthetical comment is in order. It is indeed possible (as was pointed out to us by the referee) to have a situation where

- (1) geodesics t_{1i} and t_{2i} in $\mathbb{H}^3$ start and end outside large compacts
- (2) and the geodesic a_i also lies outside large compacts.
- (3) But the geodesic σ_i joining the endpoint of b_i do intersect a given compact K_o .

To ensure this, the geodesics t_{1i} and t_{2i} must lie very close to σ_i for a very long period of time, and hence must themselves come close to K_o . This is prevented in our situation by the definite progress of $\underline{t}_i$ (and hence t_{1i} or t_{2i}).

Returning to the proof, we summarize by saying that t_{1i} (and similarly t_{2i}) lie at a large distance from p . Since s_i is uniformly bounded in length, it follows that b_i lies at a large distance from p .

Since $\mathbb{H}^3$ is δ -hyperbolic for some $\delta > 0$, it follows that the geodesic joining the end-points of b_i (and hence a_i which has the same end-points) lies in a 2δ -neighborhood of b_i .

At this stage we invoke the existence theorem for Cannon-Thurston maps, Theorem 4.19. Since a_i ’s converge to λ and the hyperbolic geodesics joining the end-points of a_i exit all compact sets, it follows that $\partial i(u) = \partial i(v)$, where u, v denote the boundary points of λ . The Proposition follows. $\square$

Generalization to arbitrary finitely generated Kleinian groups: Any finitely generated Kleinian group is geometrically tame ([Ago04] [GC06] [Thu80] Ch. 9) and has finitely many ends. Observe that the proof of the above Proposition used the freeness of G only at the stage of applying Theorem 4.19. The same proof goes through verbatim for freely decomposable Kleinian groups with degenerate ends. The only modification to the above proof is that we consider one end of the manifold M at a time (and the pigeon-hole principle) along with Theorem 4.20 in place of Theorem 4.19 to obtain the following Proposition.

Proposition 5.3. EL is CT - General Case *Let G be a finitely generated freely decomposable Kleinian group. Let u, v be either ideal end-points of a leaf of an ending lamination of G , or ideal boundary points of a complementary ideal polygon. Then $\partial i(u) = \partial i(v)$.*

5.2. CT leaves are EL leaves. As usual we deal first with free degenerate groups without parabolics. We restate Theorem 2.13 in a form that we shall use. Recall that $M \setminus \text{Int}(H)$ is homeomorphic to $S \times [0, \infty)$ and is bi-Lipschitz homeomorphic to an end M_S of a simply degenerate hyperbolic manifold without accidental parabolics [Bow05] [BCM14]. Hence by Theorem 2.13 we have the following.

Theorem 5.4. [Mj14b] *Let S, M_S be as above. Then the inclusion of universal covers $\tilde{j} : \tilde{S} \rightarrow \tilde{M}_S$ extends continuously to the boundary. Further, pre-images of points on the boundary are precisely ideal boundary points of a leaf of the ending lamination Λ_{ELS} of M_S , or ideal boundary points of a complementary ideal polygon whenever the Cannon-Thurston map is not one-to-one.*

We identify the Cayley graph Γ of the free group with a subset of $\tilde{H} \subset \tilde{M}$, viz. the orbit of a base-point joined by edges. The next Theorem is one of the main Theorems of this paper.

Theorem 5.5. *Let G be a free degenerate Kleinian group without parabolics. Let $i : \Gamma_G \rightarrow \mathbb{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbb{H}^3$. Then i extends continuously to a map $\hat{i} : \widehat{\Gamma}_G \rightarrow \mathbb{D}^3$, where $\widehat{\Gamma}_G$ denotes the (Gromov) hyperbolic compactification of Γ_G . Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial \Gamma_G$ of Γ_G .*

Then $\partial i(a) = \partial i(b)$ for $a \neq b \in \partial \Gamma$ iff a, b are either ideal end-points of a leaf of an ending lamination of G , or ideal boundary points of a complementary ideal polygon.

Proof. By Theorem 4.19 the inclusion $i : \Gamma \rightarrow \tilde{M}$ extends continuously to a map between the Gromov compactifications $\hat{i} : \widehat{\Gamma} \rightarrow \mathbb{D}^3$. Let ∂i denote the values of the above continuous extension to the boundary. Suppose $\partial i(a) = \partial i(b)$. Λ_{EL} is the ending lamination of M regarded as a subset of S . Let Λ_{ELG} denote Λ_{EL} lifted to $S_G = \partial \tilde{H}$, which is a cover of S . We want to show that a, b are the end-points of a leaf of Λ_{ELG} . Suppose $(a, b)_\Gamma$ is the bi-infinite geodesic from a to b in $\Gamma \subset \tilde{M}$. Assume without loss of generality that (a, b) passes through $1 \in \Gamma$. Let a_k, b_k be points on (a, b) such that $a_k \rightarrow a$ and $b_k \rightarrow b$. Let $\overline{a_k b_k}$ denote the geodesic in $\tilde{M}$ joining a_k, b_k . By continuity of the Cannon-Thurston map (Theorem 4.19) there exists $N(k) \rightarrow \infty$ as $k \rightarrow \infty$ such that $\overline{a_k b_k}$ lies outside an $N(k)$ ball about $1 \in \Gamma \subset \tilde{M}$, where radius is measured in the hyperbolic metric on $\tilde{M}$. Isotoping $\overline{a_k b_k}$ slightly, we can assume without loss of generality that it meets $\Gamma \subset \tilde{M}$ only at its end-points (since Γ is one dimensional). We can further isotope $\overline{a_k b_k}$ rel. endpoints by a bounded amount (depending on the Hausdorff distance between $\tilde{H}$ and $\Gamma \subset \tilde{H}$) such that $\overline{a_k b_k} \cap S_G = \{c_k, d_k\}$, where.

- 1) $c_k, d_k \in S_G = \partial \tilde{H}$ with $d(a_k, c_k)$ and $d(b_k, d_k)$ uniformly bounded (independent of k)
- 2) if $\overline{c_k d_k}$ denotes the subpath of $\overline{a_k b_k}$ between c_k, d_k then (modifying $N(k)$ by an additive constant if necessary) $\overline{c_k d_k}$ lies outside an $N(k)$ ball about $1 \in \Gamma \subset \tilde{M}$.

3) $\overline{c_k d_k}$ intersects $\tilde{H}$ only at the endpoints c_k, d_k .

Thus $\overline{a_k b_k}$ is a concatenation of three pieces, $\overline{a_k c_k}$, $\overline{c_k d_k}$, $\overline{d_k b_k}$, where $\overline{a_k c_k}$ and $\overline{d_k b_k}$ are uniformly bounded in length and lie in $\tilde{H}$, whereas $\overline{c_k d_k}$ lies in $\tilde{M} \setminus \text{Int}(\tilde{H})$.

Let $[c_k, d_k]_G$ denote the geodesic in the intrinsic metric on S_G which is homotopic (rel. endpoints) to $\overline{c_k d_k}$ in $\tilde{M} \setminus \text{Int}(\tilde{H})$. Since G is free, we can assume that its Cayley graph is a tree and (since $\tilde{H}$ is quasi-isometric to Γ) $[c_k, d_k]_G$ passes through a point $o_k \in S_G$ at a uniformly bounded distance from $1 \in \Gamma$.

Recall that $\tilde{S}(\subset \tilde{M}_S)$ is the universal cover of S inside the universal cover of the end M_S . Lift $[c_k, d_k]_G$ to some geodesic $\widetilde{c_k, d_k}(\subset \tilde{S})$ in the intrinsic metric on $\tilde{S}$. Further assume that there exists some fixed $o \in \tilde{S}$ such that the corresponding lift o'_k of o_k lies in a uniformly bounded neighborhood of o . Let $(\widetilde{c_k d_k})^\sim$ denote the corresponding lift of $\overline{c_k d_k}$ having the same endpoints as $\widetilde{c_k, d_k}$ (such a choice is possible as $[c_k, d_k]_G$ and $\overline{c_k d_k}$ are homotopic rel. endpoints in the complement of $\text{Int}(\tilde{H})$ in $\tilde{M}$). It follows that $(\widetilde{c_k d_k})^\sim$ lies outside an $N(k)$ -ball about o'_k in $\tilde{M}_S$. Hence (modifying $N(k)$ by a further additive constant if necessary), $(\widetilde{c_k d_k})^\sim$ lies outside an $N(k)$ -ball about $o \in \tilde{M}_S$. Therefore, by the existence of Cannon-Thurston maps for $j : \tilde{S} \rightarrow \tilde{M}_S$ (Theorem 5.4) it follows that if $c_\infty d_\infty$ denotes any subsequential limit of the segments $\widetilde{c_k, d_k}$ on $\tilde{S}$, then $\partial j(c_\infty) = \partial j(d_\infty)$ and hence again by Theorem 5.4 c_∞, d_∞ are end-points of leaves (or vertices of a complementary ideal polygon) of the ending lamination Λ_{ELS} of $S \subset M_S$. Finally, since c_∞, d_∞ is a bi-infinite geodesic passing through a bounded neighborhood of o , it projects to a leaf (or diagonal of a complementary ideal polygon) of Λ_{ELG} in S_G . Such leaves are also well-defined as leaves of the ending lamination Λ_{ELH} , i.e. as leaves of the ending lamination of M regarded as an element of the Masur domain, cf. Theorem 4.4. We have thus finally shown that $\Lambda_{CT} \subset \Lambda_{ELH}$. Combining this with Proposition 5.2 and Theorem 4.19 we have the Theorem. $\square$

Note that in the proof of Theorem 5.5 we have used freeness of G to conclude only two things:

- 1) The manifold M has exactly one end.
- 2) The path λ in $\tilde{H}$ can be isotoped off the Cayley graph of G embedded in $\tilde{H}$.

To prove an analogue of Theorem 5.5 for arbitrary finitely generated Kleinian groups we continue with the notation that M is a hyperbolic manifold with augmented Scott core H . Then M has finitely many ends. We first note, that if $\lambda = (a_\infty, b_\infty)$ is a CT leaf then there exist $a_n \rightarrow a_\infty$ and $b_n \rightarrow b_\infty$ such that the geodesic realizations μ_n of $[a_n, b_n]$ in $\tilde{M}$ leave arbitrarily large compact sets. We may assume that $\tilde{M} \setminus \tilde{H}$ consists of lifts of the ends of M to $\tilde{M}$. Each μ_n intersects finitely many such lifts of ends and hence has subsegments $\mu_{n1}, \dots, \mu_{nk}$, where each μ_{ni} lies in a lift $\widetilde{H \cup E_i}$ for some end E_i of M . Assume that such a decomposition is minimal (i.e. k is the minimal possible for μ_n). Then, since μ_n leaves arbitrarily large compact sets, so must each μ_{ni} . Further, each μ_{ni} has end-points in $\tilde{H}$. It follows that there will be at least one of the μ_{ni} 's - call it ν_n - such that

- (a) ν_n is contained entirely in one of these lifts of the ends
- (b) Endpoints c_n, d_n of ν_n lie on $\tilde{H}$
- (c) $c_n \rightarrow c_\infty$ and $d_n \rightarrow d_\infty$, where c_∞, d_∞ lie in the boundary of $\tilde{H}$.
- (d) Finally, by considering all segments ν_n (the non-uniqueness of ν_n is used at this stage) satisfying properties (1)-(3), there exists a finite sequence $a_\infty = a_0, \dots, a_n = b_\infty$ such that each pair (a_i, a_{i+1}) arises as a limiting pair c_∞, d_∞ as in (3).

We may therefore assume for the time being that μ_n lies in precisely one of the lifts of the ends E of M . If $S^h = H \cap E$ be its boundary then the ending lamination lies in the boundary of the (relatively) hyperbolic group $j_*(\pi_1(S^h))$ (hyperbolic relative to the cusp groups if any), where $j : S^h \rightarrow M$ is inclusion.

Fact (2) now goes through for arbitrary finitely generated Kleinian groups, as the inclusion of the augmented Scott core into M is a homotopy equivalence (in fact a deformation retract) and we are only interested in leaves which are limits of segments whose geodesic realizations lie inside the lift of a fixed end.

With this modification, and with Theorem 4.20 in place, the proof of Theorem 5.5 goes through for arbitrary finitely generated Kleinian groups provided a Minsky model obtains (for instance if M has no parabolics, cf. Appendix 7). However, owing to Item (d) above, the statement is a bit more involved.

Theorem 5.6. *Let G be a finitely generated Kleinian group. Let $M = \mathbf{H}^3/G$ and assume that each degenerate end of M admits a Minsky model (for instance if M has no parabolics). Let $i : \Gamma_G \rightarrow \mathbf{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbf{H}^3$. Then i extends continuously to a map $\hat{i} : \widehat{\Gamma}_G \rightarrow \mathbb{D}^3$, where $\widehat{\Gamma}_G$ denotes the (relative) hyperbolic compactification of Γ_G . Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial\Gamma_G$ of Γ_G .*

Let E be a degenerate end of $N^h = \mathbf{H}^3/G$ and $\tilde{E}$ a lift of E to $\tilde{N}^h$ and let M_{gf} be an augmented Scott core of N^h . Then the ending lamination $\mathcal{L}_E$ for the end E lifts to a lamination on $\tilde{M}_{gf} \cap \tilde{E}$. Each such lift $\mathcal{L}$ of the ending lamination of a degenerate end defines a relation $\mathcal{R}_\mathcal{L}$ on the (Gromov) hyperbolic boundary $\partial\tilde{M}_{gf}$ (equal to the relative hyperbolic boundary $\partial\Gamma_G$ of Γ_G), given by $a\mathcal{R}_\mathcal{L}b$ iff a, b are end-points of a leaf of $\mathcal{L}$. Let $\{\mathcal{R}_i\}_i$ be the entire collection of relations on $\partial\tilde{M}_{gf}$ obtained this way. Let $\mathcal{R}$ be the transitive closure of the union $\bigcup_i \mathcal{R}_i$. Then $\partial i(a) = \partial i(b)$ iff $a\mathcal{R}b$.

6. APPLICATIONS

In this section we shall first mention a couple of applications of the main Theorems of this paper. Finally we indicate an extension of the Sullivan-McMullen dictionary between complex dynamics and Kleinian groups.

6.1. Primitive Stable Representations. In [Min09] Minsky introduced and studied primitive stable representations, an open set of $PSl_2(\mathbb{C})$ characters of a non-abelian free group, on which the action of the outer automorphism group is properly discontinuous, and which is strictly larger than the set of discrete, faithful convex-cocompact (i.e. Schottky) characters.

In [Min09] Minsky also conjectured that
A discrete faithful representation of F is primitive-stable if and only if every component of the ending lamination is blocking.

Using the structure of the Cannon-Thurston map for handlebody groups, Jeon, Kim, Lecuire and Ohshika [JKLO14] have solved this conjecture. We sketch their proof for degenerate free groups without parabolics with associated representation denoted by ρ . They show that for the ending lamination Λ_E of a degenerate free group without parabolics, $Wh(\Lambda_E, \Delta)$ is connected and has no cutpoints. Then they argue by contradiction. If ρ is not primitive stable, then there exists a sequence of primitive cyclically reduced elements w_n such that $\rho(\cdots w_n w_n w_n \cdots)$ is not an n -quasi-geodesic. After passing to a subsequence, w_n and hence $\cdots w_n w_n w_n \cdots$ converges to a bi-infinite geodesic w_∞ in the Cayley graph with two distinct endpoints w_+, w_- in the Gromov boundary of F identified by the Cannon-Thurston map. It follows from a Lemma due to Whitehead that w_n cannot be primitive for large n , a contradiction.

6.2. Discreteness of Commensurators. In [LLR11] and [Mj11], the main results of this paper are used to prove that commensurators of finitely generated, infinite covolume, Zariski dense Kleinian groups are discrete. The proof proceeds by showing that commensurators preserve the structure of point pre-images of Cannon-Thurston maps.

6.3. Extending the Sullivan-McMullen Dictionary. A celebrated theorem of Yoccoz in Complex Dynamics (see Hubbard [Hub93], or Milnor [Mil00]) proves the local connectivity of certain Julia sets using a technique called ‘puzzle pieces’, which consists of a decomposition of a complex domain into pieces each of which under iteration by a quadratic map converges to a single point. The dynamical system can then be regarded as a semigroup $\mathbb{Z}_+$ of transformations acting on a complex domain.

Split components can be regarded as a 3-dimensional analogue of puzzle pieces. Let us try to justify this analogy. Suppose there is a group G acting cocompactly on $\mathbb{H}^3$. Let $H \subset G$ be a subgroup. Let G/H denote the coset space. Then what we would want as the right analogue is that if one takes a sequence of elements g_i going to infinity in the coset space, the iterates of the convex hull of the limit set of H converge to a point in the limit sphere. Thus going to infinity in the coset space G/H would be the right Kleinian groups analogue of going to infinity in the semigroup $\mathbb{Z}_+$ of transformations acting on a complex domain.

In the context of this paper, H would be the fundamental group of a split component. However, for us G is a surface Kleinian group and does not act cocompactly on $\mathbb{H}^3$. We think of the quotient space $\mathbb{H}^3/H$ as parametrizing the set of normal directions to the split component. The graph metric gives a combinatorial distance on $\mathbb{H}^3/H$ and we think of $(\mathbb{H}^3/H, d_G)$ as the analogue of the semigroup $\mathbb{Z}_+$. Thus, instead of going to infinity by iteration in the semigroup $\mathbb{Z}_+$, we go to infinity in the graph metric. Further, the analogue of the requirement that iterates go to infinity, is that the visual diameter goes to zero as we move to infinity in the graph metric. This is ensured by hyperbolic quasiconvexity, and also follows easily from **graph quasiconvexity**. Note that **graph quasiconvexity** is a statement that gives uniform shrinking of visual diameter to zero as one goes to infinity.

Thus we extend the Sullivan-McMullen dictionary (see [Sul85], [McM98]) between Kleinian groups and complex dynamics by suggesting the following analogy:

- (1) *Puzzle pieces* are analogous to *split components*
- (2) *Convergence to a point under iteration* is analogous to *graph quasiconvexity*

One issue that gets clarified by the above analogy is a point raised by McMullen in [McM01]. McMullen indicates that though the Julia set $J(P_\theta)$, where

$$P_\theta(z) = e^{2\pi i\theta}z + z^2$$

need not be locally connected in general by a result of Sullivan [Sul83], the limit sets of punctured torus groups are nevertheless locally connected. Local connectivity of Julia sets would therefore not be the right analogue of local connectivity of limit sets in this setup. Instead we look at the *techniques* for proving local connectivity of limit sets vis-a-vis the *techniques* for proving local connectivity of Julia sets. Thus, by proposing the analogy between puzzle pieces and split components as above, this issue is to an extent clarified. In short, the analogy is in the technique rather than in the result.

An analogue of the $\mathbb{Z}_+$ dynamical system may also be extracted from the split geometry model. Note that each block corresponds to a splitting of the surface group, and hence an action on a tree. As $i \rightarrow \infty$, the split blocks B_i^s and hence the induced splittings also go to infinity, converging to a **free action of the surface group on an $\mathbb{R}$ -tree dual to the ending lamination**. Thus iteration of the quadratic function corresponds to taking a sequence of splittings of the surface group converging to a (particular) action on an $\mathbb{R}$ -tree.

Problem: The building of the Minsky model and its bi-Lipschitz equivalence to a hyperbolic manifold [Min10] [BCM12] gives rise to a speculation that there should be a purely combinatorial way of doing much of the work. Bowditch's rendering [Bow11], [Bow05] of the Minsky, Brock-Canary-Minsky results is a step in this direction. This paper brings out the possibility that the whole thing should be doable purely in terms of actions on trees. Of course there is an action of the surface group on a tree dual to a pants decomposition. So we do have a starting point. However, one ought to be able to give a purely combinatorial description, *ab initio*, in terms of a sequence of actions of surface groups on trees converging to an action on an $\mathbb{R}$ -tree. This would open up the possibility of extending these results (including those of this paper) to other hyperbolic groups with infinite automorphism groups, notably free groups.

7. APPENDIX: MODEL MANIFOLD FOR COMPRESSIBLE BOUNDARY

In this Appendix, for the sake of completeness, we provide a sketch of a bi-Lipschitz Minsky model for general M without parabolics, following work and ideas of several authors, notably Brock, Bromberg and Souto. No claim to originality is made here. Our aim is modest: to reduce the problem of existence of such a model to the case with incompressible boundary discussed in [BCM12] and hence conclude the existence of a bi-Lipschitz model geometry in general. We shall focus on a degenerate end E such that $\partial E = S$ is compressible. We would like to prove that E has a bi-Lipschitz model as in [BCM12]. By passing to the cover of M corresponding to the end E (i.e. the cover of M corresponding to the image of

$\pi_1(E)$ in $\pi_1(M)$), we might as well **assume that M is a compression body**. We sketch the proof below in the case when $S(= \partial E)$ is closed. The case when S has parabolics, and hence M has rank one cusps, is technically quite a bit more involved and we refer the reader to [BBES03, p. 148] where the presence of accidental parabolics is addressed.

The model geometry for the case where E has bounded geometry has been treated in detail in [Mos03, Bow02, Ohs98] (see also [Mit98]) and so we assume that E has unbounded geometry. The main idea is to isolate E and reduce to the incompressible case. The point is to show that the asymptotic geometry of E is bi-Lipschitz to an incompressible simply degenerate end.

Step 1: Existence of Disk-busting curves: We consider a sequence of simple closed curves c_i on S whose geodesic realizations exit E and have length going to zero. We note first that for large enough i , these curves nontrivially intersect all compressing disks with boundary in S . If not, there exists compressing disk D_i with $\partial D_i \subset S$ such that $\partial D_i \cap c_i = \emptyset$. Normalizing c_i by its length and taking a limit in $\mathcal{PML}(S)$, it would follow that the (limiting) ending lamination does not lie in the Masur Domain, a contradiction (cf. the definition of the Masur domain after Theorem 4.2). Hence, for all large enough i , the curves c_i are disk-busting.

Hence if we drill out the geodesic realizations of c_i from E , and equip the resulting manifold with a complete hyperbolic structure with a rank 2 cusp where c_i has been drilled (while fixing the end-invariants), the new end E_i is incompressible. Then, by [BCM12], E_i has a model geometry. We would be done if we could establish that E and E_i are bi-Lipschitz homeomorphic, as this would transfer the bi-Lipschitz model of E_i to E . Instead of drilling, we shall perform the closely related construction of grafting below.

Step 2: Grafting Constructions: In [Bro07, BB04, BBES03, BB11], two kinds of grafting constructions are used. We call these up-grafting and down-grafting below.

The standard construction [Bro07] (Section 4, where grafting an infinite annulus is described) involves cutting the manifold open along a semi-infinite cylinder and gluing in a wedge. Thus one *up-grafts* an annulus along the geodesic realization of c_i in M as in [Bro07, BB04] to get a cone manifold N_i with cone angle 4π along the geodesic realizations of c_i .

The other kind of grafting (which we call down-grafting) is described in detail in [Bro07] (see Section 4 of the paper, especially Theorem 4.2)³. We give a quick summary. Take an annulus going out the end based at the curve c_0 . We pass to the cover of E associated to the subsurface obtained by taking the complement of the curve. For ease of exposition, assume that c_0 is non-separating. In the cover the annulus will have two isometric lifts. Cut the cover along both of them and then glue them together. The result is a cone-manifold M_{c_0} that is homeomorphic to $S \times \mathbb{R}$.

To ensure that (up or down) grafting is possible, one needs two conditions:

- a) that the geodesic is unknotted,
- b) that the geodesic is sufficiently small. This allows the relevant technical tools of [Bro07], [BB04] (Section 5, especially Theorem 5.1 which proves the existence of

³I am grateful to Ken Bromberg for explaining this construction to me.

the geometrically finite cone manifold), to go through.

The first condition follows from the second thanks to work of Otal [Ota95, Ota03] (See also [Sou08]). Shortness of curves is guaranteed by unbounded geometry of E .

Step 3: Double Grafting and Convergence: Choose sufficiently short (short enough to allow grafting in Step 2 to go through) disk-busting curves c_0 and c_i ($i \in \mathbb{N}$). Along c_0 perform down-grafting to obtain a cone-manifold M_{c_0} that is homeomorphic to $S \times \mathbb{R}$. As a result, one end (the up end) will be the original unbounded geometry end. The down end will be geometrically finite. This process effectively isolates the "up" end (isometric to E) with the other end of M_{c_0} being geometrically finite. We need to show that E is bi-Lipschitz to the end of a simply degenerate manifold.

Along the c_i curves now perform the standard (up) grafting construction on M_{c_0} described in Step 2 above to obtain quasi-Fuchsian cone manifolds N_i . Smooth quasi-Fuchsian manifolds M_i may be obtained from N_i by decreasing the cone angle to 2π [Bro07].

The main technical Theorems of Brock and Bromberg on inflexibility [BB11, Theorem 3.6], [BB14, Theorems 1.1, 1.2] now guarantee that the cone deformations do not change the hyperbolic structure much away from the boundary. More precisely, fix an arbitrary compact core $\mathcal{M}$ of M_{c_0} and assume that all the curves c_i have geodesic realizations outside $\mathcal{M}$. Then the inflexibility Theorem [BB11, Theorem 3.6] shows that the cone deformation from N_i to M_i has small effect in the thick part of $\mathcal{M}$ (more precisely, the harmonic strain field decays exponentially with distance from the boundary). It follows that the hyperbolic structures on N_i and M_i when restricted to $\mathcal{M}$ are $(1 + \epsilon_i)$ -biLipschitz to each other, where $\epsilon_i \rightarrow 0$ as $i \rightarrow \infty$.

Now, in the cone-manifolds N_i , between the two curves c_0 and c_i , there is a larger and larger product region that is isometric to a product region in the original end E . Therefore (from the inflexibility theorem above) the sequence of smooth quasifuchsian manifolds M_i converges to a simply degenerate manifold M_∞ whose degenerate end E_∞ is bilipschitz to the original end E . Since E_∞ has a model geometry by [Min10, BCM12], so does E .

REFERENCES

- [Ago04] I. Agol. Tameness of hyperbolic 3-manifolds. *preprint*, *arXiv:math.GT/0405568*, 2004.
- [BB04] J. Brock and K. Bromberg. Density of Geometrically Finite Kleinian Groups. *Acta Mathematica* 192, pages 33–93, 2004.
- [BB11] J. Brock and K. Bromberg. Geometric inflexibility and 3-manifolds that fiber over the circle. *J. Topol.* 4, no. 1, pages 1–38, 2011.
- [BB14] J. Brock and K. Bromberg. Geometric inflexibility of hyperbolic cone-manifolds. *arXiv:1412.4635*, to appear in the *Proceedings of the 2014 MSJ-SI*, 2014.
- [BBES03] J. Brock, K. Bromberg, R. Evans, and J. Souto. Tameness on the boundary and Ahlfors' measure conjecture. *Publ. Math. I.H.E.S.* 98, pages 145–166, 2003.
- [BCM12] J. F. Brock, R. D. Canary, and Y. N. Minsky. The Classification of Kleinian surface groups II: The Ending Lamination Conjecture. *Ann. of Math.* 176 (1), pages 1–149, 2012.
- [BCM14] J. F. Brock, R. D. Canary, and Y. N. Minsky. The Classification of finitely generated Kleinian groups. *in preparation*, 2014.
- [Bon86] F. Bonahon. Bouts de varietes hyperboliques de dimension 3. *Ann. Math. vol. 124*, pages 71–158, 1986.

- [Bow02] B. H. Bowditch. Stacks of hyperbolic spaces and ends of 3 manifolds. *preprint, Southampton*, 2002.
- [Bow05] B. H. Bowditch. Model geometries for hyperbolic manifolds. *preprint, Southampton*, 2005.
- [Bow11] B. H. Bowditch. The ending lamination theorem. *preprint, Warwick*, 2011.
- [Bow12] B. H. Bowditch. Relatively hyperbolic groups. *Internat. J. Algebra Comput.* 22, no. 3, 2012.
- [Bro07] K. Bromberg. Projective structures with degenerate holonomy and the Bers density conjecture. *Ann. of Math. (2)* 166, no. 1, pages 77–93, 2007.
- [Can93] R. D. Canary. Ends of hyperbolic 3 manifolds. *J. Amer. Math. Soc.*, pages 1–35, 1993.
- [CT85] J. Cannon and W. P. Thurston. Group Invariant Peano Curves. *preprint, Princeton*, 1985.
- [CT07] J. Cannon and W. P. Thurston. Group Invariant Peano Curves. *Geometry and Topology vol 11*, pages 1315–1356, 2007.
- [DM16] S. Das and M. Mj. Semiconjugacies between relatively hyperbolic boundaries. *Groups Geom. Dyn.* 10, no. 2, pages 733–752, 2016.
- [Far98] B. Farb. Relatively hyperbolic groups. *Geom. Funct. Anal.* 8, pages 810–840, 1998.
- [Flo80] W. J. Floyd. Group Completions and Limit Sets of Kleinian Groups. *Invent. Math. vol.57*, pages 205–218, 1980.
- [GC06] D. Gabai and D. Calegari. Shrink-wrapping and the Taming of Hyperbolic 3-manifolds. *Jour. Amer. Math. Soc.* 19, no. 2, pages 385–446, 2006.
- [Hem76] J. Hempel. 3 manifolds. *Princeton University Press*, 1976.
- [Hub93] J. H. Hubbard. Local Connectivity of Julia sets and Bifurcation Loci: three Theorems of J.-C. Yoccoz. In L. R. Goldberg and A. G. Philips, Editors, *Topological Methods in Modern Mathematics, Publish or Perish Inc.*, pages 467–511, 1993.
- [JKLO14] W. Jeon, I. Kim, C. Lecuire, and K. Ohshika. Primitive stable representations of free Kleinian groups. *Israel J. Math.* 199, no. 2, pages 841–866, 2014.
- [Kla99] E. Klarreich. Semiconjugacies between Kleinian group actions on the Riemann sphere. *Amer. J. Math* 121, pages 1031–1078, 1999.
- [KS03] G. Kleineidam and J. Souto. Ending laminations in the masur domain. in *Kleinian groups and hyperbolic 3-manifolds (Warwick, 2001)*, ed. by Y. Komori, V. Markovic and C. Series, *London Math. Soc. Lecture Notes* 299, pages 105–129, 2003.
- [LLR11] C. J. Leininger, D. D. Long, and A. W. Reid. Commensurators of non-free finitely generated Kleinian groups. *Algebraic and Geometric Topology* 11, pages 605–624, 2011.
- [Luf78] E. Luft. Actions of the homeotopy group of an orientable 3-dimensional handlebody. *Math. Ann.* 234, no. 3, pages 279–292, 1978.
- [McM98] C. T. McMullen. Renormalization and 3-Manifolds which Fiber over the Circle. *Princeton University Press*, 1998.
- [McM01] C. T. McMullen. Local connectivity, Kleinian groups and geodesics on the blow-up of the torus. *Invent. math.*, 97:95–127, 2001.
- [Mil00] J. Milnor. Local Connectivity of Julia sets: Expository Lectures. In Tan Lei, Editor, *The Mandelbrot Set, Theme and Variations*, CUP, pages 67–116, 2000.
- [Min09] Y. N. Minsky. On dynamics of $Out(F_n)$ on $PSL_2(C)$ characters. *preprint, arxiv:0906.3491*, 2009.
- [Min10] Y. N. Minsky. The Classification of Kleinian surface groups I: Models and bounds. *Ann. of Math.* 171, pages 1–107, 2010.
- [Mit98] M. Mitra. Cannon-Thurston Maps for Trees of Hyperbolic Metric Spaces. *Jour. Diff. Geom.* 48, pages 135–164, 1998.
- [Miy02] H. Miyachi. Semiconjugacies between actions of topologically tame Kleinian groups. *preprint*, 2002.
- [Mj11] M. Mj. On Discreteness of Commensurators. *Geometry and Topology* 15, pages 331–350, 2011.
- [Mj14a] M. Mj. Cannon-Thurston Maps for Surface Groups. *Ann. of Math.*, 179(1), pages 1–80, 2014.
- [Mj14b] M. Mj. Ending Laminations and Cannon-Thurston Maps, with an appendix by S. Das and M. Mj. *Geom. Funct. Anal.* 24, pages 297–321, 2014.
- [MM86] D. McCullough and A. Miller. Homeomorphisms of 3-manifolds with compressible boundary. *Mem. Amer. Math. Soc.* 61, no. 344, pages xii+100 pp., 1986.

- [Mos03] L. Mosher. Stable Teichmüller quasigeodesics and ending laminations. *Geom. Topol.* 7, pages 33–90, 2003.
- [MP89] J. D. McCarthy and A. Papadopoulos. Dynamics on Thurston’s Sphere of Projective Measured Foliations. *Comment. Math. Helv.* 64, pages 133–166, 1989.
- [MP11] M. Mj and A. Pal. Relative Hyperbolicity, Trees of Spaces and Cannon-Thurston Maps. *Geom. Dedicata* 151, pages 59–78, 2011.
- [Ohs98] K. Ohshika. Rigidity and topological conjugates of topologically tame kleinian groups. *Trans. Amer. Math. Soc.* 350, no. 10, pages 3989–4022, 1998.
- [Ota88] J. P. Otal. Courants géodésiques et produit libres. *Thèse d’Etat, Université Paris-Sud, Orsay*, 1988.
- [Ota95] J. P. Otal. Sur le nouage des géodésiques dans les variétés hyperboliques. *C. R. Acad. Sci. Paris Ser. I Math.* 320, no. 7, pages 847–852, 1995.
- [Ota03] J. P. Otal. Les géodésiques fermées d’une variété hyperbolique en tant que noeuds. In *Kleinian Groups and Hyperbolic 3-Manifolds. London Math. Soc.*, 2003.
- [Sou06] J. Souto. Cannon-Thurston Maps for Thick Free Groups. *preprint*, 2006.
- [Sou08] J. Souto. Short geodesics in hyperbolic compression bodies are not knotted. *preprint*, 2008.
- [Sul83] D. Sullivan. Conformal dynamical systems. *Geometric Dynamics, LNM 1007, Springer-Verlag*, pages 725–752, 1983.
- [Sul85] D. Sullivan. Quasiconformal Homeomorphisms and Dynamics I: Solution of the Fatou-Julia problem on Wandering Domains. *Annals of Math.* 122, pages 401–418, 1985.
- [Suz77] S. Suzuki. On homeomorphisms of a 3-dimensional handlebody. *Canad. J. Math.* 29, no. 1, pages 111–124, 1977.
- [Thu80] W. P. Thurston. The Geometry and Topology of 3-Manifolds. *Princeton University Notes*, 1980.
- [Thu82] W. P. Thurston. Three dimensional manifolds, Kleinian groups and hyperbolic geometry. *Bull. Amer. Math. Soc.*, pages 357–382, 1982.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI-400005, INDIA

E-mail address: mahan.mj@gmail.com; mahan@math.tifr.res.in