# Symmetries for the Ablowitz-Ladik hierarchy: I. Four-potential case

Da-jun Zhang<sup>\*</sup>, Shou-ting Chen

Department of Mathematics, Shanghai University, Shanghai 200444, P.R. China

#### Abstract

In the paper we first investigate symmetries of isospectral and non-isospectral fourpotential Ablowitz-Ladik hierarchies. We express these hierarchies in the form of  $u_{n,t} = L^m H^{(0)}$ , where *m* is an arbitrary integer (instead of a nature number) and *L* is the recursion operator. Then by means of the zero-curvature representations of the isospectral and non-isospectral flows, we construct symmetries for the isospectral equation hierarchy as well as non-isospectral equation hierarchy, respectively. The symmetries, respectively, form two centerless Kac-Moody-Virasoro algebras. The recursion operator *L* is proved to be hereditary and a strong symmetry for this isospectral equation hierarchy. Besides, we make clear for the relation between four-potential and two-potential Ablowitz-Ladik hierarchies. The even order members in the four-potential Ablowitz-Ladik hierarchies together with their symmetries and algebraic structures can be reduced to two-potential case. The reduction keeps invariant for the algebraic structures and the recursion operator for two potential case becomes  $L^2$ .

**Key words:** Ablowitz-Ladik hierarchies; isospectral and non-isospectral flows; symmetries; Lie algebra.

**PACS:** 02.30.Ik, 05.45.Yv

# 1 Introduction

As is well known, many physically interesting problems are modeled by nonlinear differentialdifference equations, such as the Toda lattice, Volterra lattice and discrete nonlinear Schrödinger equation. Since 1970s discrete systems have received considerable attention from variety of aspects (e.g., [1–4]), such as Inverse Scattering Transform, bilinear method, Sato's approach, symmetry analysis and so on. One of famous discrete spectral problems is given by Ablowitz and Ladik [1,5] which is now referred to as the Ablowitz-Ladik (AL) spectral problem. This spectral problem, coupled with different time evolution parts, has provided Lax integrabilities for many discrete soliton systems, such as integrable discrete nonlinear Schrödinger equation [6], discrete mKdV equation and so forth.

There are two types of the AL spectral problems, which contains two potentials  $\{Q_n, R_n\}$ and four potentials  $\{Q_n, R_n, S_n, T_n\}$ , respectively. The two-potential one is the direct discretization (cf. [6]) of the famous continuous AKNS-ZS spectral problem [7], and besides solutions,

<sup>\*</sup>Corresponding author. E-mail: djzhang@staff.shu.edu.cn

the related Hamiltonian structures, constraint flows, nonlinearization, Darboux transformation, conservation laws, symmetries and Lie algebra structures have been studied (cf. [8–17]). The four-potential AL spectral problem is more complicated than the two-potential case because of containing two more potentials and its unsymmetrical matrix form. Cheng [18] transformed the four-potential AL spectral problem to a bit simple form which could be further related to the two-potential case, but the relation (connecting two and four potentials) given in [18] is nothing helpful for discussions of Hamiltonian structures and symmetries. Recently, based on Cheng's transformation, Geng and Dai [19] separated the four-potential AL spectral matrix into two symmetrical two-potential AL spectral matrices. This makes it possible to construct a four-potential hierarchy (e.g. [19]) and its recursion operator and then investigate more characteristics of integrability.

Infinite symmetries act as an important characteristic for integrable systems [20]. Symmetryanalysis is also a powerful approach to finding exact solutions for nonlinear systems [21, 22]. In this paper we focus on symmetries and their Lie algebras for the four-potential isospectral and non-isospectral AL hierarchies. We will first derive positive and negative order isospectral and non-isospectral flows. These flows share a same recursion operator L and can be respectively uniformed as  $K^{(m)} = L^m K^{(0)}$  and  $\sigma^{(m)} = L^m \sigma^{(0)}$ , where m is an arbitrary integer (instead of a nature number in most of cases). Then we can imbed these flows into their zero-curvature equations by means of functional derivatives. The resulting expressions, which we refer to as zero-curvature representations, have been shown to be powerful in constructing symmetries for Lax integrable systems (cf. [10,15,23–27]). We will derive algebraic relations for isospectral and non-isospectral flows and then derive symmetries and their Lie algebras for not only isospectral AL hierarchy but also non-isospectral hierarchy. Both algebras are the type of centerless Kac-Moody-Virasoro algebra. The recursion operator L is hereditary and a strong symmetry for the isospectral hierarchy.

A natural question is whether all these structures w.r.t. four potentials  $(Q_n, R_n, S_n, T_n)$ , including hierarchies, recursion operator L, symmetries and algebras admit a closed reduction w.r.t. two potentials  $(Q_n, R_n)$  by directly taking  $(S_n, T_n) = (0, 0)$ . This is true for those even order members in the four-potential isospectral and non-isospectral hierarchies, and the new recursion operator becomes  $L^2$ . We will discuss the reduction in the paper.

This paper is the first part of our series investigations which consist of two parts. In Part II we will focus on symmetries of the integrable discrete nonlinear Schrödinger equation and discrete AKNS hierarchy. The integrable discrete nonlinear Schrödinger equation consists of positive and negative order flows which correspond to a central-difference discretization for a continuous second order derivative. We will also give a recursion operator which generates discrete AKNS hierarchies. The obtained symmetry algebras are not centerless Kac-Moody-Virasoro type. The structure changes will also be explained in Part II.

The present paper is organized as follows. Sec.2 contains some basic notations and backgrounds on the AL spectral problem. Sec.3 derives four-potential isospectral and non-isospectral flows and their zero-curvature representations. In Sec.4 we derive symmetries and their algebraic structures for both four-potential isospectral and non-isospectral hierarchies. In Sec.5 we discuss reduction relation between four-potential case and two-potential case. There are also two Appendix sections. Sec.A lists out first few equations in four-potential AL hierarchies and their Lax pairs, and Sec.B is a theorem obtained in Ref. [11] which we give here for self-containedness.

### 2 Basic notations and backgrounds

Let us first introduce some basic notations and notions which have been used for discussing symmetries of discrete systems (cf. [11, 15, 27]).

Assume that  $u_n \doteq u(t,n) = (u^{(1)}, u^{(2)}, u^{(3)}, u^{(4)})^T$  is a four-dimensional vector field, where  $u^{(i)} = u^{(i)}(t,n), 1 \le i \le 4$ , are all functions defined over  $\mathbb{R} \times \mathbb{Z}$  and vanish rapidly as  $|n| \to \infty$ . By  $\mathcal{V}_4$  we denote a linear space consisting of all vector fields  $f = (f^{(1)}, f^{(2)}, f^{(3)}, f^{(4)})^T$ , where each  $f^{(i)}$  is a function of u(t,n) and its shifts  $u(t,n+j), j \in \mathbb{Z}$ , satisfying  $f^{(i)}(u(t,n))|_{u_n=0} = 0$ , and each  $f^{(i)}$  is  $C^{\infty}$  differentiable w.r.t. t and n, and  $C^{\infty}$ -Gauteaux differentiable w.r.t.  $u_n$ . Here the Gateaux (or Fréchet) derivative of  $f \in \mathcal{V}_4$  (or f an operator on  $\mathcal{V}_4$ ) in the direction  $g \in \mathcal{V}_4$  is defined as

$$f'[g] = \frac{d}{d\epsilon} \Big|_{\epsilon=0} f(u+\epsilon g).$$
(2.1)

By means of the Gateaux derivative one can define a Lie product for any  $f, g \in \mathcal{V}_4$  as

$$[\![f,g]\!] = f'[g] - g'[f].$$
(2.2)

We also define a Laurent matrix polynomials space  $Q_2(z)$  composed by all  $2 \times 2$  matrices  $Q = Q(z, u(t, n)) = (q_{ij}(z, u(t, n)))_{2 \times 2}$ , where all the  $\{q_{ij}\}$  are Laurent polynomials of z. Two subspaces of  $Q_2(z)$  we will need are

$$\mathcal{Q}_2^+(z) = \{ Q \in \mathcal{Q}_2(z) | \text{ the lowest degree of } z \ge 0 \}, \mathcal{Q}_2^-(z) = \{ Q \in \mathcal{Q}_2(z) | \text{ the highest degree of } z \le 0 \}.$$

We note that in a similar way we can define spaces  $\mathcal{V}_s$ ,  $\mathcal{Q}_m(z)$  and  $\mathcal{Q}_m^{\pm}(z)$  (cf. [11]).

In general a discrete evolution equation arises from the compatibility of a pair discrete linear problems  $^{\rm 1}$ 

$$\Phi_{n+1} = U_n(z, u(t, n))\Phi_n, \quad \Phi_{n,t} = V_n(z, u(t, n))\Phi_n,$$
(2.3)

where  $\Phi_n$  is a wave function,  $U_n$  is a spectral matrix with spectral parameter z and potential vector u(t,n) while  $V_n$  is a matrix governing time evolution. The compatibility condition, also called discrete zero-curvature equation, reads

$$U_{n,t} = (EV_n)U_n - U_n V_n. (2.4)$$

Here and in the following E is a shift operator defined as  $E^j f(n) = f(n+j)$  for  $j \in \mathbb{Z}$ . Suppose that the corresponding nonlinear evolution equation is

$$u_{n,t} = K(u_n). \tag{2.5}$$

Then by means of the Gateaux derivative the flow  $K(u_n)$  can be embedded into the zerocurvature equation (2.4) as the following,

$$U'_{n}[K(u_{n})] = (EV_{n})U_{n} - U_{n}V_{n} - U_{n,z}z_{t},$$
(2.6)

which is usually called the zero-curvature representation of the flow  $K(u_n)$ .

<sup>&</sup>lt;sup>1</sup> Actually, they are semi-discrete.

For the nonlinear evolution equation (2.5),  $\sigma(u_n) \in \mathcal{V}_4$  is its symmetry if  $\sigma_t = K'[\sigma]$ , i.e.,

$$\frac{\tilde{\partial}\sigma}{\tilde{\partial}t} = \llbracket K, \sigma \rrbracket, \tag{2.7}$$

where by  $\frac{\partial \sigma}{\partial t}$  we specially denote the derivative of  $\sigma$  w.r.t. t explicitly included in  $\sigma$ , (for example,  $\frac{\partial \sigma}{\partial t} = u_n$  if  $\sigma = tu_n + u_{n+1}$ ).

Next, let us recall some backgrounds on four-potential AL hierarchy. The four-potential AL spectral problem reads [1,5]

$$\psi_{1,n+1} = \lambda \psi_{1,n} + Q_n \psi_{2,n} + S_n \psi_{2,n+1}, \psi_{2,n+1} = \lambda^{-1} \psi_{2,n} + R_n \psi_{1,n} + T_n \psi_{1,n+1},$$
(2.8)

where  $\lambda$  is a spectral parameter and  $Q_n, R_n, S_n, T_n$  are four potential functions of n and t. When  $S_n = T_n = 0$  (2.8) reduces to the two-potential AL spectral problem, i.e.,

$$\psi_{1,n+1} = \lambda \psi_{1,n} + Q_n \psi_{2,n}, \psi_{2,n+1} = \lambda^{-1} \psi_{2,n} + R_n \psi_{1,n},$$
(2.9)

which is a discrete version of the AKNS-ZS spectral problem (cf. [6]).

An alternative (matrix) form of (2.8) is [1]

$$\Psi_{n+1} = \frac{1}{\Lambda_n} \begin{pmatrix} z^2 + S_n R_n & Q_n + z^{-2} S_n \\ z^2 T_n + R_n & z^{-2} + T_n Q_n \end{pmatrix} \Psi_n, \quad \Psi_n = \begin{pmatrix} \psi_{1,n} \\ \psi_{2,n} \end{pmatrix},$$
(2.10)

where  $\Lambda_n = 1 - S_n T_n$  and we have substituted  $z^2$  for  $\lambda$ . This form can be gauge-transformed to [18]

$$\Phi_{n+1} = U_n \Phi_n, \quad U_n = \begin{pmatrix} z^2 + S_n R_n & zQ_n + z^{-1}S_n \\ zT_n + z^{-1}R_n & z^{-2} + T_nQ_n \end{pmatrix}, \quad \Phi_n = \begin{pmatrix} \phi_{1,n} \\ \phi_{2,n} \end{pmatrix},$$
(2.11)

where  $\Phi_n$  and  $\Psi_n$  are related through

$$\Phi_n = \rho(z)\Psi_n \prod_{i=n}^{+\infty} \Lambda_i^{-1}, \quad \rho(z) = \begin{pmatrix} z^{\frac{1}{2}} & 0\\ 0 & z^{-\frac{1}{2}} \end{pmatrix}.$$
 (2.12)

Here on  $U_n$  we impose a condition

$$Q_n R_n + S_n T_n \neq 0 \tag{2.13}$$

so that  $U'_n$  is an injective homomorphism.

Suppose that the time evolution of  $\Phi_n$  is

$$\Phi_{n,t} = V_n \Phi_n, \qquad V_n = \begin{pmatrix} A_n & B_n \\ C_n & D_n \end{pmatrix}.$$
(2.14)

Then the compatibility condition with (2.11) yields

$$U_{n,t} = (EV_n)U_n - U_n V_n. (2.15)$$

Usually the above discrete zero-curvature equation contributes a discrete nonlinear evolution equation hierarchy with four potentials and their recursion operator, but this is not as easy as in two-potential case (related to (2.9), cf. [9–11, 15]). However, the spectral matrix  $U_n$  can be separated into [19]

$$U_n = U_n^{(2)} U_n^{(1)}, \quad U_n^{(1)} = \begin{pmatrix} z & Q_n \\ R_n & z^{-1} \end{pmatrix}, \quad U_n^{(2)} = \begin{pmatrix} z & S_n \\ T_n & z^{-1} \end{pmatrix}.$$
 (2.16)

Then (2.15) holds if [19]

$$U_{n,t}^{(1)} = \widehat{V}_n U_n^{(1)} - U_n^{(1)} V_n, \qquad U_{n,t}^{(2)} = (EV_n) U_n^{(2)} - U_n^{(2)} \widehat{V}_n, \qquad (2.17)$$

where  $\widehat{V}_n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}$ . In fact,  $U_{n,t} - (EV_n)U_n + U_nV_n = (U_{n,t}^{(2)} - (EV_n)U_n^{(2)} + U_n^{(2)}\widehat{V}_n)U_n^{(1)} + U_n^{(2)}(U_{n,t}^{(1)} - \widehat{V}_nU_n^{(1)} + U_n^{(1)}V_n).$ 

Thus, one can consider the two auxiliary systems given in (2.17), where each of  $U_n^{(j)}$  contains two potentials. Recently, starting from (2.17) Geng and Dai [19] derived a four-potential AL hierarchy and their recursion relation and considered Hamiltonian structures and nonlinearization of Lax pair.

Noting that in the AL spectral problem (2.8) the spectral parameter  $\lambda$  appears symmetrically w.r.t. positive and negative powers, it is then understood that the recursion operator and its inverse can be derived symmetrically. So are the negative order and positive order hierarchies. In [15], starting from (2.9) we have expressed isospectral and non-isospectral two-potential AL hierarchies in the form of  $u_{n,t} = L^m H^{(0)}$ , where m is an arbitrary integer (instead of a nature number) and L is the recursion operator. This is also true for four-potential case. In the next section we will derive four-potential AL hierarchies from the two auxiliary systems given in (2.17).

### 3 AL hierarchies and zero-curvature representations

Now we derive isospectral and non-isospectral AL hierarchies and their recursion operator. The procedure is quite like the one given in [15]. In addition, we will express the obtained isospectral and non-isospectral flows in terms of zero-curvature equation, by means of which we will prove that the recursion operator is hereditary and a strong symmetry of the isospectral hierarchy.

### 3.1 Isospectral hierarchy

The explicit form of the auxiliary systems (2.17) is

$$z^{-1}z_t = a_n - A_n + R_n b_n z^{-1} - C_n Q_n z^{-1}, aga{3.1a}$$

$$Q_{n,t} = b_n z^{-1} - B_n z + Q_n (a_n - D_n), \qquad (3.1b)$$

$$R_{n,t} = c_n z - C_n z^{-1} + R_n (d_n - A_n), \qquad (3.1c)$$

$$z(z^{-1})_t = d_n - D_n + Q_n c_n z - R_n B_n z;$$
(3.1d)

$$z^{-1}z_t = A_{n+1} - a_n + T_n B_{n+1} z^{-1} - c_n S_n z^{-1},$$
(3.1e)

$$S_{n,t} = B_{n+1}z^{-1} - b_n z + S_n(A_{n+1} - d_n),$$
(3.1f)

$$T_{n,t} = C_{n+1}z - c_n z^{-1} + T_n(D_{n+1} - a_n),$$
(3.1g)

$$z(z^{-1})_t = D_{n+1} - d_n + S_n C_{n+1} z - T_n b_n z.$$
(3.1h)

From (3.1a), (3.1d), (3.1e) and (3.1h) one can get

$$A_n = (E-1)^{-1} (S_n c_n z^{-1} + Q_n C_n z^{-1} - T_n B_{n+1} z^{-1} - R_n b_n z^{-1}) + 2n z^{-1} z_t + A_0, \qquad (3.2a)$$
$$a_n = (E-1)^{-1} (S_n c_n z^{-1} + Q_n C_n z^{-1} - T_n B_{n+1} z^{-1} - R_n b_n z^{-1})$$

$$+Q_n C_n z^{-1} - R_n b_n z^{-1} + (2n+1)z^{-1} z_t + A_0,$$
(3.2b)

$$D_n = (E-1)^{-1} (R_n B_n z + T_n b_n z - S_n C_{n+1} z - Q_n c_n z) + 2nz(z^{-1})_t + D_0,$$

$$d_n = (E-1)^{-1} (R_n B_n z + T_n b_n z - S_n C_{n+1} z - Q_n c_n z) + R_n B_n z - Q_n c_n z$$
(3.2c)

$$d_n = (E-1)^{-1} (R_n B_n z + T_n b_n z - S_n C_{n+1} z - Q_n c_n z) + R_n B_n z - Q_n c_n z + (2n+1) z (z^{-1})_t + D_0.$$
(3.2d)

Here  $A_0 = A_n|_{u_n=0} - 2nz^{-1}z_t$  and  $D_0 = D_n|_{u_n=0} - 2nz(z^{-1})_t$  where  $u_n = (Q_n, R_n, S_n, T_n)^T$ . Thus (3.1) simplifies to

$$u_{n,t} = (zL_1 + z^{-1}L_2) \begin{pmatrix} -B_n \\ C_n \\ -b_n \\ c_n \end{pmatrix} + (A_0 - D_0) \begin{pmatrix} Q_n \\ -R_n \\ S_n \\ -T_n \end{pmatrix} + z^{-1}z_t \begin{pmatrix} (4n+1)Q_n \\ -(4n+1)R_n \\ (4n+3)S_n \\ -(4n+3)T_n \end{pmatrix}, \quad (3.3)$$

where

$$L_{1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -R_{n}^{2} & 0 & 0 & \gamma_{n}^{2} \\ S_{n}R_{n} & 0 & 1 & S_{n}Q_{n} \\ -T_{n}R_{n} & \pi_{n}^{2}E & -T_{n}^{2} & -T_{n}Q_{n} \end{pmatrix} + \begin{pmatrix} Q_{n} \\ -R_{n} \\ S_{n} \\ -T_{n} \end{pmatrix} (E-1)^{-1}(R_{n}, S_{n}E, T_{n}, Q_{n}), \quad (3.4a)$$

$$\begin{pmatrix} 0 & Q_{n}^{2} & -\gamma_{n}^{2} & 0 \\ 0 & Q_{n}^{2} & -\gamma_{n}^{2} & 0 \end{pmatrix} \begin{pmatrix} Q_{n} \\ -T_{n} \end{pmatrix}$$

$$L_{2} = \begin{pmatrix} 0 & -1 & 0 & 0\\ -\pi_{n}^{2}E & S_{n}Q_{n} & R_{n}S_{n} & S_{n}^{2}\\ 0 & -T_{n}Q_{n} & -T_{n}R_{n} & -1 \end{pmatrix} + \begin{pmatrix} -R_{n}\\ S_{n}\\ -T_{n} \end{pmatrix} (E-1)^{-1}(T_{n}E,Q_{n},R_{n},S_{n}), \quad (3.4b)$$

in which  $\gamma_n = \sqrt{1 - Q_n R_n}$ ,  $\pi_n = \sqrt{1 - S_n T_n}$ . One can verify that the inverse operators of  $L_1$  and  $L_2$  are

$$L_{1}^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ \frac{T_{n-1}R_{n-1}E^{-1}}{\gamma_{n-1}^{2}} & \frac{T_{n-1}Q_{n-1}E^{-1}}{\gamma_{n-1}^{2}} & \frac{T_{n-1}^{2}E^{-1}}{\pi_{n-1}^{2}} & \frac{E^{-1}}{\pi_{n-1}^{2}} \\ -\frac{S_{n}R_{n}}{\gamma_{n}^{2}} & -\frac{S_{n}Q_{n}}{\gamma_{n}^{2}} & 1 & 0 \\ \frac{R_{n}^{2}}{\gamma_{n}^{2}} & \frac{1}{\gamma_{n}^{2}} & 0 & 0 \end{pmatrix} \\ - \begin{pmatrix} Q_{n} \\ -T_{n-1}E^{-1} \\ S_{n} \\ -R_{n} \end{pmatrix} (E-1)^{-1} \begin{pmatrix} R_{n} \\ \gamma_{n}^{2} \\ \gamma_{n}^{2}$$

$$L_{2}^{-1} = \begin{pmatrix} -\frac{S_{n-1}R_{n-1}E^{-1}}{\gamma_{n-1}^{2}} & -\frac{S_{n-1}Q_{n-1}E^{-1}}{\gamma_{n-1}^{2}} & -\frac{E^{-1}}{\pi_{n-1}^{2}} & -\frac{S_{n-1}^{2}E^{-1}}{\pi_{n-1}^{2}} \\ 0 & -1 & 0 & 0 \\ -\frac{1}{\gamma_{n}^{2}} & -\frac{Q_{n}^{2}}{\gamma_{n}^{2}} & 0 & 0 \\ \frac{T_{n}R_{n}}{\gamma_{n}^{2}} & \frac{T_{n}Q_{n}}{\gamma_{n}^{2}} & 0 & -1 \end{pmatrix} \\ - \begin{pmatrix} S_{n-1}E^{-1} \\ -R_{n} \\ Q_{n} \\ -T_{n} \end{pmatrix} (E-1)^{-1} \begin{pmatrix} R_{n}, Q_{n}, T_{n}, S_{n} \\ \gamma_{n}^{2}, \gamma_{n}^{2}, \frac{T_{n}}{\pi_{n}^{2}}, \frac{S_{n}}{\pi_{n}^{2}} \end{pmatrix}.$$
(3.5b)

To derive isospectral AL hierarchy, we need to take  $z_t \equiv 0$  in (3.3) and expand  $(B_n, C_n, b_n, c_n)^T$ as

$$\begin{pmatrix} B_n \\ C_n \\ b_n \\ c_n \end{pmatrix} = \sum_{j=0}^m \begin{pmatrix} B_n^{(j)} \\ C_n^{(j)} \\ b_n^{(j)} \\ c_n^{(j)} \end{pmatrix} z^{2(m-j)+1}, \qquad m = 0, 1, 2, \cdots.$$
(3.6)

Then by setting  $(B_n^{(0)}, C_n^{(0)}, b_n^{(0)}, c_n^{(0)})^T = (0, 0, 0, 0)^T$ ,  $A_0 = -D_0 = A_n|_{u_n=0} = -D_n|_{u_n=0} = \frac{1}{2}z^{2m}$ , and comparing the coefficients of the same powers of z in (3.3) we get

$$\begin{pmatrix} Q_n \\ R_n \\ S_n \\ T_n \end{pmatrix}_{t_m} = (1 - \delta_{0,m}) L_2 \begin{pmatrix} -B_n^{(m)} \\ C_n^{(m)} \\ -b_n^{(m)} \\ c_n^{(m)} \end{pmatrix} + \delta_{0,m} \begin{pmatrix} Q_n \\ -R_n \\ S_n \\ -T_n \end{pmatrix},$$
(3.7a)

$$\begin{pmatrix} -B_n^{(j+1)} \\ C_n^{(j+1)} \\ -b_n^{(j+1)} \\ c_n^{(j+1)} \end{pmatrix} = -L_1^{-1}L_2 \begin{pmatrix} -B_n^{(j)} \\ C_n^{(j)} \\ -b_n^{(j)} \\ c_n^{(j)} \end{pmatrix}, \qquad j = 1, 2, \cdots, m-1,$$
(3.7b)

$$\begin{pmatrix} -B_n^{(1)} \\ C_n^{(1)} \\ -b_n^{(1)} \\ c_n^{(1)} \end{pmatrix} = -L_1^{-1} \begin{pmatrix} Q_n \\ -R_n \\ S_n \\ -T_n \end{pmatrix},$$
(3.7c)

where the subindex m for t indicates the order of the expansion (3.6) as well as the order of member in isospectral hierarchy. This further yields a isospectral hierarchy

$$u_{n,t_m} = K^{(m)} = L^m K^{(0)}, \qquad m = 0, 1, 2, \cdots,$$
(3.8)

where

$$K^{(0)} = (Q_n, -R_n, S_n, -T_n)^T, (3.9)$$

and the recursion operator L is defined by

$$L = -L_2 L_1^{-1} = \begin{pmatrix} -S_n R_n & -S_n Q_n & \gamma_n^2 & -Q_n^2 E^{-1} \\ 0 & 0 & 0 & E^{-1} \\ \pi_n^2 E & -S_n^2 & -S_n R_n & -S_n Q_n E^{-1} \\ \frac{\pi_n^2 R_n^2}{\gamma_n^2} & \frac{1 - Q_n R_n S_n T_n}{\gamma_n^2} & T_n R_n & T_n Q_n E^{-1} \end{pmatrix}$$
$$- \begin{pmatrix} Q_n \\ -R_n \\ S_n \\ -T_n \end{pmatrix} (E - 1)^{-1} (T_n E, S_n, R_n, Q_n E^{-1})$$
$$- \begin{pmatrix} S_n \gamma_n^2 \\ -T_{n-1} \gamma_n^2 \\ Q_{n+1} \pi_n^2 E \\ -R_n \pi_n^2 \end{pmatrix} (E - 1)^{-1} \left( \frac{R_n}{\gamma_n^2}, \frac{Q_n}{\gamma_n^2}, \frac{T_n}{\pi_n^2}, \frac{S_n}{\pi_n^2} \right).$$
(3.10)

If we expand  $(B_n, C_n, b_n, c_n)^T$  in another direction, i.e.,

$$\begin{pmatrix} B_n \\ C_n \\ b_n \\ c_n \end{pmatrix} = \sum_{j=m}^0 \begin{pmatrix} B_n^{(j)} \\ C_n^{(j)} \\ b_n^{(j)} \\ c_n^{(j)} \end{pmatrix} z^{2(m-j)-1}, \qquad m = 0, -1, -2, \cdots,$$
(3.11)

and take  $(B_n^{(0)}, C_n^{(0)}, b_n^{(0)}, c_n^{(0)})^T = (0, 0, 0, 0)^T$ ,  $A_0 = -D_0 = A_n|_{u_n=0} = -D_n|_{u_n=0} = \frac{1}{2}z^{2m}$ , we can have relation

$$\begin{pmatrix} Q_n \\ R_n \\ S_n \\ T_n \end{pmatrix}_{t_m} = (1 - \delta_{0,m}) L_1 \begin{pmatrix} -B_n^{(m)} \\ C_n^{(m)} \\ -b_n^{(m)} \\ c_n^{(m)} \end{pmatrix} + \delta_{0,m} \begin{pmatrix} Q_n \\ -R_n \\ S_n \\ -T_n \end{pmatrix}, \quad (3.12a)$$
$$\begin{pmatrix} -B_n^{(j)} \\ C_n^{(j)} \\ -b_n^{(j)} \\ c_n^{(j)} \end{pmatrix} = -L_2^{-1} L_1 \begin{pmatrix} -B_n^{(j+1)} \\ C_n^{(j+1)} \\ -b_n^{(j+1)} \\ c_{n+1}^{(j)} \end{pmatrix}, \quad j = -2, -3, \cdots, m, \quad (3.12b)$$

$$\begin{pmatrix} -B_n^{(j)} \\ C_n^{(j)} \\ -b_n^{(j)} \\ c_n^{(j)} \end{pmatrix} = -L_2^{-1}L_1 \begin{pmatrix} -B_n^{(j+1)} \\ C_n^{(j+1)} \\ -b_n^{(j+1)} \\ c_{n+1}^{(j)} \end{pmatrix}, \qquad j = -2, -3, \cdots, m,$$
(3.12b)

$$\begin{pmatrix} -B_n^{(-1)} \\ C_n^{(-1)} \\ -b_n^{(-1)} \\ c_n^{(-1)} \end{pmatrix} = -L_2^{-1} \begin{pmatrix} Q_n \\ -R_n \\ S_n \\ -T_n \end{pmatrix},$$
(3.12c)

and further get a negative order isospectral hierarchy

$$u_{n,t_m} = K^{(m)} = L^m K^{(0)}, \qquad m = 0, -1, -2, \cdots,$$
 (3.13)

where  $K^{(0)}$  and L are given by (3.9) and (3.10) respectively.

Obviously, (3.8) and (3.13) can be jointed together and written as a uniformed isospectral AL hierarchy

$$u_{n,t_m} = K^{(m)} = L^m K^{(0)}, \qquad m \in \mathbb{Z}.$$
 (3.14)

In Appendix A we will list out the first few isospectral equations and their related Lax pairs.

### 3.2 Non-isospectral hierarchy

For the non-isospectral case, we suppose the time evolution of spectral parameter z follows  $z_{t_m} = \frac{1}{2}z^{2m+1}$  for any given  $m \in \mathbb{Z}$ . We can first expand  $(B_n, C_n, b_n, c_n)^T$  as (3.6) and still take  $(B_n^{(0)}, C_n^{(0)}, b_n^{(0)}, c_n^{(0)})^T = (0, 0, 0, 0)^T$  but  $A_0 = -D_0 = 0$ . Then we can get a recursion relation which is similar to (3.7) but in this non-isospectral case (3.7a) and (3.7c) are replaced by

$$\begin{pmatrix} Q_n \\ R_n \\ S_n \\ T_n \end{pmatrix}_{t_m} = (1 - \delta_{0,m}) L_2 \begin{pmatrix} -B_n^{(m)} \\ C_n^{(m)} \\ -b_n^{(m)} \\ c_n^{(m)} \end{pmatrix} + \delta_{0,m} \begin{pmatrix} (2n + \frac{1}{2})Q_n \\ -(2n + \frac{3}{2})S_n \\ (2n + \frac{3}{2})S_n \\ -(2n + \frac{3}{2})T_n \end{pmatrix}$$
$$\begin{pmatrix} -B_n^{(1)} \\ C_n^{(1)} \\ -b_n^{(1)} \\ c_n^{(1)} \end{pmatrix} = -L_1^{-1} \begin{pmatrix} (2n + \frac{1}{2})Q_n \\ -(2n + \frac{1}{2})R_n \\ (2n + \frac{3}{2})S_n \\ -(2n + \frac{3}{2})T_n \end{pmatrix} .$$

It then follows that a positive order non-isospectral hierarchy is

$$u_{n,t_m} = \sigma^{(m)} = L^m \sigma^{(0)}, \qquad m = 0, 1, 2, \cdots,$$
 (3.15)

where

$$\sigma^{(0)} = \begin{pmatrix} (2n + \frac{1}{2})Q_n \\ -(2n + \frac{1}{2})R_n \\ (2n + \frac{3}{2})S_n \\ -(2n + \frac{3}{2})T_n \end{pmatrix},$$
(3.16)

and L is the recursion operator given by (3.10). After a discussion for a negative order expansion of  $(B_n, C_n, b_n, c_n)^T$  and deriving a negative order non-isospectral hierarchy, one can finally reach to a uniformed non-isospectral AL hierarchy:

$$u_{n,t_m} = \sigma^{(m)} = L^m \sigma^{(0)}, \qquad m \in \mathbb{Z}.$$
 (3.17)

In Appendix A we will also list out the first few non-isospectral equations and their related Lax pairs.

### 3.3 Zero-curvature representations

We have derived isospectral hierarchy (3.14) and non-isospectral hierarchy (3.17). Suppose that their Lax pairs are respectively

$$\Phi_{n+1} = U_n \Phi_n, \quad \Phi_{n,t_m} = G_n^{(m)} \Phi_n, \tag{3.18}$$

and

$$\Phi_{n+1} = U_n \Phi_n, \quad \Phi_{n,t_m} = W_n^{(m)} \Phi_n, \tag{3.19}$$

where  $m \in \mathbb{Z}$  and  $U_n$  is defined in (2.11). Then in isospectral case the zero-curvature equation which is related to (3.18) is

$$U_{n,t_m} = (EG_n^{(m)})U_n - U_n G_n^{(m)}, (3.20)$$

and in non-isospectral case

$$U_{n,t_m} = (EW_n^{(m)})U_n - U_n W_n^{(m)}.$$
(3.21)

Noticing the definition (2.1) for a Gateaux derivative,  $U_{n,t_m}$  can be rewritten as

$$U_{n,t_m} = U'_n[u_{n,t_m}] + U_{n,z} \cdot z_{t_m}, \qquad (3.22)$$

by means of which we have

**Proposition 3.1.** The isospectral flows  $\{K^{(m)}\}\$  and non-isospectral flows  $\{\sigma^{(m)}\}\$  admit zero curvature representations

$$U'_{n}[K^{(m)}] = (EG_{n}^{(m)})U_{n} - U_{n}G_{n}^{(m)}, \qquad (3.23)$$

$$U'_{n}[\sigma^{(m)}] = (EW_{n}^{(m)})U_{n} - U_{n}W_{n}^{(m)} - \frac{1}{2}z^{2m+1}U_{n,z}, \qquad (3.24)$$

where  $m \in \mathbb{Z}$ ,  $K^{(m)}$  and  $\sigma^{(m)} \in \mathcal{V}_4$ ,  $G_n^{(m)}$  and  $W_n^{(m)} \in \mathcal{Q}_2(z)$  and satisfy

$$G_n^{(m)}|_{u_n=0} = \frac{z^{2m}}{2} \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix}, \quad W_n^{(m)}|_{u_n=0} = z^{2m} \begin{pmatrix} n & 0\\ 0 & -n \end{pmatrix}.$$
 (3.25)

Besides, noting that  $U'_n$  is an injective homomorphism when  $Q_n R_n + S_n T_n \neq 0$ , we have (cf. [11, 15])

Lemma 3.1. The matrix equation

$$U'_{n}[X_{n}] = (EV_{n})U_{n} - U_{n}V_{n}, \quad X_{n} \in \mathcal{V}_{4}, \quad V_{n} \in \mathcal{Q}_{2}(z) \text{ and } V_{n}|_{u_{n}=0} = 0$$
(3.26)

has only zero solutions  $X_n = 0$  and  $V_n = 0$ .

This lemma and the zero-curvature representations (3.23) and (3.24) will play important roles in constructing symmetries and determining their algebraic structures.

### 3.4 Recursion operator and hereditary and strong symmetry

The flows  $\{K^{(m)}\}, \{\sigma^{(m)}\}\)$  and their recursion relation can also be derived from the following way.

Firstly, we start from the matrix equations

$$U'_{n}[X_{n}] = (EG_{n})U_{n} - U_{n}G_{n}, \qquad (3.27)$$

$$U'_{n}[Z_{n}] = (EW_{n})U_{n} - U_{n}W_{n} - U_{n,z} \cdot \frac{z}{2}, \qquad (3.28)$$

where the unknowns are  $X_n, Z_n \in \mathcal{V}_4$  and  $G_n, W_n \in \mathcal{Q}_2(z)$ . Obviously, both of the two equations admit non-zero solution pairs  $\{X_n, G_n\}$  and  $\{Z_n, W_n\}$  (see (3.23) and (3.24) for m = 0). Secondly, for equation

$$U'_{n}[X_{n} - z^{\alpha}Y_{n}] = (EV_{n})U_{n} - U_{n}V_{n}$$
(3.29)

where  $X_n, Y_n \in \mathcal{V}_4$  and  $V_n \in \mathcal{Q}_2(z)$ , one can find that when  $\alpha = 2$ , for any given  $Y_n \neq 0 \in \mathcal{V}_4$ , there exist unique solutions  $X_n \in \mathcal{V}_4$ ,  $V_n \in \mathcal{Q}_2(z)$  where  $V_n|_{u_n=0} = 0$ . Thus  $X_n$  and  $Y_n$  are connected by a map:<sup>2</sup>

$$L: \quad X_n = LY_n. \tag{3.30}$$

The map, L, is nothing but the recursion operator (3.10). Then, thirdly, let  $Y_n = K^{(0)}$  or  $\sigma^{(0)}$  in (3.29), one can get  $K^{(1)}$  or  $\sigma^{(1)}$  by taking  $\alpha = 2$  and  $K^{(-1)}$  or  $\sigma^{(-1)}$  by taking  $\alpha = -2$ . Repeating the procedure one can generate all the isospectral flows  $\{K^{(m)}\}$  and non-isospectral flows  $\{\sigma^{(m)}\}$ .

Besides, with the recursion operator L in hand, (3.29) becomes

$$U'_{n}[X - z^{2}LX] = (EV_{n})U_{n} - U_{n}V_{n}.$$
(3.31)

Then, in the light of Theorem 1 in Ref. [11] (also see Appendix B), we immediately have

**Proposition 3.2.** The recursion operator L is hereditary and a strong symmetry<sup>3</sup> for the isospectral AL hierarchy (3.14).

### 4 Symmetries and Lie algebras

In this section, we construct two types of symmetries for the isospectral and non-isospectral AL hierarchies, respectively. To do that, let us first look at the algebraic structure of flows  $\{K^{(m)}\}\$  and  $\{\sigma^{(m)}\}\$ . We note that the proofs for this section are quite similar to those in [15] and here we skip them.

### 4.1 Algebra of flows

Making use of the identity [21]

$$U'_{n}[\llbracket f, g \rrbracket] = (U'_{n}[f])'[g] - (U'_{n}[g])'[f], \quad \forall f, g \in \mathcal{V}_{4},$$
(4.1)

from the zero-curvature representations (3.23) and (3.24) we can derive the following relations:

<sup>&</sup>lt;sup>2</sup>If  $\alpha = -2$  then the map is  $X = L^{-1}Y$ .

<sup>&</sup>lt;sup>3</sup> For the definitions of a hereditary operator and a strong symmetry operator, one can refer to [28].

**Lemma 4.1.** The isospectral flows  $\{K^{(m)}\}$  and non-isospectral flows  $\{\sigma^{(m)}\}$  satisfy

$$U'_{n}[\llbracket K^{(m)}, K^{(s)} \rrbracket] = (E < G_{n}^{(m)}, G_{n}^{(s)} >) U_{n} - U_{n} < G_{n}^{(m)}, G_{n}^{(s)} >,$$
(4.2a)

$$U'_{n}[\llbracket K^{(m)}, \sigma^{(s)} \rrbracket] = (E < G_{n}^{(m)}, W_{n}^{(s)} >) U_{n} - U_{n} < G_{n}^{(m)}, W_{n}^{(s)} >,$$
(4.2b)

$$U_n'[\llbracket\sigma^{(m)}, \sigma^{(s)}\rrbracket] = (E < W_n^{(m)}, W_n^{(s)} >)U_n - U_n < W_n^{(m)}, W_n^{(s)} > -\frac{1}{2}(m-s)U_{n,z}z^{2(m+s)+1},$$
(4.2c)

where

$$\langle G_n^{(m)}, G_n^{(s)} \rangle = G_n^{(m)\prime}[K^{(s)}] - G_n^{(s)\prime}[K^{(m)}] + [G_n^{(m)}, G_n^{(s)}],$$
(4.3a)

$$\langle G_n^{(m)}, W_n^{(s)} \rangle = G_n^{(m)\prime}[\sigma^{(s)}] - W_n^{(s)\prime}[K^{(m)}] + [G_n^{(m)}, W_n^{(s)}] + \frac{1}{2}G_{n,z}^{(m)}z^{2s+1},$$
(4.3b)

$$\langle W_{n}^{(m)}, W_{n}^{(s)} \rangle = W_{n}^{(m)\prime}[\sigma^{(s)}] - W_{n}^{(s)\prime}[\sigma^{(m)}] + [W_{n}^{(m)}, W_{n}^{(s)}] + \frac{1}{2}W_{n,z}^{(m)}z^{2s+1} - \frac{1}{2}W_{n,z}^{(s)}z^{2m+1}.$$
(4.3c)

This lemma can be proved via a similar procedure as in [15] and here we skip the proof. Based on the lemma we then come up with a algebra for the flows  $\{K^{(m)}\}$  and  $\{\sigma^{(m)}\}$  (cf. [15]).

**Lemma 4.2.** The isospectral and non-isospectral flows,  $\{K^{(m)}\}$  and  $\{\sigma^{(m)}\}$ , compose an infinitedimensional Lie algebra  $\mathcal{F}$  through the Lie product  $\llbracket, \cdot \rrbracket$  and possess the following relations

$$[\![K^{(m)}, K^{(s)}]\!] = 0, (4.4a)$$

$$\llbracket K^{(m)}, \sigma^{(s)} \rrbracket = m K^{(m+s)}, \tag{4.4b}$$

$$[\![\sigma^{(m)}, \sigma^{(s)}]\!] = (m-s)\sigma^{(m+s)}.$$
(4.4c)

### 4.2 Symmetries for the isospectral and non-isospectral AL hierarchies

**Theorem 4.1.** Any given member  $u_{n,t_m} = K^{(m)}$  in the isospectral four-potential AL hierarchy (3.14) possesses the following two sets of symmetries, i.e.,

$$\{K^{(s)}\}$$
 and  $\{\tau^{(m,s)} = mt_m K^{(m+s)} + \sigma^{(s)}\}, s \in \mathbb{Z}.$  (4.5)

These symmetries form a centerless Kac-Moody-Virasoro (KMV) algebra  $\mathcal{S}$  with the following structure

$$\llbracket K^{(l)}, K^{(s)} \rrbracket = 0, \tag{4.6a}$$

$$\llbracket K^{(l)}, \tau^{(m,s)} \rrbracket = l K^{(l+s)}, \tag{4.6b}$$

$$\llbracket \tau^{(m,s)}, \tau^{(m,l)} \rrbracket = (s-l)\tau^{(m,s+l)}.$$
(4.6c)

Obviously, the Lie algebras  $\mathcal{F}$  and  $\mathcal{S}$  are respectively generated by the following elements

{
$$\sigma^{(1)}$$
 (or  $\sigma^{(-1)}$ ),  $\sigma^{(2)}$ ,  $\sigma^{(-2)}$ ,  $K^{(1)}$  (or  $K^{(-1)}$ )}; (4.7)

{
$$\tau^{(m,1)}$$
 (or  $\tau^{(m,-1)}$ ),  $\tau^{(m,2)}$ ,  $\tau^{(m,-2)}$ ,  $K^{(1)}$  (or  $K^{(-1)}$ )}. (4.8)

**Theorem 4.2.** Any given member  $u_{n,t_m} = \sigma^{(m)}$  in the non-isospectral four-potential AL hierarchy (3.17) has two sets of symmetries, i.e.,

$$\eta^{(m,s)} = \sum_{j=0}^{s} C_s^j (mt_m)^{s-j} \sigma^{(m-jm)} \qquad (s = 0, 1, 2, \cdots),$$
(4.9a)

$$\gamma^{(m,s)} = \sum_{j=0}^{s} C_s^j (mt_m)^{s-j} K^{(-jm)} \quad (s = 0, 1, 2, \cdots),$$
(4.9b)

which we call  $\eta$ -symmetries and  $\gamma$ -symmetries, respectively. Here  $C_s^j = \frac{s!}{j!(s-j)!}$ . These symmetries form a centerless KMV algebra  $\mathcal{H}$  with the following structure

$$[\eta^{(m,s)}, \eta^{(m,l)}] = (l-s)m\eta^{(m,s+l-1)},$$
(4.10a)

$$[\![\gamma^{(m,s)}, \gamma^{(m,l)}]\!] = 0, \tag{4.10b}$$

$$[\![\eta^{(m,s)}, \gamma^{(m,l)}]\!] = lm\gamma^{(m,s+l-1)}.$$
(4.10c)

The algebra can be generated by

$$\{\eta^{(m,0)}, \eta^{(m,3)}, \gamma^{(m,1)}\}.$$
 (4.11)

### 4.3 Relations between the recursion operator and flows

**Theorem 4.3.** The isospectral flows  $\{K^{(m)}\}$ , non-isospectral flows  $\{\sigma^{(m)}\}\$  and recursion operator L satisfy the relations

$$L'[K^{(m)}] - [K^{(m)'}, L] = 0, (4.12)$$

$$L'[\sigma^{(m)}] - [\sigma^{(m)'}, L] - L^{m+1} = 0, \qquad (4.13)$$

where  $m \in \mathbb{Z}$  and [A, B] = AB - BA.

Utilizing the relations one can also derive symmetries (4.5) and their algebra structure (4.6) by means of inductive approach, as in [29]. In that way the non-isospectral flow  $\sigma^{(0)}$  will play the role of a master symmetry [30].

### 5 Reduction to two-potential case

It is possible to reduce isospectral, non-isospectral AL hierarchies and their symmetries from four-potential case to two-potential case.

### **5.1** Reduction of $(S_n, T_n) = (0, 0)$

#### 5.1.1 Spectral problem

Let us start from the four-potential AL-spectral problem (2.11). Taking  $(S_n, T_n) = (0, 0)$  in (2.11) yields

$$\Phi_{n+1} = \overline{U}_n \Phi_n, \quad \overline{U}_n = \begin{pmatrix} z^2 & zQ_n \\ z^{-1}R_n & z^{-2} \end{pmatrix}, \quad \Phi_n = \begin{pmatrix} \phi_{1,n} \\ \phi_{2,n} \end{pmatrix}, \tag{5.1}$$

but this is not the canonical form (2.9) yet. Next we introduce a gauge transformation

$$\Psi_n = \begin{pmatrix} z^{-1} & 0\\ 0 & 1 \end{pmatrix} \Phi_n, \quad \Psi_n = \begin{pmatrix} \psi_{1,n}\\ \psi_{2,n} \end{pmatrix},$$
(5.2)

under which (5.1) becomes

$$\Psi_{n+1} = M_n \Psi_n, \quad M_n = \begin{pmatrix} z^2 & Q_n \\ R_n & z^{-2} \end{pmatrix},$$
(5.3)

which is the canonical form (2.9) in the light of  $z^2 = \lambda$ .

### 5.1.2 Closeness discussion

For convenience we introduce some notations. Let

$$\overline{u}_n = \begin{pmatrix} Q_n \\ R_n \end{pmatrix}, \quad \overline{K}^{(m)} = \begin{pmatrix} K_1^{(m)} \\ K_2^{(m)} \end{pmatrix}_{(S_n, T_n) = (0, 0)}, \quad \overline{\sigma}^{(m)} = \begin{pmatrix} \sigma_1^{(m)} \\ \sigma_2^{(m)} \end{pmatrix}_{(S_n, T_n) = (0, 0)}, \quad (5.4)$$

where  $K_j^{(m)}$  and  $\sigma_j^{(m)}$  are the *j*-th elements of  $K^{(m)}$  and  $\sigma^{(m)}$ . Noting that the reduction  $(S_n, T_n) = (0, 0)$  yields  $u_n|_{(S_n, T_n) = (0, 0)} = (\overline{u}_n^T, 0, 0)^T$ , it is necessary to discuss whether the reduction is closed, i.e., whether

$$K^{(m)}|_{(S_n,T_n)=(0,0)} = \begin{pmatrix} \overline{K}^{(m)} \\ 0 \\ 0 \end{pmatrix}, \ \sigma^{(m)}|_{(S_n,T_n)=(0,0)} = \begin{pmatrix} \overline{\sigma}^{(m)} \\ 0 \\ 0 \end{pmatrix}.$$
 (5.5)

Next we will see this is true for  $m = 2h, h \in \mathbb{Z}$ .

**Theorem 5.1.** Under the reduction  $(S_n, T_n) = (0, 0)$ , those even order members in the fourpotential AL hierarchies (3.14) and (3.17) reduce to two-potential isospectral AL-hierarchies

$$\overline{u}_{n,t_{2h}} = \overline{K}^{(2h)} = \overline{L}^h \overline{K}^{(0)}, \qquad h \in \mathbb{Z}$$
(5.6)

and

$$\overline{u}_{n,t_{2h}} = \overline{\sigma}^{(2h)} = \overline{L}^h \overline{\sigma}^{(0)}, \qquad h \in \mathbb{Z},$$
(5.7)

where

$$\overline{K}^{(0)} = (Q_n, -R_n)^T, \quad \overline{\sigma}^{(0)} = (2n + \frac{1}{2})(Q_n, -R_n)^T,$$
(5.8)

and the recursion operator  $\overline{L}$  is

$$\overline{L} = \begin{pmatrix} E & 0\\ 0 & E^{-1} \end{pmatrix} + \begin{pmatrix} -Q_n E\\ R_n \end{pmatrix} (E-1)^{-1} (R_n E, Q_n E^{-1}) + \gamma_n^2 \begin{pmatrix} -Q_{n+1}E\\ R_{n-1} \end{pmatrix} (E-1)^{-1} \begin{pmatrix} R_n\\ \gamma_n^2, \frac{Q_n}{\gamma_n^2} \end{pmatrix}.$$
(5.9)

*Proof.* First, look at (3.9) and (3.16). When m = 0, the reduction  $(S_n, T_n) = (0, 0)$  is obviously closed and the resulting flows  $\overline{K}^{(0)}$  and  $\overline{\sigma}^{(0)}$  are as given in (5.8). Meanwhile, by direct calculation we find

$$L^2|_{(S_n,T_n)=(0,0)} = \begin{pmatrix} \overline{L} & \mathbf{0} \\ \mathbf{0} & H \end{pmatrix}$$

where

$$H = \begin{pmatrix} \gamma_{n+1}^2 E & -Q_{n+1}^2 \\ R_n^2 & (1+Q_n R_n) E^{-1} \end{pmatrix} - 2 \begin{pmatrix} Q_{n+1}E \\ -R_n \end{pmatrix} (E-1)^{-1} (R_n, Q_n E^{-1}).$$

That means the closeness is valid for those even order members of four-potential AL hierarchies (3.14) and (3.17) when we take  $(S_n, T_n) = (0, 0)$ . Thus the proof is completed.

#### 5.1.3 Flows under gauge transformation

With the closeness in hand, let us make a comparison for the present results and those in [15]. Isospectral flows and recursion operator are exactly same but non-isospectral flows are different. The basic non-isospectral flow  $\hat{\sigma}^{(0)}$  given in [15] is

$$\widehat{\sigma}^{(0)} = (2n+1)(Q_n, -R_n)^T, \tag{5.10}$$

and the difference from  $\overline{\sigma}^{(0)}$  is

$$\overline{\sigma}^{(0)} - \widehat{\sigma}^{(0)} = -\frac{1}{2} (Q_n, -R_n)^T.$$
(5.11)

To understand the difference on non-isospectral flows, we go back to the gauge transformation (5.2).

Obviously, the Lax pair of  $\overline{u}_{n,t_0} = \overline{\sigma}^{(0)}$  is composed by (5.1) and

$$\Phi_{n,t_0} = \overline{V}_n^{(0)} \Phi_n, \qquad \overline{V}_n^{(0)} = V_n^{(0)}|_{(S_n,T_n) = (0,0)}, \tag{5.12}$$

of which the zero-curvature equation reads

$$\overline{U}_{n,t_0} = (E\overline{V}_n^{(0)})\overline{U}_n - \overline{U}_n\overline{V}_n^{(0)}.$$
(5.13)

Under the gauge transformation (5.2), (5.12) turns out to be

$$\Psi_{n,t_0} = \hat{V}_n^{(0)} \Psi_n \tag{5.14}$$

where

$$\widehat{V}_{n}^{(0)} = \widehat{N}_{n}^{(0)} - \begin{pmatrix} z^{-1}z_{t} & 0\\ 0 & 0 \end{pmatrix}, \quad z_{t} = \frac{z}{2}, \quad \widehat{N}_{n}^{(0)} = \begin{pmatrix} z^{-1} & 0\\ 0 & 1 \end{pmatrix} \overline{V}_{n}^{(0)} \begin{pmatrix} z & 0\\ 0 & 1 \end{pmatrix}.$$
(5.15)

Meanwhile the zero-curvature equation (5.13) is transformed to

$$M_{n,t_0} - (E\widehat{V}_n^{(0)})M_n + M_n\widehat{V}_n^{(0)} = 0, (5.16)$$

i.e.,

$$M_{n,t_0} - (E\hat{N}_n^{(0)})M_n + M_n\hat{N}_n^{(0)} = \frac{1}{2} \begin{pmatrix} 0 & -Q_n \\ R_n & 0 \end{pmatrix}.$$
 (5.17)

Compared with [15] we find that the l.h.s. of (5.17) equating to zero yields the non-isospectral equation  $\overline{u}_{n,t_0} = \widehat{\sigma}^{(0)}$  while the r.h.s. just gives the difference (5.11). Now we can see that it is just the time-dependent multiplier  $\begin{pmatrix} z^{-1} & 0 \\ 0 & z^{-1} \end{pmatrix}$  in the transformation

Now we can see that it is just the time-dependent multiplier  $\begin{pmatrix} z^{-1} & 0 \\ 0 & 1 \end{pmatrix}$  in the transformation (5.2) contributes the extra term (5.11) for the non-isospectral flow  $\overline{\sigma}^{(0)}$ , but in isospectral case this multiplier adds nothing new since in this turn z is independent of time. However, the extra term in (5.11) is nothing but  $\frac{1}{2}\overline{K}^{(0)}$ , which means the obtained non-isospectral flow  $\overline{\sigma}^{(2h)}$  is only a summation of  $\widehat{\sigma}^{(h)}$  and  $-\frac{1}{2}\overline{K}^{(2h)}$ , where  $\widehat{\sigma}^{(h)} = \overline{L}^h \widehat{\sigma}^{(0)}$  is the non-isospectral flow derived in [15].

#### 5.2 Symmetries

The closeness also guarantees reduction of symmetries and their algebraic structures. From Theorem 4.1 and 4.2 one can directly have

**Theorem 5.2.** Any given member  $\overline{u}_{n,t_{2h}} = \overline{K}^{(2h)}$  in the isospectral two-potential AL hierarchy (5.6) can have two sets of symmetries

$$\{\overline{K}^{(2s)}\}$$
 and  $\{\overline{\tau}^{(2h,2s)} = 2ht_{2h}\overline{K}^{(2h+2s)} + \overline{\sigma}^{(2s)}\}, s \in \mathbb{Z},$  (5.18)

which form a centerless KMV algebra  $\overline{\mathcal{S}}$  with structure

$$\llbracket \overline{K}^{(2l)}, \overline{K}^{(2s)} \rrbracket = 0, \tag{5.19a}$$

$$\llbracket \overline{K}^{(2l)}, \overline{\tau}^{(2h,2s)} \rrbracket = 2l \overline{K}^{(2l+2s)}, \tag{5.19b}$$

$$[\![\overline{\tau}^{(2h,2s)},\overline{\tau}^{(2h,2l)}]\!] = 2(s-l)\overline{\tau}^{(2h,2s+2l)},$$
(5.19c)

and generators

$$\{\overline{\tau}^{(2h,2)} \text{ (or } \overline{\tau}^{(2h,-2)}), \overline{\tau}^{(2h,4)}, \overline{\tau}^{(2h,-4)}, \overline{K}^{(2)} \text{ (or } \overline{K}^{(-2)})\}.$$
 (5.20)

**Theorem 5.3.** Any given member  $\overline{u}_{n,t_{2h}} = \overline{\sigma}^{(2h)}$  in the non-isospectral two-potential AL hierarchy (5.7) can have two sets of symmetries

$$\overline{\eta}^{(2h,s)} = \sum_{j=0}^{s} C_s^j (2ht_{2h})^{s-j} \overline{\sigma}^{(2h-2jh)} \qquad (s=0,1,2,\cdots),$$
(5.21a)

$$\overline{\gamma}^{(2h,s)} = \sum_{j=0}^{s} C_s^j (2ht_{2h})^{s-j} \overline{K}^{(-2jh)} \quad (s = 0, 1, 2, \cdots),$$
(5.21b)

which form a centerless KMV algebra  $\overline{\mathcal{H}}$  with structure

$$\llbracket \overline{\eta}^{(2h,s)}, \overline{\eta}^{(2h,l)} \rrbracket = 2(l-s)h\overline{\eta}^{(2h,s+l-1)},$$
(5.22a)

$$[\![\overline{\gamma}^{(2h,s)}, \overline{\gamma}^{(2h,l)}]\!] = 0, \qquad (5.22b)$$

$$[\![\overline{\eta}^{(2h,s)}, \overline{\gamma}^{(2h,l)}]\!] = 2lh\overline{\gamma}^{(2h,s+l-1)}, \qquad (5.22c)$$

and generators

$$\{\overline{\eta}^{(2h,0)}, \overline{\eta}^{(2h,3)}, \overline{\gamma}^{(2h,1)}\}.$$
 (5.23)

# 6 Conclusions

We have derived isospectral and non-isospectral four-potential AL hierarchies and their recursion operator. Both hierarchies have been shown to have infinitely many symmetries and Lie algebras which belong to centerless KMV algebras, respectively. These structures are derived by means of zero-curvature representations of flows. What is special is that these structures cover both positive and negative order four-potential AL hierarchies, and each member in the non-isospectral AL hierarchy also possesses two sets of symmetries which compose a centerless KMV algebra too. These two points make the algebraic structure of the AL hierarchies quite different from many other Lax integrable systems. This is due to the symmetrical form (in terms of z and  $z^{-1}$ ) of the AL spectral matrix (see (2.11)). Besides, it is now clear for the relation between four-potential and two-potential AL hierarchies. All the even order members in the four-potential isospectral and non-isospectral hierarchies can be reduced to two-potential case by directly taking  $(S_n, T_n) = (0, 0)$ . The new recursion operator for two-potential case is  $L^2$ . Meanwhile, the reduction keeps the algebraic structures of symmetries invariant. The procedure and main results are possible to apply to Ablowitz-Ladik systems with vector potentials and this will be discussed elsewhere. This paper will be continued by the second part in which we will discuss symmetries and recursion operator for the integrable discrete nonlinear Schrödinger equation and the discrete AKNS hierarchy.

# Acknowledgement

This project is supported by the National Natural Science Foundation of China (10671121) and Shanghai Leading Academic Discipline Project (No.J50101).

# References

- M.J. Ablowitz, J.F. Ladik, Nonlinear differential-difference equations, J. Math. Phys., 16 (1975) 598-603.
- [2] R. Hirota, Nonlinear partial difference equations. I. A difference analogue of the Korteweg-de Vries equation, J. Phys. Soc. Jpn., 43 (1977) 1424-33.
- [3] E. Date, M. Jimbo, T. Miwa, Method for generating discrete soliton equations. I, J. Phys. Soc. Jpn., 51 (1982) 4116-24.
- [4] B.A. Kuperschmidt, Discrete Lax Equations and Differential-difference Calculus, Astérisque, No. 123 (1985).
- [5] M.J. Ablowitz, J.F. Ladik, Nonlinear differential-difference equations and Fourier analysis, J. Math. Phys., 17 (1976) 1011-8.

- [6] M.J. Ablowitz, B. Prinari, A.D. Trubatch, Discrete and Continuous Nonlinear Schrödinger Systems, Cambridge Univ. Press, 2004.
- [7] M.J. Ablowitz, D.J. Kaup, A.C. Newell, H. Segur, Nonlinear-evolution equations of physical significance, Phys. Rev. Lett., 31 (1973) 125-7.
- [8] X.G. Geng, Darboux transformation of the discrete Ablowitz-Ladik eigenvalue problem, Acta Math. Sci., 9 (1989) 21-6.
- [9] Y.B. Zeng, S.R. Wojciechowski, Restricted flows of the Ablowitz-Ladik hierarchy and their continuous limits, J. Phys. A: Math. Gen., 28 (1995) 113-34.
- [10] K.M. Tamizhmani, W.X. Ma, Master symmetries from Lax operators for certain lattice soliton hierarchies, J. Phys. Soc. Jpn., 69 (2000) 351-61.
- [11] D.J. Zhang, D.Y. Chen, Hamiltonian structure of discrete soliton systems, J. Phys. A: Math. Gen., 35 (2002) 7225-41.
- [12] D.J. Zhang, D.Y. Chen, The conservation laws of some discrete soliton systems, Chaos, Solitons & Fractals, 14 (2002) 573-9.
- [13] X.G. Geng, H.H. Dai, C.W. Cao, Algebro-geometric constructions of the discrete Ablowitz-Ladik flows and applications, J. Math. Phys., 44 (2003) 4573-88.
- [14] V.E. Vekslerchik, Implementation of the Bäcklund transformations for the Ablowitz-Ladik hierarchy, J. Phys. A: Math. Gen., 39 (2006) 6933-53.
- [15] D.J. Zhang, T.K. Ning, J.B. Bi, D.Y. Chen, New symmetries for the Ablowitz-Ladik hierarchies, Phys. Lett. A, 359 (2006) 458-66.
- [16] X.G. Geng, H.H. Dai, J.Y. Zhang, Decomposition of the discrete Ablowitz-Ladik hierarchy, Stud. Appl. Math., 118 (2007) 281-312.
- [17] F. Gesztesy, H. Holden, J. Michor, G. Teschl, Local conservation laws and the Hamiltonian formalism for the Ablowitz-Ladik hierarchy, Stud. Appl. Math., 120 (2008) 361-423.
- [18] Y. Cheng, On discrete Ablowitz-Ladik eigenvalue problem, Sci. Sinica Ser. A, 29 (1986), 582-94.
- [19] X.G. Geng, H.H. Dai, Nonlinearization of the Lax pairs for discrete Ablowitz-Ladik hierarchy, J. Math. Anal. Appl., 327 (2007) 829-53.
- [20] A.S. Fokas, Symmetries and integrability, Stud. Appl. Math., 77 (1987) 253-99.
- [21] P.J. Olver, Applications of Lie Groups to Differential Equations, Springer-Verlag, New York, 1993.
- [22] G.W. Bluman, S.C. Anco, Symmetry and Integration Methods for Differential Equations, Springer-Verlag, New York, 2002.
- [23] W.X. Ma, K symmetries and  $\tau$  symmetries of evolution equations and their Lie algebras, J. Phys. A: Math. Gen., 23 (1990) 2707-16.
- [24] D.Y. Chen, H.W. Zhang, Lie algebraic structure for the AKNS system, J. Phys. A: Math. Gen., 24 (1991) 377-83.
- [25] D.Y. Chen, D.J. Zhang, Lie algebraic structures of (1+1)-dimensional Lax integrable systems, J. Math. Phys., 37 (1996) 5524-38.
- [26] D.Y. Chen, H.W. Xin, D.J. Zhang, Lie algebraic structures of some (1+2)-dimensional Lax integrable systems, Chaos Solitons Fractals, 15 (2003) 761-70.

- [27] W.X. Ma, B. Fuchssteiner, Algebraic structure of discrete zero curvature equations and master symmetries of discrete evolution equations, J. Math. Phys., 40 (1999) 2400-18.
- [28] B. Fuchssteiner, A.S. Fokas, Symplectic structures, their Bäcklund transformations and hereditary symmetries, Phys. D, 4 (1981) 47-66.
- [29] C. Tian, Symmetries, in Soliton Theory and Its Applications, Ed. C.H. Gu, Springer-Verleg, Berlin, 1996.
- [30] B. Fuchssteiner, Mastersymmetries, higher order time-dependent symmetries and conserved densities of nonlinear evolution equations, Prog. Theor. Phys., 70 (1983) 1508-22.

# A The first few flows and their Lax pairs

For the isospectral four-potential AL hierarchy (3.14), when m = 1 we have

$$u_{n,t_1} = K^{(1)} = \begin{pmatrix} (1 - Q_n R_n) S_n \\ -(1 - Q_n R_n) T_{n-1} \\ (1 - S_n T_n) Q_{n+1} \\ -(1 - S_n T_n) R_n \end{pmatrix}.$$
 (A.1)

In its Lax pair

$$V_n^{(1)} = \begin{pmatrix} -\frac{1}{2}Q_n T_{n-1} + \frac{1}{2}z^2 & Q_n z \\ T_{n-1}z & \frac{1}{2}Q_n T_{n-1} - \frac{1}{2}z^2 \end{pmatrix}.$$
 (A.2)

For m = -1, we have

$$u_{n,t_{-1}} = K^{(-1)} = \begin{pmatrix} (1 - Q_n R_n) S_{n-1} \\ -(1 - Q_n R_n) T_n \\ (1 - S_n T_n) Q_n \\ -(1 - S_n T_n) R_{n+1} \end{pmatrix},$$
 (A.3)

and

$$V_n^{(-1)} = \begin{pmatrix} -\frac{1}{2}R_nS_{n-1} + \frac{1}{2}z^{-2} & -S_{n-1}z^{-1} \\ -R_nz^{-1} & \frac{1}{2}R_nS_{n-1} - \frac{1}{2}z^{-2} \end{pmatrix}.$$
 (A.4)

For the non-isospectral four-potential AL hierarchy (3.17), when m = 1 we have

$$u_{n,t_1} = \sigma^{(1)} = \begin{pmatrix} (2n + \frac{3}{2})(1 - Q_n R_n)S_n - Q_n(E - 1)^{-1}(Q_{n+1}T_n + S_n R_n) \\ -(2n - \frac{1}{2})(1 - Q_n R_n)T_{n-1} + R_n(E - 1)^{-1}(Q_{n+1}T_n + S_n R_n) \\ (2n + \frac{5}{2})(1 - S_n T_n)Q_{n+1} - S_n(E - 1)^{-1}(Q_{n+1}T_n + S_n R_n) - S_n^2 R_n \\ -(2n + \frac{1}{2})(1 - S_n T_n)R_n + T_n(E - 1)^{-1}(Q_{n+1}T_n + S_n R_n) + S_n T_n R_n \end{pmatrix}.$$
(A.5)

In its Lax pair

$$V_n^{(1)} = \begin{pmatrix} -(2n+\frac{1}{2})Q_nT_{n-1} - (E-1)^{-1}(T_{n-1}Q_n + R_nS_n) + nz^2 & (2n+\frac{1}{2})Q_nz \\ (2n-\frac{1}{2})T_{n-1}z & -nz^2 \end{pmatrix}.$$
 (A.6)

For m = -1,

$$u_{n,t_{-1}} = \sigma^{(-1)} = \begin{pmatrix} (2n - \frac{1}{2})(1 - Q_n R_n)S_{n-1} - Q_n(E - 1)^{-1}(R_{n+1}S_n + T_n Q_n) \\ -(2n + \frac{3}{2})(1 - Q_n R_n)T_n + R_n(E - 1)^{-1}(R_{n+1}S_n + T_n Q_n) \\ (2n + \frac{1}{2})(1 - S_n T_n)Q_n - S_n(E - 1)^{-1}(R_{n+1}S_n + T_n Q_n) - Q_n S_n T_n \\ -(2n + \frac{5}{2})(1 - S_n T_n)R_{n+1} + T_n(E - 1)^{-1}(R_{n+1}S_n + T_n Q_n) + T_n^2 Q_n \end{pmatrix},$$
(A.7)

and

satisfying

$$V_n^{(-1)} = \begin{pmatrix} nz^{-2} & -(2n-\frac{1}{2})S_{n-1}z^{-1} \\ -(2n+\frac{1}{2})R_nz^{-1} & (2n+\frac{1}{2})R_nS_{n-1} + (E-1)^{-1}(S_{n-1}R_n + T_nQ_n) - nz^{-2} \end{pmatrix}.$$
(A.8)

# B Theorem 1 in Ref. [11]

**Theorem B.1.** Suppose that the linear problem

$$\Phi_{n+1} = M_n \Phi_n \tag{B.1}$$

satisfies the following conditions: (1). matrix equation

$$M'_n[X] = (EN_n)M_n - M_nN_n \tag{B.2}$$

possesses a unique couple of nonzero solutions  $X(u(t,n)) \in \mathcal{V}_s$  and  $N_n = N(z, u(t,n)) \in \mathcal{Q}_m(z)$ satisfying  $[M_n, N_n]|_{u_n=0} = 0$ , (2). for any given  $Y(u(t,n)) \neq 0 \in \mathcal{V}_s$ , there exist solutions  $X(u(t,n)) \in \mathcal{V}_s$  and  $N_n \in \mathcal{Q}_m(z)$ 

$$M'_{n}[X - z^{\alpha}Y] = (EN_{n})M_{n} - M_{n}N_{n}, \quad N_{n}|_{u_{n}=0} = 0,$$
(B.3)

where  $\alpha$  is a constant related to (B.1), then the following results hold: (1). there exist the following Lax representations for isospectral flows  $\{K^{(l)}\}$ :

$$M'_{n}[K^{(l)}] = (EN_{n}^{(l)})M_{n} - M_{n}N_{n}^{(l)}, \quad l = 0, 1, \cdots,$$
(B.4)

(2). there exists a unique recursion operator L such that

$$u_{n,t} = K^{(l)} = L^l K^{(0)}, \quad l = 0, 1, \cdots,$$
 (B.5)

and L is a strong and hereditary symmetry of each member in the above hierarchy.