SYMMETRIC APPROXIMATIONS OF PSEUDO-BOOLEAN FUNCTIONS WITH APPLICATIONS TO INFLUENCE INDEXES

JEAN-LUC MARICHAL AND PIERRE MATHONET

ABSTRACT. We introduce an index for measuring the influence of the kth smallest variable on a pseudo-Boolean function. This index is defined from a weighted least squares approximation of the function by linear combinations of order statistic functions. We give explicit expressions for both the index and the approximation and discuss some properties of the index. Finally, we show that this index subsumes the concept of system signature in engineering reliability and that of cardinality index in decision making.

1. INTRODUCTION

Boolean and pseudo-Boolean functions play a central role in various areas of applied mathematics such as cooperative game theory, engineering reliability, and decision making (where fuzzy measures and fuzzy integrals are often used). In these areas indexes have been introduced to measure the importance of a variable or its influence on the function under consideration (see, e.g., [3, 7]). For instance, the concept of importance of a player in a cooperative game has been studied in various papers on *values* and *power indexes* starting from the pioneering works by Shapley [13] and Banzhaf [1]. These power indexes were rediscovered later in system reliability theory as Barlow-Proschan and Birnbaum measures of importance (see, e.g., [10]).

In general there are many possible influence/importance indexes and they are rather simple and natural. For instance, a cooperative game on a finite set $[n] = \{1, \ldots, n\}$ of *players* is a set function $v: 2^{[n]} \to \mathbb{R}$ with $v(\emptyset) = 0$, which associates with any *coalition* of players $S \subseteq [n]$ its *worth* v(S). The *Banzhaf value* of player $i \in [n]$ in the game v is then defined as

(1)
$$\phi_B(v,i) = \frac{1}{2^{n-1}} \sum_{S \subseteq [n] \setminus \{i\}} \left(v(S \cup \{i\}) - v(S) \right) = \frac{1}{2^{n-1}} \sum_{S \neq i} v(S) - \frac{1}{2^{n-1}} \sum_{S \neq i} v(S).$$

Thus, $\phi_B(v, i)$ is the average of the marginal contributions of player *i* to all coalitions $S \subseteq [n] \setminus \{i\}$, or the difference between the average worth over all coalitions $S \ni i$ and the average worth over all coalitions $S \not\equiv i$. Considering weighted averages instead of symmetric averages gives rise to various *probabilistic values* (see [14]), including the Shapley value and weighted Banzhaf values (see [8]).

The choice of a suitable influence/importance index depends on the practical problem under consideration and is usually made by considering the properties

Date: February 27, 2012.

²⁰¹⁰ Mathematics Subject Classification. Primary 41A10, 93E24; Secondary 62G30, 90B25, 91A12.

Key words and phrases. Pseudo-Boolean function, least squares approximation, symmetric function, cooperative game theory, system reliability, system signature, cardinality index.

that the index should satisfy. This is why many indexes have been characterized axiomatically. Besides these characterizations, it is well known in statistics that one can measure the influence of a variable using linear regression. This approach was applied successfully to pseudo-Boolean functions by Hammer and Holzman [5], who showed that the Banzhaf value appears as the coefficients of the linear terms of the standard least squares approximation of a game (or its corresponding pseudo-Boolean function) by a function of degree at most 1. Weighted versions of this least squares approach were also considered to characterize the Shapley value [4] and weighted Banzhaf values [8].

Slightly different influence indexes emerged in certain applications where it is not the influence of a variable on a function that is to be measured but rather the influence of adding a variable to a given subset of variables. For instance, considering a system made up of n interconnected components with independent and identically distributed (i.i.d.) lifetimes, Samaniego [11, 12] defined the *signature* of the system as the n-tuple (s_1, \ldots, s_n) where s_k is the probability that the kth failure causes the system to fail. Due to the i.i.d. assumption, the signature only depends on the (Boolean) structure function of the system. Thus, the number s_k can be interpreted as a measure of the influence on the structure function of adding a kth element to the set of failed components. Another example of such a measure of influence was introduced by Yager [15] in the context of fuzzy measures. Considering a fuzzy measure on an n-set X, he introduced the *cardinality index* as the n-tuple (C_0, \ldots, C_{n-1}) , where C_k is the average gain in certitude one gets when going from a k-subset to a (k + 1)-subset.

In this paper we show that the concepts of system signature and cardinality index are special instances of a more general notion of influence index: the *influence index* of the kth smallest variable on a pseudo-Boolean function. We define this index by considering the least squares approximation of a given function by a linear combination of order statistic functions. Such linear combinations are particularly suitable for encoding the influence that we want to measure and are exactly the symmetric (i.e., invariant under a permutation of the variables) pseudo-Boolean functions (see Proposition 1). Here we consider the general framework of *arbitrarily weighted* least squares approximations. In Section 2 we give explicit expressions for the approximation and discuss some of its properties. In Sections 3 and 4 we introduce our influence index and show how it subsumes the concepts of system signature and cardinality index. We also show how this index can be used in cooperative game theory to define a new influence index.

We employ the following notation throughout the paper. We denote by \mathbb{B} the two-element set $\{0,1\}$. For any $\mathbf{x} \in \mathbb{B}^n$, we set $|\mathbf{x}| = \sum_{i=1}^n x_i$. For any $S \subseteq [n] = \{1, \ldots, n\}$, we denote by $\mathbf{1}_S$ the *n*-tuple whose *i*th coordinate is 1, if $i \in S$, and 0, otherwise (with the particular cases $\mathbf{0} = \mathbf{1}_{\emptyset}$ and $\mathbf{1} = \mathbf{1}_{[n]}$).

Through the usual identification of the elements of \mathbb{B}^n with the subsets of [n], a pseudo-Boolean function $f:\mathbb{B}^n \to \mathbb{R}$ can be equivalently described by a set function $v_f:2^{[n]} \to \mathbb{R}$. We simply write $v_f(S) = f(\mathbf{1}_S)$. To avoid cumbersome notation, we henceforth use the same symbol to denote both a given pseudo-Boolean function and its underlying set function, thus writing $f:\mathbb{B}^n \to \mathbb{R}$ or $f:2^{[n]} \to \mathbb{R}$ interchangeably.

Recall that if the B-valued variables x_1, \ldots, x_n are rearranged in ascending order of magnitude $x_{(1)} \leq \cdots \leq x_{(n)}$, then $x_{(k)}$ is called the *kth order statistic* and the function $os_k: \mathbb{B}^n \to \mathbb{B}$, defined as $os_k(\mathbf{x}) = x_{(k)}$, is the *kth order statistic function*. We then have $os_k(\mathbf{x}) = 1$, if $\sum_{i=1}^n x_i \ge n - k + 1$, and 0, otherwise. As a matter of convenience, we also formally define $os_0 \equiv 0$ and $os_{n+1} \equiv 1$. An *L*-statistic function is a linear combination of the functions os_1, \ldots, os_n while a shifted *L*-statistic function is a linear combination of the functions os_1, \ldots, os_{n+1} .

2. Symmetric approximations

In this section we present and solve the problem of approximation of pseudo-Boolean functions by shifted L-statistic functions and discuss a few properties of the approximations.

Recall that any *n*-ary pseudo-Boolean function f can always be represented by a multilinear polynomial of degree at most n (see [6]). More precisely, f can always be written in the form

(2)
$$f(\mathbf{x}) = \sum_{S \subseteq [n]} f(S) \prod_{i \in S} x_i \prod_{i \in [n] \setminus S} (1 - x_i).$$

By expanding the second product, we see that this polynomial can be further simplified into

$$f(\mathbf{x}) = \sum_{S \subseteq [n]} a_f(S) \prod_{i \in S} x_i,$$

where the set function $a_f: 2^{[n]} \to \mathbb{R}$, called the *Möbius transform* of f, is defined by

$$a_f(S) = \sum_{T \subseteq S} (-1)^{|S| - |T|} f(T)$$

Denote by $F(\mathbb{B}^n)$ the vector space of *n*-ary pseudo-Boolean functions and by $F_S(\mathbb{B}^n)$ the subspace of symmetric *n*-ary pseudo-Boolean functions. It is clear that a function $f \in F(\mathbb{B}^n)$ is symmetric if and only if it is *cardinality-based*, i.e., it satisfies the property f(S) = f(T) for every $S, T \subseteq [n]$ such that |S| = |T|. Equivalently, there exists a unique function $\overline{f}: \{0, 1, \ldots, n\} \to \mathbb{R}$ such that $f(S) = \overline{f}(|S|)$.

The following proposition shows that the shifted *L*-statistic functions are precisely those pseudo-Boolean functions that are symmetric.

Proposition 1. A pseudo-Boolean function is symmetric if and only if it is a shifted L-statistic function.

Proof. The class of *n*-ary shifted *L*-statistic functions is clearly a subspace of $F_S(\mathbb{B}^n)$. Since each of these spaces has dimension n + 1, they must coincide. \Box

Given a weight function $w: \mathbb{B}^n \to]0, \infty[$ and a function $f \in F(\mathbb{B}^n)$, we define the best symmetric approximation of f with respect to w as the unique function $f_L \in F_S(\mathbb{B}^n)$ that minimizes the weighted squared distance

$$||f-g||^2 = \sum_{\mathbf{x}\in\mathbb{B}^n} w(\mathbf{x})(f(\mathbf{x})-g(\mathbf{x}))^2$$

among all symmetric functions $g \in F_S(\mathbb{B}^n)$. Since $\|\cdot\|$ is the norm associated with the inner product

$$\langle f,g\rangle = \sum_{\mathbf{x}\in\mathbb{B}^n} w(\mathbf{x})f(\mathbf{x})g(\mathbf{x}),$$

the solution f_L of this approximation problem exists and is uniquely determined by the orthogonal projection of f onto $F_S(\mathbb{B}^n)$. We then write $f_L = A(f)$.

We will henceforth assume (without loss of generality) that the weights are multiplicatively normalized so that $\sum_{\mathbf{x} \in \mathbb{B}^n} w(\mathbf{x}) = 1$. Although this assumption is not necessary for most of the results, it will enable us to interpret w as a probability distribution and make use of certain concepts in probability theory.

Definition 2. For every $f \in F(\mathbb{B}^n)$, define $\overline{f}: \{0, 1, \dots, n\} \to \mathbb{R}$ as

(3)
$$\overline{f}(s) = E(f(\mathbf{x}) | |\mathbf{x}| = s) = \frac{\sum_{|\mathbf{x}|=s} w(\mathbf{x}) f(\mathbf{x})}{\sum_{|\mathbf{x}|=s} w(\mathbf{x})} .^{1}$$

We also formally define $\overline{f}(-1) = 0$.

The next theorem gives an explicit expression for A(f).

Theorem 3. The best symmetric approximation of $f \in F(\mathbb{B}^n)$ is given by

(4)
$$A(f) = \sum_{j=1}^{n+1} c_j \operatorname{os}_j$$

where $c_j = \overline{f}(n-j+1) - \overline{f}(n-j)$ for every $j \in [n+1]$.

Proof. Since $F_S(\mathbb{B}^n)$ is spanned by the n + 1 functions $os_1, \ldots, os_n, os_{n+1}$, the projection A(f) is characterized by the conditions

(5)
$$\langle f - A(f), \operatorname{os}_i \rangle = 0$$
 $(i \in [n+1]),$

that is,

(6)
$$\sum_{|\mathbf{x}| \ge n-i+1} w(\mathbf{x}) (f(\mathbf{x}) - A(f)(\mathbf{x})) = 0 \qquad (i \in [n+1]).$$

We observe that the system (6) remains equivalent if we replace the inequality $|\mathbf{x}| \ge n - i + 1$ with the equality. Using (4), we then obtain

(7)
$$\left(\sum_{j=i}^{n+1} c_j\right) \left(\sum_{|\mathbf{x}|=n-i+1} w(\mathbf{x})\right) = \sum_{|\mathbf{x}|=n-i+1} w(\mathbf{x}) f(\mathbf{x}) \qquad (i \in [n+1])$$

We finally obtain the result by using (3) and subtracting equation i+1 from equation i.

We now provide alternative expressions for A(f) as a shifted L-statistic function and symmetric multilinear polynomials. Observing first that $\overline{f}(0) = f(0)$ and then using (4) and (5) for i = n + 1, we obtain

(8)
$$A(f) = f(\mathbf{0}) + \sum_{j=1}^{n} c_j \operatorname{os}_j = \langle f, 1 \rangle + \sum_{j=1}^{n} c_j (\operatorname{os}_j - \langle \operatorname{os}_j, 1 \rangle),$$

where $c_j = \overline{f}(n-j+1) - \overline{f}(n-j)$ for every $j \in [n]$. Then, using (3), (4), and (7), we obtain

(9)
$$A(f)(S) = \sum_{j=n-|S|+1}^{n+1} c_j = E(f(\mathbf{x}) | |\mathbf{x}| = |S|) = \overline{f}(|S|) \quad (S \subseteq [n])$$

so that by (2) we obtain immediately

$$A(f)(\mathbf{x}) = \sum_{S \subseteq [n]} \overline{f}(|S|) \prod_{i \in S} x_i \prod_{i \in [n] \setminus S} (1 - x_i) = \sum_{S \subseteq [n]} \Delta_k^{|S|} \overline{f}(k)|_{k=0} \prod_{i \in S} x_i,$$

¹Although \overline{f} depends explicitly on w, we use this notation for it is consistent with that introduced for cardinality-based set functions. Note also that, in the special case when the weight function w is symmetric, $\overline{f}(s)$ clearly reduces to $\binom{n}{s}^{-1} \sum_{|\mathbf{x}|=s} f(\mathbf{x})$.

where $\Delta_k^s \overline{f}(k)|_{k=0} = \sum_{t=0}^s {s \choose t} (-1)^{s-t} \overline{f}(t).$

We now examine the effect of a permutation of the variables of f on the symmetric approximation A(f). Let S_n denote the symmetric group on [n]. A permutation $\pi \in S_n$ acts on a pseudo-Boolean function $f \in F(\mathbb{B}^n)$ by $\pi(f)(x_1, \ldots, x_n) = f(x_{\pi(1)}, \ldots, x_{\pi(n)})$. A permutation $\pi \in S_n$ is said to be a symmetry of $f \in F(\mathbb{B}^n)$ if $\pi(f) = f$.

Proposition 4. If $\pi \in S_n$ is a symmetry of the weight function w, then for every $f \in F(\mathbb{B}^n)$ we have $A(\pi(f)) = A(f)$ and $\|\pi(f) - A(f)\| = \|f - A(f)\|$.

Proof. If π is a symmetry of w, then clearly it is an isometry of $F(\mathbb{B}^n)$, that is, $\langle \pi(f), \pi(g) \rangle = \langle f, g \rangle$. Now, if $g \in F_S(\mathbb{B}^n)$, then by (5), we have $\langle \pi(f), g \rangle = \langle \pi(f), \pi(g) \rangle = \langle f, g \rangle = \langle A(f), g \rangle$, which shows that $A(\pi(f)) = A(f)$. We prove the second equality similarly since $\|\pi(f) - A(f)\|^2 = \|\pi(f) - \pi(A(f))\|^2$. \Box

With any pseudo-Boolean function $f \in F(\mathbb{B}^n)$, we can associate the symmetric function $\operatorname{Sym}(f) = \frac{1}{n!} \sum_{\pi \in S_n} \pi(f)$. We then have the following result.

Corollary 5. If the weight function w is symmetric, then for every $f \in F(\mathbb{B}^n)$ we have Sym(f) = A(Sym(f)) = A(f).

Proof. The first equality follows from the symmetry of Sym(f). The second one follows from Proposition 4 and the linearity of the projector A.

We end this section by analyzing the effect of dualization of f on the symmetric approximation A(f). The dual of a function $f \in F(\mathbb{B}^n)$ is the function $f^d \in F(\mathbb{B}^n)$ defined by $f^d(\mathbf{x}) = f(\mathbf{0}) + f(\mathbf{1}) - f(\mathbf{1}_{[n]} - \mathbf{x})$.

Proposition 6. If the weight function w satisfies $w(\mathbf{1}_{[n]} - \mathbf{x}) = w(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{B}^n$, then for every $f \in F(\mathbb{B}^n)$ we have $A(f^d) = A(f)^d$.

Proof. By (5) and (9), we have $A(f)(1)-f(1) = A(f)(0)-f(0) = \langle A(f), 1 \rangle - \langle f, 1 \rangle = 0$. From these equalities, we obtain

$$\langle f^d - A(f)^d, g^d \rangle = \langle (f - A(f))^d, g^d \rangle = \langle f - A(f), g \rangle = 0$$

for every function $g \in F_S(\mathbb{B}^n)$. The result then follows.

3. Influence of the kth smallest variable

Following Hammer and Holzman's approach [5], to measure the influence of the kth smallest variable $x_{(k)}$ on a pseudo-Boolean function $f \in F(\mathbb{B}^n)$, it is natural to define an index $I: F(\mathbb{B}^n) \times [n] \to \mathbb{R}$ as $I(f,k) = c_k$, where c_k is defined in Theorem 3.²

Definition 7. Let $I: F(\mathbb{B}^n) \times [n] \to \mathbb{R}$ be defined as $I(f,k) = \overline{f(n-k+1)} - \overline{f(n-k)}$.

Thus we have defined an influence index from an elementary approximation (projection) problem. Conversely, the following result shows that A(f) is the unique function of $F_S(\mathbb{B}^n)$ that preserves the average value and the influence index of f.

Proposition 8. A function $g \in F_S(\mathbb{B}^n)$ is the best symmetric approximation of $f \in F(\mathbb{B}^n)$ if and only if $\langle f, 1 \rangle = \langle g, 1 \rangle$ and I(f,k) = I(g,k) for all $k \in [n]$.

²We observe that, by definition, this index remains invariant under normalization of w.

Proof. The necessity is trivial (use Eq. (5) for i = n + 1). To prove the sufficiency, observe that any $q \in F_S(\mathbb{B}^n)$ satisfying the assumptions of the proposition is of the form

$$g = g(\mathbf{0}) + \sum_{j=1}^{n} I(g, j) \operatorname{os}_{j} = g(\mathbf{0}) + \sum_{j=1}^{n} I(f, j) \operatorname{os}_{j}$$

We then have $g(\mathbf{0}) + \sum_{j=1}^{n} I(f, j) \langle os_j, 1 \rangle = \langle g, 1 \rangle = \langle f, 1 \rangle$. Using (8), we finally obtain g = A(f).

The next proposition reassembles several properties of the index I(f,k). These properties follow easily from the definition of the index and the properties of the approximations.

Proposition 9. Let $k \in [n]$ and let $w: \mathbb{B}^n \to [0, \infty]$ be a weight function. Then

- (i) The map $f \mapsto I(f,k)$ is linear.
- (ii) If π is a symmetry of w, then $I(\pi(f), k) = I(f, k)$ for every $f \in F(\mathbb{B}^n)$.
- (iii) If w is symmetric, then I(f,k) = I(Sym(f),k) for every $f \in F(\mathbb{B}^n)$.
- (iv) If w satisfies $w(\mathbf{1}_{[n]} \mathbf{x}) = w(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{B}^n$, then $I(f^d, k) = I(f, n-k+1)$ for every $f \in F(\mathbb{B}^n)$. (v) We have $\sum_{j=1}^n I(f,j) = f(1) - f(0)$.

It is a well-known fact of linear algebra that a linear map on a finite dimensional inner product space can be expressed as an inner product with a fixed vector. The next proposition gives the explicit form of such a vector for $I(\cdot, k)$. To this extent, for every $k \in [n]$ we introduce the function $g_k: \mathbb{B}^n \to \mathbb{R}$ as $g_k(\mathbf{x}) = \Delta_k(d_k \Delta_k os_{k-1})$, where $d_k = -1 / \sum_{|x|=n-k+1} w(\mathbf{x})$.

Proposition 10. For every $f \in F(\mathbb{B}^n)$ and every $k \in [n]$, we have $I(f,k) = \langle f, g_k \rangle$.

Proof. We have $d_{k+1}(f, \Delta_k os_k) = d_{k+1} \sum_{|\mathbf{x}|=n-k} w(\mathbf{x}) f(\mathbf{x}) = -\overline{f}(n-k)$, which leads immediately to the result.

Proposition 10 shows that the index I(f,k) is the covariance of the random variables f and g_k . Indeed, we have $I(f,k) = E(fg_k) = cov(f,g_k) + E(f)E(g_k)$, where $E(g_k) = \langle 1, g_k \rangle = I(1, k) = 0$. From the usual interpretation of the concept of covariance, we see that an element $\mathbf{x} \in \mathbb{B}^n$ makes a positive contribution to I(f,k) whenever the values of $f(\mathbf{x}) - E(f)$ and $g_k(\mathbf{x}) - E(g_k) = g_k(\mathbf{x})$ have the same sign. Note that $g_k(\mathbf{x})$ is positive whenever $x_{(k)}$ is greater than the value $(d_{k+1}x_{(k+1)} + d_kx_{(k-1)})/(d_{k+1} + d_k)$, which lies in the range of $x_{(k)}$ when the other order statistics are fixed at **x**.

4. Two special cases: Cardinality index and system signature

We now show that the *cardinality index* and *system signature* are particular instances of our influence index.

4.1. The cardinality index of a fuzzy measure. A fuzzy measure on the finite set X = [n] is a nondecreasing set function $\mu: 2^X \to [0,1]$ satisfying the boundary conditions $\mu(\emptyset) = 0$ and $\mu(X) = 1$. For any subset $S \subseteq X$, the number $\mu(S)$ can be interpreted as the certitude that we have that a variable will take on its value in the set $S \subseteq X$. In this context, Yager [15] introduced the *cardinality index* associated with a fuzzy measure μ as the *n*-tuple (C_0, \ldots, C_{n-1}) where C_k is the average gain in certitude that we obtain by adding an arbitrary element to an arbitrary k-subset, that is,

$$C_{k} = \frac{1}{(n-k)\binom{n}{k}} \sum_{|S|=k} \sum_{x \notin S} (\mu(S \cup \{x\}) - \mu(S)).$$

We observe that this expression, which resembles the Banzhaf value (1), could be used in cooperative game theory to measure the marginal contribution of an additional player to a k-coalition. It is also clear that this index can be written as

$$C_{k} = \frac{1}{\binom{n}{k+1}} \sum_{|S|=k+1} \mu(S) - \frac{1}{\binom{n}{k}} \sum_{|S|=k} \mu(S),$$

which shows that we have $C_k = I(\mu, n - k) = I(\mu^d, k + 1)$ in the special case when the weight function w defining the index I is symmetric.

4.2. System signatures in engineering reliability. Consider a system consisting of n interconnected components. When the components have continuous and i.i.d. lifetimes X_1, \ldots, X_n , the signature of the system is defined as the n-tuple $(s_1, \ldots, s_n) \in [0, 1]^n$ with $s_k = \Pr(T = X_{(k)})$, where T denotes the system lifetime. That is, s_k is the probability that the kth failure causes the system to fail (for a recent reference, see [12]). It was proved [2] that

$$s_{k} = \frac{1}{\binom{n}{n-k+1}} \sum_{|\mathbf{x}|=n-k+1} \phi(\mathbf{x}) - \frac{1}{\binom{n}{n-k}} \sum_{|\mathbf{x}|=n-k} \phi(\mathbf{x}),$$

where $\phi: \mathbb{B}^n \to \mathbb{B}$ is the structure function of the system. Thus, in view of Definitions 2 and 7, we have $s_k = I(\phi, k)$ in the special case where the weight function wis symmetric. Interestingly, the identity $s_k = I(\phi, k)$ still holds in the non-i.i.d. case if we define the weight function w as the (non-normalized) relative quality function

$$w(S) = \Pr\left(\max_{i \in [n] \setminus S} X_i < \min_{j \in S} X_j\right)$$

for which we have $\sum_{|\mathbf{x}|=s} w(\mathbf{x}) = 1$ for all $s \in [n]$ (see [9]). Therefore s_k can be obtained from a weighted least squares approximation problem of the structure function and can always be interpreted as the influence on the system of the component that has the kth smallest lifetime.

Acknowledgments

This research was supported by the internal research project F1R-MTH-PUL-09MRDO of the University of Luxembourg.

References

- J. Banzhaf. Weighted voting doesn't work : A mathematical analysis. *Rutgers Law Review*, 19:317–343, 1965.
- [2] Ph. J. Boland. Signatures of indirect majority systems. J. Appl. Prob., 38:597-603, 2001.
- [3] J. Bourgain, J. Kahn, G. Kalai, Y. Katznelson, and N. Linial. The influence of variables in product spaces. Isr. J. Math., 77(1-2):5564, 1992.
- [4] A. Charnes, B. Golany, M. Keane, and J. Rousseau. Extremal principle solutions of games in characteristic function form: core, Chebychev and Shapley value generalizations. In *Econometrics of planning and efficiency*, volume 11 of *Adv. Stud. Theoret. Appl. Econometrics*, pages 123–133. Kluwer Acad. Publ., Dordrecht, 1988.
- [5] P. Hammer and R. Holzman. Approximations of pseudo-Boolean functions; applications to game theory. Z. Oper. Res., 36(1):3–21, 1992.

- [6] P. Hammer and S. Rudeanu. Boolean methods in operations research and related areas. Berlin-Heidelberg-New York: Springer-Verlag, 1968.
- [7] J.-L. Marichal. The influence of variables on pseudo-Boolean functions with applications to game theory and multicriteria decision making. *Discrete Appl. Math.*, 107(1-3):139–164, 2000.
- [8] J.-L. Marichal and P. Mathonet. Weighted Banzhaf power and interaction indexes through weighted approximations of games. Eur. J. of Operational Research, 211(2):352–358, 2011.
- [9] J.-L. Marichal and P. Mathonet. Extensions of system signatures to dependent lifetimes: Explicit expressions and interpretations. J. of Multivariate Analysis, 102(5):931–936, 2011.
- [10] K.G. Ramamurthy. *Coherent structures and simple games*. Dordrecht-Boston-London: Kluwer Academic Publisher, 1990.
- [11] F.J. Samaniego. On closure of the IFR class under formation of coherent systems. *IEEE Trans. Reliability*, 34:69–72, 1985.
- [12] F.J. Samaniego. System signatures and their applications in engineering reliability. Int. Series in Operations Research & Management Science, 110. New York: Springer, 2007.
- [13] L. Shapley. A value for n-person games. In Contributions to the Theory of Games II (Annals of Mathematics Studies 28), pages 307–317. Princeton University Press, 1953.
- [14] R. J. Weber. Probabilistic values for games. In *The Shapley value*, pages 101–119. Cambridge Univ. Press, Cambridge, 1988.
- [15] R. Yager. On the cardinality index and attitudinal character of fuzzy measures Int. J. Gen. Syst., 31(3):303–329, 2002.

Mathematics Research Unit, FSTC, University of Luxembourg, 6, rue Coudenhove-Kalergi, L-1359 Luxembourg, Grand Duchy of Luxembourg.

E-mail address: jean-luc.marichal[at]uni.lu

MATHEMATICS RESEARCH UNIT, FSTC, UNIVERSITY OF LUXEMBOURG, 6, RUE COUDENHOVE-KALERGI, L-1359 LUXEMBOURG, GRAND DUCHY OF LUXEMBOURG. *E-mail address*: p.mathonet[at]ulg.ac.be