

Transversality Conditions for Higher Order Infinite Horizon Discrete Time Optimization Problems

Dapeng CAI ¹ and Takashi Gyoshin NITTA ²

¹*Corresponding author. Institute for Advanced Research, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8601, Japan;*
²*Department of Mathematics, Faculty of Education, Mie University, Kurimamachiya 1577, Tsu, 514-8507, Japan*

Abstract

In this paper, we examine higher order difference problems:
 $\max_{\mathbf{c}} \sum_{t=0}^{\infty} U(\mathbf{c}(t), \mathbf{c}(t+1), \dots, \mathbf{c}(t+N-1), t)$. Using the “squeezing” argument, we derive both Euler’s condition and the transversality condition. In order to derive the two conditions, two needed assumptions are identified. A counterexample, in which the transversality condition is not satisfied without the two assumptions, is also presented.

Keywords: Keywords: Transversality condition; Dynamic optimization; Infinite horizon; Higher order difference problems

1. INTRODUCTION

In this paper, we consider the following reduced form model

$$(1) \quad \begin{cases} \max_{\mathbf{c}} \sum_{t=0}^{\infty} U(\mathbf{c}(t), \mathbf{c}(t+1), \dots, \mathbf{c}(t+N-1), t) \\ \text{subject to } \mathbf{c}(0) = \mathbf{c}_0, \\ \forall t \geq 0, (\mathbf{c}(t), \mathbf{c}(t+1), \dots, \mathbf{c}(t+N-1)) \in X(t) \subset (\mathbb{R}^n)^N, \end{cases}$$

where $N \in \mathbb{N}$, U is a real-valued N th-order continuously differentiable function, and $\mathbf{c} \equiv (c_1, c_2, \dots, c_n)$ is N th-order continuously differentiable.¹ Notice that the objective functional of (1) can be infinite. [7] considers the continuous time first order differential problems: $v(\mathbf{x}(t), \dot{\mathbf{x}}(t), t)$. It generalizes the results of [5, 6, 10, 12]. So far, the most general form of the transversality conditions for continuous time version of problem (1) is presented in [11], which extends the first order case considered in [7](Theorem 3.2) to higher order cases. [7] was later extended to the discrete time stochastic case by [8]. In this paper, we aim to extend these results to deterministic higher order difference problems, using the “squeezing” argument.

The application of higher order difference problems can be widely found in economics. In particular, they appear in the discussion concerning the overlapping generations models. A satisfactory examination of the individuals’ marriage and fertility decisions would necessitate the division of the representative agent’s lifetime to multiple periods, instead of only two periods, young and old. However, as argued in [4], the properties of a model with two-period-lived agents cannot be readily extended to n -period-lived agents. To consider the n -period-lived agents case, transversality conditions for higher order difference problems would be imperative.

We first use the “squeezing” argument to derive both Euler’s condition and the transversality condition for higher order difference problems, showing the argument needs two imperative assumptions. These two assumptions constitute the discrete time version of Assumption 1 and 2 in [11]. We then provide a counterexample, in which the transversality condition is not satisfied without the two assumptions. Because Assumption 1 and 2 are satisfied when a discounting factor is incorporated into the model, our transversality conditions also generalize the results obtained in the presence of discounting. For approaches on

¹ Normally, U is defined on $(\mathbb{R}^n)^N \times \mathbb{R}$. The domain of U is denoted by $X(t)$, in included in $(\mathbb{R}^n)^N$, for all t .

how to explicitly construct the optimal solutions to the undiscounted infinite horizon optimization problems, see [2, 3].

2. DERIVATION OF THE TRANSVERSALITY CONDITIONS

Suppose that the optimal path to (1) exists and is given by $\mathbf{c}^*(t)$, optimal in the sense of an overtaking criterion to be defined below. We perturb it with N th-order continuously differentiable curves $\mathbf{q}(t)$,

$$(2) \quad \mathbf{c}(t) = \mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t).$$

We define

$$(3) \quad \begin{aligned} V(\varepsilon, T) = \inf_{T \leq T'} \sum_{t=0}^{T'} [& U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \mathbf{c}^*(t+1) + \varepsilon \cdot \mathbf{q}(t+1), \\ & \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ & - U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)]. \end{aligned}$$

In this paper, [1]'s notion of weak maximality is used as our optimality criterion. We assume that there exists an optimal path that satisfy the weak maximality criterion, which is defined as: an attainable path ($\mathbf{c}^*(t)$) is optimal if no other attainable path overtakes it²:

$$(4) \quad \begin{aligned} \lim_{T \rightarrow +\infty} \inf_{T \leq T'} \sum_{t=0}^{T'} [& U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \mathbf{c}^*(t+1) + \varepsilon \cdot \mathbf{q}(t+1), \\ & \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ & - U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)] \leq 0. \end{aligned}$$

Let $V(\varepsilon) = \lim_{T \rightarrow \infty} V(\varepsilon, T)$. Differentiating it with respect to ε , we have

$$(5) \quad \begin{aligned} & \lim_{\varepsilon \rightarrow +0} \frac{V(\varepsilon)}{\varepsilon} \\ & = \lim_{\varepsilon \rightarrow +0} \lim_{T \rightarrow \infty} \inf_{T \leq T'} \sum_{t=0}^{T'} \frac{1}{\varepsilon} [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \mathbf{c}^*(t+1) + \varepsilon \cdot \mathbf{q}(t+1), \\ & \quad \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ & \quad - U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)]. \end{aligned}$$

Let $\lim_{\varepsilon \rightarrow +0} \frac{V(\varepsilon)}{\varepsilon} \equiv \Omega$. Generally, $\frac{d}{d\varepsilon} \lim_{T \rightarrow \infty} f(\varepsilon, T) = \lim_{T \rightarrow \infty} \frac{d}{d\varepsilon} f(\varepsilon, T)$ only when $\lim_{T \rightarrow \infty} \frac{d}{d\varepsilon} f(\varepsilon, T)$ converges uniformly for ε ([9]). We assume

² [1] shows that such a path exists once two assumptions are satisfied.

Assumption 1. Assume Ω converges uniformly for ε when $T \rightarrow \infty$

Assume Assumption 1, we can then restate (5) as

$$\begin{aligned} \Omega &= \lim_{T \rightarrow \infty} \lim_{\varepsilon \rightarrow +0} \inf_{T \leq T'} \frac{1}{\varepsilon} \\ &\quad \times \sum_{t=0}^{T'} [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ &\quad - U(\mathbf{c}^*(t), \dots, \mathbf{c}^*(t+N-1), t)]. \end{aligned} \quad (6)$$

We also assume

Assumption 2. We assume for any $T > 0$,

$$\begin{aligned} \inf_{T \leq T'} \sum_{t=0}^{T'} \frac{1}{\varepsilon} [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ - U(\mathbf{c}^*(t), \dots, \mathbf{c}^*(t+N-1), t)] \end{aligned}$$

converges uniformly for ε .

As in [11], a precise interpretation of Assumption 2 can be given as follows: Let

$$\begin{aligned} A(T, \varepsilon) &= \sum_{t=0}^{T'} \frac{1}{\varepsilon} [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ &\quad - U(\mathbf{c}^*(t), \dots, \mathbf{c}^*(t+N-1), t)]. \end{aligned}$$

Then there exists a sequence $A(T'_n, \varepsilon)$ for each $\varepsilon > 0$, so that $\lim_{n \rightarrow \infty} A(T'_n, \varepsilon) = \inf_{T \leq T'} A(T', \varepsilon)$, uniformly for ε , that is, the sequence is uniformly convergence for ε .

Assumptions 1 and 2 extend Assumption 3.1 in [7]. When Assumptions 1 and 2 are satisfied, then $\lim_{\varepsilon \rightarrow +0}$ and $\inf_{T \leq T'}$ can be interchanged, and equality (6) can then restated as

$$\begin{aligned} \Omega &= \lim_{T \rightarrow \infty} \inf_{T \leq T'} \lim_{\varepsilon \rightarrow +0} \frac{1}{\varepsilon} \\ &\quad \times \sum_{t=0}^{T'} [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ &\quad - U(\mathbf{c}^*(t), \dots, \mathbf{c}^*(t+N-1), t)]. \end{aligned} \quad (7)$$

Because T' is finite uniformly for ε , if

$$\sum_{t=0}^{T'} \frac{1}{\varepsilon} [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) - U(\mathbf{c}^*(t), \dots, \mathbf{c}^*(t+N-1)), t]$$

exists, (7) is then rewritten as

$$\begin{aligned} \Omega &= \lim_{T \rightarrow \infty} \inf_{T \leq T'} \sum_{t=0}^{T'} \frac{1}{\varepsilon} \\ &\quad \times [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ &\quad - U(\mathbf{c}^*(t), \dots, \mathbf{c}^*(t+N-1), t)]. \end{aligned} \tag{8}$$

From the differentiability of U , we have

$$\begin{aligned} &\lim_{\varepsilon \rightarrow +0} \frac{1}{\varepsilon} [U(\mathbf{c}^*(t) + \varepsilon \cdot \mathbf{q}(t), \dots, \mathbf{c}^*(t+N-1) + \varepsilon \cdot \mathbf{q}(t+N-1), t) \\ &\quad - U(\mathbf{c}^*(t), \dots, \mathbf{c}^*(t+N-1), t)] \\ &= \sum_{i=1}^n \left[\frac{\partial U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)}{\partial c_i(t)} q_i(t) \right. \\ &\quad + \frac{\partial U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)}{\partial c_i(t+1)} q_i(t+1) \\ &\quad \left. + \dots + \frac{\partial U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)}{\partial c_i(t+N-1)} q_i(t+N-1) \right]. \end{aligned}$$

Hence,

$$\begin{aligned} \Omega &= \lim_{T \rightarrow \infty} \inf_{T \leq T'} \sum_{t=0}^{T'} \sum_{i=1}^n \left[\frac{\partial U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)}{\partial c_i(t)} q_i(t) \right. \\ &\quad + \frac{\partial U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)}{\partial c_i(t+1)} q_i(t+1) + \dots \\ &\quad \left. + \frac{\partial U(\mathbf{c}^*(t), \mathbf{c}^*(t+1), \dots, \mathbf{c}^*(t+N-1), t)}{\partial c_i(t+N-1)} q_i(t+N-1) \right]. \end{aligned}$$

We derive

$$\begin{aligned}
& \sum_{t=0}^{T'} \left(\sum_{i=1}^n \left(\frac{\partial U(t)}{\partial c_i(t)} q_i(t) + \cdots + \frac{\partial U(t)}{\partial c_i(t+N-1)} q_i(t+N-1) \right) \right) \\
&= \sum_{i=1}^n \left\{ \frac{\partial U(0)}{\partial c_i(0)} q_i(0) + \frac{\partial (U(0) + U(1))}{\partial c_i(1)} q_i(1) + \cdots \right. \\
&\quad + \frac{\partial (U(0) + \cdots + U(N-2))}{\partial c_i(t)} q_i(N-2) \\
&\quad + \sum_{t=N-1}^{T'} \frac{\partial (U(t-N+1) + \cdots + U(t))}{\partial c_i(t)} q_i(t) \\
&\quad + \left. \left(\frac{\partial (U(T'-N+2) + \cdots + U(T'))}{\partial c_i(T'+1)} \right) q_i(T'+1) + \cdots \right. \\
(9) \quad & \left. + \frac{\partial U(T')}{\partial c_i(T'+N-1)} q_i(T'+N-1) \right\}.
\end{aligned}$$

Hence, Euler's condition is

$$\begin{aligned}
& \frac{\partial U(0)}{\partial c_i(0)} = 0, \\
& \frac{\partial (U(0) + U(1))}{\partial c_i(1)} = 0, \\
& \dots\dots, \\
& \frac{\partial (U(0) + \cdots + U(N-2))}{\partial c_i(t)} = 0, \\
(10) \quad & \frac{\partial (U(t-N+1) + \cdots + U(t))}{\partial c_i(t)} = 0, \text{ for } N-1 \leq t \leq T',
\end{aligned}$$

which extends the standard Euler's condition, and the transversality condition is given by

$$\begin{aligned}
& \lim_{T' \rightarrow \infty} \inf_{T \leq T'} \sum_{i=1}^n \left[\frac{\partial (U(T'-N+2) + \cdots + U(T'))}{\partial c_i(T'+1)} q_i(T'+1) \right. \\
(11) \quad & \left. + \cdots + \frac{\partial (U(T'))}{\partial c_i(T'+N-1)} q_i(T'+N-1) \right] \leq 0.
\end{aligned}$$

Note that when $\varepsilon \rightarrow^- 0$, the argument is the same:

$$(11') \quad \lim_{T \rightarrow \infty} \sup_{T \leq T'} \sum_{i=1}^n \left[\frac{\partial (U(T' - N + 2) + \cdots + U(T'))}{\partial c_i(T' + 1)} q_i(T' + 1) \right. \\ \left. + \cdots + \frac{\partial (U(T'))}{\partial c_i(T' + N - 1)} q_i(T' + N - 1) \right] \geq 0.$$

Next, we consider the linkage between our result and that in [7]. We fix $0 < \bar{\alpha} < 1$ and $\alpha : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, C^∞ , $\alpha(0) = 0, \dots, \alpha^{(n-1)}(0) = 0$, $\alpha(t) = \bar{\alpha}$, $t \geq 1$. We let $\varepsilon \rightarrow^+ 0$. Let $q(t) = \alpha c^*(t)$, then (11) is modified to

$$\lim_{T \rightarrow \infty} \inf_{T \leq T'} \sum_{i=1}^n \left[\frac{\partial (U(T' - N + 2) + \cdots + U(T'))}{\partial c_i(T' + 1)} \alpha c_i^*(T' + 1) \right. \\ \left. + \cdots + \frac{\partial (U(T'))}{\partial c_i(T' + N - 1)} \alpha c_i^*(T' + N - 1) \right] \\ = \bar{\alpha} \lim_{T \rightarrow \infty} \inf_{T \leq T'} \sum_{i=1}^n \left[\frac{\partial (U(T' - N + 2) + \cdots + U(T'))}{\partial c_i(T' + 1)} c_i^*(T' + 1) \right. \\ \left. + \cdots + \frac{\partial (U(T'))}{\partial c_i(T' + N - 1)} c_i^*(T' + N - 1) \right] \\ (12) \quad \leq 0.$$

Because $\bar{\alpha} > 0$, we then have

$$\lim_{T \rightarrow \infty} \inf_{T \leq T'} \sum_{i=1}^n \left[\frac{\partial (U(T' - N + 2) + \cdots + U(T'))}{\partial c_i(T' + 1)} c_i^*(T' + 1) \right. \\ \left. + \cdots + \frac{\partial (U(T'))}{\partial c_i(T' + N - 1)} c_i^*(T' + N - 1) \right] \leq 0. \\ (13)$$

which is an extension of [7]'s transversality condition.

3. A COUNTEREXAMPLE

We proceed to show that Assumption 1 and 2 are imperative in the sense that (11) becomes invalid if one of them is violated. We consider the following simple counterexample:

$$(14) \quad U(c(t), c(t+1), c(t+2), t) = (c(t) - \alpha)^2 + \beta c(t+1) + \gamma c(t+2),$$

where $\alpha > 0$, $\beta > 0$, $\gamma > 0$, and the initial values $c(0) = c_0$, $c(1) = c_1$ are given. From (10), we see that Euler's condition is given by

$$(15) \quad \begin{aligned} \frac{\partial U(0)}{\partial c(0)} &= 0, \\ \frac{\partial (U(0) + U(1))}{\partial c(1)} &= 0, \\ \frac{\partial (U(t-2) + U(t-1) + U(t))}{\partial c(t)} &= 0, 2 \leq t \leq T', \end{aligned}$$

which implies

$$\begin{aligned} (15') \quad & t = 0, \quad 2(c(0) - \alpha) = 0, \\ (15'') \quad & t = 1, \quad 2(c(1) - \alpha) + \beta = 0, \\ (15''') \quad & t = 2, \quad 2(c(2) - \alpha) + \beta + \gamma = 0, \\ (15'''') \quad & t = 3, \quad 2(c(3) - \alpha) + \beta + \gamma = 0, \\ & \dots\dots\dots, \\ (15''''') \quad & t = T', \quad 2(c(T') - \alpha) + \beta + \gamma = 0. \end{aligned}$$

Thus, we have $c(2) = c(3) = \dots = c(T') = \alpha - \frac{\beta + \gamma}{2}$.

Choosing a p so that $p(0) = 0$ and $p(t) > 0$, there exists $T_0 > 0$, $p(t)$ is a constant $p_\infty > 0$ when $t \geq T_0$.

From (14), we see that

$$(16) \quad \frac{\partial (U(T'-1) + U(T'))}{\partial c(t+1)} q(T'+1) = (\gamma + \beta)q(T'+1) \leq 0,$$

$$(17) \quad \frac{\partial (U(T'))}{\partial c(t+2)} q(T'+2) = \gamma q(T'+2) \leq 0.$$

Hence, we have arrived at a contradiction to (11).

Next, we show that Assumption 1 is violated, which causes this contradiction. We consider $U(c^*(t) + \varepsilon p(t), c^*(t+1) + \varepsilon p(t+1), c^*(t+2) + \varepsilon p(t+2)) - U(c^*(t), c^*(t+1), c^*(t+2))$. Substituting $c^*(t) = \alpha - \frac{\beta + \gamma}{2}$ into it,

we have

$$\begin{aligned}
& U(c^*(t) + \varepsilon p(t), c^*(t+1) + \varepsilon p(t+1), c^*(t+2) + \varepsilon p(t+2)) \\
& \quad - U(c^*(t), c^*(t+1), c^*(t+2)) \\
&= (c^*(t) + \varepsilon p(t) - \alpha)^2 + \beta(c^*(t+1) + \varepsilon p(t+1)) + \gamma(c^*(t+2) + \varepsilon p(t+2)) \\
& \quad - ((c^*(t) - \alpha)^2 + \beta c^*(t+1) + \gamma c^*(t+2)) \\
& \quad (18) \\
&= \left(\varepsilon p(t) - \left(\frac{\beta + \gamma}{2} \right) \right)^2 + \varepsilon(\beta p(t+1) + \gamma p(t+2)) - \left(\frac{\beta + \gamma}{2} \right)^2.
\end{aligned}$$

Hence,

$$\begin{aligned}
& \inf_{T \leq T'} \sum_{t=0}^{T'} \frac{\left(\varepsilon p(t) - \left(\frac{\beta + \gamma}{2} \right) \right)^2 + \varepsilon\beta p(t+1) + \varepsilon\gamma p(t+2) - \left(\frac{\beta + \gamma}{2} \right)^2}{\varepsilon} \\
&= \inf_{T \leq T'} \sum_{t=0}^{T'} (\varepsilon p(t)^2 - (\beta + \gamma)p(t) + \beta p(t+1) + \gamma p(t+2)) \\
&= \inf_{T \leq T'} \left(\varepsilon \sum_{t=0}^{T'} p(t)^2 + \beta p(T'+1) + \gamma p(T'+2) \right) \\
&= \inf_{T \leq T'} \left(\varepsilon \sum_{t=0}^{T'} (p(t)^2) \right) + \beta p_\infty + \gamma p_\infty \\
& \quad (19) \\
&= \varepsilon \sum_{t=0}^T (p(t)^2) + \beta p_\infty + \gamma p_\infty.
\end{aligned}$$

Ω is the limit of (18) when $T \rightarrow \infty$, $\varepsilon \rightarrow 0$. However,

$$\text{because } \lim_{\varepsilon \rightarrow 0} \lim_{T \rightarrow \infty} \left(\varepsilon \sum_{t=0}^T (p(t)^2) + \beta p_\infty + \gamma p_\infty \right) = \infty,$$

$$\text{whereas } \lim_{T \rightarrow \infty} \lim_{\varepsilon \rightarrow 0} \left(\varepsilon \sum_{t=0}^T (p(t)^2) + \beta p_\infty + \gamma p_\infty \right) = \beta p_\infty + \gamma p_\infty, \text{ we see}$$

that Ω does not converge uniformly for ε when $T \rightarrow \infty$. Hence, Assumption 1 is violated and (11) is also not satisfied.

4. CONCLUSION

This paper gives the two assumptions that would be imperative when examining infinite horizon discrete time optimization problems in which

the objective functions are unbounded. Our results generalizes the results of [5, 6, 7, 10, 12] ($N = 1$) to higher order difference problems. Specifically, when $N = 1$, our transversality condition is the discrete time version of that of [7] (Theorem 3.2). Moreover, our Assumption 1 and 2 also constitute the discrete time version of the Assumptions in [11]. As in [11], our Assumption 1 and 2 obviously hold when a discounting factor is incorporated into the model. In this sense, this paper also extends the transversality conditions examined in the presence of discounting.

REFERENCES

- [1] W.A. Brock, On existence of weakly maximal programmes in a multi-sector economy, *Review of Economic Studies* 37 (1970), pp. 275–280.
- [2] D. Cai, and T.G. Nitta, Optimal solutions to the infinite horizon problems: Constructing the optimum as the limit of the solutions for the finite horizon problems. *Nonlinear Analysis: Theory, Methods and Applications*(2009), doi:10.1016/j.na.2009.03.066.
- [3] D. Cai, and T.G. Nitta, Limit of the solutions for the finite horizon problems as the optimal solution to the infinite horizon optimization problems, *Journal of Difference Equations and Applications*(2010), doi:10.1080/10236190902953763.
- [4] D. De La Croix, and P. Michel, *A Theory of Economic Growth: Dynamics and Policy in Overlapping Generations*, Cambridge University Press, 2002.
- [5] I. Ekeland, and J.A. Scheinkman, Transversality conditions for some infinite horizon discrete time optimization problems, *Mathematics of Operations Research* 11(1986), pp.216-229.
- [6] T. Kamishigashi, A simple proof of Ekeland and Scheinkman’s result on the the necessity of a transversality condition, *Economic Theory* 15(2000), pp.463-468.
- [7] T. Kamishigashi, Necessity of transversality conditions for infinite horizon problems, *Econometrica* 69(2001), pp.995-1012.
- [8] T. Kamishigashi, Necessity for transversality conditions for stochastic problems, *Journal of Economic Theory* 109(2003), 140-149.
- [9] S. Lang, *Undergraduate Analysis*. 2nd ed., Springer, New York, 1997.
- [10] P. Michel, Some clarifications on the transversality condition, *Econometrica* 58(1990), 705-723.
- [11] R. Okumura, D. Cai, and T.G. Nitta, Transversality conditions for infinite horizon optimality: Higher order differential problems, *Nonlinear Analysis: Theory, Methods and Applications*(2009), doi:10.1016/j.na.2009.02.110.
- [12] M.L. Weitzman, Duality theory for infinite horizon convex models, *Management Science* 19 (1973), pp.783-789.