



Departament de Física
Grup de Física Teòrica

Efficiency in Quantum Key Distribution Protocols using entangled Gaussian states

Carles Rodó Sarró

Treball de recerca del programa de Doctorat de Física de la Universitat Autònoma de Barcelona, realitzat per Carles Rodó Sarró sota la direcció de la Dra. Anna Sanpera i Trigueros.

Bellaterra, Abril de 2008

La Dra. Anna Sanpera i Trigueros, Investigadora ICREA al departament de Física Teòrica de la Facultat de Ciències de la Universitat Autònoma de Barcelona, CERTIFICA: Que la present memòria, que porta per títol: “Efficiency in Quantum Key Distribution”, ha estat realitzada sota la seva direcció pel Carles Rodó i Sarró i que constitueix el seu Treball de Recerca del programa de Doctorat de Física.

Bellaterra, Abril de 2008

Anna Sanpera i Trigueros

Vull donar els meus agraïments, en primer lloc a l'Albert Bramon qui, farà ja uns anys, em va començar mostrant els misteris de la Física Quàntica en aquelles tan ben cronometrades classes i, qui, em va facilitar poder treballar amb l'Anna Sanpera.

Vull agrair, com no, la inagotable paciència i l'esforç que ha posat en mi l'Anna. No vull oblidar la generositat i lo molt que he après col·laborant amb el Gerardo Adesso.

També vull recordar molt especialment als meus ex-companys amb qui més temps he passat, el Javi i l'Àlex per les variopintes discussions de Física i pels bons moments compartits fora de la feina. També a les noves “promeses” Marc i Joan Antoni.

Als amics d'arreu, Rubén, Dani, Àlex, Ignasi, Marina, Mara, Sito, Xavi, Pepe, Cristina, Hugo, Edu, Jaime, Pedro, Shona, Mirka, Guillem, Felix, Deivid, Chechu, Christian, ...

Finalment agrair a la família que sempre m'ha recolzat, al meu pare per haver-me motivat a encuriosir-me per la Física i sobretot a la persona amb qui més m'he barallat, la meva mare, però qui ho ha donat tot sempre per mi, i qui més es mereix els meus agraïments.

Carles

Resum

L'estudi de l'eficiència en els protocols de Criptografia és una tasca clau per a la seva futura implementació tecnològica ja que els recursos físics dels quals es disposa experimentalment no són il·limitats. En aquesta tesina s'estudia l'eficiència en els protocols de distribució de claus quàntiques basats en l'entrellaçament quàntic. Usant estats de variable discreta i destil·lant entrellaçament, sempre és possible destil·lar singlets i així extreure correlacions clàssiques perfectes. En canvi, mitjançant estats de variable contínua com ho són els modes de la llum, la destil·lació no és possible mitjançant només operacions Gaussians. No obstant, mostrem que, utilitzant estats Gaussians de llum bipartits i l'entrellaçament en les quadratures del camp electromagnètic d'aquests estats, sí és possible extreure seqüències de bits correlacionats, necessàries per destil·lar claus aleatòries segures de forma eficient. Obtenim que, usant estats mescla Gaussians NPPT simètrics i operacions Gaussians es poden distribuir claus quàntiques segures davant d'atacs individuals d'un espia. Davant d'atacs finits coherents no tots els estats permeten una extracció segura. Mostrem quins estats sí són segurs davant d'atacs finits coherents i l'eficiència amb la qual es poden extreure aquestes correlacions clàssiques davant d'atacs individuals. Finalment s'analitza la relació entre l'eficiència, l'entrellaçament i la puresa per estats Gaussians simètrics. Concloem que un entrellaçament gran (més correlacions entre els modes) implica un guany en eficiència a l'hora d'extreure bits ben correlacionats. No hi ha una relació un a un donat que estats amb el mateix entrellaçament però diferents pureses donen lloc a diferent eficiència. El que succeeix és que estats amb una puresa alta afavoreixen un augment en l'eficiència donat que estem desentrellaçant l'espia, però aquests estats són ineficients quan finalment realitzem Classical Advantage Distillation on, per extreure la clau aleatòria segura un alt grau de correlacions és essencial.

Contents

1	Introduction	1
1.1	Classical Cryptography	2
1.1.1	Vernam cypher	2
1.1.2	Public key distribution: The RSA algorithm	4
1.2	The solution to the distribution of the key	5
1.2.1	”Prepare and Measure“ scheme	5
1.2.2	”Entanglement based“ scheme	6
1.3	Efficiency in Quantum Key Distribution (QKD) protocols	7
1.4	Outline	9
2	Continuous Variable formalism	11
2.1	Continuous Variable system	11
2.2	Canonical Commutation Relations	12
2.3	Phase-space	13
2.3.1	Phase space geometry	13
2.3.2	Symplectic operations	13
2.4	Probability Distribution functions	15
2.4.1	Quantum states	16
2.4.2	Properties of the Wigner distribution	17
2.4.3	The generating function of a Classical probability distribution	18
2.4.4	The generating function of a quasi-probability distribution	19
2.5	Gaussian states	20
2.5.1	Displacement Vector (DV) and Covariance Matrix (CM)	20
2.5.2	Hilbert space, phase space and DV&CM connection	22
2.6	Multipartite states and entanglement	25
2.6.1	Bipartite Gaussian states	25
2.6.2	Entanglement of Gaussian states	26
2.7	Appendix of integrals	28
3	Efficiency in QKD protocols with entangled Gaussian states	29
3.1	State of the problem: QKD with entangled Gaussian states	29
3.2	Efficient QKD with entangled Gaussian states	31

4	Summary and Conclusions	37
	Bibliography	40

Chapter 1

Introduction

Rolf Landauer, known for his remarkable contributions to the theory of electrical conductivity, mesoscopic phenomena, and the physics of information, during the 1960s, as the director of IBM Research's Solid State Science Division persuaded the idea that information was rather than an abstract concept a physical one.

Nowadays the Quantum Information science (QI) has become one of the most active fields of research. QI research spreads in two main areas, Quantum Communication and Quantum Computation tasks. One of the most important topics of Quantum Communication deals with the study of secure communications. Thus Cryptography, based in protocols that exploit Quantum Mechanics properties, plays a central role in Quantum Communication.

Cryptography refers to strategies which permit the secure communication between two distant parties (traditionally denoted by Alice and Bob) that wish to communicate secretly. So its purpose is to design new communication algorithms being sure that secrecy is preserved. In Classical Cryptography there is only one cryptographic protocol, known as the ideal Vernam cypher method, which is absolutely secure. The Vernam cypher consists of a random secret key (private key) shared between Alice (the sender) and Bob (the receiver) used to encode and decode messages. However, this method suffers from two drawbacks. First, if Alice wants to communicate N bits to Bob, they will need to have a random secret key in advance with at least N bits. Moreover, this key can only be used once to make the method unbreakable. Second, the key must originally be exchanged by hand before the communication to keep secrecy and it is essential that the key is totally random. To solve the second issue, classical ways exist to distribute the key among them being this distribution partially secure. This problem is known as Key Distribution.

In Classical Cryptography the problem of Key Distribution can be solved partially by designing new algorithms which permit the distribution of the secret key in a "practical secure" way. The price to pay is that absolute security is not achievable. Nevertheless as security relies on the fact that to decrypt one needs to invert difficult mathematical operations, the time to decrypt the key is long, and thus practical security is achievable.

The distribution problem is classically solved by using Public Key Cryptosystems. They work as follows: Alice and Bob possess, in advance, a common secret key. This secret key can be used many times to distribute among them, several private keys that they are going to use to encrypt messages through *e.g.* the Vernam cypher method. Any time Alice (the sender) wants to distribute a private key to encrypt later messages, she only needs to make one key publicly available, referred as the public key. From this public key any receiver can extract the private key, but only Bob (the receiver in possession of the secret key) can extract it in an efficient way. Any receiver without the secret key needs to decrypt a problem which has non-polynomial (NP) complexity. This way of distributing a secure key can be done even when the key's length is much smaller than the message and even if the key is used several times.

Nowadays bank's security, electronic commerce and the internet are mostly based on one of these algorithms, the RSA Cryptography algorithm proposed in 1977 by Rivest, Shamir, and Adleman from the MIT. Security in RSA algorithm relies on the fact that to decrypt the algorithm one needs to solve the factorisation problem which is an NP problem. Even now, there is no efficient algorithm known to solve the factorisation problem. This means that even though computational resources increase constantly, one simply needs to exploit the NP character of the algorithm to make the solution harder to find. Moreover, the key can eventually be redistributed and changed.

The "Quantum Computer" arises here first as a menace for Classical Key Distribution methods and then as a the solution for the security in Cryptography. Based on the Quantum nature of the microscopic world, this new generation of computers, still in a theoretical stage, are known to be able to solve hard mathematical problems rapidly. In 1994 Peter Shor proposed a Quantum protocol to solve the factorisation problem in an efficient way, known as the Shor's algorithm. If such a computer can be realised, current cryptographic protocols will not be anymore secure.

Can Quantum Mechanics then offer a solution for a secure Cryptography method? The answer is yes. Based on the intrinsic nature of the microscopic world, Quantum Mechanics permits to perform Cryptography in an unconditional secure way. At present, Quantum Cryptography is the only real implementation of QI. In this work, we will present one way of performing efficient Quantum Cryptography using entangled Gaussian states of light and standard optical devices (Gaussian operations).

1.1 Classical Cryptography

1.1.1 Vernam cypher

In Classical Cryptography there is a protocol called Vernam cypher which is absolutely secure. This protocol is the best and most well known classical private key (or ONE-TIME-PAD) cryptosystem. In order to achieve security in communications between two parties (Alice and Bob) with the Vernam cypher protocol, one needs to have a private key, in advance, possessed by Alice and Bob exclusively of at least the

same length (# of bits) as the message to be encoded. This key has to be random and in possession of Alice and Bob only, and so a Classical Key Distribution is a pre-condition to achieve this absolute security. If this is the case, then Cryptography is going to be absolutely secure and works as follows.

If Alice wants to encode a message m with the key k , she only needs to perform the following operation between the message and the key, and send the encoded message e to Bob:

$$\text{Enc}_k(m) = m \oplus k = e. \quad (1.1)$$

Only Bob who has also the key can decode the message (invert the operation) because the key Alice has used is random. Thus, he only needs to use the key again and perform the following operation to the encoded message e to retrieve the original message m :

$$\text{Dec}_k(e) = \text{Dec}_k[\text{Enc}_k(m)] = e \oplus k = d = m. \quad (1.2)$$

Let us illustrate Vernam cypher with an specific example. Alice wants to communicate to Bob, in a secure way, a message m (in a binary string) of *e.g.* 9 bits. They share the key k of the same size as the message (9 bits). Alice encodes her message by applying a *XOR* (exclusive *OR*)¹ operation between the message m and the key k .

In the following table we summarise the *XOR* operation

$$\begin{array}{c|cc} \oplus & 0 & 1 \\ \hline 0 & 0 & 1 \\ 1 & 1 & 0 \end{array} \quad (1.3)$$

As a result Alice has the encoded message e , that will send to Bob in a public way.

message	$m = 010011101$
key	$k = 110100011$
encoded message	$e = 1000111110$

Then Bob wants to readout the message, and thus, performs the inverse operation (which is again a *XOR*) between the encoded message e and the key k .

As a result Bob has the decoded message d that coincides with the message Alice wanted to communicate to him.

encoded message	$e = 1000111110$
key	$k = 110100011$
decoded message	$d = 010011101$

¹Also known as *AND* or $\oplus \bmod (2)$.

1.1.2 Public key distribution: The RSA algorithm

Thus far, Classical Cryptography has not solved the distribution of the private key needed to perform Vernam cypher encryption. This distribution can be done in a public and practical secure way with the RSA algorithm. Nowadays the RSA algorithm is the most commonly used algorithm of public key. It is a Classical Key Distribution algorithm that permits together with the Vernam cypher method to perform secure communication as long as the factorisation problem is unsolved. Let us illustrate how it works with the following example.

i) The sender, say Alice, chooses two "big" different prime numbers, say $p = 61$ and $q = 53$ and computes its product $n = p q = 3233$ and also the following quantity $\phi = (p - 1)(q - 1) = 3120$.

ii) She chooses a positive integer l smaller and coprime with ϕ , in the example $l = 17$.

iii) As a private key, Alice gives to Bob the number k such that $k l = 1 \bmod (\phi)$, take *e.g.* $k = 2753$. At the same time Alice makes public l and n , what we call public key.

iv) With the public key (l and n) anyone can encrypt a message m and send it to Bob, but only Bob who is in possession of the private key k is able to decrypt the message. This method can thus be used to perform Classical Key Distribution. Any time Alice wants to communicate with Bob, she sends a secure key encrypted with l and n and only Bob will be able to retrieve it. Once Bob has the secure key, Alice can send messages to Bob *via* the Vernam cypher using this secure key.

v) Encryption proceeds as follows. Alice wants to distribute a key encoded in a message $m = 123$. She uses the public key and computes the encryption $\text{Enc}_{l,n}(m) = m^l \bmod (n) = e = 855$.

vi) Bob now wants to decrypt the message e to extract a secure key, so he calculates $\text{Dec}_{k,n}(e) = \text{Dec}_{k,n}[\text{Enc}_{l,n}(m)] = e^k \bmod (n) = d = m = 123$. Bob is the only one in possession of the private key k and so the only one that can decrypt the message e to find the secure key Alice is going to use with the Vernam cypher to communicate securely with him.

The private key Alice and Bob share can be used more than once to distribute secure keys, in such a way that, to perform Vernam cypher we no longer need to share a long private key because we can distribute many of them in a secure way. Security relies in the fact that, from the encrypted message e , and the public keys l and n it is very difficult to find the private key k (and so m) or the original two prime numbers q and p , even in the case we are reusing the key k . This is because factorisation is a NP problem, whose efficient solution is not known yet. Security thus, in the RSA algorithm, relies on the fact that with the current computation resources, NP problems cannot be solved efficiently.

1.2 The solution to the distribution of the key

What Quantum Cryptography offers is an absolutely secure distribution of a random key which combined with the Vernam cypher guarantees completely secure Cryptography. Thus, the Quantum Cryptography problem is in fact the problem of distributing a secure random key, *i.e.* the Quantum Key Distribution (QKD) problem.

Quantum Cryptography relies on the possibility of establishing a secret random key between two distant parties traditionally denoted by Alice and Bob. If the key is securely distributed, the algorithms used to encode and decode any message can be made public without compromising security. The key consists typically in a random sequence of bits which both, Alice and Bob, share as a string of classically correlated data. The superiority of Quantum Cryptography comes from the fact that the laws of Quantum Mechanics permit to the legitimate users (Alice and Bob) to infer if an eavesdropper has monitored the distribution of the key and has gained information about it. If this is the case, Alice and Bob will both agree in withdrawing the key and will start the distribution of a new one. In contrast, Classical Key Distribution, no matter how difficult the distribution from a technological point of view is, can always be intercepted by an eavesdropper without Alice and Bob realising it.

In Quantum Cryptography two seemingly independent main schemes exist for QKD. The first, the "Prepare and Measure" scheme, originally proposed by C.H. Bennett and G. Brassard in 1988 and known as BB84 [1], does not use entangled states shared between Alice and Bob and the key is established by sending non-entangled quantum states between the parties and communicating classically. Security is guaranteed by the Quantum nature of the measurements. The second scheme ("Entanglement based"), uses as a resource shared entanglement, like the one originally proposed by A. Ekert in 1991 known as Ekert91 [2], where indeed entanglement is explicitly distributed and the security is guaranteed by Bell's theorem. However, the two schemes have been shown to be completely equivalent [3], and specifically entanglement stands as a precondition for any secure key distribution [4].

1.2.1 "Prepare and Measure" scheme

BB84 permits a secure distribution of a secret key. The protocol does not avoid an eavesdropper from intercepting the key, but it lets Alice and Bob know if the key has been intercepted, and so that it can be discarded. Security relies in the Quantum nature of the measurement. We sketch here the steps of the protocol.

i) Alice prepares a secret sequence of random bits and encodes them in the state of a spin-1/2 system (or the polarisation of photons) by choosing randomly between two bases (Z and X). Alice encodes $|\pm\rangle$ ($|\pm\rangle_x$) according to 0/1 in base Z (0/1 in base X). Then she sends Bob the states she has prepared. As an example:

Alice random bits	0	1	1	0	0	1	1	0	0
Alice random bases	Z	X	X	X	Z	X	Z	X	X
Alice final states	$ +\rangle$	$ -\rangle_x$	$ -\rangle_x$	$ +\rangle_x$	$ +\rangle$	$ -\rangle_x$	$ -\rangle$	$ +\rangle_x$	$ +\rangle_x$

ii) Bob receives the states and measures in another random choice of bases. The outcome of the measurements is going to be retained as the bits received.

Alice final states	$ +\rangle$	$ -\rangle_x$	$ -\rangle_x$	$ +\rangle_x$	$ +\rangle$	$ -\rangle_x$	$ -\rangle$	$ +\rangle_x$	$ +\rangle_x$
Bob random bases	X	Z	X	X	X	X	X	X	X
Bob final states	$ +\rangle_x$	$ +\rangle$	$ -\rangle_x$	$ +\rangle_x$	$ +\rangle_x$	$ -\rangle_x$	$ -\rangle_x$	$ +\rangle_x$	$ +\rangle_x$
Bob received bits	0	0	1	0	0	1	1	0	0

iii) Bob communicates to Alice his choice of basis in a public way.

iv) Alice identifies the set of bits for which they have performed the measurement in the same basis, *i.e.* outcomes 3, 4, 6, 8 and 9. Alice and Bob discard the set of data in which they did not agree (the rest).

v) Bob sends part of his data (received bits) to Alice by a public channel. Alice checks the correlation between the data.

vi) If the error rate is less than 25% Alice deduces that there is not an eavesdropper present and she communicates it to Bob. Alice and Bob use the set of remaining data as a private key to encrypt messages with Vernam cypher.

1.2.2 "Entanglement based" scheme

Other protocols exist, which demand as a fundamental resource, shared entanglement between Alice and Bob. In the same way as in BB84, these protocols permit a secure distribution of a secret key. This can be done as far as the protocol ensures if there has been an interception of the key. Security relies in the Quantum correlations *i.e.* entanglement like in Ekert91. We sketch here the steps of this well-known protocol below.

i) The first step consists on distributing (along the z direction) singlet states of a spin-1/2 system (or polarisations of photons) between Alice and Bob. Thus Alice and Bob share many copies of a Bell state $|\Psi^-\rangle = \frac{1}{\sqrt{2}}(|0\rangle|1\rangle - |1\rangle|0\rangle)$.

ii) Alice and Bob are going to measure in the $x - y$ plane in one of the three directions given by unit vectors $\vec{A}_i = (\cos \phi_i^A, \sin \phi_i^A)$ and $\vec{B}_j = (\cos \phi_j^B, \sin \phi_j^B)$ respectively, where the azimuthal angles are fixed to $\phi_i^A = (0, \pi/4, \pi/2)^i$ and to $\phi_j^B = (\pi/4, \pi/2, 3\pi/4)^j$. Each time they will choose the basis randomly and independently for each pair of incoming particles.

iii) The quantity

$$\mathcal{E}(\vec{A}_i, \vec{B}_j) = \mathcal{P}_{++}(\vec{A}_i, \vec{B}_j) + \mathcal{P}_{--}(\vec{A}_i, \vec{B}_j) - \mathcal{P}_{+-}(\vec{A}_i, \vec{B}_j) - \mathcal{P}_{-+}(\vec{A}_i, \vec{B}_j) \quad (1.4)$$

is the correlation coefficient of the measurements performed by Alice and Bob along $\vec{A}_i$ and by Bob along $\vec{B}_j$. Here $\mathcal{P}_{\pm\pm}(\vec{A}_i, \vec{B}_j)$ denotes the probability that result ± 1

has been obtained along $\vec{A}_i$ and ± 1 along $\vec{B}_j$. Straightforward calculations give rise to

$$\begin{aligned}\mathcal{P}_{++}(\vec{A}_i, \vec{B}_j) &= \frac{1}{2} \sin^2(\phi_i^A - \phi_j^B), \\ \mathcal{P}_{--}(\vec{A}_i, \vec{B}_j) &= \frac{1}{2} \sin^2(\phi_i^A - \phi_j^B), \\ \mathcal{P}_{+-}(\vec{A}_i, \vec{B}_j) &= \frac{1}{2} \cos^2(\phi_i^A - \phi_j^B), \\ \mathcal{P}_{-+}(\vec{A}_i, \vec{B}_j) &= \frac{1}{2} \cos^2(\phi_i^A - \phi_j^B),\end{aligned}$$

thus according to the Quantum rules $\mathcal{E}(\vec{A}_i, \vec{B}_j) = -\vec{A}_i \vec{B}_j = \cos(\phi_i^A - \phi_j^B)$. We see that whenever they choose the same orientation Quantum Mechanics predicts total anticorrelation in the outcomes *i.e.* $\mathcal{E}(\vec{A}_i, \vec{B}_j) = -1$.

iv) Here we define a quantity composed of those correlation coefficients for which Alice and Bob have measured in different directions,

$$\mathcal{S} = |\mathcal{E}(\vec{A}_1, \vec{B}_1) + \mathcal{E}(\vec{A}_3, \vec{B}_3) - \mathcal{E}(\vec{A}_1, \vec{B}_3) + \mathcal{E}(\vec{A}_3, \vec{B}_1)|. \quad (1.5)$$

Again, Quantum Mechanics requires, $\mathcal{S} = 2\sqrt{2} > 2$.

v) After the transmission has taken place, Alice and Bob can announce in public the orientations they have chosen for each measurements and divide them into two separated groups. A first group for which they coincide and a second group for which they do not. The second group of outcomes is made public and it is used to establish the value of $\mathcal{S}$.

The CHSH (Clauser, Horne, Shimony, and Hold) inequalities, a generalisation of Bell inequalities, asserts that $\mathcal{S} \leq 2$. But Quantum Mechanics, and in particular Bell states, violate CHSH inequalities. If this is the case, *i.e.* the value of $\mathcal{S}$ that they find is exactly $2\sqrt{2}$, they now that their states have not been disturbed and so the first group of outcomes, that are random, are totally anticorrelated and can be converted into a secret string of bits. Later on they can use this string as a private key to encrypt messages with Vernam cypher.

1.3 Efficiency in Quantum Key Distribution (QKD) protocols

As already mentioned, Quantum Key Distribution, refers to specific Quantum strategies which permit the secure distribution of a secret key between two parties that wish to communicate secretly. Quantum Cryptography has been proved unconditionally secure in ideal scenarios and has been successfully implemented using quantum states with finite (discrete) as well as infinite (continuous) degrees of freedom. We have analysed the efficiency of QKD protocols that use as a resource entangled Gaussian states and Gaussian operations only. In this framework, it has already been shown that QKD is possible [5] but the issue of its efficiency was not considered. We have propose a figure of merit (the efficiency E) to quantify the number of classical correlated bits that can be used to distill a key from a sample of N entangled states. We have related the efficiency of the protocol to the entanglement and purity of the states shared between the parties.

Notice that if Alice and Bob share a collection of distillable entangled states, they can always obtain a smaller number of maximally entangled states from which they can establish a secure key [6]. The number of singlets (maximally entangled states) that can be extracted from a quantum state using only Local Operations and Classical Communication (LOCC) is referred to as the Entanglement of Distillation E_D . In order to establish a key, another important concept is the number of secret bits K_D , that can be extracted from a quantum state using LOCC. As a secret bit can always be extracted from maximally entangled state, $E_D \leq K_D$. Furthermore, there are quantum states which cannot be distilled but in spite of being entangled, *i.e.*, have $E_D = 0$. They are usually referred to as bound entangled states since its entanglement is bound to the state. Nevertheless, for some of those states it has been shown that $K_D \neq 0$ and thus they can be used to establish a secret key [7].

A particular case of states that cannot be “distilled” by “normal” procedures are continuous variables Gaussian states, *e.g.*, coherent, squeezed and thermal states of light. By “normal” procedures we mean operations that preserve the Gaussian character of the state (Gaussian operations). They correspond *e.g.* to beam splitters, phase shifts, mirrors, squeezers, etc. Thus, in the Gaussian scenario all entangled Gaussian states possess bound entanglement. Quantum Cryptography with Gaussian states using Gaussian operations has been experimentally implemented using “Prepare and Measure” schemes with either squeezed or coherent states [8, 9, 10]. Those schemes do not demand entanglement between the parties.

Navascués *et al.* [5] have shown that it is also possible using only Gaussian operations to extract a secret key *à la* Ekert91 from entangled Gaussian states, in spite the fact that these states are not distillable. In other words, it has been proven that in the Gaussian scenario all entangled Gaussian states fulfil $GK_D > 0$ (where the letter G stands for Gaussian) while $GE_D = 0$.

The way to proceed in this scheme *i.e.*, how can, Alice and Bob extract a list of classically correlated bits from a set of symmetric 1×1 entangled modes goes as follows:

- i) they agree on a value $x_0 > 0$,
- ii) Alice(Bob) measures the quadrature of each of her(his) modes $\hat{X}_A(\hat{X}_B)$,
- iii) they make public the modulus of their outcomes, but not the sign and accept only outputs such that $|x_A| = |x_B| = x_0$,
- iv) they associate *e.g.*, the classical value 0(1) to $x_i = +x_0(-x_0)$, $i = A, B$ and thus establish a list of classically correlated bits.

From there, they can apply Classical Advantage Distillation [11] to establish the secret key. This protocol is secure against individual eavesdropper attacks. As the protocol is based on output coincidences of the measurements of the quadratures which, by definition, are operators with a continuous spectrum, this protocol has zero efficiency.

1.4 Outline

Efficiency is a key issue for any experimental implementation of Quantum Cryptography since available resources are not unlimited. Since it is possible to extract a secret key *à la* Eckert91 from entangled Gaussian states in the Gaussian scenario one important question is to address the efficiency.

In fact, the protocol suffers from an efficiency problem because the success probability of the protocol is vanishingly small. Here we study the consequences of relaxing the conditions to a more realistic scenario. We assume that Alice and Bob can extract a list of sufficiently correlated classical bits obtained by accepting measurement outputs that do not coincide but are bound within a range. We ask ourselves which is the possibility that Alice and Bob can still distribute the key in a secure way under individual and finite coherent attacks.

We find that there always exists a finite interval which the protocol can be implemented successfully. The length of this interval depends on the entanglement and on the purity of the shared states, and increases with increasing entanglement.

In Chapter 2 we review the formalism of Continuous Variable (CV) systems focussing on Gaussian states. It will be shown that Gaussian states admit an easy mathematical description based on phase space Wigner functions. We will introduce also the basic ingredients to describe also systems with entanglement.

In Chapter 3 we will present first a simple academic protocol that permits to extract a quantum key from an entangled continuous variable system. Differently from discrete systems, Gaussian entangled states cannot be distilled with Gaussian operations. However, as we will show, it is possible to extract perfectly correlated classical bits to establish a secret key between the sender and receiver. We will then demonstrate that this protocol, properly modified, can be made efficient and can be implemented with present technology [12].

Finally, in Chapter 4, we conclude summarising our results and giving some general conclusions and future directions of research.

Chapter 2

Continuous Variable formalism

This Chapter is intended exclusively to describe Continuous Variable systems and to provide the mathematical framework to analyse the problem of QKD in Continuous Variable systems. We will focus then on Gaussian Continuous Variable states, which describe among others, coherent, squeezed and thermal states of light. Presently these states are the preferred resources in experiments of QI using Continuous Variable systems. For further background information the interested reader is referred to [13, 14, 15, 16].

2.1 Continuous Variable system

A system corresponds to a continuous variable system if it possesses two canonical conjugated degrees of freedom *i.e.* there exist two observables that fulfil the Canonical Commutation Relations (CCR). The CCR for two canonical observables $\hat{q}$ and $\hat{p}$ read ¹

$$[\hat{q}, \hat{p}] = i\mathbb{I}. \quad (2.1)$$

It is a direct consequence that they possess a continuous spectra and act in an infinite dimensional Hilbert space.

As examples of CV systems we can think as the position-momentum of a massive particle, the quadratures of an electromagnetic field or the collective spin of a polarised ensemble of atoms. In all of the three examples above there exist two observables fulfilling (2.1). As we will show, they obey the standard bosonic commutation relations and so we call these systems bosonic modes. We can deal with several modes, and in this case ordering the operators by canonical pairs by $\hat{R}^T = (\hat{q}_1, \hat{p}_1, \hat{q}_2, \hat{p}_2, \dots, \hat{q}_N, \hat{p}_N)$ we can compactly state CCR as

$$[\hat{R}_i, \hat{R}_j] = i(J_N)_{ij} \quad (2.2)$$

¹Quadrature operators are chosen dimensionless in such a way that $\hbar$ is not going to appear in any formula.

where $i, j = 1, 2, \dots, 2N$ and $J_N = \oplus_{i=1}^N J$ accounts for all modes while J is the so-called symplectic matrix which corresponds to a antisymmetric and non-degenerate form fullfilling (i) $\forall \eta, \zeta \in \mathbb{R}^{2N} : \langle \eta | J | \zeta \rangle = -\langle \zeta | J | \eta \rangle$ and (ii) $\forall \eta : \langle \eta | J | \zeta \rangle = 0 \Rightarrow \zeta = 0$. In the appropriate choice of basis (canonical coordinates) the symplectic matrix is brought into the standard form $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$

2.2 Canonical Commutation Relations

The CCR (2.1) are related with the classical Poisson brackets *via* the 1st quantisation transcription: $\{\mathcal{A}, \mathcal{B}\}_{pp} \equiv \sum_{\mu} (\frac{\partial \mathcal{A}}{\partial Q_{\mu}} \frac{\partial \mathcal{B}}{\partial P_{\mu}} - \frac{\partial \mathcal{B}}{\partial Q_{\mu}} \frac{\partial \mathcal{A}}{\partial P_{\mu}}) \rightarrow -i[\hat{A}, \hat{B}] \equiv -i(\hat{A}\hat{B} - \hat{B}\hat{A})$ and $\mathcal{A} \rightarrow \hat{A}$. They can be also described by using the annihilation and creation operators $\hat{a}_{\mu}$ and $\hat{a}_{\mu}^{\dagger}$ which obey the standard bosonic commutation relations

$$[\hat{a}_{\mu}, \hat{a}_{\nu}^{\dagger}] = \delta_{\mu\nu}, \quad [\hat{a}_{\mu}, \hat{a}_{\nu}] = [\hat{a}_{\mu}^{\dagger}, \hat{a}_{\nu}^{\dagger}] = 0 \quad (2.3)$$

$\mu, \nu = 1, 2, \dots, N$. The CCR in forms (2.2) and (2.3) are related by a unitary matrix $U = 1/\sqrt{2} \begin{pmatrix} \mathbb{I}_N & i\mathbb{I}_N \\ \mathbb{I}_N & -i\mathbb{I}_N \end{pmatrix}$ such that if we define $\hat{O}^T = (\hat{a}_1, \hat{a}_2, \dots, \hat{a}_N, \hat{a}_1^{\dagger}, \hat{a}_2^{\dagger}, \dots, \hat{a}_N^{\dagger})$ then $\hat{O}_i = U_{ij} \hat{R}_j$.

The representation of the CCR up to unitaries is not unique. For instance, for a single mode in the Schrödinger representation each degree of freedom is embedded in $\mathcal{H} = \mathcal{L}^2(\mathbb{R})$, while the operators $\hat{q}$ and $\hat{p}$ act multiplicative and derivative respectively

$$\left. \begin{aligned} \hat{q} &= q \\ \hat{p} &= -i\frac{\partial}{\partial q} \end{aligned} \right\} \quad (2.4)$$

but also $\hat{q} = +i\frac{\partial}{\partial p}$, $\hat{p} = p$ is equally possible. In both representations the operators are unbounded.

A way to remove ambiguities (up to unitaries) and to treat with bounded operators is using the Weyl operators. The Weyl operator is defined as

$$\hat{W}_{\zeta} \equiv e^{i\zeta^T \cdot J \cdot \hat{R}} \quad (2.5)$$

where $\zeta^T = (\zeta_1, \zeta_2, \dots, \zeta_{2N})$. Then the Weyl operators satisfies the Weyl relation

$$\hat{W}_{\zeta} \hat{W}_{\eta} = e^{-\frac{i}{2}\zeta^T \cdot J \cdot \eta} \hat{W}_{\zeta+\eta}, \quad (2.6)$$

or in an analogous way

$$\hat{W}_{\zeta} \hat{W}_{\eta} = \hat{W}_{\eta} \hat{W}_{\zeta} e^{-i\zeta^T \cdot J \cdot \eta}. \quad (2.7)$$

Theorem 2.2.1 (Stone-von Neumann theorem) *Let $\hat{W}_1$ and $\hat{W}_2$ be two Weyl systems over a finite dimensional phase space ($N < \infty$). If the two Weyl systems are strongly continuous ² and irreducible ³ then they are equivalent (up to an unitary).*

² $\forall |\psi\rangle \in \mathcal{H} : \lim_{\zeta \rightarrow 0} \|\psi\rangle - \hat{W}_{\zeta} |\psi\rangle\| = 0$.

³ $\forall \zeta \in \mathbb{R}^{2N} : [\hat{W}_{\zeta}, \hat{A}] = 0 \Rightarrow \hat{A} \propto \mathbb{I}$.

According to the theorem 2.2.1 there exists only one equivalent representation of the Weyl relation.

The Weyl operator acts in the states as a translation in the phase space (displacements $e^{i\eta\hat{p}}|q\rangle = |q - \eta\rangle$ and kicks $e^{i\zeta\hat{q}}|q\rangle = e^{i\zeta q}|q\rangle$) as it can be checked by looking to its action onto an arbitrary position-momentum operator

$$\hat{W}_\zeta^\dagger \hat{R}_i \hat{W}_\zeta = \hat{R}_i - \zeta_i \mathbb{I} \quad (2.8)$$

2.3 Phase-space

Phase space formulation of Quantum Mechanics offers a framework in which Quantum phenomena can be described using as much classical language as allowed. It appeals naturally to one's intuition and can often provide useful physical insights. Furthermore, it requires dealing only with constant number equations and not with operators, which can be of significant practical advantage. This mathematical advantage arises here from the fact that the infinite-dimensional complex Hilbert space structure which is in principle a difficult object to work with, can be mapped into the linear algebra structure of the finite-dimensional real phase space. We will extend this map (2.3.1) and how to characterise states and operations in sections (2.4) and (2.3.2) respectively.

2.3.1 Phase space geometry

A system of N canonical degrees of freedom is described classically in a $2N$ -dimensional real vector space ⁴ $V \simeq \mathbb{R}^{2N}$. Together with the symplectic form it defines a symplectic real vector space (the phase space) $\Omega \simeq \mathbb{R}^{2N}$. The phase space is naturally equipped with a complex structure and can be identified with a complex Hilbert space $\mathcal{H}_\Omega \simeq \mathbb{C}^N$. If $\langle | \rangle$ stands for the scalar product in $\mathcal{H}_\Omega$ and $\langle | \rangle_J$ for the symplectic scalar product in V their connection reads

$$\langle \eta | \zeta \rangle = \langle J\eta | \zeta \rangle_J + i \langle \eta | \zeta \rangle_J \quad (2.9)$$

where $\eta = (q, p) \in V$ while $\eta = q + ip \in \mathcal{H}_\Omega$ such that any orthonormal basis in $\mathcal{H}_\Omega$ leads to a canonical basis in V . Moreover, any unitary operator (which preserve the scalar product) acting on $\mathcal{H}_\Omega$ leads to a symplectic operation S in the phase space in such a way that the symplectic scalar product is also preserved. The inverse is also true provided that the symplectic operation commutes with the symplectic matrix J .

2.3.2 Symplectic operations

We hence define symplectic canonical transformation S as these ones which preserve the CCR and therefore leave the basic kinematic rules unchanged. That is, if we

⁴They are isomorphic (there exist a bijective morphism between the two groups).

transform our quadrature operators $\hat{R}_S = S \cdot \hat{R}$ still equation (2.2) is fulfilled. In a totally equivalent way we can define symplectic transformation as the ones which preserve the symplectic scalar product and therefore ⁵

$$S^T \cdot J \cdot S = J. \quad (2.10)$$

The set of real $2N \times 2N$ matrices S satisfying the above condition form the symplectic group $Sp(2N, \mathbb{R})$. To construct the affine symplectic group we just need to add also the phase space translations whose group generators are $\hat{G}_i^{(0)} = J_{ij} \hat{R}_j$. Apart from that, the group generators of the representation of $Sp(2N, \mathbb{R})$ which physically corresponds to the Hamiltonians which perform the symplectic transformations on the states are of the form $\frac{1}{2} \{ \hat{R}_i, \hat{R}_j \}$. This corresponds to hermitian Hamiltonians of quadratic order in the canonical operators. When rewriting them in terms of creation / annihilation operators we can divide it into two groups.

Compact generators (passive):

$$\hat{G}_{\mu\nu}^{(1)} = i \frac{(\hat{a}_\mu^\dagger \hat{a}_\nu - \hat{a}_\nu^\dagger \hat{a}_\mu)}{2}, \quad \hat{G}_{\mu\nu}^{(2)} = \frac{(\hat{a}_\mu^\dagger \hat{a}_\nu + \hat{a}_\nu^\dagger \hat{a}_\mu)}{2}$$

and non-Compact generators (active):

$$\hat{G}_{\mu\nu}^{(3)} = i \frac{(\hat{a}_\mu^\dagger \hat{a}_\nu^\dagger - \hat{a}_\nu \hat{a}_\mu)}{2}, \quad \hat{G}_{\mu\nu}^{(4)} = \frac{(\hat{a}_\mu^\dagger \hat{a}_\nu^\dagger + \hat{a}_\nu \hat{a}_\mu)}{2}.$$

The passive ones are generators which commute with all the number operators $\hat{n}_\mu \equiv \hat{a}_\mu^\dagger \hat{a}_\mu$ and so they preserve the total number, in this sense they are passive. In case our system corresponds to modes of the electromagnetic field, then, what is being preserved is the total number of photons. They can be implemented optically by only using beam splitters, phase shifts and mirrors and conversely, only using them we can implement any Hamiltonian constructed by a linear combination of the compact generators. With all the generators we then have enough to generate all the unitaries, $\hat{U}_\lambda = e^{i\lambda \cdot \hat{G}}$.

As the simplest example for, one mode ($N = 1$) we have the phase shift operator

$$\hat{U}_\theta = e^{i\theta \hat{a}^\dagger \hat{a}} \quad (2.11)$$

which amounts to the symplectic operation in phase space

$$S_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}. \quad (2.12)$$

On the other hand we have the active ones, they change the energy of the state. The most important one is the single mode squeezing operator ($N = 1$), whose unitary expression for a fixed angle $\phi = 0$ and squeezing parameter $r > 0$ reads

$$\hat{U}_r = e^{\frac{r}{2}(\hat{a}^2 - \hat{a}^{\dagger 2})} \quad (2.13)$$

⁵From now on we neglect the subscript N in symplectic matrix.

while its symplectic operation in phase space reads

$$S_r = \begin{pmatrix} e^{-r} & 0 \\ 0 & e^r \end{pmatrix}. \quad (2.14)$$

Finally phase space translations for one mode whose unitary is $\hat{U}_\alpha = e^{\alpha\hat{a}^\dagger - \alpha^*\hat{a}}$ where $\alpha = \frac{(q_0 + ip_0)}{\sqrt{2}}$, amounts to the symplectic operation in phase space as $s_\alpha = \begin{pmatrix} q_0 \\ p_0 \end{pmatrix}$. The most important non-trivial unitaries for two modes ($N = 2$) are beam splitters $\hat{U}_{BS} = e^{\frac{\theta}{2}(\hat{a}_1\hat{a}_2^\dagger - \hat{a}_1^\dagger\hat{a}_2)}$ (at fixed angle $\phi = 0$, reflectivity $R = \sin^2 \theta/2$ and transmitivity $T = \cos^2 \theta/2$) and two mode squeezings $\hat{U}_{TMS} = e^{r(\hat{a}_1\hat{a}_2 - \hat{a}_1^\dagger\hat{a}_2^\dagger)}$ that amounts to

$$S_{BS} = \begin{pmatrix} \cos \theta/2 & 0 & \sin \theta/2 & 0 \\ 0 & \cos \theta/2 & 0 & \sin \theta/2 \\ -\sin \theta/2 & 0 & \cos \theta/2 & 0 \\ 0 & -\sin \theta/2 & 0 & \cos \theta/2 \end{pmatrix} \quad (2.15)$$

and

$$S_{TMS} = \begin{pmatrix} \cosh r & 0 & \sinh r & 0 \\ 0 & \cosh r & 0 & -\sinh r \\ \sinh r & 0 & \cosh r & 0 \\ 0 & -\sinh r & 0 & \cosh r \end{pmatrix}. \quad (2.16)$$

2.4 Probability Distribution functions

One of the most important tools of the phase space formulation of Quantum Mechanics are the phase space probability distribution functions. The best known and widely used is the Wigner distribution function, but there is not a unique way of defining a quantum phase space distribution function. In fact, several distribution functions with different properties, rules of association and operator ordering can also be well defined. For instance sometimes normal ordered (P-function), antinormal ordered (Q-function), generalised antinormal ordered (Husimi-function),... distributions can be more convenient depending on the problem being considered. In this dissertation we are only going to work with the totally symmetrical ordered (Weyl ordered) one, the Wigner distribution function.

Due to the fact that a joint probability at a fixed position $\hat{q}$ and momentum $\hat{p}$ point is not allowed in Quantum Mechanics by Heisenberg uncertainty theorem, the quantum phase space distribution function should, therefore, be considered simply as a mathematical tool that facilitates quantum calculations. Joint probabilities can be negative, so that one deals with quasiprobability distributions as long as it yields a correct description of physically observable quantities.

2.4.1 Quantum states

At the level of density operators, $\hat{\rho}$ defines a quantum state iff it satisfies the following properties

$$\text{tr}\hat{\rho} = 1, \quad \hat{\rho} \geq 0 \quad [\Rightarrow \quad \hat{\rho}^\dagger = \hat{\rho}]. \quad (2.17)$$

This operator belongs to the bounded linear operators Hilbert space $\mathcal{B}(\mathcal{H})$. For pure states the ket and wave function formulation is enough to describe our state. There, a state lives in a Hilbert space $\mathbb{C}^d$ for qudits, (a system of discrete variables) or $\mathcal{L}^2(\mathbb{R}^N)$ for N modes, (a system of continuous variables).

For systems of continuous variables, the Wigner distribution function gives a complete description of the state. Given a state $\hat{\rho}$ (a single mode) we define the Wigner distribution function as ⁶

$$\mathcal{W}_\rho(q, p) = \frac{1}{\pi} \int dx \langle q+x | \hat{\rho} | q-x \rangle e^{-2ipx}. \quad (2.18)$$

This transformation is called Weyl-Fourier transformation and it gives the bridge between density operators and distribution functions. Sometimes, for computational reasons it is better to compute first the characteristic distribution function which is obtained through

$$\chi_\rho(\zeta, \eta) = \text{tr}\{\hat{\rho} \hat{W}_{(\zeta, \eta)}\}. \quad (2.19)$$

The above two distribution functions are fully equivalent in the sense of describing completely our quantum state and are related by a Symplectic-Fourier transform

$$\mathcal{W}_\rho(q, p) = \frac{1}{(2\pi)^2} \int d\zeta \int d\eta \chi_\rho(\zeta, \eta) e^{-i\zeta p + i\eta q} = \frac{1}{2\pi} \mathcal{SFT}\{\chi_\rho(\zeta, \eta)\}, \quad (2.20)$$

$$\chi_\rho(\zeta, \eta) = \int dq \int dp \mathcal{W}_\rho(q, p) e^{i\zeta p - i\eta q} = 2\pi \mathcal{SFT}^{-1}\{\mathcal{W}_\rho(q, p)\}. \quad (2.21)$$

The Weyl-Fourier transformation is invertible and it provides a way to recover our density operator from both distribution functions

$$\begin{aligned} \hat{\rho} &= \frac{1}{2\pi} \int dq \int dp \int d\zeta \int d\eta \mathcal{W}_\rho(q, p) e^{-i\zeta p + i\eta q} \hat{W}_{(-\zeta, -\eta)} = \\ &= \frac{1}{2\pi} \int d\zeta \int d\eta \chi_\rho(\zeta, \eta) \hat{W}_{(-\zeta, -\eta)}. \end{aligned} \quad (2.22)$$

At this level, $\mathcal{W}$ (and χ) defines a quantum state iff they satisfy the following properties

$$\int dq \int dp \mathcal{W}(q, p) = 1, \quad \int dq \int dp \mathcal{W}(q, p) \mathcal{W}_p(q, p) \geq 0 \quad [\Rightarrow \quad \mathcal{W}^*(q, p) = \mathcal{W}(q, p)] \quad (2.23)$$

⁶For pure states the definition gets simplified to $\mathcal{W}_\rho(q, p) = \frac{1}{\pi} \int dx e^{-2ipx} \psi^*(q-x) \psi(q+x)$.

for all pure states $\mathcal{W}_p$ and

$$\chi(0,0) = 1, \quad \sum_{i,j=1}^{2N} a_i^* a_j \chi(\zeta_j - \zeta_i) e^{\frac{i}{2}(\zeta_i^T \cdot J \cdot \zeta_j)} \geq 0 \quad [\Rightarrow \quad \chi^*(\zeta, \eta) = \chi(-\zeta, -\eta)] \quad (2.24)$$

for all $a_{i,j} \in \mathbb{R}$. This can be shown using the following theorem.

Theorem 2.4.1 (Quantum Bochner-Khinchin theorem) *For $\chi(\eta)$ to be a characteristic function of a quantum state the following conditions are necessary and sufficient*

- 1.) $\chi(0) = 1$ and $\chi(\eta)$ is continuous at $\eta = 0$,
- 2.) $\chi(\eta)$ is J -positive (symplectic-positive defined).

2.4.2 Properties of the Wigner distribution

Properties

i) *Quasidistribution:* It is real valued quasidistribution because it admit negatives values (a Quantum Mechanics signature).

ii) *T-symmetry:* It has time symmetry

$$t \rightarrow -t \iff \mathcal{W}(q, p, t) \rightarrow \mathcal{W}(q, -p, t). \quad (2.25)$$

iii) *X-symmetry:* It has space symmetry

$$q \rightarrow -q \iff \mathcal{W}(q, p, t) \rightarrow \mathcal{W}(-q, -p, t). \quad (2.26)$$

iv) *Galilei invariant:* It is Galilei invariant

$$q \rightarrow q - a \iff \mathcal{W}(q, p, t) \rightarrow \mathcal{W}(q + a, p, t). \quad (2.27)$$

v) *T-evolution:* The equation of motion for each point in the phase space is classical in the absence of forces ⁷

$$\frac{d\hat{p}}{dt} = -\frac{1}{i}[\hat{p}, \hat{H}] \iff \frac{\partial \mathcal{W}(q, p, t)}{\partial t} = -\frac{p}{m} \frac{\partial \mathcal{W}(q, p, t)}{\partial q}. \quad (2.29)$$

vi) *Bounded:* It is bounded ⁸

$$|\mathcal{W}(q, p)| \leq \frac{1}{\pi}. \quad (2.30)$$

⁷Remember that when we are speaking about states of light m has to be interpret as permittivity of vacuum ϵ_0 while there is a minus sign difference with Heisenberg's equation of motion.

$$\frac{d\hat{A}}{dt} = \frac{1}{i}[\hat{A}, \hat{H}]. \quad (2.28)$$

⁸Use Schwarz's inequality $|\langle \psi_1 | \psi_2 \rangle|^2 \leq \langle \psi_1 | \psi_1 \rangle \langle \psi_2 | \psi_2 \rangle$ at the density operator level *i.e.* $0 \leq \text{tr}(\rho_1 \rho_2)^n \leq \text{tr}(\rho_1)^n \text{tr}(\rho_2)^n$. (*no es ven be la dem*)

vii) *Normalised*: It is well normalised

$$\int dq \int dp \mathcal{W}(q, p) = 1. \quad (2.31)$$

viii) *Quantum marginal distributions*: It possesses good marginal distributions ⁹

$$\int dp \mathcal{W}(q, p) = \langle q | \hat{\rho} | q \rangle \geq 0 \quad (2.32)$$

$$\int dq \mathcal{W}(q, p) = \langle p | \hat{\rho} | p \rangle \geq 0. \quad (2.33)$$

ix) *Complete orthonormal set*: The set of functions $\mathcal{W}_{nm}(q, p)$ form a complete orthonormal set (if $\psi_n(q)$ are already a set)

$$\int dq \int dp \mathcal{W}_{nm}^*(q, p) \mathcal{W}_{n'm'}(q, p) = \frac{1}{2\pi} \delta_{nn'} \delta_{mm'} \quad (2.34)$$

$$\sum_{n,m} \mathcal{W}_{nm}^*(q, p) \mathcal{W}_{nm}(q', p') = \frac{1}{2\pi} \delta(q - q') \delta(p - p') \quad (2.35)$$

where

$$\mathcal{W}_{nm}(q, p) = \frac{1}{\pi} \int dx e^{-2ipx} \psi_n^*(q - x) \psi_m(q + x). \quad (2.36)$$

2.4.3 The generating function of a Classical probability distribution

Denoting by y (x) a random variable which can be discrete $y \in \{y_i\}$ (or continuous $x \in [a, b]$) and its corresponding (density) probability $p(y_i)$ ($p(x)$), we can establish the normalisation constrain as

$$\left. \begin{aligned} \sum_{y_i} p(y_i) &= 1 \\ \int_a^b p(x) dx &= 1 \end{aligned} \right\}. \quad (2.37)$$

Of relevant importance given a probability distribution are the following quantities:

i) *Mean value of $u(x)$* : $E[u(x)] = \int u(x) p(x) dx.$

ii) *Moment of order m respect point c of x* : $\alpha_c^m = E[(x - c)^m].$

iii) *Mean value of x* : $\mu = \alpha_0^1 = E[x].$

iv) *Standard deviation of x* : ¹⁰ $\sigma = \sqrt{\alpha_\mu^2} = \sqrt{E[(x - \mu)^2]}.$

v) *Covariance of x_i and x_j* : ¹¹ $C_{ij} = \text{cov}(x_i, x_j) = E[(x_i - \mu_i)(x_j - \mu_j)]$

where $i, j = 1, 2, \dots, 2N.$

⁹For pure state they correspond to the square modulus of the wave function in position $|\psi(q)|^2$ and in momentum $|\tilde{\psi}(p)|^2$ representation.

¹⁰Square root of the variance.

¹¹Here subindex i, j labels all the possible variables of the distribution, when they are equal, C_{ii} corresponds to the variance of the variable $x_i.$

Theorem 2.4.2 (Taylor's theorem) *Any well behaved distribution function can be reconstructed by its (in general) infinite moments.*

This theorem, of considerable importance, tell us that any distribution $p(x)$ can be retrieved only by its moments α_c^m . We define the vector d and the matrix C called mean vector and covariance matrix by

$$\left. \begin{aligned} C &= [[\text{cov}(x_i, x_j)]] \\ d &= [[\mu_i]] \end{aligned} \right\}. \quad (2.38)$$

What is more important is that d and C encode all the information of 1st and 2nd moments.

If we define the generating function of the distribution function by a Laplace transformation (provided it exists)

$$M(\eta) = \mathcal{LT}\{p(x)\} = E[e^{x\eta}] \quad (2.39)$$

all moments can be obtained by subsequently differentiating the generating function

$$\alpha_0^m = \frac{\partial^{(m)} M(\eta)}{\partial \eta^m} \Big|_{\eta=0}. \quad (2.40)$$

2.4.4 The generating function of a quasi-probability distribution

In the same way as in Classical Probability where all the moments of a distribution characterise the distribution, the Wigner quasidistribution function is fully characterised by its moments.

To adapt the classical formalism to the quantum Wigner quasidistribution function we have to introduce the following transcription $\eta \rightarrow i\eta$, $M \rightarrow \chi$, $\mathcal{LT} \rightarrow \mathcal{FT}$.

We then define the generating function of the Wigner distribution (characteristic function) by a Fourier transformation, which always exists, because the Wigner distribution is an integrable function. In general it is complex and reads

$$\chi(\eta) = \mathcal{FT}\{\mathcal{W}(x)\} = E[e^{ix\eta}] \quad (2.41)$$

then all moments can be obtained by subsequently differentiating the generating function

$$\beta_0^m = \frac{1}{i^m} \frac{\partial^{(m)} \chi(\eta)}{\partial \eta^m} \Big|_{\eta=0}. \quad (2.42)$$

Analogously, we define, given a quantum Wigner distribution function the displacement vector (DV) d (a $2N$ real vector) and the covariance matrix (CM) γ (a $2N \times 2N$ real symmetric matrix). The DV contains the information of the first moments and in general plays no role, by the space symmetry only relative DVs are of physical meaning. The CM is much more richer, it contains information (up to

second moments) about the purity, entanglement, ... The CM to describe a physical state must be symplectic-positive defined

$$\gamma + iJ \geq 0. \quad (2.43)$$

2.5 Gaussian states

Among all the CV systems Gaussian states are of greatest importance. The Gaussian distribution is simple, it is the limit of many others and appears in a great variety of different conditions. In order to give a motivation we state here one of the most important theorems (together with the Law of Large Numbers) of the Theory of the Classical Probability and Statistics, the Central Limit Theorem.

Theorem 2.5.1 (Central limit theorem) *Suppose we have n independent random variables $x_1, x_2, \dots, x_n$ which are all distributed with a mean value μ and a standard deviation σ (each of them can have different arbitrary distribution functions $p_i(x_i)$). In the limit $n \rightarrow \infty$ the arithmetic mean $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ is Gaussian (or normal) distributed with mean value μ and standard deviation $\frac{\sigma}{\sqrt{n}}$ i.e. $\bar{p}(\bar{x}) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(\bar{x}-\mu)^2}{2\sigma^2}}$.*

Another way to see the importance of Gaussian probability distributions is encoded in the following theorem

Theorem 2.5.2 (Marcinkiewicz's theorem) *If we define the cumulant generating function as $K(\eta) = \ln M(\eta)$, then, either the cumulant is a polynomial of order 2 or it has infinite terms.*

Lemma 2.5.1 (Gaussianity lemma) *As a consequence then $p(x)$ is a Gaussian (non-Gaussian) distribution iff the cumulant is a polynomial of order 2 (∞).*

In general all moments are necessary but as long as we are concern with only Gaussian distributions, 1st and 2nd moments are sufficient, in fact all other higher moments can be rewritten in terms of them. This is a consequence of the theorem 2.5.2.

2.5.1 Displacement Vector (DV) and Covariance Matrix (CM)

An important class of quantum states are Gaussian states. They can be defined as all quantum states whose Wigner distribution function is Gaussian. Thus the DV and the CM are enough to describe them. Analogously to the classical case, it is straightforward to obtain the moments of order β_0^m of a distribution by differentiating the generating function. Computing 1st and 2nd moments, through (2.19) we get ¹²

¹²Notice: $(\frac{\partial}{\partial \eta_i} e^{i\eta^T \cdot \hat{R}})|_{\eta_i=0} = i\hat{R}_i$ and $(\frac{\partial^2}{\partial \eta_i \partial \eta_j} e^{i\eta^T \cdot \hat{R}})|_{\eta_{i,j}=0} = \frac{1}{2}[\hat{R}_i, \hat{R}_j] - \hat{R}_i \hat{R}_j = -\frac{1}{2}\{\hat{R}_i, \hat{R}_j\}$ where we have used Cambell-Hausdorff formula $e^{\hat{A}+\hat{B}} = e^{\hat{A}} e^{\hat{B}} e^{-\frac{1}{2}[\hat{A}, \hat{B}]}$ (when $[\hat{A}, \hat{B}] \propto \mathbb{I}$).

$$\beta_{0,i}^1 = -i \frac{\partial}{\partial \eta_i} \chi(\eta)|_{\eta=0} = \text{tr}(\hat{\rho} \hat{R}'_i) \quad (2.44)$$

$$\beta_{0,ij}^2 = (-i)^2 \frac{\partial^2}{\partial \eta_i \partial \eta_j} \chi(\eta)|_{\eta=0} = \frac{1}{2} \text{tr}(\hat{\rho} \{\hat{R}'_i, \hat{R}'_j\}) = \text{tr}(\hat{\rho} \hat{R}'_i \hat{R}'_j) - \frac{i}{2} J_{ij} \quad (2.45)$$

where $\hat{R}'_i = J \hat{R}_i$.

Finally we define the DV and the CM as ¹³

$$d_i = \text{tr}(\hat{\rho} \hat{R}_i) \quad (2.46)$$

$$\begin{aligned} \gamma_{ij} &= \text{tr}(\hat{\rho} \{\hat{R}_i - d_i \hat{\mathbb{I}}, \hat{R}_j - d_j \hat{\mathbb{I}}\}) = 2 \text{tr}[\hat{\rho} (\hat{R}_i - d_i \hat{\mathbb{I}}) (\hat{R}_j - d_j \hat{\mathbb{I}})] - i J_{ij} = \\ &= 2 \text{Re}\{\text{tr}[\hat{\rho} (\hat{R}_i - d_i \hat{\mathbb{I}}) (\hat{R}_j - d_j \hat{\mathbb{I}})]\} \end{aligned} \quad (2.47)$$

It is important to remark here that symplectic operations at the level of the DV and CM act in such a way that any unitary $\hat{U}_S$ maps to the following transformation $\gamma_S = S \cdot \gamma \cdot S^T$ and $d_S = d + s$ where S stands for an element of the symplectic group, while s stands for a phase space translation.

With these definitions it can be shown that the Wigner distribution of any Gaussian state can be written in terms of the DV and CM through ¹⁴

$$\mathcal{W}(\zeta) = \frac{1}{\pi^N \sqrt{\det \gamma}} e^{-(\zeta-d)^T \cdot \frac{1}{\gamma} \cdot (\zeta-d)} \quad (2.48)$$

while its symplectic-Fourier transform reads

$$\chi(\eta) = e^{i\eta^T \cdot J \cdot d - \eta^T \cdot J^T \frac{1}{4} J \cdot \eta} = e^{i\eta^T \cdot d' - \eta^T \cdot \frac{\gamma'}{4} \cdot \eta} \quad (2.49)$$

where $d'_i = J_{ij} d_j$ and $\gamma'_{ij} = J_{ik}^T \gamma_{kl} J_{lj}$.

Theorem 2.5.3 (Minimum uncertainty states theorem) *Equality in Heisenberg's uncertainty theorem is attained iff the state is a pure Gaussian state i.e. a rotated squeezed coherent state, $|\psi\rangle = \hat{U}_\theta \hat{U}_r \hat{U}_\alpha |0\rangle$.*

All pure Gaussian states of one mode, characterised by its γ (and if necessary by d), can be obtained from the vacuum state by an arbitrary displacement+squeezing+rotation in the phase space. These states, by theorem 2.5.3, are minimum uncertainty states.

¹³For pure states $d_i = \langle \hat{R}_i \rangle_\rho$ and $\gamma_{ij} = \langle \{\hat{R}_i - d_i \hat{\mathbb{I}}, \hat{R}_j - d_j \hat{\mathbb{I}}\} \rangle_\rho = \langle \{\hat{R}_i, \hat{R}_j\} \rangle_\rho - 2 \langle \hat{R}_i \rangle_\rho \langle \hat{R}_j \rangle_\rho$, where, by the anticommutator definition, we see a factor 2 of difference with the classical analog and so $\gamma_{ij} \sim 2C_{ij}$. The diagonal terms can be rewritten in terms of the uncertainties as $\gamma_{ii} = 2(\Delta R_i)_\rho^2$ where as usual $(\Delta A)_\psi = \sqrt{\langle \hat{A}^2 \rangle_\psi - (\langle \hat{A} \rangle_\psi)^2}$.

¹⁴We see here that from (2.30) $\max[\mathcal{W}(\zeta)] = \mathcal{W}(d) = \frac{1}{\pi^N \sqrt{\det \gamma}} \leq \frac{1}{\pi^N}$ where the equality holds for pure states only.

Instead, mixed Gaussian states of one mode can be all obtained from a thermal state by an arbitrary displacement+squeezing+rotation.

As the cornerstone examples of Gaussian states, we have the vacuum, coherent, squeezed and thermal states.

**Vacuum:* $|0\rangle$

$$\gamma_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad d_0 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (2.50)$$

**(Pure) Coherent:* ¹⁵ $|\alpha\rangle = \hat{\mathcal{D}}(\alpha)|0\rangle = \hat{U}_\alpha|0\rangle$

$$\gamma_\alpha = S_\alpha \gamma_0 S_\alpha^T = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad d_\alpha = d_0 + s_\alpha = \begin{pmatrix} q_0 \\ p_0 \end{pmatrix}, \quad (2.51)$$

where $\alpha = \alpha_R + i\alpha_I = \frac{q_0 + ip_0}{\sqrt{2}}$.

**(Pure) Squeezed:* $|r\rangle = \hat{\mathcal{S}}(r)|0\rangle = \hat{U}_r|0\rangle$

$$\gamma_r = S_r \gamma_0 S_r^T = \begin{pmatrix} e^{-2r} & 0 \\ 0 & e^{2r} \end{pmatrix}, \quad d_r = d_0 + s_r = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (2.52)$$

**(Mixed) Thermal:* $\hat{\rho}_\beta = \frac{1}{\pi M} \int d^2\alpha |\alpha\rangle\langle\alpha| e^{-|\alpha|^2/M}$

$$\gamma_\beta = \begin{pmatrix} 2M+1 & 0 \\ 0 & 2M+1 \end{pmatrix}, \quad d_\beta = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (2.53)$$

where $M = \frac{1}{e^\beta - 1} \geq 0$ being β the inverse temperature.

2.5.2 Hilbert space, phase space and DV&CM connection

We have already shown how to describe quantum states and operations at the different levels *i.e.* Hilbert space, phase space and DV&CM. Two main connections are needed still to perform calculations in the phase space: the ordering of the operators and the metric between them.

The Weyl association rule tells us about the ordering operators. Provided we are using with the Wigner distribution, which is symmetrical ordered, when working with observables we have to take into account that as we are in the phase space and we have avoided its operator character we have to symmetrise them. The way we have to symmetrise is

$$e^{i\zeta\hat{q}+i\eta\hat{p}} \longrightarrow: e^{i\zeta\hat{q}+i\eta\hat{p}} := e^{i\zeta\hat{q}+i\eta\hat{p}} \longleftarrow e^{i\zeta q+i\eta p}, \quad (2.54)$$

where $: :$ stands for the symmetrical order. In general for a polynomial on q and p

¹⁵A Coherent state can alternatively be defined as the eigenstate of the annihilation operator, $\hat{a}|\alpha\rangle = \alpha|\alpha\rangle$. Coherent states form an *overcomplete* non-orthogonal ($\langle\alpha|\alpha'\rangle = \exp[-(|\alpha|^2 + |\alpha'|^2)/2 + \alpha^*\alpha']$) set base ($\frac{1}{\pi} \int d^2\alpha |\alpha\rangle\langle\alpha| = \mathbb{I}$) of vectors of the Hilbert space.

$$\hat{q}^n \hat{p}^m \longrightarrow \hat{q}^n \hat{p}^m := \frac{1}{2^n} \sum_{r=0}^n \binom{n}{r} \hat{q}^r \hat{p}^m \hat{q}^{n-r} = \frac{1}{2^m} \sum_{r=0}^m \binom{m}{r} \hat{p}^r \hat{q}^n \hat{p}^{m-r} \longleftrightarrow q^n p^m. \quad (2.55)$$

As an example take the observable $\mathcal{Q}P$, its quantum associated operator is of course $\hat{q}\hat{p}$. We know that $\hat{q}$ and $\hat{p}$ do not commute but in the phase space qp and pq are functionally treated in the same way. Imagine we need to find its average value, we have then to remove the ambiguity by totally symmetrising. The recipe is $\hat{q}\hat{p} \longrightarrow \hat{q}\hat{p} := \frac{\hat{q}\hat{p} + \hat{p}\hat{q}}{2} \longleftrightarrow qp$. And so the average to be performed is

$$\langle \mathcal{Q}P \rangle = \langle \hat{q}\hat{p} \rangle_\rho = \langle qp + i/2 \rangle_{\mathcal{W}} \quad (2.56)$$

because

$$\hat{q}\hat{p} = \frac{\hat{q}\hat{p} + \hat{p}\hat{q}}{2} + i/2. \quad (2.57)$$

More important and relevant averages concern the moments which can be obtained *via* the Wigner distribution as

$$d_i = \text{tr}(\hat{\rho} \hat{R}_i) = \int d^{2N} \zeta [\zeta_i] \mathcal{W}(\zeta) \quad (2.58)$$

$$\gamma_{ij} = \text{tr}(\hat{\rho} \{\hat{R}_i - d_i \hat{\mathbb{I}}, \hat{R}_j - d_j \hat{\mathbb{I}}\}) = \int d^{2N} \zeta [2(\zeta_i - d_i)(\zeta_j - d_j)] \mathcal{W}(\zeta). \quad (2.59)$$

Theorem 2.5.4 (Quantum Parseval theorem) *Let $\hat{W}_\zeta$ be a strongly continuous and irreducible Weyl system acting on the Hilbert space $\mathcal{H}_\Omega$ with phase space Ω . Then $\hat{A} \mapsto \text{tr}\{\hat{A}\hat{W}_\zeta\}$, with $\zeta \in \Omega$, is an isometric map from the Hilbert space $\mathcal{H}$ (Hilbert-Schmidt operators) onto the Hilbert space $\mathcal{L}^2(\Omega)$ (square-integrable measurable functions on Ω) such that*

$$\text{tr}(\hat{A}^\dagger \hat{B}) = \frac{1}{(2\pi)^N} \int d^{2N} \zeta \text{tr}\{\hat{A}\hat{W}_\zeta\}^* \text{tr}\{\hat{B}\hat{W}_\zeta\}. \quad (2.60)$$

This theorem is of capital importance, it follows from it how to compute the scalar product between operators

$$\begin{aligned} \text{tr}\{\hat{A}^\dagger \hat{B}\} &= \frac{1}{(2\pi)^N} \int d^{2N} \eta A^{X^*}(\eta) B^X(\eta) = \\ &= \frac{1}{(2\pi)^N} \int d^{2N} \zeta A^{\mathcal{W}}(\zeta) B^{\mathcal{W}}(\zeta), \end{aligned} \quad (2.61)$$

the trace of an operator ¹⁶

$$\text{tr}\{\hat{A}\} = A^X(0,0) = \frac{1}{(2\pi)^N} \int d^{2N} \zeta A^{\mathcal{W}}(\zeta), \quad (2.62)$$

¹⁶Use that $\mathbb{I}^{\mathcal{W}} = 1$ and $\mathbb{I}^X = (2\pi)^N \delta^{(2N)}(\eta)$ computed from eq. (2.66) and eq. (2.65).

and the expectation value of an observable

$$\begin{aligned}\langle \hat{A} \rangle_\rho &= \text{tr}\{\hat{\rho}\hat{A}\} = \frac{1}{(2\pi)^N} \int d^{2N}\eta \chi^*(\eta) A^\chi(\eta) = \\ &= \int d^{2N}\zeta \mathcal{W}(\zeta) A^\mathcal{W}(\zeta).\end{aligned}\tag{2.63}$$

To justify the above expression we just need to define properly the Fourier-Weyl transform as

$$\begin{aligned}\hat{A} &= \mathcal{F}\mathcal{W}\mathcal{T}\{A^\chi(\eta)\} = \frac{1}{(2\pi)^N} \int d^{2N}\eta A^\chi(\eta) \hat{W}_{-\eta} = \\ &= \frac{1}{(2\pi)^{2N}} \int d^{2N}\eta \int d^{2N}\zeta A^\mathcal{W}(\zeta) e^{i\zeta^T \cdot J \cdot \eta} \hat{W}_{-\eta},\end{aligned}\tag{2.64}$$

and its inverse

$$A^\chi(\eta) = \mathcal{F}\mathcal{W}\mathcal{T}^{-1}\{\hat{A}\} = \text{tr}\{\hat{A}\hat{W}_\eta\},\tag{2.65}$$

$$A^\mathcal{W}(\bar{\zeta}, \bar{\eta}) = 2^N \int d^N\lambda \langle \bar{\zeta} + \lambda | \hat{A} | \bar{\zeta} - \lambda \rangle e^{-2i\bar{\eta}\lambda}\tag{2.66}$$

where $\bar{\zeta}^T = (\zeta_1, \zeta_2, \dots, \zeta_N)$ idem for $\bar{\eta}$.

$$A^\mathcal{W}(\zeta) = \mathcal{S}\mathcal{F}\mathcal{T}\{A^\chi(\eta)\}^{17} = \frac{1}{(2\pi)^N} \int d^{2N}\eta A^\chi(\eta) e^{-i\zeta^T \cdot J \cdot \eta},\tag{2.67}$$

$$A^\chi(\eta) = \mathcal{S}\mathcal{F}\mathcal{T}^{-1}\{A^\mathcal{W}(\zeta)\}^{18} = \frac{1}{(2\pi)^N} \int d^{2N}\zeta A^\mathcal{W}(\zeta) e^{i\zeta^T \cdot J \cdot \eta}.\tag{2.68}$$

An important concept in Quantum Information is the fidelity $\mathcal{F}$ between quantum states. The one we adopt here is the so called Bures-Uhlmann fidelity and it is defined as follows

$$\mathcal{F}(\hat{\rho}_1, \hat{\rho}_2) = [\text{tr}\sqrt{\hat{\rho}_1^{1/2} \hat{\rho}_2 \hat{\rho}_1^{1/2}}]^2.\tag{2.69}$$

It is symmetric and normalised between 1 (equal states) and 0 (orthogonal states). Its definition is simplified when one of the two states is pure (say $\hat{\rho}_1$), in this case it converges to the Hilbert-Schmidt fidelity

$$\mathcal{F}(\hat{\rho}_1, \hat{\rho}_2) = \text{tr}(\hat{\rho}_1 \hat{\rho}_2) = \langle \psi_1 | \hat{\rho}_2 | \psi_1 \rangle.\tag{2.70}$$

In case both states are pure, then, the fidelity becomes simply the overlap between the two states

$$\mathcal{F}(\hat{\rho}_1, \hat{\rho}_2) = |\langle \psi_1 | \psi_2 \rangle|^2.\tag{2.71}$$

¹⁷Notice that $A^\mathcal{W} = (2\pi)^N \mathcal{W}$ if $\hat{A} = \hat{\rho}$ see eq. (2.18) (for normalisation convenience).

¹⁸Notice that $A^\chi = \chi$ if $\hat{A} = \hat{\rho}$ see eq. (2.19).

It is useful here to use theorem 2.5.4 to evaluate the Hilbert-Schmidt fidelity between two Gaussian state (at least when one is pure)¹⁹

$$\begin{aligned}\mathcal{F}(\hat{\rho}_1, \hat{\rho}_2) &= \text{tr}(\hat{\rho}_1 \hat{\rho}_2) = \left(\frac{1}{2\pi}\right)^N \int d^{2N} \eta \chi_1^*(\eta) \chi_2(\eta) = (2\pi)^N \int d^{2N} \zeta \mathcal{W}_1(\zeta) \mathcal{W}_2(\zeta) = \\ &= \frac{1}{\sqrt{\det(\frac{\gamma_1 + \gamma_2}{2})}} e^{-d^T (\frac{1}{\gamma_1 + \gamma_2}) d}\end{aligned}\tag{2.72}$$

where $\gamma_{1(2)}$ and $d_{1(2)}$ belongs to $\hat{\rho}_{1(2)}$, while $d = d_2 - d_1$.

Another important concept in Quantum Information is the purity $\mathcal{P}$ of a quantum state. The purity is defined as follows

$$\mathcal{P}(\hat{\rho}) = \text{tr}(\hat{\rho}^2)\tag{2.73}$$

It is normalised between 1 (pure states) and 0 (maximally mixed states). Also here using theorem 2.5.4 we can evaluate the purity of a Gaussian state²⁰

$$\mathcal{P}(\hat{\rho}) = (2\pi)^N \int d^{2N} \zeta [\mathcal{W}(\zeta)]^2 = \frac{1}{\sqrt{\det \gamma}}\tag{2.74}$$

2.6 Multipartite states and entanglement

If we want to treat the entanglement of a quantum state, first we need to introduce multipartite states. In general we have to extend the Hilbert space. At the level of density operators this means that we have to “tensor product” $\otimes$, the Hilbert space of each party *i.e.* $\mathcal{H} = \bigotimes_{k=1}^N \mathcal{H}_k$. The covariance matrix structure for Gaussian states turns to be simplified to a “direct sum” $\oplus$, of each party’s associated phase space *i.e.* $\Omega = \bigoplus_{k=1}^N \Omega_k$. This is reminiscent of the Quantum Parseval theorem, which transforms tensor product between density matrices to products of Wigner functions (and Characteristic functions) and at the same time direct sums of covariance matrices.

Therefore, the advantage of using Gaussian states is that we fully describe a state by a finite dimensional 2×2 matrix instead of its infinite dimensional density matrix. Additionally, dimensionality of the phase space increases slower, as dimensions are added instead of multiplied.²¹

2.6.1 Bipartite Gaussian states

Any bipartite Gaussian state can be written in a block structure as $\gamma = \begin{pmatrix} A & C \\ C^T & B \end{pmatrix}$, where $A = A^T$ and $B = B^T$.

¹⁹The second and third equality is true for all CV states.

²⁰The first equality is true for all CV states.

²¹Remember that $\dim(\hat{\rho}_1 \otimes \hat{\rho}_2) = \dim(\hat{\rho}_1) \dim(\hat{\rho}_2)$ while $\dim(\gamma_1 \oplus \gamma_2) = \dim(\gamma_1) + \dim(\gamma_2)$.

Lemma 2.6.1 (Normal form) *Every 1×1 mode Gaussian state can be transformed (by two local symplectic transformations) to*

$$\gamma = \begin{pmatrix} \lambda_a & 0 & k_x & 0 \\ 0 & \lambda_a & 0 & k_p \\ k_x & 0 & \lambda_b & 0 \\ 0 & k_p & 0 & \lambda_b \end{pmatrix}. \quad (2.75)$$

If we define the four invariants of an arbitrary state $\mathcal{A} = \det A$, $\mathcal{B} = \det B$, $\mathcal{C} = \det C$ and $\Upsilon = \det \gamma$, then the following holds

$$\begin{aligned} \lambda_a &= \sqrt{\mathcal{A}} \\ \lambda_b &= \sqrt{\mathcal{B}} \\ k_x &= \frac{1}{2}(\alpha - \sqrt{\alpha^2 - 4\mathcal{C}}) \\ k_p &= \frac{1}{2}(\alpha + \sqrt{\alpha^2 - 4\mathcal{C}}) \\ \alpha &= \sqrt{\left(\frac{(\sqrt{\mathcal{A}\mathcal{B}} + \mathcal{C})^2 - \Upsilon}{\sqrt{\mathcal{A}\mathcal{B}}}\right)} \end{aligned} \quad (2.76)$$

Lemma 2.6.2 (Standard form) *Every 1×1 mode Gaussian state can be transformed (by local quasi-free symplectic transformations) to*

$$\gamma = \begin{pmatrix} \lambda_a & 0 & k_x & 0 \\ 0 & \lambda_a & 0 & -k_p \\ k_x & 0 & \lambda_b & 0 \\ 0 & -k_p & 0 & \lambda_b \end{pmatrix}, \quad (2.77)$$

where $\lambda_a, \lambda_b \geq 1$ and $k_x \geq |k_p|$.

A Gaussian state in the Standard form is called symmetric if $\lambda_a = \lambda_b$, and fully symmetric if it is symmetric and in addition $k_x = k_p$.

2.6.2 Entanglement of Gaussian states

For discrete variable systems an important separability criteria based on the partial transpose (time reversal) exists.

Lemma 2.6.3 (NPPT Peres criteria) *Given a bipartite state $\hat{\rho}$, if it has non-positive partial transpose ($\hat{\rho}^{T_A} \not\geq 0 \Rightarrow \hat{\rho}^{T_B} \not\geq 0$), then $\hat{\rho}$ is entangled.*

Lemma 2.6.4 (NPPT Horodecki criteria) *In $\mathbb{C}^2 \otimes \mathbb{C}^2$ and $\mathbb{C}^2 \otimes \mathbb{C}^3$ given a bipartite state $\hat{\rho}$, it is entangled iff it has non-positive partial transpose ($\hat{\rho}^{T_A} \not\geq 0 \Rightarrow \hat{\rho}^{T_B} \not\geq 0$).*

For continuous variable states, Peres criteria also holds while Horodecki criteria is true provided our state is composed of $1 \times N$ modes. In particular for Gaussian states, time reversal is very easy to implement at the covariance matrix level. If $\hat{T}$ is the reversal operator then $S_T = \theta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ is the symplectic operations in phase space. So we can rewrite the lemma 2.6.4 for Gaussian states as

Lemma 2.6.5 (NPPT Horodecki's criteria) *For $1 \times N$ modes given a bipartite Gaussian state γ , it is entangled iff it has non-positive partial transpose ($\theta_A \gamma \theta_A^T + iJ \not\geq 0 \Rightarrow \theta_B \gamma \theta_B^T + iJ \not\geq 0$).*

Concerning entanglement measures we use as an entanglement measure for pure state the Entropy of entanglement and for mixed ones the Logarithmic negativity.

(Pure states) Entropy of entanglement:

$$E_S(\hat{\rho}) = S(\hat{\rho}_A) = -\text{tr}(\hat{\rho}_A \log \hat{\rho}_A), \quad (2.78)$$

where S is the von Neumann Entropy ²² $S(\hat{\rho}) = -\text{tr}(\hat{\rho} \log \hat{\rho})$, and $\hat{\rho}_A$ is the trace over B defined as $\hat{\rho}_A = \text{tr}_B(\hat{\rho})$. For any state it reduces (in terms of the Schmidt coefficients) to

$$E_S(\hat{\rho}) = -\sum_{i=0}^{\infty} C_i^2 \log C_i^2, \quad (2.79)$$

while for Gaussian states,

$$E_S(\gamma) = -\sum_{i=1}^{N_A} \left[\left(\frac{\mu_i + 1}{2} \right) \log \left(\frac{\mu_i + 1}{2} \right) - \left(\frac{\mu_i - 1}{2} \right) \log \left(\frac{\mu_i - 1}{2} \right) \right], \quad (2.80)$$

where $\{\mu_i\} = \text{spec}(-iJ\gamma_A)$.

(Mixed states) Logarithmic negativity (additive monotone):

$$E_N(\hat{\rho}) = LN(\hat{\rho}) = \log \|\hat{\rho}^{TA}\|_1, \quad (2.81)$$

where $\|\cdot\|_1$ is the trace norm defined as $\|\hat{\rho}\|_1 = \text{tr}|\hat{\rho}| = \text{tr}\sqrt{\hat{\rho}^T \hat{\rho}} = \sum \text{singular values}(\hat{\rho})$. For Gaussian states,

$$E_N(\gamma) = -\sum_{i=1}^N \log[\min(\tilde{\mu}_i, 1)], \quad (2.82)$$

where $\{\tilde{\mu}_i\} = \text{spec}(-iJ\gamma^{TA})$.

²²The logarithm is in base 2.

2.7 Appendix of integrals

Gaussian integrals:

$$\int e^{-\zeta^T \cdot A \cdot \zeta} d^{2N} \zeta = \frac{\pi^N}{\sqrt{\det A}} \quad (2.83)$$

$$\int e^{-\zeta^T \cdot A \cdot \zeta - b^T \cdot \zeta} d^{2N} \zeta = \frac{\pi^N}{\sqrt{\det A}} e^{b^T \cdot \frac{1}{4A} \cdot b} \quad (2.84)$$

Non-Gaussian integrals:

$$\int \zeta_i^{m_i} \zeta_j^{m_j} \zeta_k^{m_k} \dots e^{-\zeta^T \cdot A \cdot \zeta} d^{2N} \zeta = (-1)^{m_i+m_j+m_k+\dots} \frac{\pi^N}{\sqrt{\det A}} \frac{\partial^{m_i}}{\partial b_i^{m_i}} \frac{\partial^{m_j}}{\partial b_j^{m_j}} \frac{\partial^{m_k}}{\partial b_k^{m_k}} \dots e^{b^T \cdot \frac{1}{4A} \cdot b} \Big|_{b_i=0, b_j=0, b_k=0, \dots} \quad (2.85)$$

$$\begin{aligned} & \int \zeta_i^{m_i} \zeta_j^{m_j} \zeta_k^{m_k} \dots e^{-\zeta^T \cdot A \cdot \zeta - c^T \cdot \zeta} d^{2N} \zeta = \\ & = (-1)^{m_i+m_j+m_k+\dots} \frac{\pi^N}{\sqrt{\det A}} \frac{\partial^{m_i}}{\partial b_i^{m_i}} \frac{\partial^{m_j}}{\partial b_j^{m_j}} \frac{\partial^{m_k}}{\partial b_k^{m_k}} \dots e^{(b+c)^T \cdot \frac{1}{4A} \cdot (b+c)} \Big|_{b_i=0, b_j=0, b_k=0, \dots} \end{aligned} \quad (2.86)$$

Useful integrals:

$$\int \mathcal{W}(\zeta) d^{2N} \zeta = 1 \quad (2.87)$$

$$\int (\mathcal{W}(\zeta))^2 d^{2N} \zeta = \frac{1}{(2\pi)^N \sqrt{\det \gamma}} \quad (2.88)$$

$$\int \chi(\eta) d^{2N} \eta = \frac{(4\pi)^N}{\sqrt{\det \gamma}} e^{-d^T \frac{1}{\gamma} d} \quad (2.89)$$

$$\int (\chi(\eta))^2 d^{2N} \eta = \frac{(2\pi)^N}{\sqrt{\det \gamma}} e^{-d^T \frac{4}{\gamma} d} \quad (2.90)$$

$$\int |\chi(\eta)|^2 d^{2N} \eta = \frac{(2\pi)^N}{\sqrt{\det \gamma}} \quad (2.91)$$

Chapter 3

Efficiency in QKD protocols with entangled Gaussian states

Efficiency is a key issue in any real implementation of a cryptographic protocol since the physical resources are not unlimited. We will first show that Quantum Key Distribution is possible with an "Entanglement based" scheme with NPPT symmetric Gaussian states in spite of the fact that these systems cannot be distilled with Gaussian operations (they are all bound entangled). In this work we analyze the secrecy properties of Gaussian states under Gaussian operations. Although such operations are useless for quantum distillation, we prove that it is possible to distill efficiently a secret key secure against finite coherent attacks from sufficiently entangled Gaussian states with non-positive partial transposition. Moreover, all such states allow for efficient key distillation, when the eavesdropper is assumed to perform individual attacks before the reconciliation process. In section (3.1) we present the academic protocol [5], while in section (3.2) we present the way to perform QKD with in the protocol in an efficient way.

3.1 State of the problem: QKD with entangled Gaussian states

Before presenting the protocol it is important to notice that Gaussian states always admit a purification. Thus, any mixed Gaussian state of N modes can be expressed as the reduction of a pure Gaussian state of $2N$ modes of the form:

$$\gamma_{2N} = \begin{pmatrix} \gamma_N & C_N \\ C_N^T & \theta_N \gamma_N \theta_N^T \end{pmatrix}, \quad C_N = J_N \sqrt{-(J_N \gamma_N)^2 - \mathbb{I}} \theta_N, \quad \theta_N = \bigoplus_{i=1}^N \theta,$$

such that the mixed state can be obtained after tracing out N modes from γ_{2N} . Here $\theta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, which is the momentum reflection in phase-space, is the associated symplectic operation.

For what follows it is also important the fact that any NPPT Gaussian state can be mapped by Gaussian Local Operations and Classical Communication (GLOCC) to an NPPT symmetric state of 1×1 modes *i.e.* preserving the amount of entanglement.

As the last remark, to deal with the content of the entanglement in Gaussian states we are going to use the negativity. As it was stated in section (2.6.2), in terms of CMs and for 1×1 and $1 \times N$ modes of bipartite Gaussian states the PPT criterion, which tells us that a state $\hat{\rho}$ is entangled if and only if it has non positive partial transposition, reads $\theta_A \gamma \theta_A^T + iJ < 0$.

With all the formalism at hand we now review the main steps of the protocol used in [5]. Without loosing generality, and by virtue of the above properties of Gaussian states, one should only consider the case in which Alice and Bob share many copies of a quantum system of 1×1 symmetric NPPT Gaussian state $\hat{\rho}_{AB}$. To extract a list of classically correlated bits to establish a secret key, each party measures the quadratures of her/his mode $\hat{X}_{A,B}$ and accepts only those outputs $x_{A,B}$ for which both parties have a consistent result $|x_A| = |x_B| = x_0$. With probability $p(i, j)$, each party associates the classical bit $i = 0(1)$ to her/his outcome $+x_0(-x_0)$. The probability that their symbols do not coincide is given by $\epsilon_{AB} = (\sum_{i \neq j} p(i, j)) / (\sum_{i, j} p(i, j))$. Having fixed a string of M classical correlated values, they can apply Classical Advantage Distillation [11]. To this aim, Alice generates a random bit b and encodes her string of M classical bits into a vector $\vec{b}$ of length M such that $b_{Ai} + b_i = b \pmod{2}$. Bob checks that for his symbols all results $b_{Bi} + b_i = b' \pmod{2}$ are consistent, and in this case accepts the bit b . The new error probability is given by

$$\epsilon_{AB,M} = \frac{(\epsilon_{AB})^M}{(1 - \epsilon_{AB})^M + (\epsilon_{AB})^M} < \left(\frac{\epsilon_{AB}}{1 - \epsilon_{AB}} \right)^M, \quad (3.1)$$

which tends to zero for sufficiently large M . The most general scenario for eavesdropping is to assume that Eve has access to the states before their distribution. Hence, the states that Alice and Bob share correspond to the reduction of a pure 4-mode state. We consider two types of attacks: (i) individual (or incoherent) attack, where Eve performs individual measurements, possibly non-Gaussian, over her set of states and (ii) finite coherent (or collective) attack, where Eve waits until the distribution has been performed, and, decides, which collective measurement gives her more information on the final key. Now, security with respect to individual attacks from the eavesdropper Eve, can be established if

$$\left(\frac{\epsilon_{AB}}{1 - \epsilon_{AB}} \right)^M < |\langle e_{++} | e_{--} \rangle|^M, \quad (3.2)$$

where $|e_{\pm\pm}\rangle$ denotes the state of Eve once Alice and Bob have projected their states onto $|\pm x_0\rangle$. Notice that Eve can gain information if the overlap between her states after Alice and Bob have measured coincident results is sufficiently small. The above inequalities come from the fact that in the case of individual attacks the error on

Eve's estimation of the final bit b is bound from below by a term proportional to $|\langle e_{++}|e_{--}\rangle|^M$ [5]. Therefore, Alice and Bob can establish a key if

$$\frac{\epsilon_{AB}}{1 - \epsilon_{AB}} < |\langle e_{++}|e_{--}\rangle|. \quad (3.3)$$

In [5] it was shown that any 1×1 NPPT state fulfils the above inequality and thus any NPPT Gaussian state can be used to establish a secure key in front of individual eavesdropper attacks. If we assume that Eve performs more powerful attacks, namely finite coherent attacks, then security is only guaranteed if the much more restrictive condition

$$\frac{\epsilon_{AB}}{1 - \epsilon_{AB}} < |\langle e_{++}|e_{--}\rangle|^2 \quad (3.4)$$

is fulfilled. This new inequality is violated by some NPPT states. Notice that this implies that the analysed protocol is not good for these states in this more general scenario. Nevertheless, using the recent techniques of [17], one can find states for which the presented protocol allows to extract common bits secure against this attack.

3.2 Efficient QKD with entangled Gaussian states

Let us now present our results. Notice that since security relies on the fact that Alice and Bob have better correlations than the information the eavesdropper can learn about their state, perfect correlation is not a requirement to establish a secure key. We denote Alice's outputs by x_{0A} and we calculate which are the outputs Bob can accept so that the correlation established between Alice and Bob outputs can be used to extract a secret bit.

We use the standard form of a bipartite 1×1 mode Gaussian state,

$$\gamma_{AB} = \begin{pmatrix} \lambda_A & 0 & c_x & 0 \\ 0 & \lambda_A & 0 & -c_p \\ c_x & 0 & \lambda_B & 0 \\ 0 & -c_p & 0 & \lambda_B \end{pmatrix} \quad (3.5)$$

with $\lambda_{A,B} \geq 1$, and $c_x \geq |c_p| \geq 0$ (we can shift the displacement vector to 0). We shall deal with mixed symmetric states and so $\lambda_A = \lambda_B = \lambda$. The positivity condition reads $(\lambda - c_x)(\lambda + c_p) \geq 1$, while the entanglement NPPT condition is given by $(\lambda - c_x)(\lambda - c_p) < 1$. As in [5], we impose that the global state including Eve is pure (she has access to all degrees of freedom outside Alice and Bob) while the mixed symmetric state, shared by Alice and Bob is just its reduction,

$$\gamma_{ABE} = \begin{pmatrix} \gamma_{AB} & C \\ C^T & \theta \gamma_{AB} \theta^T \end{pmatrix}, \quad (3.6)$$

$$C = J_{AB} \sqrt{-(J_{AB} \gamma_{AB})^2 - \mathbb{I}_2} \theta_{AB} = \begin{pmatrix} 0 & -X & 0 & -Y \\ -X & 0 & -Y & 0 \\ 0 & -Y & 0 & -X \\ -Y & 0 & -X & 0 \end{pmatrix}, \quad (3.7)$$

$$\theta_{AB} = \theta_A \oplus \theta_B, \quad J_{AB} = J_A \oplus J_B, \quad (3.8)$$

where

$$X = \frac{\sqrt{a+b} + \sqrt{a-b}}{2},$$

$$Y = \frac{\sqrt{a+b} - \sqrt{a-b}}{2},$$

and $a = \lambda^2 - c_x c_p - 1$, $b = \lambda(c_x - c_p)$.

Performing a measurement with uncertainty σ , the probability that Alice finds $\pm|x_{0A}|$ while Bob finds $\pm|x_{0B}|$, is given by the overlap between the state of Alice and Bob, $\hat{\rho}_{AB}$, and a pure product state $\hat{\rho}_{A,i} \otimes \hat{\rho}_{B,j}$ (with $i, j = 0, 1$) of Gaussians centred at $\pm|x_{0A}|(\pm|x_{0B}|)$ respectively with σ width (notice $\hat{\rho}_{A,0} \equiv | +|x_{0A}| \rangle \langle +|x_{0A}| |$). We use here the Hilbert-Schmidt fidelity for calculation, which gives:

$$\begin{aligned} p(0,0) &= p(1,1) = \text{tr}[\hat{\rho}_{AB}(\hat{\rho}_{A,0} \otimes \hat{\rho}_{B,0})] = \\ &= (2\pi)^4 \int d^4 \zeta_{AB} \mathcal{W}_{\rho_{AB}}(\zeta_{AB}) \mathcal{W}_{\rho_{A,0} \otimes \rho_{B,0}}(\zeta_{AB}) = \\ &= K(\sigma) \exp \left(\frac{2|x_{0A}||x_{0B}|c_x - (\lambda + \sigma^2)(x_{0A}^2 + x_{0B}^2)}{(\lambda + \sigma^2)^2 - c_x^2} \right), \end{aligned} \quad (3.9)$$

for the probability that their symbols do coincide and,

$$p(0,1) = p(1,0) = K(\sigma) \exp \left(\frac{-2|x_{0A}||x_{0B}|c_x - (\lambda + \sigma^2)(x_{0A}^2 + x_{0B}^2)}{(\lambda + \sigma^2)^2 - c_x^2} \right), \quad (3.10)$$

for the probability that they do not coincide, where

$$K(\sigma) = \frac{4\sigma^2}{\sqrt{(\lambda + \sigma^2)^2 - c_x^2} \sqrt{(\lambda\sigma^2 + 1)^2 - c_p^2\sigma^4}}. \quad (3.11)$$

Their error probability for $\sigma \rightarrow 0$ reads

$$\epsilon_{AB} = \lim_{\sigma \rightarrow 0} \frac{\sum_{i \neq j} p(i,j)}{\sum_{i,j} p(i,j)} = \frac{1}{1 + \exp \left(\frac{4c_x|x_{0A}||x_{0B}|}{\lambda^2 - c_x^2} \right)}. \quad (3.12)$$

Let us calculate the state of Eve $|e_{\pm\pm}\rangle$ after Alice has projected onto $|\pm|x_{0A}|\rangle$ and Bob onto $|\pm|x_{0B}|\rangle$:

$$\gamma_{++} = \gamma_{--} = \begin{pmatrix} \gamma_x & 0 \\ 0 & \gamma_x^{-1} \end{pmatrix}, \quad \gamma_x = \begin{pmatrix} \lambda & c_x \\ c_x & \lambda \end{pmatrix}, \quad (3.13)$$

$$d_{\pm\pm} = \mp \begin{pmatrix} 0 \\ 0 \\ A\delta x_0 - B\Delta x_0 \\ A\delta x_0 + B\Delta x_0 \end{pmatrix}, \quad (3.14)$$

where $A = \frac{\sqrt{a+b}}{\lambda+c_x}$, $B = \frac{\sqrt{a-b}}{\lambda-c_x}$, $\Delta x_0 = |x_{0B}| - |x_{0A}|$ and $\delta x_0 = |x_{0B}| + |x_{0A}|$. The overlap between the two states of Eve is given by:

$$|\langle e_{++}|e_{--}\rangle|^2 = \exp\left(\frac{-4}{\lambda^2 - c_x^2} \left[\left(\frac{x_{0A}^2 + x_{0B}^2}{2}\right) (\lambda^2 - c_x^2 - 1)\lambda + |x_{0A}||x_{0B}| (c_x - c_p(\lambda^2 - c_x^2)) \right]\right). \quad (3.15)$$

Substituting Eqs. (3.12) and (3.15) into (3.3) one can check, after some algebra, that the last inequality reduces to:

$$\left(\frac{x_{0A}^2 + x_{0B}^2}{2}\right) (\lambda^2 - c_x^2 - 1)\lambda + |x_{0A}||x_{0B}| (-c_x - c_p(\lambda^2 - c_x^2)) < 0. \quad (3.16)$$

Notice that condition (3.16) imposes both, restrictions on the parameters defining the state (λ, c_x, c_p) , and on the outcomes of the measurements (x_{0A}, x_{0B}) . The constraints on the state parameters are equivalent to demand that the state is NPPT and satisfies

$$(\lambda - c_x)(\lambda + c_x) \geq 1. \quad (3.17)$$

Nevertheless, as $c_x \geq c_p$, any positive state fulfils this condition. Hence for any NPPT symmetric state, there exists, for a given x_{0A} , a range of values of x_{0B} such that secret bits can be extracted (Eq. (3.3) is fulfilled). This range is given by

$$\Delta x_0 = |x_{0B}| - |x_{0A}| \in \mathfrak{D}_\alpha = \left[\frac{2}{-\sqrt{\alpha} - 1}, \frac{2}{\sqrt{\alpha} - 1} \right] |x_{0A}|, \quad (3.18)$$

where

$$\alpha = \left(\frac{c_x - \lambda}{c_x + \lambda}\right) \left[\frac{1 - (\lambda + c_x)(\lambda + c_p)}{1 - (\lambda - c_x)(\lambda - c_p)} \right]. \quad (3.19)$$

After Alice communicates $|x_{0A}|$ to Bob, he will accept only measurement outputs within the above interval. The interval Δx_0 is well defined if $\alpha \geq 1$, which equals to fulfil Eq. (3.17). Notice also that the interval is not symmetric around $|x_{0A}|$ because the probabilities calculated in Eqs. (3.9) and (3.10) do depend on this value in a non-symmetric way. The length D_α of the interval of valid measurements outputs for Bob is given by

$$D_\alpha = \frac{4\sqrt{\alpha}}{\alpha - 1} |x_{0A}|. \quad (3.20)$$

It can be observed that maximal $D_\alpha \rightarrow \infty$ ($\alpha = 1$) corresponds to the case when Alice and Bob share a pure state (Eve is disentangled from the system) and thus condition (3.3) is always fulfilled. On the other hand, any mixed NPPT symmetric state ($\alpha > 1$) admits a finite D_α . This ensures a *finite* efficiency on establishing a secure secret key in front of individual attacks.

If we assume that Eve performs more powerful attacks, namely finite coherent attacks, then security is only guaranteed if [5]:

$$\frac{\epsilon_{AB}}{1 - \epsilon_{AB}} < |\langle e_{++} | e_{--} \rangle|^2. \quad (3.21)$$

This condition is more restrictive than (3.3). With a similar calculation as before we obtain that now security is not guaranteed for all mixed entangled symmetric NPPT states, but only for those that also satisfy:

$$\lambda - (\lambda + c_x)(\lambda - c_x)(\lambda - c_p) > 0. \quad (3.22)$$

For such states, and given a measurement result x_{0A} of Alice, Bob will only accept outputs within the range:

$$\Delta x_0 = |x_{0B}| - |x_{0A}| \in \mathfrak{D}_\beta = \left[\frac{2}{-\sqrt{\beta} - 1}, \frac{2}{\sqrt{\beta} - 1} \right] |x_{0A}|, \quad (3.23)$$

where

$$\beta = \frac{2\lambda(\lambda^2 - c_x^2 - 1)}{\lambda - (\lambda + c_x)(\lambda - c_x)(\lambda - c_p)} \geq 1. \quad (3.24)$$

As before, $\beta \geq 1$ is fulfilled by conditions (3.17) and (3.22).

Let us now focus on the efficiency issue. We define the efficiency $E(\gamma_{AB})$ of the protocol for a given state γ_{AB} , as the average probability of obtaining a classically correlated bit. Explicitly,

$$E(\gamma_{AB}) = \int_{\Delta x_0 \in \mathfrak{D}} dx_{0A} dx_{0B} (1 - \epsilon_{AB}) \text{tr}(\hat{\rho}_{AB} |x_{0A}, x_{0B}\rangle \langle x_{0A}, x_{0B}|). \quad (3.25)$$

The marginal distribution in phase-space is easily computed by integrating the corresponding Wigner function in momentum space [16]:

$$\begin{aligned} \text{tr}(\hat{\rho}_{AB} |x_{0A}, x_{0B}\rangle \langle x_{0A}, x_{0B}|) &= \int \int dp_A dp_B \mathcal{W}_{\rho_{AB}}(\zeta_{AB}) = \\ &= \frac{\exp\left(\frac{2c_x x_{0A} x_{0B} - \lambda(x_{0A}^2 + x_{0B}^2)}{\lambda^2 - c_x^2}\right)}{\pi \sqrt{\lambda^2 - c_x^2}}, \end{aligned} \quad (3.26)$$

but the final expression of Eq. (3.25) has to be calculated numerically. Note that if Alice and Bob share as a resource M identical states (NPPT state for individual attacks, and NPPT fulfilling condition (3.22) for finite coherent attacks), the number of classically correlated bits that can be extracted from them is $\sim M \times E(\gamma_{AB})$. The efficiency Eq. (3.25) increases with increasing D and decreasing ϵ_{AB} . In particular, for the protocol given in [5], $D = 0$, and therefore $E(\gamma_{AB}) = 0$ for any state.

We investigate now the dependence of $E(\gamma_{AB})$ with the entanglement of the NPPT mixed symmetric state used for the protocol as well as with the purity of the




Figure 3.1: Protocol efficiency (quantified by $E(\gamma_{AB})$) versus the entanglement measured by logarithmic negativity $\text{LN}(\gamma_{AB})$. The shading from black to white corresponds to purity from zero to one.

state. As a measure of the entanglement between Alice and Bob we compute the logarithmic negativity

$$\text{LN}(\gamma_{AB}) = \log_2 \left(\frac{1}{\sqrt{(\lambda - c_x)(\lambda - c_p)}} \right) > 0. \quad (3.27)$$

In Fig. 3.1, we display the efficiency of the protocol (assuming individual attacks) versus entanglement shared between Alice and Bob for different states γ_{AB} . There is not a one-to-one correspondence between $E(\gamma_{AB})$ and entanglement, since states with the same entanglement can have different purity, which can lead to different efficiency. This is so because there are two favourable scenarios to fulfil Eq. (3.3). The first one is to demand large correlations so that the relative error ϵ_{AB} of Alice and Bob is small. The second scenario happens when Alice and Bob share a state with high purity, *i.e.*, Eve is very disentangled. In this case, independently of the error ϵ_{AB} , Eq. (3.3) can be fulfilled more easily.

Despite the fact that efficiency generally increases with increasing entanglement, this enhancement, as depicted in the figure, is a complex function of the parameters involved. Nevertheless, one can see that there exist an entanglement threshold (around $\text{LN}(\gamma_{AB}) \simeq 0.2$) below which the protocol efficiency diminishes drastically no matter how mixed are the states shared between Alice and Bob.

It is also illustrative to examine the dependence of α (which determines the interval length D_α) on the entanglement of the states shared by Alice and Bob. In Fig. 3.2 we plot the logarithmic negativity of a given state versus the parameter α . States with the same entanglement but different purity are associated to quite different values of α . Nevertheless states with high entanglement permit a large interval length (small α) and, thus, high efficiency.




Figure 3.2: Entanglement of the states shared between Alice and Bob measured in terms of the logarithmic negativity $\text{LN}(\gamma_{AB})$ as a function of the parameter $\alpha(\gamma_{AB})$ under individual attacks. The shading from black to white corresponds to purity from zero to one.

In both, Fig. 3.1 and Fig. 3.2, we have observed that states with different entanglement give the same efficiency. However it is important to pointed out that to extract the key's bits, Classical Advantage Distillation [11] stills needs to be performed. The efficiency of Maurer's protocol, strongly increases with decreasing ϵ_{AB} , and, therefore, the states with higher entanglement will provide a higher key rate.

Chapter 4

Summary and Conclusions

Efficiency is a key issue for any experimental implementation of Quantum Cryptography since available resources are not unlimited. Here, we have shown that the sharing of entangled Gaussian variables and the use of only Gaussian operations permits efficient Quantum Key Distribution against individual and finite coherent attacks.

We have used the fact that all mixed NPPT symmetric states can be used to extract secret bits under individual attacks whereas under finite coherent attacks an additional condition has to be fulfilled. We have introduced a figure of merit (the efficiency E) to quantify the number of classical correlated bits that can be used to distill a key from a sample of M entangled states. We have observed that this quantity grows with the entanglement shared between Alice and Bob. This relation it is not one-to-one due to the fact that states with less entanglement but with more purity (eavesdropper more disentangled) can be equally efficient. Nevertheless we have point out that, these states would be inefficient, when performing the Classical Advantage Distillation of the key.

Finally, we would like to remark that our study is not restricted to Quantum Key Distribution protocols, but can be extended to any other protocol that uses as a resource entangled continuous variables to establish a set of classically correlated bits between distant parties, see *e.g.* [18].

In [18] an efficient solution of the Byzantine Agreement problem (detectable broadcast) in the continuous variable scenario with multipartite entangled Gaussian states and Gaussian operations (homodyne detection) is presented. In a cryptographic context, detectable broadcast refers to distributed protocols in which some of the participants might have malicious intentions and could try to sabotage the distributed protocol inducing the honest parties to take contradictory actions between them. Entanglement is used in the protocol to distribute classical private random variables with a specific correlation between the players, in such a way that any malicious manipulation of the data can be detected by all honest parties allowing them to abort the protocol. We discuss realistic implementations of the protocol, which consider the possibility of having inefficient homodyne detectors, not perfectly

correlated outcomes, and noise in the preparation of the resource states. The proposed protocol is proven to be robust and efficiently applicable under such non-ideal conditions.

Following [19], it is known that, in spite of their exceptional role within the space of all continuous variables states, in fact, Gaussian states are not the best candidates as resources to perform Quantum tasks. In general, any continuous, strongly, super-additive functional acting on any given covariance matrix is minimised by Gaussian states. This is the case for all entanglement measures fulfilling the above conditions *e.g.* the distillable entanglement or the entanglement of formation. In this sense Gaussian states are extremal. With this idea on mind, naturally one could try to perform QKD with non-Gaussian states. Following the presented protocol here one should expect an enhancement on the efficiency on the key distribution with non-Gaussian states. Gaussian states possess an easy mathematical description at the covariance matrix level while for non-Gaussian states this description is not complete. Nevertheless, the Wigner distribution function formalism presented here, allows to perform the needed calculations to study QKD with non-Gaussian states, in a very similar way.

Also, for non-Gaussian states there are no computable entanglement measures while these are well established in the case of Gaussian states. Thus, one might think that a way to quantify entanglement in non-Gaussian states can be accomplished by relating the efficiency of distilling correlated bits with the entanglement of non-Gaussian states [20]. From the experimental point of view, there already exist several groups which actually have succeeded in producing non-Gaussian states like photon-subtracted, states that up to now lack of a complete well characterisation.

Bibliography

- [1] C. H. Bennett and G. Brassard. Quantum cryptography: Public key distribution and coin tossing. *Proceedings of IEEE International Conference on Computers, Systems, and Signals Processing, Bangalore, India (IEEE, New York, 1984)*, page 175, (1984).
- [2] A. Ekert. Quantum cryptography based on bell's theorem. *Phys. Rev. Lett.*, **67**:661, (1991).
- [3] C. H. Bennett, G. Brassard, and N. D. Mermin. Quantum cryptography without bell's theorem. *Phys. Rev. Lett.*, **68**:557, (1992).
- [4] M. Curty, M. Lewenstein, and N. Lütkenhaus. Entanglement as a precondition for secure quantum key distribution. *Phys. Rev. Lett.*, **92**:217903, (2004).
- [5] M. Navascués, J. Bae, J. I. Cirac, M. Lewestein, A. Sanpera, and A. Acín. Quantum key distillation from gaussian states by gaussian operations. *Phys. Rev. Lett.*, **94**:010502, (2005).
- [6] D. Deutsch, A. Ekert, R. Jozsa, S. Popescu C. Macchiavello, and A. Sanpera. Quantum privacy amplification and the security of quantum cryptography over noisy channels. *Phys. Rev. Lett.*, **77**:2818, (1996).
- [7] K. Horodecki, M. Horodecki, P. Horodecki, and J. Oppenheim. Secure key from bound entanglement. *Phys. Rev. Lett.*, **94**:160502, (2005).
- [8] D. Gottesman and J. Preskill. Secure quantum key distribution using squeezed states. *Phys. Rev.*, **63**:022309, (2001).
- [9] F. Grosshans and P. Grangier. Continuous variable quantum cryptography using coherent states. *Phys. Rev. Lett.*, **88**:057902, (2002).
- [10] Ch. Silberhorn, T. C. Ralph, N. Lütkenhaus, and G. Leuchs. Continuous variable quantum cryptography: Beating the 3 db loss limit. *Phys. Rev. Lett.*, **89**:167901, (2002).
- [11] U. M. Maurer. Secret key agreement by public discussion from common information. *IEEE Trans. Inf. Theory*, **39**:733, (1993).

-
- [12] C. Rodó, O. Romero-Isart, K. Eckert, and A. Sanpera. Efficiency in quantum key distribution protocols with entangled gaussian states. *Open Syst. Inf. Dyn.*, **14**:69, (2007).
 - [13] S. L. Braunstein and P. van Loock. Quantum information with continuous variables. *Rev. Mod. Phys.*, **77**:513, (2005).
 - [14] P. van Loock. Quantum communication with continuous variables. *Fortschr. Phys.*, **50**:1177, (2002).
 - [15] W-M. Zhang, Da H. Feng, and R. Gilmore. Coherent states: Theory and some applications. *Rev. Mod. Phys.*, **62**:867, (1990).
 - [16] H. W. Lee. Theory and application of the quantum phase-space distribution functions. *Phys. Rep.*, **259**:147, (1995).
 - [17] M. Christandl, R. Renner, and A. Ekert. A generic security proof for quantum key distribution. *ArXiv:quant-ph/0402131*, (2004).
 - [18] R. Neigovzen, C. Rodó, G. Adesso, and A. Sanpera. Continuous variable solution for byzantine agreement. *Phys. Rev. A*, **77**:062307, (2008).
 - [19] M. M. Wolf, G. Giedke, and J. I. Cirac. Extremality of gaussian quantum states. *Phys. Rev. Lett.*, **96**:080502, (2006).
 - [20] C. Rodó, G. Adesso, and A. Sanpera. Operational quantification of continuous variable correlations. *Phys. Rev. Lett.*, **100**:110505, (2008).