Renormalization Group Functions for the Radiative Symmetry Breaking Scheme with Multi Mass Scale

Chungku Kim

(Dated: February 23, 2024)

We obtain the renormalization group(RG) functions for the O(N) scalar field theory and the Higgs-Yukawa field theory with the Coleman-Weinberg mechanism in which the symmetry breaking occurs radiatively by using the method proposed previously in case of the neutral scalar field theory.

PACS numbers: 11.15.Bt, 12.38.Bx

The electroweak(EW) model with the radiative symmetry breaking(RSB) scheme [1] in which the scalar field does not have a tree level mass and the symmetry breaking(SB) occurs from the effective potential [2] has been studied extensively [3] due to its predictive power for the magnitude of the Higgs boson mass. In order to investigate the higher-order perturbative corrections, we need the renormalization group(RG) function [4] in case of the RSB scheme. Recently, we have obtained the RG function for the neutral scalar field theory with the RSB scheme [5] by two different procedures. Among these, the one step procedure can be used in case of the theory which have an effective potential depending on multi-mass scales [6]. In this procedure, we first obtain the effective potential for the RSB scheme by a finite transform of both the coupling constant and the classical field from the well known effective potential in the minimal subtraction(MS) scheme. Then we obtain the resulting change of the RG functions due to these transforms.

In this paper, we will apply the one step procedure proposed in [5] to the case of the O(N) scalar field theory and the Higgs-Yukawa field theory which have an effective potential depending on multi-mass scales and have obtained the two-loop effective potential and the corresponding RG functions in the RSB scheme.

A. O(N) Scalar Field Theory

The effective potential of the O(N) scalar field theory in the MS scheme has the form [7]

$$V^{MS}(\lambda,\phi,\mu) = \phi^{4} \left[\frac{1}{24}\lambda + \kappa\lambda^{2} \left(\frac{1}{16}L_{2} + \frac{n}{144}L_{1} - \frac{3}{32} - \frac{n}{96}\right) + \kappa^{2}\lambda^{3} \left(\frac{3}{32}L_{2}^{2} + \frac{n}{48}L_{2}L_{1} + \frac{n^{2}}{864}L_{1}^{2} - \left(\frac{5}{16} + \frac{5n}{144}\right)L_{2} - \left(\frac{13n}{432} + \frac{n^{2}}{432}\right)L_{1} + \frac{11}{32} + \frac{1}{2}\Omega(1) + \frac{29n}{432} + \frac{5n}{54}\Omega(3) + \frac{n^{2}}{864}\right) + \dots\right],$$
(1)

where $\kappa = (16\pi^2)^{-1}$, n = N - 1, l is the order of the loop and

$$L_1 \equiv \log\left(\frac{\lambda\phi^2}{6\mu^2}\right) \text{ and } L_2 \equiv \log\left(\frac{\lambda\phi^2}{2\mu^2}\right)$$
 (2)

and $\Omega(x)$ is given by

$$\Omega(x) = \frac{\sqrt{x(4-x)}}{x+2} \int_0^\theta \ln(2\sin t) dt \ (\sin\theta = \frac{\sqrt{x}}{2}) \tag{3}$$

Since the effective potential is independent of the renormalization mass scale μ , it satisfies the renormalization group equation

$$\left[\mu\frac{\partial}{\partial\mu} + \beta_{MS}(\lambda)\frac{\partial}{\partial\lambda} + \gamma_{MS}(\lambda)\phi\frac{\partial}{\partial\phi}\right]V^{MS}(\lambda,\phi,\mu) = 0 \tag{4}$$

where the RG functions β_{MS} and γ_{MS} are given by [4]

$$\beta^{MS}(\lambda) = \mu \frac{d\lambda}{d\mu} = \kappa (3 + \frac{n}{3})\lambda^2 - \kappa^2 (\frac{17}{3} + n)\lambda^3 + \dots \equiv \sum_{l=1}^{\infty} b_l^{MS} \kappa^l \lambda^{l+1}$$
(5)

arXiv:1008.4843v1 [hep-ph] 28 Aug 2010

and

$$\gamma^{MS}(\lambda) = \frac{\mu}{\phi} \frac{d\phi}{d\mu} = -\kappa^2 \left(\frac{1}{12} + \frac{n}{36}\right) \lambda^2 + \dots \equiv \sum_{l=1}^{\infty} g_l^{MS} \kappa^l \lambda^l \tag{6}$$

In order to obtain the effective potential for the RSB scheme from $V^{MS}(\lambda, \phi, \mu)$, let us perform a finite transformations of the coupling constant λ and the classical field ϕ to $\tilde{\lambda}$ and $\tilde{\phi}$ as $\lambda = \lambda(\tilde{\lambda})$ and $\phi = \tilde{\phi} f(\tilde{\lambda})$ respectively so that the resulting effective potential for the RSB scheme $V^R(\tilde{\lambda}, \tilde{\phi}, \mu)$ has the form

$$V^{R}(\widetilde{\lambda},\widetilde{\phi},\mu) = V^{MS}(\lambda(\widetilde{\lambda}),\widetilde{\phi}f(\widetilde{\lambda}),\mu) = \sum_{l=0}^{\infty} \kappa^{l} \widetilde{\phi}^{4} \sum_{n=0}^{l} a_{l,n}(\widetilde{\lambda}) \log^{n}\left(\frac{\widetilde{\phi}^{2}}{\mu^{2}}\right) \equiv \sum_{l=0}^{\infty} \kappa^{l} V_{l}^{R}(\widetilde{\lambda},\widetilde{\phi},\mu)$$
(7)

and satisfies the renormalization condition

$$\left[\frac{d^4 V^R(\tilde{\lambda}, \tilde{\phi}, \mu)}{d\tilde{\phi}^4}\right]_{\tilde{\phi}=\mu} = \tilde{\lambda}.$$
(8)

Then the corresponding change of the RG function due to these transformations are given by [5]

$$\beta^{R}(\widetilde{\lambda}) = \mu \frac{\partial \widetilde{\lambda}}{\partial \mu} = \frac{\partial \widetilde{\lambda}}{\partial \lambda} \mu \frac{\partial \lambda}{\partial \mu} = \frac{\beta^{MS}(\lambda(\widetilde{\lambda}))}{\frac{\partial \lambda(\widetilde{\lambda})}{\partial \widetilde{\lambda}}} \equiv \sum_{l=1}^{\infty} \kappa^{l} b_{l}^{R}(\widetilde{\lambda}).$$
(9)

and

$$\gamma^{R}(\widetilde{\lambda}) = \frac{\mu}{\widetilde{\phi}} \frac{d\widetilde{\phi}}{d\mu} = \frac{1}{\widetilde{\phi}} \left[\frac{\mu}{f(\widetilde{\lambda})} \frac{\partial \phi}{\partial \mu} + \phi \mu \frac{\partial \widetilde{\lambda}}{\partial \mu} \frac{\partial f^{-1}(\widetilde{\lambda})}{\partial \widetilde{\lambda}} \right]$$
$$= \gamma^{MS}(\lambda(\widetilde{\lambda})) + \beta^{R}(\widetilde{\lambda}) f(\widetilde{\lambda}) \frac{\partial f^{-1}(\widetilde{\lambda})}{\partial \widetilde{\lambda}} \equiv \sum_{l=1}^{\infty} \kappa^{l} g_{l}^{R}(\widetilde{\lambda}).$$
(10)

Note that the *l*-loop of order transformation $\lambda = \lambda(\widetilde{\lambda})$ and $\phi = \widetilde{\phi} f(\widetilde{\lambda})$ determine the (l+1)-loop order coefficient of the RG functions $\beta_R(\widetilde{\lambda})$ and $\gamma_R(\widetilde{\lambda})$.

Now, let us apply the above procedure to obtain the two-loop effective potential and the RG functions in the RSB scheme. The general form of the $O(\kappa)$ transforms can be written as

$$\lambda(\tilde{\lambda}) = \tilde{\lambda} + \kappa \tilde{\lambda}^2 (c_{110}l_1 + c_{101}l_2 + c_{100}) f(\tilde{\lambda}) = 1 + \kappa \tilde{\lambda} (d_{110}l_1 + d_{101}l_2 + d_{100})$$
(11)

where $l_1 \equiv \log(\frac{\tilde{\lambda}}{6})$ and $l_2 \equiv \log(\frac{\tilde{\lambda}}{2})$. By demanding that the one-loop effective potential have the form (7), we obtain

$$c_{110} = -4d_{110} - \frac{n}{6}, c_{101} = -4d_{101} - \frac{3}{2}$$
 (12)

By substituting Eq.(11) into Eq.(10) we can see that if d_{110}, d_{101} and d_{100} are not zero then the $g_R^{(2)}(\tilde{\lambda})$ which is the $O(\kappa^2)$ coefficient of the RG function $\gamma^R(\tilde{\lambda})$ depend on l_1 and l_2 . Hence, in order to obtain simple results for the two-loop RG functions, we choose $d_{110} = d_{101} = d_{100} = 0$. Finally, by demanding that $V^R(\tilde{\lambda}, \tilde{\phi}, \mu)$ satisfies the renormalization condition given in Eq.(8), we obtain

$$c_{100} = -4\left(1 + \frac{n}{9}\right) \tag{13}$$

Next, let us write the $O(\kappa^2)$ transformation as

$$\lambda(\widetilde{\lambda}) = \widetilde{\lambda} - \kappa \widetilde{\lambda}^2 (\frac{3}{2}l_1 + \frac{n}{6}l_2 + 4(1 + \frac{n}{9})) + \kappa^2 \widetilde{\lambda}^3 (c_{220}l_1^2 + c_{211}l_1l_2 + c_{202}l_2^2 + c_{210}l_1 + c_{201}l_2 + c_{200}) + \dots$$

$$f(\widetilde{\lambda}) = 1 + \kappa^2 \widetilde{\lambda}^2 (d_{220}l_1^2 + d_{211}l_1l_2 + d_{202}l_2^2 + d_{210}l_2 + d_{201}l_2 + d_{200}) + \dots$$
(14)

By substituting Eqs.(14) into (7) we can see that by choosing

$$c_{220} = \frac{n^2}{36} - 4d_{220}, \ c_{211} = \frac{n}{2} - 4d_{211}, \ c_{202} = \frac{9}{4} - 4d_{202} ,$$

$$c_{210} = -4d_{210} + \frac{4n^2}{27} + \frac{14n}{9} \text{ and } c_{201} = -4d_{201} + \frac{5n}{3} + 15$$
(15)

we can make those terms that depend on l_1^2 , $l_1 l_2$, l_2^2 , l_1 and l_2 in the two-loop effective potential vanish. Moreover, by investigating $b_R^{(3)}(\tilde{\lambda})$ and $g_R^{(3)}(\tilde{\lambda})$ which is the three-loop coefficients of the RG functions β^R and γ^R , we can see that those terms that depend on l_1^2 , $l_1 l_2$ and l_2^2 can be made to zero by choosing $d_{220} = d_{211} = d_{202} = 0$ and that those terms that depend on l_1 and l_2 does not vanish by any choice of d_{210} and d_{201} . This means that the RG functions becomes a polynomial of the coupling constant only up to the $O(\kappa^2)$. Finally, by demanding the renormalization condition given in Eq.(8), we obtain

$$c_{200} = -4d_{200} - \frac{20n\Omega(2)}{9} + \frac{202n}{27} + \frac{139}{4} + \frac{113n^2}{324} - 12\Omega(1)$$
(16)

As a result, up to two loop order, we obtain the RG functions and effective potential in RSB scheme as

$$\beta^R(\widetilde{\lambda}) = \kappa(3+\frac{n}{3})\widetilde{\lambda}^2 + \kappa^2(-\frac{7}{6} + \frac{1}{18}n^2)\widetilde{\lambda}^3 + \cdots$$
(17)

$$\gamma^R(\widetilde{\lambda}) = -\kappa^2 (\frac{1}{12} + \frac{n}{36})\widetilde{\lambda}^2 + \cdots$$
(18)

and

$$V^{R}(\tilde{\lambda}, \tilde{\phi}, \mu) = \tilde{\phi}^{4} [\frac{1}{24}\tilde{\lambda} + \kappa \tilde{\lambda}^{2} \{ (\frac{1}{16} + \frac{n}{144})L - \frac{25}{96} - \frac{25n}{864} \} + \kappa^{2} \tilde{\lambda}^{3} \{ (\frac{3}{32} + \frac{n}{48} + \frac{n^{2}}{864})L^{2} - (\frac{13}{16} + \frac{19n}{108} + \frac{11n^{2}}{1296})L + \frac{55}{24} + \frac{635n}{1296} + \frac{85n^{2}}{3888} \} + O(\kappa^{3})].$$

$$(19)$$

Note that up to this order $V^R(\tilde{\lambda}, \tilde{\phi}, \mu)$ coincide with the effective potential given in Ref.[8] and satisfies the RG equation

$$[\mu \frac{\partial}{\partial \mu} + \beta^R(\widetilde{\lambda}) \frac{\partial}{\partial \widetilde{\lambda}} + \gamma^R(\widetilde{\lambda}) \widetilde{\phi} \frac{\partial}{\partial \widetilde{\phi}}] V^R(\widetilde{\lambda}, \widetilde{\phi}, \mu) = 0$$
⁽²⁰⁾

B. Higgs-Yukawa field theory

The effective potential of the Higgs-Yukawa field theory which is the O(4) symmetric scalar field theory coupled to the Dirac fermion with the Yukawa coupling constant h in the MS scheme is given by [9]:

$$V^{MS}(\lambda, h, \phi, \mu) = \phi^{4} [\frac{1}{24}\lambda + \kappa (\frac{\lambda^{2}}{48}\log(\frac{G}{\mu^{2}}) + \frac{\lambda^{2}}{16}\log(\frac{H}{\mu^{2}}) - \frac{3h^{2}}{4}\log(\frac{T}{\mu^{2}}) - \frac{\lambda^{2}}{8} + \frac{9h^{2}}{8}) + \kappa^{2} [-\frac{\lambda^{2}\phi^{2}}{12}(\hat{I}(H, H, H) + \hat{I}(G, H, H)) + \frac{\lambda}{8}(\hat{J}(H, H) + \hat{J}(H, G) + \hat{J}(G, G)) + 3h^{2}((2T - \frac{H}{2})\hat{I}(H, T, T) - \frac{T}{2}\hat{I}(G, T, T) + (T - G)\hat{I}(G, T, 0) + \hat{J}(T, T) - \hat{J}(H, T) - 2\hat{J}(G, T)] + O(\kappa^{3}),$$
(21)

where

$$H = \frac{\lambda \phi^2}{2}, G = \frac{\lambda \phi^2}{6}, T = \frac{h^2 \phi^2}{2}$$
(22)

and

$$\widehat{I}(x,y,z) = 2(x\log(\frac{x}{\mu^2}) + y\log(\frac{y}{\mu^2}) + z\log(\frac{z}{\mu^2})) - \frac{1}{2}(y+z-x)\log(\frac{y}{\mu^2})\log(\frac{z}{\mu^2}) - \frac{1}{2}(z+x-y)\log(\frac{z}{\mu^2})\log(\frac{x}{\mu^2}) - \frac{1}{2}(x+y-z)\log(\frac{x}{\mu^2})\log(\frac{y}{\mu^2}) - \frac{5}{2}(x+y+z) - \frac{1}{2}\xi(x,y,z)$$
(23)

and

$$\widehat{J}(x,y) = xy(1 - \log(\frac{x}{\mu^2}) - y\log(\frac{y}{\mu^2}) + \log(\frac{x}{\mu^2})\log(\frac{y}{\mu^2}))$$
(24)

The $\xi(x, y, z)$ is a function independent of the logarithms and the detailed form of this function is given in [9]. Since the effective potential is independent of the renormalization mass scale μ , it satisfies the renormalization group equation

$$\left[\mu\frac{\partial}{\partial\mu} + \beta_{\lambda}^{MS}\frac{\partial}{\partial\lambda} + \beta_{h}^{MS}\frac{\partial}{\partialh} + \gamma^{MS}(\lambda)\phi\frac{\partial}{\partial\phi}\right]V^{MS}(\lambda, h, \phi, \mu) = 0$$
(25)

where the RG functions β_{MS} and γ_{MS} are given by [4]

$$\beta_{\lambda}^{MS} = \mu \frac{d\lambda}{d\mu} = \kappa (4\lambda^2 + 12\lambda h^2 - 36h^4) + \kappa^2 (-\frac{26}{3}\lambda^3 - 24\lambda^2 h^2 - 3\lambda h^4 + 180h^6) + \dots$$
(26)

$$\beta_{h^2}^{MS} = \mu \frac{dh^2}{d\mu} = 9\kappa h^4 + \kappa^2 (\frac{1}{3}\lambda^2 h^2 - 4\lambda h^4 - 24h^6) + \dots$$
(27)

and

$$\gamma^{MS} = \frac{\mu}{\phi} \frac{d\phi}{d\mu} = -3\kappa h^2 + \kappa^2 (-\frac{1}{6}\lambda^2 + \frac{27}{4}h^4) \cdots$$
(28)

Since the effective potential depends on two different coupling constants λ and h, let us generalize the previous results (Eqs.(9) and (10)) which correspond to a single coupling constant λ to the case where several coupling constants λ_i (i = 1, ..., N) exist and consider the finite transforms of the coupling constants λ_i and the classical field ϕ as

$$\lambda_i = \lambda_i(\widetilde{\lambda}_j) \ (i, j = 1, .., p) \tag{29}$$

and

$$\phi = \widetilde{\phi} f(\widetilde{\lambda}_j) \tag{30}$$

Then the RG of the coupling constants in RSB scheme can be obtained as

$$\beta_i^R(\widetilde{\lambda}) = \mu \frac{d\widetilde{\lambda}_i}{d\mu} = \sum_{j=1}^p \frac{\partial\widetilde{\lambda}_i}{\partial\lambda_j} \mu \frac{d\lambda_j}{d\mu} = \sum_{j=1}^p (M^{-1})_{ij} \beta_j^{MS}(\lambda(\widetilde{\lambda})) \equiv \sum_{l=1}^\infty \kappa^l b_l^R(\widetilde{\lambda}).$$
(31)

where $(M)_{ij} \equiv \frac{\partial \lambda_i}{\partial \lambda_i}$ which can be obtained from (29). The RG of the classical field ϕ in RSB scheme can be obtained by a simple extension of the Eq.(10) as

$$\gamma^{R}(\widetilde{\lambda}) = \mu \frac{d\widetilde{\phi}}{d\mu} = \gamma^{MS}(\lambda(\widetilde{\lambda})) + \sum_{i=1}^{p} \beta_{i}^{R}(\widetilde{\lambda}) f(\widetilde{\lambda}) \frac{\partial f^{-1}(\widetilde{\lambda})}{\partial \widetilde{\lambda}_{i}} \equiv \sum_{l=1}^{\infty} \kappa^{l} g_{l}^{R}(\widetilde{\lambda}).$$
(32)

As in case of the O(N) symmetric scalar field theory, we have searched the finite transformation of the coupling constants and the classical field by requiring that the one-loop effective potential take the form eq.(7), satisfy the renormalization condition given in eq.(8) and that the coefficients of the logarithms $L \equiv \log\left(\frac{\tilde{\phi}^2}{\mu^2}\right)$ of the two loop effective potential does not depend on $\log\left(\tilde{\lambda}\right)$ and $\log(\tilde{h}^2)$. The resulting one-loop order transformation was not unique and one possible solution was

$$\lambda = \tilde{\lambda} + \kappa \left[\left(-\frac{3}{2} l_1 - \frac{1}{2} l_2 - \frac{16}{3} \right) \tilde{\lambda}^2 + (18 l_3 + 48) \tilde{h}^4 \right]$$

$$h^2 = \tilde{h}^2 + \kappa \left[\left(\frac{1}{4} l_1 + \frac{1}{12} l_2 - \frac{1}{6} \right) \tilde{\lambda}^2 + \left(-\frac{3}{2} l_1 - \frac{1}{2} l_2 + 2 l_3 \right) \tilde{\lambda} \tilde{h}^2 - \frac{3}{2} \tilde{h}^4 \right]$$

$$f = 1$$
(33)

where $l_1 \equiv \log(\tilde{\lambda}_6)$, $l_2 \equiv \log(\tilde{\lambda}_2)$ and $l_3 \equiv \log(\tilde{h}^2/2)$. Then, by using the equations (21) and (30)-(33), the effective potential and the RG functions in the RSB scheme can be determined up to the two loop order as

$$V^{R}(\tilde{\lambda},\tilde{h},\tilde{\phi},\mu) = \phi^{4}[\frac{1}{24}\tilde{\lambda} + \kappa(\frac{\tilde{\lambda}^{2}}{12}L - \frac{3}{4}\tilde{h}^{2}L - \frac{25}{72}\tilde{\lambda}^{2} + \frac{25}{8}\tilde{h}^{2}) + \kappa^{2}[(\frac{1}{6}\tilde{\lambda}^{3} + \frac{1}{4}\tilde{\lambda}^{2}\tilde{h}^{2} - \frac{3}{2}\tilde{\lambda}\tilde{h}^{4} - \frac{9}{8}\tilde{h}^{6})L^{2} + (-\frac{17}{12}\tilde{\lambda}^{3} - \frac{3}{4}\tilde{\lambda}^{2}\tilde{h}^{2} + \frac{23}{2}\tilde{\lambda}\tilde{h}^{4} + 6\tilde{h}^{6})L + \frac{25}{24}\tilde{\lambda}^{3} + \frac{5}{24}\tilde{\lambda}^{2}\tilde{h}^{2} - \frac{365}{12}\tilde{\lambda}\tilde{h}^{4} - \frac{95}{8}\tilde{h}^{6}] + O(\kappa^{3}), \quad (34)$$

$$\beta_{\widetilde{\lambda}}^{R} = \mu \frac{\partial \widetilde{\lambda}}{\partial \mu} = \kappa (4\widetilde{\lambda}^{2} + 12\widetilde{\lambda}\widetilde{h}^{2} - 36\widetilde{h}^{4}) + \kappa^{2} [-\frac{8}{3}\widetilde{\lambda}^{3} + 76\widetilde{\lambda}^{2}\widetilde{h}^{2} - 75\widetilde{\lambda}\widetilde{h}^{4} - 2706\widetilde{h}^{6} + (3\widetilde{\lambda}^{3} - 18\widetilde{\lambda}^{2}\widetilde{h}^{2})l_{1} + (\widetilde{\lambda}^{3} - 6\widetilde{\lambda}^{2}\widetilde{h}^{2})l_{2} + (24\widetilde{\lambda}^{2}\widetilde{h}^{2} - 18\widetilde{\lambda}\widetilde{h}^{4})l_{3}],$$

$$(35)$$

$$\beta_{\tilde{h}^{2}}^{R} = \mu \frac{\partial \tilde{h}^{2}}{\partial \mu} = 9\kappa \tilde{h}^{4} + \kappa^{2} [\frac{16}{3} \tilde{\lambda}^{2} \tilde{h}^{2} + 2\tilde{\lambda} \tilde{h}^{4} - \frac{165}{2} \tilde{h}^{6} + (-2\tilde{\lambda}^{3} + \frac{9}{2} \tilde{\lambda}^{2} \tilde{h}^{2} + \frac{45}{2} \tilde{\lambda} \tilde{h}^{4} - 54\tilde{h}^{6}) l_{1} \\ + (-\frac{2}{3} \tilde{\lambda}^{3} + \frac{3}{2} \tilde{\lambda}^{2} \tilde{h}^{2} + \frac{15}{2} \tilde{\lambda} \tilde{h}^{4} - 18\tilde{h}^{6}) l_{2} + (-8\tilde{\lambda}^{2} \tilde{h}^{2} - 6\tilde{\lambda} \tilde{h}^{4} + 72\tilde{h}^{6}) l_{3}]$$
(36)

and

$$\gamma^{R}(\widetilde{\lambda},\widetilde{h}) = \mu \frac{\partial \widetilde{\lambda}}{\partial \mu} = -3\kappa \widetilde{h}^{2} + \kappa^{2} [\frac{1}{3}\widetilde{\lambda}^{2} + \frac{27}{4}\widetilde{h}^{4} + (-\frac{3}{4}\widetilde{\lambda}^{2} + \frac{9}{2}\widetilde{\lambda}\widetilde{h}^{2})l_{1} + (-\frac{1}{4}\widetilde{\lambda}^{2} + \frac{3}{2}\widetilde{\lambda}\widetilde{h}^{2})l_{2} + (-6\widetilde{\lambda}\widetilde{h}^{2} + \frac{9}{2}\widetilde{h}^{4})l_{3}].$$
(37)

where the L independent term of the $O(\kappa^2)$ term of $V^R(\lambda, \tilde{h}, \tilde{\phi}, \mu)$ was fixed by the renormalization condition given in (8). One can check that up to $O(\kappa^2)$, $V^R(\lambda, \tilde{h}, \tilde{\phi}, \mu)$ satisfies the RG equation

$$[\mu \frac{\partial}{\partial \mu} + \beta^{R}(\widetilde{\lambda}) \frac{\partial}{\partial \widetilde{\lambda}} + \beta^{R}(\widetilde{h}) \frac{\partial}{\partial \widetilde{h}} + \gamma^{R}(\widetilde{\lambda}) \widetilde{\phi} \frac{\partial}{\partial \widetilde{\phi}}] V^{R}(\widetilde{\lambda}, \widetilde{h}, \widetilde{\phi}, \mu) = 0$$
(38)

Form this equation, one can see that the coefficient of the L in the $O(\kappa^2)$ term of $V^R(\lambda, \tilde{h}, \tilde{\phi}, \mu)$ is related to the combination $b_2^R(\lambda) + 4g_2^R(\lambda)$. Hence, although both $b_2^R(\lambda)$ and $g_2^R(\lambda)$ depend on l_i terms, the combination $b_2^R(\lambda) + 4g_2^R(\lambda)$ does not depend on l_i terms so that the coefficient of the L in the $O(\kappa^2)$ term of $V^R(\lambda, \tilde{h}, \tilde{\phi}, \mu)$ is independent of l_i terms. In the case of the Higgs-Yukawa theory which contains three different mass scales in the effective potential, one can check that the l_i terms always appear in the $O(\kappa^2)$ RG functions even with the most general form of the finite transformation of the coupling constants and the classical field. The appearance of the logarithms in the coefficients of the RG functions also happens in case of the multi-scale renormalization[10]. As a result of these logarithms, the RG running of the coupling constants and the classical field will be different compared to the case of the usual RG functions which is a polynomial of the coupling constants and this is in progress.

ACKNOWLEDGMENTS

This research was supported by the Institute of the Basic Science Research Center.

- [1] S. Coleman and E.Weinberg, Phys. Rev. D7, 1888 (1973).
- [2] For a review and references, see M. Sher, Phys. Rep. 89, 273 (1989).
- [3] V. Elias, R.B.Mann, D.G.C.McKeon and T.G.Stelle, Phys. Rev. Lett. 91, 251601 (2003); Nucl. Phys. B678, 147 (2004);
 Phys. Rev. D72, 037902 (2005); F.A.Chishtie, V. Elias, R.B.Mann, D.G.C.McKeon and T.G.Stelle, Nucl. Phys. B743 ,104 (2006).
- [4] H. Kleinert, J. Neu, V. Schulte-Frohlinde, K. G. Chetyrkin, S. A. Larin, Phys. Lett. B272, 39(1991); ibid. 319, 545(1993).
- [5] C.K.Kim, Phys. Lett. B680, 104 (2009).
- [6] B. Kastening, Phys. Lett. B283, 287 (1992); M. Bando, T. Kugo, N. Maekawa and H. Nakano, Phys. Lett. B301,89 (1993).
- [7] C. Ford, D. R.T. Jones, Phys. Lett. B274, 409 (1992), 285, 399(E) (1992).
- [8] R. Jackiw, Phys. Rev. D9, 1686 (1974).
- [9] C.Ford, D.R.T.Jones, P.W.Stephenson and M.B.Einhorn, Nucl. Phys. B395, 17(1993).
- [10] C.Ford and C. Wiesendanger, Phys. Lett. B398, 342(1997).