

The Lambert Way to Gaussianize heavy tailed data with the inverse of Tukey's h transformation as a special case

Georg M. Goerg

Department of Statistics, Carnegie Mellon University
Pittsburgh, PA 15213, USA
gmg@stat.cmu.edu, www.stat.cmu.edu/~gmg

February 5, 2022

1 I present a parametric, bijective transformation to generate heavy tail versions Y of
2 arbitrary RVs $X \sim F_X$. The tail behavior of the so-called *heavy tail Lambert* $W \times F_X$ RV
3 Y depends on a tail parameter $\delta \geq 0$: for $\delta = 0$, $Y \equiv X$, for $\delta > 0$ Y has heavier tails than
4 X . For X being Gaussian, this meta-family of heavy-tailed distributions reduces to Tukey's
5 h distribution. Lambert's W function provides an explicit inverse transformation, which
6 can be estimated by maximum likelihood. This inverse can remove heavy tails from data,
7 and also provide analytical expressions for the cumulative distribution (cdf) and probability
8 density function (pdf). As a special case, these yield explicit formulas for Tukey's h pdf
9 and cdf - to the author's knowledge for the first time in the literature. Simulations and
10 applications to S&P 500 log-returns and solar flares data demonstrate the usefulness of the
11 introduced methodology.

12 The R package [LambertW](#) implementing the presented methodology is publicly available
13 at [CRAN](#).

14	Contents	
15	1 Introduction	1
16	1.1 Multivariate Extensions	4
17	1.2 Box-Cox Transformation to Remove Heavy Tails	5
18	2 Generating Heavy Tails Using Transformations	5
19	2.1 Tukey's h Distribution	6
20	2.2 Heavy Tail Lambert W Random Variables	7
21	2.3 Inverse Transformation: "Gaussianize" Heavy-Tailed Data	8
22	2.4 Distribution and Density	10
23	2.5 Quantile Function	12
24	3 Tukey's h distribution: Gaussian input	13
25	3.1 Tukey's h versus student's t	14
26	4 Parameter Estimation	15
27	4.1 Maximum Likelihood Estimation (MLE)	15
28	4.1.1 Properties of The MLE For The Heavy Tail Parameter	17
29	4.2 Iterative Generalized Method of Moments (IGMM)	18
30	5 Simulations	19
31	5.1 Estimating δ Only	19
32	5.2 Estimating All Parameters Jointly	20
33	5.3 Discussion of the Simulations	23
34	6 Applications	23
35	6.1 Estimating Location of a Cauchy With The Sample Mean	24
36	6.2 Heavy Tails in Finance: S&P 500 Case Study	25
37	6.2.1 Gaussian Fit to Returns	26
38	6.2.2 Heavy Tail Fit to Returns	27
39	6.2.3 "Gaussianizing" Returns	28
40	6.2.4 Gaussian MLE for Gaussianized Data	28
41	6.3 Removing Power Law From Solar Flare Counts	29
42	7 Discussion and Outlook	31
43	References	33
44	A Auxiliary Results and Properties	39
45	A.1 Inverse Transformation $W_\delta(z)$	39
46	A.2 Penalty $\log R(\delta z_i)$ for Standard Gaussian Input	41
47	A.3 Gaussian log-Likelihood at $W_\delta(z)$	42

48	B Proofs	42
49	B.1 Inverse transformation	42
50	B.2 Cdf and pdf	43
51	B.3 MLE for δ	44
52	C Details on IGMM	46
53	D Simulation Details	49

1 Introduction

Statistical theory and practice are both tightly linked to Gaussianity. In theory, many methods require Gaussian data or noise: i) regression often assumes Gaussian errors; ii) pattern recognition for images often model noise as a Gaussian random field (Achim, Tsakalides, and Bezerianos, 2003); iii) many time series models are based on Gaussian white noise (Brockwell and Davis, 1998; Engle, 1982; Granger and Joyeux, 2001).

In all these cases, a model $\mathcal{M}_{\mathcal{N}}$, parameter estimates and their standard errors, and other properties, are then studied – all based on the ideal(istic) assumption of Gaussianity.

In practice, however, data/noise often exhibits asymmetry and heavy tails; for example wind speed data (Field, 2004), human dynamics

(Vázquez, Oliveira, Dezső, Goh, Kondor, and Barabási, 2006), or Internet traffic data (Gidlund and Debernardi, 2009) – just to name a few. Particularly notable examples are financial data (Cont, 2001; Kim and White, 2003) and speech signals (Aysal and Barner, 2006), which almost exclusively exhibit heavy tails. Thus a model $\mathcal{M}_{\mathcal{N}}$ developed for the Gaussian case does not necessarily provide accurate inference anymore.

One way to overcome this shortcoming is to replace $\mathcal{M}_{\mathcal{N}}$ with a new model $\mathcal{M}_G$, where G is a heavy tail distribution: i) regression with Cauchy errors (Smith, 1973); ii) image denoising for α -stable noise (Achim et al., 2003); iii) forecasting long memory processes with heavy tail innovations (Ilow, 2000; Palma and Zevallos, 2011), or ARMA modeling of electricity loads with hyperbolic noise (Nowicka-Zagrajek and Weron, 2002).

While such fundamental approaches are attractive from a theoretical perspective, they can become unsatisfactory from a practical viewpoint. Many successful statistical models assume Gaussianity, their theory is very well understood, and many algorithms are implemented for the simple – and often much faster – Gaussian case. Thus developing models based on an entirely unrelated distribution G is like throwing out the (Gaussian) baby with the bathwater.

It would be very useful to transform a Gaussian RV X to a heavy-tailed RV Y and vice




Figure 1: Schematic view of the heavy tail Lambert $W \times F_X$ framework. (left) Latent input $X \sim F_X$: $H_\tau(X)$ from (6) transforms (solid arrows) X to $Y \sim \text{Lambert } W \times F_X$ and generates heavy tails. (right) Observed heavy tail world Y and $\mathbf{y}$: (1) use $W_\tau(\cdot)$ to back-transform $\mathbf{y}$ to latent “Normal” $\mathbf{x}_\tau$, (2) use model $\mathcal{M}_\mathcal{N}$ of your choice (regression, time series models, hypothesis testing, etc.) for inference on $\mathbf{x}_\tau$, and (3) convert results back to the original “heavy-tailed world” of $\mathbf{y}$.

84 versa, and thus rely on knowledge - and software - for the well-understood Gaussian case,
85 while still capturing heavy tails in the data. Optimally such a transformation should: a) be
86 bijective; b) include Normality as a special case for hypothesis testing; and c) be parametric
87 so the optimal transformation can be estimated efficiently.

88 Figure 1 illustrates this pragmatic approach: researchers can make their observations
89 $\mathbf{y}$ as Gaussian as possible ($\mathbf{x}_\tau$) before making inference based on their favorite Gaussian
90 model $\mathcal{M}_\mathcal{N}$. This avoids the development of - or the data analysts waiting for - a whole
91 new theory of $\mathcal{M}_G$ and new implementations based on a particular heavy-tailed distribution
92 G , while still improving statistical inference on heavy-tailed data $\mathbf{y}$. For example, consider
93 $\mathbf{y} = (y_1, \dots, y_{500})$ from a standard Cauchy distribution $\mathcal{C}(0, 1)$ in Fig. 2a: modeling heavy
94 tails by a transformation makes it even possible to Gaussianize this Cauchy sample (Fig.

95 **2c**). This “nice” data $\mathbf{x}_\tau$ can then be subsequently analyzed with common techniques. For
 96 example, the location can now be estimated using the sample average (Fig. **2d**). For details
 97 see Section **6.1**.

98

99 Liu, Lafferty, and Wasserman (2009) use a semi-parametric approach, where Y has a
 100 *nonparanormal* distribution if $f(Y) \sim \mathcal{N}(\mu, \sigma^2)$ where $f(\cdot)$ is an increasing smooth func-
 101 tion; they estimate $f(\cdot)$ using non-parametric methods. This leads to a greater flexibility
 102 in the distribution of Y , but it suffers from two drawbacks: i) non-parametric methods
 103 have slower convergence rates and thus need large samples, and ii) for identifiability of
 104 $f(\cdot)$, $\mathbb{E}f(Y) \equiv \mathbb{E}Y$ and $\mathbb{V}f(Y) \equiv \mathbb{V}Y$ must hold. While **i**) is inherent to non-parametric
 105 methods, point **ii**) requires Y to have finite mean and variance, which is especially limiting
 106 for heavy-tailed data where this condition is often not met. Thus here we use parametric
 107 transformations which do not rely on restrictive identifiability conditions and also work well
 108 for small sample sizes.

109

110 The main contributions of this work are three-fold: a) following Goerg (2011) I introduce
 111 a meta-family of heavy tail Lambert $W \times F_X$ distributions with Tukey’s h (Hoaglin, 2006)
 112 as a special case; b) I present a bijective transformation to “Gaussianize” heavy-tailed data
 113 (Section **2**); and c) I also provide simple expressions for the cumulative distribution function
 114 (cdf) $G_Y(y)$ and probability density function (pdf) $g_Y(y)$ - also for Tukey’s h -, which can
 115 be easily implemented in statistics software (Section **2.4**).

116 To the author’s knowledge analytic expressions for Tukey’s h cdf and pdf are presented
 117 here (Section **3**) for the first time in the literature. Section **4** introduces a methods of
 118 moments estimator and studies the maximum likelihood estimator (MLE). Section **5** shows
 119 their finite sample properties.

120 As has been shown in many case studies, Tukey’s h distribution (heavy tail Lambert
 121 $W \times$ Gaussian) is useful to model data with unimodal, heavy-tailed densities. Section **6**
 122 not only confirms this finding for S&P 500 log-returns, but also demonstrates the benefits
 123 of removing heavy tails for exploratory data analysis: Gaussianizing γ -ray intensity data
 124 reveals a bimodal density, which even non-parametric estimators fail to detect if heavy tails




Figure 2: Gaussianizing a standard Cauchy sample. For (d) $\tau^{(n)}$ was estimated for each fixed $n = 5, \dots, 500$, before Gaussianizing $(y_1, \dots, y_n)$.

are not removed. Finally, we discuss the new methodology and future work in Section 7. All proofs are given in the Supplementary Material, Appendix B. Computations, figures, and simulations were done in R (R Development Core Team, 2010). The R package LambertW is publicly available on CRAN.

1.1 Multivariate Extensions

While this work focuses on univariate case, multivariate extensions of the presented methods can be defined component-wise – analogously to the multivariate version of Tukey’s h distribution (Field and Genton, 2006). While this may not make the transformed RVs jointly Gaussian, it still provides a good starting point for more well-behaved multivariate modeling.

1.2 Box-Cox Transformation to Remove Heavy Tails

A popular method to deal with skewed, high variance data is the Box-Cox transformation

$$\mathbf{y}_\lambda = \begin{cases} \frac{\mathbf{y}^\lambda - 1}{\lambda} & \text{if } \lambda > 0 \\ \log \mathbf{y} & \text{if } \lambda = 0. \end{cases} \quad (1)$$

The parameter λ can be chosen by MLE. However one major limitation of (1) is the non-negativity constraint on $\mathbf{y}$, which prohibits its use in many applications. To avoid this limitation it is common to shift the data, $\tilde{\mathbf{y}} = \mathbf{y} + |\min(\mathbf{y})| \geq 0$. However, as Fig. 2b shows applying the Box-Cox transformation to the Cauchy sample¹ completely fails. Furthermore, this restricts Y to a half-open interval $[c, \infty)$ and is not desirable if the underlying process can occur on the entire real line, since it undermines statistical inference for yet unobserved data. See Sakia (1992) for a more detailed discussion and the Box-Cox transformation in general.

Furthermore, the main purpose of the Box-Cox transformation is to stabilize variance (Blaylock, Salathe, and Green, 1980; Lawrance, 1987; Tsiotas, 2007) and remove right tail skewness (Goncalves and Meddahi, 2011); a lower empirical kurtosis is merely a by-result of the variance stabilization. In contrast, the Lambert W framework is designed to model heavy-tailed RVs and remove heavy tails from data, and has no difficulties with negative values.

2 Generating Heavy Tails Using Transformations

Random variables exhibit heavy tails if more mass than for a Gaussian RV lies at the outer end of the density support. A RV Z has a tail index a if its cdf satisfies $1 - F_Z(z) \sim L(z)z^{-a}$, where $L(z)$ is a slowly varying function at infinity, i.e. $\lim_{z \rightarrow \infty} \frac{L(tz)}{L(z)} = 1$ for all $t > 0$ (Baek and Phipras, 2010).² The heavy tail index a is an important characteristic of Z ; for example, only moments up to order a exist.

¹We use $\tilde{\mathbf{y}} = \mathbf{y} + |\min(\mathbf{y})| + 1$ and use `boxcox` from the MASS R package; $\hat{\lambda} = 0.37$.

²There are various similar definitions of heavy/fat/long tails; for this work these differences are not essential.




Figure 3: Transformation and inverse transformation for $\delta_\ell = 0$ and $\delta_r = 1/10$: identity on the left (same tail behavior) and a heavy-tailed transformation in the right tail of input U .

2.1 Tukey's h Distribution

A parametric transformation is the basis of Tukey's h RVs (Hoaglin, 2006)

$$Z = U \exp\left(\frac{h}{2}U^2\right), \quad h \geq 0, \quad (2)$$

where U is standard Normal RV and h is the heavy tail parameter. The RV Z has tail parameter $a = 1/h$ (Morgenthaler and Tukey, 2000) and reduces to the Gaussian for $h = 0$. Morgenthaler and Tukey (2000) extend the h distribution to the skewed, heavy-tailed family of hh RVs

$$Z = \begin{cases} U \exp\left(\frac{\delta_\ell}{2}U^2\right), & \text{if } U \leq 0, \\ U \exp\left(\frac{\delta_r}{2}U^2\right), & \text{if } U > 0, \end{cases} \quad (3)$$

where again $U \sim \mathcal{N}(0,1)$. Here $\delta_\ell \geq 0$ and $\delta_r \geq 0$ shape the left and right tail of Z , respectively; thus transformation (3) can model skewed and heavy-tailed data - see Fig. 3a. For simplicity let $H_\delta(u) := u \exp\left(\frac{\delta}{2}u^2\right)$.

Despite their great flexibility they are not popular in statistical practice, because the inverse of (2) or (3) has not been found. Consequently, no closed-form expressions for the cdf or pdf are available. Although Morgenthaler and Tukey (2000) express the pdf of (2) as ($h \equiv \delta$)

$$g_Z(z) = \frac{f_U(H_\delta^{-1}(z))}{H'_\delta(H_\delta^{-1}(z))}, \quad (4)$$

they fall short of explicitly specifying $H_\delta^{-1}(z)$. So far this inverse has been considered an-

162 alytically intractable (Field, 2004), or was only numerically approximated (Fischer, 2010;
 163 Headrick, Kowalchuk, and Sheng, 2008). Thus parameter inference relied on matching em-
 164 pirical and theoretical quantiles (Field, 2004; Morgenthaler and Tukey, 2000), or by the
 165 method of moments (Headrick et al., 2008). Only recently Headrick et al. (2008) provided
 166 numerical approximations. Hence, a closed form, analytically tractable pdf that can be
 167 computed efficiently is essential for a wide-spread use of Tukey’s h (& variants).

168

169 In this work I present this long sought explicit inverse, which is readily available in
 170 standard statistics software. For ease of notation and concision main results are shown for
 171 $\delta_\ell = \delta_r = \delta$; analogous results for $\delta_\ell \neq \delta_r$ will be stated without details.

172 2.2 Heavy Tail Lambert W Random Variables

173 Tukey’s h transformation (2) is strongly related to the approach taken by Goerg (2011) to
 174 introduce skewness in continuous RVs $X \sim F_X(x)$. In particular, if $Z \sim$ Tukey’s h , then
 175 $Z^2 \sim$ skewed Lambert $W \times \chi_1^2$ with skew parameter $\gamma = h$.

176 Adapting the skew Lambert $W \times F_X$ input/output idea³ (see Fig. 1), Tukey’s h RVs can
 177 be generalized to *heavy-tailed Lambert $W \times F_X$ RVs*.

Definition 2.1. Let U be a continuous RV with cdf $F_U(u | \beta)$, pdf $f_U(u | \beta)$, and parameter vector β . Then,

$$Z = U \exp\left(\frac{\delta}{2}U^2\right), \quad \delta \in \mathbb{R}, \quad (5)$$

178 is a non-central, non-scaled heavy tail Lambert $W \times F_X$ RV with parameter vector $\theta =$
 179 (β, δ) , where δ is the tail parameter.

180 Tukey’s h distribution results for U being a standard Gaussian $\mathcal{N}(0, 1)$.

Definition 2.2. For a continuous location-scale family RV $X \sim F_X(x | \beta)$ define a location-scale heavy-tailed Lambert $W \times F_X$ RV

$$Y = \left\{ U \exp\left(\frac{\delta}{2}U^2\right) \right\} \sigma_x + \mu_x, \quad \delta \in \mathbb{R}, \quad (6)$$

³Most concepts and methods from the skew Lambert $W \times F_X$ case transfer one-to-one to the heavy tail Lambert W RVs presented here. Thus for the sake of concision I refer to Goerg (2011) for details and background information on the Lambert W framework.

181 with parameter vector $\theta = (\beta, \delta)$, where $U = (X - \mu_x)/\sigma_x$.

182 The input is not necessarily Gaussian but can be any other location-scale continuous RV,
 183 e.g., from a uniform distribution: $X \sim U(a, b)$.

Definition 2.3. Let $X \sim F_X(x/s \mid \beta)$ be a continuous scale-family RV, with standard deviation σ_x ; let $U = X/\sigma_x$. Then,

$$Y = X \exp\left(\frac{\delta}{2}U^2\right), \quad \delta \in \mathbb{R}, \quad (7)$$

184 is a scaled heavy-tailed Lambert $W \times F_X$ RV with parameter $\theta = (\beta, \delta)$.

185 Let $\tau := (\mu_x(\beta), \sigma_x(\beta), \delta)$ define transformation (6).⁴ If $X \in (-\infty, \infty)$, then for all $\delta \geq 0$
 186 also the location-scale $Y \in (-\infty, \infty)$. For a scale family $X \in [0, \infty)$ also the scale Lambert
 187 $W \times F_X$ RV $Y \in [0, \infty)$.

188

189 The shape parameter δ (= Tukey's h) governs the tail behavior of Y : for $\delta > 0$ values
 190 further away from μ_x are increasingly emphasized, leading to a heavy-tailed version of
 191 $F_X(x)$; for $\delta = 0$, $Y \equiv X$; and for $\delta < 0$ values far away from the mean are mapped
 192 back again closer to μ_x . Thus heavy tail Lambert $W \times F_X$ RVs generalize $X \sim F_X(x)$ to
 193 heavy-tailed versions of itself, $Y \sim G_Y(y)$, with a reduction to X for $\delta = 0$.

194 The Lambert W formulation of heavy tail modeling is more general than Tukey's h
 195 distribution as X can have any distribution $F_X(x)$, not necessarily Gaussian (Fig. 4).

196 **Remark 2.4** (Only non-negative δ). Although $\delta < 0$ leads to interesting properties of Y ,
 197 it yields a non-bijective transformation and thus to parameter-dependent support and non-
 198 unique input. Thus for the remainder of this work I tacitly assume $\delta \geq 0$, unless stated
 199 otherwise.

200 2.3 Inverse Transformation: “Gaussianize” Heavy-Tailed Data

201 Transformation (6) is bijective and its inverse can be obtained via Lambert's W function,
 202 which is the inverse of $z = u \exp(u)$, i.e., that function which satisfies $W(z) \exp(W(z)) = z$.

⁴For non-central, non-scale input set $\tau = (0, 1, \delta)$; for scale-family input $\tau = (0, \sigma_x, \delta)$.

Lambert’s W has been studied extensively in mathematics, physics, and other areas of science (Corless, Gonnet, Hare, and Jeffrey, 1996; Rosenlicht, 1969; Valluri, Jeffrey, and Corless, 2000), and is implemented in the GNU Scientific Library (GSL) (Galassi, Davies, Theiler, Gough, Jungman, Alken, Booth, and Rossi, 2011). Only very recently it received attention in the statistics literature (Goerg, 2011; Jodrá, 2009; Pakes, 2011; Rathie and Silva, 2011). It has many useful properties (see Appendix A and Corless et al. (1996)), in particular $W(z)$ is bijective for $z \geq 0$.

Lemma 2.5. *The inverse transformation of (6) is*

$$W_\tau(Y) := W_\delta \left(\frac{Y - \mu_x}{\sigma_x} \right) \sigma_x + \mu_x = U \sigma_x + \mu_x = X, \quad (8)$$

where

$$W_\delta(z) := \text{sgn}(z) \left(\frac{W(\delta z^2)}{\delta} \right)^{1/2}, \quad (9)$$

and $\text{sgn}(z)$ is the sign of z . $W_\delta(z)$ is bijective for all $\delta \geq 0$ and all $z \in \mathbb{R}$.

Lemma 2.5 gives for the first time an analytic, bijective inverse of Tukey’s h transformation: $H_\delta^{-1}(y)$ of Morgenthaler and Tukey (2000) is now analytically available as (8). Bijectivity implies that for any data $\mathbf{y}$ and parameter τ , the exact input $\mathbf{x}_\tau = W_\tau(\mathbf{y}) \sim F_X(x)$ can be obtained.

In view of the importance and popularity of Gaussianity, we clearly want to back-transform heavy-tailed data to a Gaussian rather than yet another heavy-tailed distribution. Typically tail behavior of RVs are compared by their kurtosis $\gamma_2(X) = \mathbb{E}(X - \mu_x)^4 / \sigma_x^4$, which for a Gaussian RV equals 3. Hence for the future when we “normalize $\mathbf{y}$ ” we can not only subtract the mean, and divide by the standard deviation, but also transform it to $\mathbf{x}_\tau$ with $\hat{\gamma}_2(\mathbf{x}_\tau) = 3$ – a “Normalization” in the true sense of the word (see Fig. 2c).

This data-driven view of the Lambert W framework can also be useful for kernel density estimation (KDE), where multivariate data is often pre-scaled to unit-variance, so the same bandwidth can be used in each dimension (Hwang, Lay, and Lippman, 1994; Wasserman, 2007). Thus “normalizing” the Lambert Way might likely also improve KDE for heavy-tailed data (see also Maiboroda and Markovich, 2004; Markovich, 2005).

Corollary 2.6 (Inverse transformation for Tukey's hh). *The inverse transformation of (3) is*

$$W_{\delta_\ell, \delta_r}(z) = \begin{cases} W_{\delta_\ell}(z), & \text{if } z \leq 0, \\ W_{\delta_r}(z), & \text{if } z > 0. \end{cases} \quad (10)$$

Figure 3b shows $W_{\delta_\ell, \delta_r}(z)$ for $\delta_\ell = 0$ and $\delta_r = 1/10$. The transformation in Fig. 3a generates a right heavy tail version of U (x-axis) by stretching only the positive axis (y-axis). By definition $W_{\delta_\ell, \delta_r}(z)$ removes the heavier right tail in Z (positive y-axis). Figure 3c shows how $W_\delta(z)$ operates for various degrees of heavy tails and $z \in [0, 3]$. If δ is close to zero, then also $W_\delta(z) \approx z$; for larger δ , the inverse maps z to (much) smaller u .

Remark 2.7 (Generalized transformation). *Transformation (2) can be generalized to*

$$Z = U \exp\left(\frac{\delta}{2\alpha} (U^2)^\alpha\right), \quad \alpha > 0. \quad (11)$$

The inner term U^2 guarantees bijectivity for all $\alpha > 0$. The inverse is

$$W_{\delta, \alpha}(z) := \text{sgn}(z) \left(W\left(\frac{\delta z^{2\alpha}}{\delta}\right) \right)^{\frac{1}{2\alpha}}. \quad (12)$$

For comparison with Tukey's h I consider $\alpha = 1$ only. For $\alpha = 1/2$ transformation (11) is closely related to skewed Lambert $W \times F_X$ distributions.

2.4 Distribution and Density

For ease of notation let

$$z = \frac{y - \mu_x}{\sigma_x}, \quad u = W_\delta(z), \quad \text{and } x = W_\tau(y) = u\sigma_x + \mu_x. \quad (13)$$

Theorem 2.8 (Distribution and Density of Y). *The cdf and pdf of a location-scale heavy tail Lambert $W \times F_X$ RV Y equal*

$$G_Y(y \mid \beta, \delta) = F_X(W_\delta(z)\sigma_x + \mu_x \mid \beta) \quad (14)$$




Figure 4: Pdf (top) and cdf (bottom) of a heavy-tail (a) “non-central, non-scaled”, (b) “scale”, and (c and d) “location-scale” Lambert $W \times F_X$ RV Y for various degrees of heavy tails (color, dashed lines).

and

$$g_Y(y | \beta, \delta) = f_X \left(W_\delta \left(\frac{y - \mu_x}{\sigma_x} \right) \sigma_x + \mu_x | \beta \right) \cdot \frac{W_\delta \left(\frac{y - \mu_x}{\sigma_x} \right)}{\frac{y - \mu_x}{\sigma_x} \left[1 + W \left(\delta \left(\frac{y - \mu_x}{\sigma_x} \right)^2 \right) \right]}. \quad (15)$$

Clearly, $G_Y(y | \beta, \delta = 0) = F_X(y | \beta)$ and $g_Y(y | \beta, \delta = 0) = f_X(y | \beta)$, since $\lim_{\delta \rightarrow 0} W_\delta(z) = z$ and $\lim_{\delta \rightarrow 0} W(\delta z^2) = 0$ for all $z \in \mathbb{R}$.

For scale family or non-central, non-scale input set $\mu_x = 0$ or $\mu_x = 0, \sigma_x = 1$.

The explicit formula (15) allows a fast computation and theoretical analysis of the likelihood, which is essential for – either frequentist or Bayesian – statistical inference. Detailed properties of (15) are given in Section 4.1.

Figure 4 shows (14) and (15) for various $\delta \geq 0$ with for four different input $X \sim F_X(x | \beta)$: for $\delta = h = 0$ the input equals the output (solid black); for larger δ the tails of $G_Y(y | \theta)$ and $g_Y(y | \theta)$ get heavier (dashed colored).

Corollary 2.9. *The cdf and pdf of Z in (3) equal*

$$G_Z(z \mid \boldsymbol{\beta}, \delta_\ell, \delta_r) = \begin{cases} G_Z(z \mid \boldsymbol{\beta}, \delta_\ell), & \text{if } z \leq 0, \\ G_Z(z \mid \boldsymbol{\beta}, \delta_r), & \text{if } z > 0, \end{cases} \quad (16)$$

and

$$g_Z(z \mid \boldsymbol{\beta}, \delta) = \begin{cases} g_Z(z \mid \boldsymbol{\beta}, \delta_\ell), & \text{if } z \leq 0, \\ g_Z(z \mid \boldsymbol{\beta}, \delta_r), & \text{if } z > 0. \end{cases} \quad (17)$$

2.5 Quantile Function

Quantile fitting has been the standard technique to estimate μ_x , σ_x , and δ of Tukey’s h . In particular, the median of Y and X are equal. Thus for symmetric, location-scale family input the sample median of $\mathbf{y}$ is a robust estimate for μ_x for any $\delta \geq 0$ (see also Section 5). General quantiles can be computed via (Hoaglin, 2006)

$$y_\alpha = u_\alpha \exp\left(\frac{\delta}{2} u_\alpha^2\right) \sigma_x + \mu_x, \quad (18)$$

where $u_\alpha = W_\delta(z_\alpha)$ are the α -quantiles of $F_U(u)$. As quantiles of U are typically tabulated, or easily available in software packages, (18) can be computed very efficiently using u_α and τ .

This simple conversion can be especially useful for education: teaching heavy-tailed statistics in introductory courses soon becomes too difficult using e.g., Cauchy or α -stable distributions. Yet, transforming data via Lambert’s W , using previously learned methods for the Gaussian case, and then transforming the inference back to the “heavy-tailed world” – e.g., transforming quantiles using (18) – is straightforward. Thus the Lambert $W \times F_X$ framework can promote heavy-tailed statistics in introductory courses.

3 Tukey's h distribution: Gaussian input

For Gaussian input Lambert $W \times F_X$ equals Tukey's h , which has been studied extensively.

[Dutta and Babbal \(2002\)](#) show that

$$\mathbb{E}Z^n = \begin{cases} 0, & \text{if } n \text{ is odd and } n < \frac{1}{\delta}, \\ \frac{n!(1-n\delta)^{\frac{-(n+1)}{2}}}{2^{n/2}(n/2)!}, & \text{if } n \text{ is even and } n < \frac{1}{\delta}, \\ \nexists, & \text{if } n > \frac{1}{\delta}, \end{cases} \quad (19)$$

which in particular implies ([Headrick et al., 2008](#))

$$\mathbb{E}Z = \mathbb{E}Z^3 = 0, \text{ if } \delta < 1 \text{ and } 1/3, \text{ respectively} \quad (20)$$

$$\text{and } \mathbb{E}Z^2 = \frac{1}{(1-2\delta)^{3/2}}, \text{ if } \delta < \frac{1}{2}, \quad \mathbb{E}Z^4 = 3\frac{1}{(1-4\delta)^{5/2}}, \text{ if } \delta < \frac{1}{4}. \quad (21)$$

Thus the kurtosis of Y equals (see Fig. 5)

$$\gamma_2(\delta) = 3\frac{(1-2\delta)^3}{(1-4\delta)^{5/2}} \text{ for } \delta < 1/4. \quad (22)$$

For $\delta = 0$, (21) and (22) reduce to the familiar Gaussian values.

Corollary 3.1. *The cdf of Tukey's h equals*

$$G_Y(y \mid \mu_x, \sigma_x, \delta) = \Phi\left(\frac{W_\tau(y) - \mu_x}{\sigma_x}\right), \quad (23)$$

where $\Phi(u)$ is the cdf of a standard Normal. The pdf equals (for $\delta > 0$)

$$g_Y(y \mid \mu_x, \sigma_x, \delta) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1+\delta}{2} W_\delta\left(\frac{y-\mu_x}{\sigma_x}\right)^2\right) \cdot \frac{1}{1 + W\left(\delta\left(\frac{y-\mu_x}{\sigma_x}\right)^2\right)} \quad (24)$$

Proof. Take $X \sim \mathcal{N}(\mu_x, \sigma_x^2)$ in Theorem 2.8. □

Section 4.1 studies functional properties of (24) in more detail.




Figure 5: Comparing moments of Lambert $W \times$ Gaussian and student's t .

3.1 Tukey's h versus student's t

Student's t_ν distribution with ν degrees of freedom is often used to model heavy-tailed data (Wong, Chan, and Kam, 2009; Yan, 2005), as its tail index equals ν . Thus the n th moment of a student t RV T exists if $n < \nu$. In particular,

$$\mathbb{E}T = \mathbb{E}T^3 = 0 \text{ if } \nu < 1 \text{ or } < 3, \quad \mathbb{E}T^2 = \frac{\nu}{\nu - 2} = \frac{1}{1 - \frac{2}{\nu}} \text{ if } \frac{1}{\nu} < \frac{1}{2}, \quad (25)$$

and kurtosis

$$\gamma_2(\nu) = 3 \frac{\nu - 2}{\nu - 4} = 3 \frac{1 - 2\frac{1}{\nu}}{1 - 4\frac{1}{\nu}} \text{ if } \frac{1}{\nu} < \frac{1}{4}. \quad (26)$$

Comparing (26) and (21) with (22) and (25) shows a natural association between $1/\nu$ and δ and a close similarity between the first four moments of student's t and Tukey's h (Fig. 5). By continuity and monotonicity, the first four moments of a location-scale t distribution can always be exactly matched by a corresponding location-scale Lambert $W \times$ Gaussian. Thus if student's t is used to model heavy tails, and not as the true distribution of a test statistic, it might be worthwhile to also fit heavy tail Lambert $W \times$ Gaussian distributions for an equally valuable “second opinion”. For example, a parallel analysis on S&P 500 log-returns in Section 6.2 leads to divergent inference regarding the existence of fourth moments. Additionally, the Lambert W approach allows to Gaussianize and thus reveal hidden patterns in the data; patterns that can be easily overseen in presence of heavy tails (Section 6.3).

4 Parameter Estimation

For a sample of N independent identically distributed (i.i.d.) observations $\mathbf{y} = (y_1, \dots, y_N)$ from transformation (6), $\theta = (\beta, \delta)$ has to be estimated from the data. Due to the lack of a closed form pdf of Y , this has been typically done by matching quantiles or a method of moments estimator (Field, 2004; Headrick et al., 2008; Morgenthaler and Tukey, 2000). These inefficient methods can now be replaced by the – fast and usually efficient – maximum likelihood estimator (MLE) using the pdf in (15). Rayner and MacGillivray (2002) introduce a numerical MLE procedure based on quantile functions, but they conclude that “sample sizes significantly larger than 100 should be used to obtain reliable estimates through maximum likelihood”. Simulations in Section 5 show that log-likelihood maximization with the Lambert W methodology converges quickly and is accurate even for sample sizes as small as $N = 10$.

4.1 Maximum Likelihood Estimation (MLE)

For an i.i.d. sample $\mathbf{y} \sim g_Y(y | \beta, \delta)$ the log-likelihood function equals

$$\ell(\theta | \mathbf{y}) = \sum_{i=1}^N \log g_Y(y_i | \beta, \delta). \quad (27)$$

The MLE is that $\theta = (\beta, \delta)$ which maximizes (27), i.e.

$$\hat{\theta}_{MLE} = \left(\hat{\beta}, \hat{\delta} \right)_{MLE} = \arg \max_{\beta, \delta} \ell(\beta, \delta | \mathbf{y}). \quad (28)$$

Since $g_Y(y_i | \beta, \delta)$ is a function of $f_X(x_i | \beta)$, the MLE depends on the specification of the input density. Eq. (27) can be decomposed as

$$\ell(\beta, \delta | \mathbf{y}) = \ell(\beta | \mathbf{x}_\tau) + \mathcal{R}(\tau | \mathbf{y}), \quad (29)$$

where

$$\ell(\beta | \mathbf{x}_\tau) = \sum_{i=1}^N \log f_X \left(W_\delta \left(\frac{y_i - \mu_x}{\sigma_x} \right) \sigma_x + \mu_x | \beta \right) = \sum_{i=1}^N \log f_X(\mathbf{x}_\tau | \beta) \quad (30)$$




Figure 6: Log-likelihood decomposition for Lambert $W \times F_X$ distributions.

is the log-likelihood of the back-transformed data $\mathbf{x}_\tau$ and

$$\mathcal{R}(\tau | \mathbf{y}) = \sum_{i=1}^n \log R(\mu_x, \sigma_x, \delta | y_i), \quad (31)$$

where

$$R(\mu_x, \sigma_x, \delta | y_i) = \frac{W_\delta \left(\frac{y_i - \mu_x}{\sigma_x} \right)}{\frac{y_i - \mu_x}{\sigma_x} \left[1 + \delta \left(W_\delta \left(\frac{y_i - \mu_x}{\sigma_x} \right) \right)^2 \right]}. \quad (32)$$

Note that $R(\mu_x, \sigma_x, \delta | y_i)$ only depends on $\mu_x(\beta)$ and $\sigma_x(\beta)$ (and δ), but not necessarily on every coordinate of β .

Decomposition (29) shows the difference between the exact MLE $(\hat{\beta}, \hat{\delta})$ based on $\mathbf{y}$ and the approximate MLE $\hat{\beta}_{\mathbf{x}_\tau}$ based on $\mathbf{x}_\tau$ alone: if we knew $\tau = (\mu_x, \sigma_x, \delta)$ beforehand, then we could back-transform $\mathbf{y}$ to $\mathbf{x}_\tau$ and estimate $\hat{\beta}_{\mathbf{x}_\tau}$ from $\mathbf{x}_\tau$ (maximize (30) with respect to β). In practice, however, τ must also be estimated and this enters the likelihood via the additive term $\mathcal{R}(\tau | \mathbf{y})$. A little calculation shows that for any $y_i \in \mathbb{R}$, $\log R(\mu_x, \sigma_x, \delta | y_i) \leq 0$ if $\delta \geq 0$, with equality if and only if $\delta = 0$. Thus $\mathcal{R}(\tau | \mathbf{y})$ can be interpreted as a penalty for transforming the data. Maximizing (29) faces a trade-off between transforming the data to follow $f_X(x | \beta)$ (and thus increasing $\ell(\beta | \mathbf{x}_{\hat{\tau}})$) versus the penalty of a more extreme transformation (and thus decreasing $\mathcal{R}(\tau | \mathbf{y})$) – see Fig. 6b.

Figure 6a shows a contour plot of $R(\mu_x = 0, \sigma_x = 1, \delta | y)$ as a function of δ and $y = z$. The penalty for transforming the data increases (in absolute value) either if δ gets larger (for fixed y) or for larger y (for fixed δ). In both cases, increasing δ makes the transformed data $W_\delta(z)$ get closer to $0 = \mu_x$, which in turn increases its input likelihood. For $\delta = 0$, the penalty disappears since input equals output; for $y = 0$ there is no penalty since $W_\delta(0) = 0$ for all δ .

Figure 6b shows a random sample ($N = 1000$) $\mathbf{z} \sim \text{Lambert } W \times \text{Gaussian}$ with $\delta = 1/3$ and the decomposition of the log-likelihood as in (29). Since $\beta = (0, 1)$ is known, the likelihood and penalty are only functions of δ . The monotonicity of the penalty (decreasing, red) and the input likelihood (increasing, green) as a function of δ is not particular to this sample, but holds true in general (see Theorem 4.1 below). This monotonicity in each component implies that their sum (black line) has a unique maximum; here $\hat{\delta}_{MLE} = 0.37$ (blue, dashed vertical line).

311

The maximization of (29) can be carried out numerically. Here I show existence and uniqueness of $\hat{\delta}_{MLE}$ assuming that μ_x and σ_x are known. Theoretical results for $\hat{\theta}_{MLE}$ remain for future work. Given the “nice” form of $g_Y(y)$ - continuous, twice differentiable,⁵ its support does not depend on the parameter, etc. - the MLE for $\theta = (\beta, \delta)$ should have the usual optimality properties (Lehmann and Casella, 1998).

4.1.1 Properties of The MLE For The Heavy Tail Parameter

Without loss of generality let $\mu_x = 0$ and $\sigma_x = 1$. In this case

$$\ell(\delta | \mathbf{z}) \propto -\frac{1}{2} \sum_{i=1}^N [W_\delta(z_i)]^2 + \sum_{i=1}^N \log \frac{W_\delta(z_i)}{z_i} - \log \left(1 + \delta [W_\delta(z_i)]^2 \right) \quad (33)$$

$$= -\frac{1+\delta}{2} \sum_{i=1}^N [W_\delta(z_i)]^2 - \sum_{i=1}^N \log \left(1 + \delta [W_\delta(z_i)]^2 \right). \quad (34)$$

Theorem 4.1 (Unique MLE for δ). *Let Z have a Lambert $W \times$ Gaussian distribution, where $\mu_x = 0$ and $\sigma_x = 1$ are assumed to be known and fixed. Also consider only the case*

⁵Assuming that $f_X(\cdot)$ is twice differentiable.

320 $\delta \in [0, \infty)$.⁶

a) If

$$\frac{\sum_{i=1}^n z_i^4}{\sum_{i=1}^n z_i^2} \leq 3, \quad (35)$$

321 then $\hat{\delta}_{MLE} = 0$.

322 If (35) does not hold, then

b) $\hat{\delta}_{MLE} > 0$ exists and is a positive solution to

$$\sum_{i=1}^N z_i^2 W'(\delta z_i^2) \left(\frac{1}{2} W_\delta(z_i)^2 - \left(\frac{1}{2} + \frac{1}{1 + W(\delta z_i^2)} \right) \right) = 0. \quad (36)$$

323 c) There is only one such δ satisfying (36), i.e. $\hat{\delta}_{MLE}$ is unique.

324 Condition (35) says that $\hat{\delta}_{MLE} > 0$ only if the data is heavy-tailed enough. Points b) and
 325 c) guarantee that there is no ambiguity in the heavy tail estimate. This is an advantage
 326 over student's t distribution, for example, which has numerical problems and local maxima
 327 for unknown (and small) ν ($\leftrightarrow$ large δ) (see also [Fernandez and Steel, 1999](#); [Liu and Rubin,](#)
 328 [1995](#)). On the contrary, $\hat{\delta}_{MLE}$ is always a global maximum.

329

330 The log-likelihood and its gradient depend on δ and $\mathbf{z}$ only via $W_\delta(\mathbf{z})$. Given the heavy
 331 tails in $\mathbf{z}$ (for $\delta > 0$) one might expect convergence issues for larger δ (e.g. expected log-
 332 likelihood, Fisher information). However, $W_\delta(Z) \sim \mathcal{N}(0, 1)$ for the true $\delta \geq 0$, and close
 333 to a standard Gaussian if $\hat{\delta}_{MLE} \approx \delta$. Thus the performance of the MLE should not get
 334 worse for large δ as long as the initial estimate is close enough to the truth. Simulations in
 335 Section 5 support this conjecture, even for $\hat{\theta}_{MLE}$.

336 4.2 Iterative Generalized Method of Moments (IGMM)

337 A disadvantage of the MLE is the mandatory a-priori specification of the input distribution.
 338 Especially for heavy-tailed data the eye is a bad judgement to choose a particular parametric

⁶While for some samples $\mathbf{z}$ the MLE also exists for $\delta < 0$, it can not be guaranteed for all $\mathbf{z}$. If $\delta < 0$ (and $z \neq 0$), then $W_\delta(z)$ is either not unique in $\mathbb{R}$ (principal and non-principal branch) or may not even have a real-valued solution.

339 $f_X(x \mid \boldsymbol{\beta})$. It would be useful to directly estimate τ , without the intermediate step of
 340 estimating θ first (and thus no distributional assumption for the input is necessary).

341 Goerg (2011) presented an estimator for τ based on iterative generalized methods of
 342 moments (IGMM). The idea of IGMM is to find a τ such that the back-transformed data
 343 $\mathbf{x}_\tau$ has desired properties, e.g., is symmetric or has kurtosis 3. An estimator for μ_x , σ_x ,
 344 and δ can be constructed completely analogously to the skewed IGMM, with the advantage
 345 that the heavy tail transformation is bijective (the skewed transformation is not). Since the
 346 algorithm is entirely analogous to the skewed case, details are given in the Supplementary
 347 Material, Appendix C.

348
 349 An advantage of IGMM is that it requires less specific knowledge about the input dis-
 350 tribution. Usually, it is also faster than the MLE. Once $\hat{\tau}_{IGMM}$ has been obtained, the
 351 back-transformed $\mathbf{x}_{\hat{\tau}_{IGMM}}$ can be used to check if X has characteristics of a known paramet-
 352 ric distribution $F_X(x \mid \boldsymbol{\beta})$. It must be noted though that testing for a particular distribution
 353 F_X are too optimistic as $\mathbf{x}_{\hat{\tau}}$ will have “nicer” properties regarding F_X than the true $\mathbf{x}$ would
 354 have. However, estimating the transformation requires only three parameters and for a large
 355 enough sample, losing three degrees of freedom should not matter for all practical purposes.

356 5 Simulations

357 This section explores finite sample properties of estimators for $\theta = (\mu_x, \sigma_x, \delta)$ and (μ_y, σ_y)
 358 under Gaussian input $X \sim \mathcal{N}(\mu_x, \sigma_x^2)$. In particular, it compares Gaussian MLE (estima-
 359 tion of μ_y and σ_y only), IGMM and Lambert W $\times$ Gaussian MLE, and - for a heavy tail
 360 competitor – the median.⁷ All results below are based on $n = 1,000$ replications.

361 5.1 Estimating δ Only

362 Here I show finite sample properties of $\hat{\delta}_{MLE}$ for $U \sim \mathcal{N}(0, 1)$, where $\mu_x = 0$ and $\sigma_x = 1$ are
 363 known and fixed. Theorem 4.1 shows that $\hat{\delta}_{MLE}$ is unique: either at the boundary $\delta = 0$
 364 or at the globally optimal solution to (36). Results in Table 1 were obtained by numerical
 365 optimization restricted to $\delta \geq 0$ ($\Leftrightarrow \log \delta \in \mathbb{R}$) using the `nlm` function in R.

⁷For IGMM, optimization was restricted to $\delta \in [0, 10]$.

Table 1: Finite sample properties of $\hat{\delta}_{MLE}$. For each N , δ was estimated $n = 1,000$ times from a random sample $\mathbf{z} \sim \text{Tukey's } h$. The left column for each δ shows bias, $\hat{\delta}_{MLE} - \delta$; each right column shows the root mean square error (RMSE) times $\sqrt{N}$.

N	$\delta = 0$		$\delta = 1/10$		$\delta = 1/3$		$\delta = 1/2$	
10	0.025	0.191	-0.017	0.394	-0.042	0.915	-0.082	1.167
50	0.013	0.187	-0.010	0.492	-0.018	0.931	-0.016	1.156
100	0.010	0.200	-0.010	0.513	-0.009	0.914	-0.006	1.225
400	0.005	0.186	-0.003	0.528	0.000	0.927	-0.004	1.211
1000	0.003	0.197	0.000	0.532	-0.001	0.928	-0.001	1.203
2000	0.003	0.217	-0.001	0.523	0.000	0.935	-0.001	1.130
N	$\delta = 1$		$\delta = 2$		$\delta = 5$			
10	-0.054	1.987	-0.104	3.384	-0.050	7.601		
50	-0.017	1.948	-0.009	3.529	0.014	7.942		
100	-0.014	2.024	-0.001	3.294	0.011	7.798		
400	0.001	1.919	-0.002	3.433	0.001	7.855		
1000	0.001	1.955	0.001	3.553	-0.001	7.409		
2000	0.001	1.896	0.000	3.508	-0.001	7.578		

Table 1 shows that the MLE is unbiased for every δ and settles down (about $N = 100$) to an asymptotic variance, which is increasing with δ . Assuming μ_x and σ_x to be known is unrealistic and thus these finite sample properties are only an indication of the behavior of the joint MLE, $\hat{\theta}_{MLE}$. Nevertheless they are very remarkable for extremely heavy-tailed data ($\delta > 1$), where standard statistical methods typically break down. One explanation in this behavior lies in the particular form of the likelihood (33) and its gradient (36) (Theorem 4.1). Although both depend on $\mathbf{z}$, they only do so through $W_\delta(\mathbf{z}) = \mathbf{u} \sim \mathcal{N}(0, 1)$. Hence as long as $\hat{\delta}_{MLE}$ is sufficiently close to the true δ , (33) and (36) are functions of almost Gaussian RVs and standard asymptotic results should still apply.

5.2 Estimating All Parameters Jointly

Here we consider the realistic scenario where μ_x and σ_x are also unknown. We consider various sample sizes ($N = 50, 100$, and 1000) and different degrees of heavy tails, $\delta \in \{0, 1/3, 1, 1.5\}$, each one representing a particularly interesting situation: i) Gaussian data (does additional - superfluous - estimation of δ affect other estimates?), ii) fourth moments do not exist anymore, iii) non-existing mean, iv) extremely heavy-tailed data – can we get

381 useful estimates at all?

Table 2: In each subtable: (first rows) average, (middle rows) proportion of estimates below truth, (bottom rows) empirical standard deviation times $\sqrt{N}$.

(a) Truly Gaussian data: $\delta = 0$

$\delta = 0$	median	Gaussian MLE		IGMM				Lambert W MLE				NA
N		μ_y	σ_y	μ_x	σ_x	δ	σ_y	μ_x	σ_x	δ	σ_y	ratio
50	0.00	0.00	0.98	0.00	0.97	0.02	0.99	0.00	0.96	0.02	0.98	0
100	0.00	0.00	0.99	0.00	0.98	0.01	1.00	0.00	0.97	0.01	0.99	0
1000	0.00	0.00	1.00	0.00	0.99	0.00	1.00	0.00	0.99	0.00	1.00	0
50	0.50	0.50	0.57	0.51	0.60	0.66	0.54	0.51	0.65	0.66	0.56	0
100	0.50	0.51	0.56	0.51	0.62	0.62	0.53	0.52	0.65	0.62	0.56	0
1000	0.50	0.49	0.52	0.49	0.62	0.56	0.52	0.49	0.63	0.56	0.52	0
50	1.24	1.01	0.72	1.01	0.76	0.21	0.73	1.02	0.78	0.26	0.72	0
100	1.25	1.02	0.70	1.02	0.76	0.23	0.70	1.03	0.78	0.26	0.70	0
1000	1.26	0.98	0.73	0.98	0.79	0.22	0.73	0.98	0.79	0.22	0.73	0

(b) No fourth moments: $\delta = 1/3$

$\delta = 1/3$	median	Gaussian MLE		IGMM				Lambert W MLE				NA
N		μ_y	σ_y	μ_x	σ_x	δ	σ_y	μ_x	σ_x	δ	σ_y	ratio
50	0.00	0.00	1.98	0.00	1.07	0.29	∞	0.00	1.01	0.33	∞	0
100	0.00	0.00	2.03	0.00	1.04	0.31	∞	0.00	1.00	0.33	∞	0
1000	0.00	0.00	2.18	0.00	1.00	0.33	2.34	0.00	1.00	0.33	2.34	0
50	0.50	0.51	0.78	0.50	0.38	0.63	0.60	0.50	0.52	0.54	0.54	0
100	0.50	0.51	0.78	0.51	0.42	0.61	0.60	0.50	0.51	0.54	0.54	0
1000	0.48	0.51	0.77	0.51	0.47	0.56	0.55	0.51	0.50	0.53	0.52	0
50	1.27	2.21	6.56	1.44	1.45	1.10	NA	1.23	1.35	1.14	NA	0
100	1.30	2.33	11.28	1.43	1.42	1.12	NA	1.19	1.34	1.09	NA	0
1000	1.23	2.25	16.76	1.39	1.45	1.20	15.97	1.17	1.33	1.08	12.30	0

(c) Non-existing mean: $\delta = 1$

$\delta = 1$	median	Gaussian MLE		IGMM				Lambert W MLE				NA
N		μ_y	σ_y	μ_x	σ_x	δ	σ_y	μ_x	σ_x	δ	σ_y	ratio
50	0.00	-0.10	24.6	-0.01	1.18	0.90	∞	0.00	1.01	0.99	∞	0
100	0.00	0.74	72.4	0.00	1.09	0.95	∞	0.00	1.01	0.99	∞	0
1000	0.00	3.84	348.1	0.00	1.01	1.00	∞	0.00	1.00	1.00	∞	0
50	0.53	0.52	1.0	0.51	0.34	0.65	1	0.51	0.52	0.52	1	0
100	0.50	0.52	1.0	0.51	0.38	0.63	1	0.50	0.53	0.53	1	0
1000	0.49	0.52	1.0	0.51	0.48	0.53	1	0.49	0.51	0.51	1	0
50	1.27	65.85	424.3	2.10	2.50	2.32	NA	1.19	1.70	2.16	NA	0
100	1.30	410.75	4050.2	2.01	2.28	2.59	NA	1.17	1.74	2.25	NA	0
1000	1.26	3307.58	104052.7	1.93	2.21	2.81	NA	1.11	1.64	2.18	NA	0

(d) Extreme heavy tails: $\delta = 1.5$

$\delta = 1.5$	median	Gaussian MLE		IGMM				Lambert W MLE				NA
N		μ_y	σ_y	μ_x	σ_x	δ	σ_y	μ_x	σ_x	δ	σ_y	ratio
50	-0.02	6.84	309	-0.02	1.23	1.37	∞	-0.01	1.00	1.49	∞	0.01
100	0.00	-51.16	3080	-0.01	1.12	1.44	∞	0.00	1.01	1.50	∞	0.00
1000	0.00	176.13	14251	0.00	1.01	1.49	∞	0.00	1.00	1.50	∞	0.00
50	0.53	0.48	1	0.51	0.34	0.64	1	0.53	0.53	0.54	1	0.01
100	0.51	0.53	1	0.54	0.37	0.61	1	0.52	0.51	0.51	1	0.00
1000	0.50	0.50	1	0.50	0.47	0.54	1	0.49	0.53	0.52	1	0.00
50	1.32	1347.71	9261	2.57	3.20	3.12	NA	1.15	1.86	2.76	NA	0.01
100	1.33	42156.28	418435	2.39	2.87	3.44	NA	1.12	1.78	2.84	NA	0.00
1000	1.26	124462.82	3903629	2.18	2.66	3.67	NA	1.11	1.80	2.85	NA	0.00

382 The convergence tolerance for IGMM was set to $tol = 1.22 \cdot 10^{-4}$. Table 5 summarizes the
383 simulation. Each sub-table is organized as follows: columns represent parameter estimates;
384 the three main rows are the average over $n = 1,000$ replications (top), the proportion of
385 estimates below the true value (middle), and the empirical standard deviation around the
386 empirical average times $\sqrt{N}$ – not around the truth (bottom).

387 The Gaussian MLE estimates σ_y directly, while IGMM and the Lambert W $\times$ Gaussian

MLE estimates δ and σ_x , which implicitly give $\hat{\sigma}_y$ through $\sigma_y(\delta, \sigma_x) = \sigma_x \cdot \frac{1}{\sqrt{(1-2\delta)^{3/2}}}$ if $\delta < 1/2$ (see (21)). For a fair comparison each sub-table also includes a column for $\hat{\sigma}_y = \hat{\sigma}_x \cdot \frac{1}{\sqrt{(1-2\hat{\delta})^{3/2}}}$. Some of these entries contain “ ∞ ”, even for $\delta < 1/2$; this occurs if at least one $\hat{\delta} \geq 1/2$.

For any $\delta < 1$, $\mu_x = \mu_y$, thus they can be directly compared. For $\delta \geq 1$, the mean does not exist; each sub-table for these δ interprets μ_y as the median.

Gaussian data: $\delta = 0$ This setting checks if imposing the Lambert W framework, even though its use is superfluous, causes a quality loss in the estimation of $\mu_y = \mu_x$ or $\sigma_y = \sigma_x$. Furthermore, critical values for $H_0 : \delta = 0$ (Gaussian tails) can be obtained. Table 2a shows that all estimators are unbiased and quickly tend to a large-sample variance. Additional estimation of δ does not affect the efficiency of $\hat{\mu}_x$ compared to estimating solely μ (both for IGMM and Lambert W $\times$ Gaussian MLE). Estimating σ_y directly by Gaussian MLE does not give better results than the Lambert W $\times$ Gaussian MLE: both are unbiased and have similar standard deviation.

No fourth moment: $\delta = 1/3$ Here $\sigma_y(\delta, \sigma_x = 1) = 2.28$, but fourth moments do not exist anymore. This results in an increasing empirical standard deviation of $\hat{\sigma}_y$ as N grows. In contrast, estimates for σ_x are not drifting off. In presence of these large heavy tails the median is much less variable than Gaussian MLE and IGMM. Yet, Lambert W $\times$ Gaussian MLE for μ_x even outperforms the median.

Non-existing mean: $\delta = 1$ Here the mean is non-finite. Thus both sample moments diverge, and their standard errors are also growing quickly. The median still provides a very good estimate for the location, but is again inferior to both Lambert W estimators, which are unbiased and seem to converge to an asymptotic variance at rate $\sqrt{N}$.

Extreme heavy tails: $\delta = 1.5$ As in Section 5.1, IGMM and Lambert W MLE continue to be unbiased even though the data is extremely heavy-tailed. Moreover, Lambert W MLE also has the smallest empirical standard deviation overall. In particular, the Lambert W MLE for μ_x has an approximately 20% lower standard deviation than the median.

415 The last column shows that for some N about 1% of the $n = 1,000$ simulations generated
 416 invalid likelihood values (NA and ∞). Here the search for the optimal δ lead into regions
 417 with a numerical overflow in the evaluation of $W_\delta(z)$. For a comparable summary, these
 418 few cases were omitted and new simulations added until a full $n = 1,000$ finite estimates
 419 were found. Since this only happened in 1% of the cases and also such heavy-tailed data
 420 is rarely encountered in practice, this numerical issue is not a real limitation in statistical
 421 practice.

422 5.3 Discussion of the Simulations

423 This simulation study confirms well-known facts about the sample average, standard devi-
 424 ation, and median and compares them to finite sample properties of the two Lambert W
 425 estimators. The median is known to be robust, which shows here as its quality does not
 426 depend on the thickness of the tails.

427 IGMM is unbiased for τ independent of the magnitude of δ . As expected the Lambert W
 428 MLE for θ has the best properties: it is unbiased for all δ , and for $\delta = 0$ it performs as well
 429 as the classic sample mean and standard deviation. For small δ it has the same empirical
 430 standard deviation as the Gaussian MLE, but a lower one than the median for large δ .

431 Hence the only advantage of estimating μ_y and σ_y by sample moments of $\mathbf{y}$ is speed;
 432 otherwise the Lambert W $\times$ Gaussian MLE is at least as good as the Gaussian MLE and
 433 clearly outperforms it for heavy-tailed data.

434 6 Applications

435 Tukey's h distribution has already proven useful to model heavy-tailed data, but parametric
 436 inference was limited to quantile fitting or methods of moments estimation (Field, 2004;
 437 Fischer, 2010; Headrick et al., 2008). Theorem 2.8 allows us to estimate θ by ML.

438 This section shows the usefulness of the presented methodology on simulated as well as
 439 real world data: i) Section 6.1 demonstrates Gaussianizing on the Cauchy sample from the
 440 Introduction; ii) Section 6.2 shows that heavy tail Lambert W $\times$ Gaussian distributions
 441 provide an excellent fit to daily S&P 500 log-return series; and iii) Section 6.3 shows how
 442 removing heavy tails reveals hidden patterns in power-law type data.

6.1 Estimating Location of a Cauchy With The Sample Mean

It is well-known that the sample mean $\bar{\mathbf{y}}$ is a poor estimate of the location parameter of a Cauchy distribution, since the sampling distribution of $\bar{\mathbf{y}}$ is again a Cauchy; in particular, its variance does not go to 0 for $n \rightarrow \infty$.

Heavy-tailed Lambert $W \times$ Gaussian distributions have similar properties to a Cauchy for $\delta \approx 1$. The mean of X equals the location of Y , due to symmetry around μ_x (for all $\delta \geq 0$) and c , respectively. Thus we can estimate τ from the Cauchy sample $\mathbf{y}$, transform $\mathbf{y}$ to $\mathbf{x}_{\hat{\tau}}$, estimate μ_x from $\mathbf{x}_{\hat{\tau}} = W_{\hat{\tau}}(\mathbf{y})$, and thus obtain an estimate of c .

The data $\mathbf{y} \sim \mathcal{C}(0, 1)$ in Fig. 2a has heavy tails with two extreme (positive) samples. A Cauchy ML fit gives $\hat{c} = 0.03(0.055)$ and $\hat{s} = 0.86(0.053)$ (standard errors in parenthesis). A Lambert $W \times$ Gaussian MLE gives $\hat{\mu}_x = 0.03(0.055)$, $\hat{\sigma}_x = 1.05(0.072)$, and $\hat{\delta} = 0.86(0.082)$. Thus both fits correctly fail to reject $\mu_x = c = 0$. Table 3a shows summary statistics on both samples. Since the Cauchy distribution does not have a well-defined mean, $\bar{\mathbf{y}} = 2.304(2.101)$ is not meaningful. However, $\mathbf{x}_{\hat{\tau}_{MLE}}$ is approximately Gaussian and we use the sample average to do inference: $\bar{\mathbf{x}}_{\hat{\tau}_{MLE}} = 0.033(0.0472)$ correctly fails to reject a zero location for $\mathbf{y}$. The transformed $\mathbf{x}_{\hat{\tau}_{MLE}}$ features additional Gaussian characteristics (symmetric, no excess kurtosis), and even the null hypothesis of Normality cannot be rejected (p-value ≥ 0.5).

Figure 2d shows the running sample average for the original sample and its Gaussianized version. For a fair comparison $\hat{\tau}_{MLE}^{(n)}$ was re-estimated cumulatively for each $n = 5, \dots, 500$, and then used to compute $(x_1, \dots, x_n)$. Even for small n the transformation works extremely well: the highly influential point around $n \approx 50$ greatly affects $\bar{\mathbf{y}}$, but has no relevant effect on $\bar{\mathbf{x}}_{\hat{\tau}_{MLE}}^{(n)}$. Overall, the sample average of the Gaussianized data has the usual good properties. And even for very small n it is already clear that the location of the underlying Cauchy distribution is approximately zero.

Although a toy example, it shows that removing (strong) heavy tails from data works and provides new, “nice” data which can then be used for more refined methods.

Table 3: Summary statistics for observed (heavy-tailed) $\mathbf{y}$ and back-transformed (Gaussianized) data $\mathbf{x}_{\hat{\tau}_{MLE}}$. ** stands for $< 10^{-16}$; * for $< 2.2 \cdot 10^{-16}$.

	(a) $\mathbf{y} \sim \mathcal{C}(0, 1)$ (Section 6.1)			(b) $\mathbf{y} = \text{S\&P 500}$ (Section 6.2)		(c) $\mathbf{y} = \text{solar flares}$ (Section 6.3)	
	$\mathbf{y}$	$\mathbf{x}_{\hat{\tau}}$	$\mathbf{x}_{\hat{\lambda}}$	$\mathbf{y}$	$\mathbf{x}_{\hat{\tau}}$	$\mathbf{y}$	$\mathbf{x}_{\hat{\tau}}$
Min	-161.59	-3.16	0	-7.11	-2.42	20	20
Max	952.95	3.81	33.18	4.99	2.23	231300	157
Mean	2.30	0.03	14.98	0.05	0.05	689.4	89.0
Median	0.04	0.04	14.96	0.04	0.04	87	87
Stdev	46.980	1.06	1.20	0.95	0.71	6520.6	27.0
Skewness	17.43	0.12	3.90	-0.30	-0.04	22.2	0.1
Kurtosis	343.34	3.21	161.75	7.70	2.93	582.1	1.9
SW	*	0.71	**	*	0.24	**	**
AD	**	0.51	**	*	0.18	**	**

6.2 Heavy Tails in Finance: S&P 500 Case Study

A lot of financial data displays negative skewness and excess kurtosis. Since financial data is in general not i.i.d., it is often modeled with a (skew) student-t distribution underlying a (generalized) auto-regressive conditional heteroskedastic (GARCH) (Bollerslev, 1986; Engle, 1982) or a stochastic volatility (SV) model (Deo, Hurvich, and Lu, 2006; Melino and Turnbull, 1990). Using the Lambert W approach we can build upon the knowledge and implications of Gaussianity (and avoid deriving properties of a GARCH or SV model with heavy-tailed innovations), and simply “Gaussianize” the reutns before fitting more complex – GARCH or SV – models.

Remark 6.1. *Time series models with Lambert $W \times$ Gaussian white noise are far beyond the scope of this work, but can be a direction of future research. Here I only consider the unconditional distribution.*

Figure 7a shows the S&P 500 log-returns with a total of $N = 2,780$ daily observations.⁸ Table 3b confirms the heavy tails (sample kurtosis 7.70), but also indicates negative skewness (-0.296). As the sample skewness $\hat{\gamma}_1(\mathbf{y})$ is very sensitive to outliers, we should test for symmetry by fitting a skewed distribution and testing its skewness parameter(s) for zero. In case of the double-tail Lambert $W \times$ Gaussian this means to test $H_0 : \delta_\ell = \delta_r = \delta$ versus $H_1 : \delta_\ell \neq \delta_r$. Since the likelihood can now be computed by (29), we can use a likelihood

⁸R package MASS, dataset SP500.




Figure 7: Lambert W Gaussianization of S&P 500 log-returns: $\hat{\tau} = (0.05, 0.70, 0.17)$. In (a) and (b): data (top left); autocorrelation function (ACF) (top right); histogram, Gaussian fit, and KDE (bottom left); Normal QQ plot (bottom right).

ratio test with one degree of freedom (3 versus 4 parameters). The log-likelihood of the double-tail Lambert W $\times$ Gaussian fit (Table 4a) equals $-3606.0 = -2972.27 + (-633.73)$ (input + penalty), while the one δ fit gives $-3606.56 = -2971.47 + (-635.09)$. Here the double tails pay a lower penalty for transforming the data, but in turn give less Gaussian transformed sample. Comparing twice their difference to a χ^2_1 distribution gives a p-value of 0.29. For comparison, a skew-t fit (Azzalini and Capitanio, 2003), with location c , scale s , shape α , and ν degrees of freedom also yields⁹ a non-significant $\hat{\alpha}$ (Table 4b). Thus both fits cannot reject symmetry.

Assume we have to make a decision if we should trade a certificate replicating the S&P 500. Since we can either buy or sell, it is not important if the average return is positive or negative, as long as it is significantly different from zero.

6.2.1 Gaussian Fit to Returns

If we ignore heavy tails and estimate (μ_y, σ_y) by Gaussian MLE, $\hat{\mu}_y = 0$ can not be rejected on a $\alpha = 1\%$ level (Table 4e). However, a plain sample average over-estimates the variance in presence of heavy tails, and thus adds bias to the test statistic.

⁹Function `st.mle` in the R package `sn`.

Table 4: MLE fits to S&P 500 $\mathbf{y}$ (a, b, c, d, e) and the Gaussianized data $\mathbf{x}_{\hat{\tau}_{MLE}}$ (f).

(a) double-tail Lambert $W \times$ Gaussian = Tukey's hh (S&P 500)					(b) skew t (S&P 500)				
	Est.	se	t	$\Pr(> t)$		Est.	se	t	$\Pr(> t)$
μ_x	0.06	0.015	3.66	0.00	c	0.10	0.061	1.65	0.10
σ_x	0.71	0.016	44.00	0.00	s	0.67	0.017	38.47	0.00
δ_ℓ	0.19	0.021	8.99	0.00	α	-0.08	0.101	-0.77	0.44
δ_r	0.16	0.019	8.24	0.00	ν	3.73	0.297	12.57	0.00
(c) Lambert $W \times$ Gaussian = Tukey's h (S&P 500)					(d) student- t (S&P 500)				
	Est.	se	t	$\Pr(> t)$		Est.	se	t	$\Pr(> t)$
μ_x	0.06	0.015	3.65	0.000	c	0.06	0.015	3.65	0.00
σ_x	0.71	0.016	43.95	0.000	s	0.67	0.017	39.51	0.00
δ	0.17	0.016	11.05	0.000	ν	3.72	0.295	12.61	0.00
(e) Gaussian (S&P 500)					(f) Gaussian ($\mathbf{x}_{\hat{\tau}_{MLE}}$)				
	Est.	se	t	$\Pr(> t)$		Est.	se	t	$\Pr(> t)$
μ_y	0.05	0.018	2.55	0.01	$\mu_{x_{\hat{\tau}}}$	0.05	0.013	3.81	0.00
σ_y	0.95	0.013	74.57	0.00	$\sigma_{x_{\hat{\tau}}}$	0.71	0.009	74.57	0.00

6.2.2 Heavy Tail Fit to Returns

Both a heavy tail Lambert $W \times$ Gaussian (Table 4c) and student- t fit (Table 4d) reject the zero mean null (p-values, 10^{-4} and $3 \cdot 10^{-5}$, respectively). The standard errors for the location parameter are essentially the same.

While location and scale estimates are almost identical, the tail estimates lead to very different conclusions: while for $\hat{\nu} = 3.71$ only moments up to order 3 exist, in the Lambert $W \times$ Gaussian case moments up to order 5 exist ($1/0.172 = 5.81$). This is especially noteworthy as many theoretical results in the (financial) time series literature rely on finite fourth moments (Mantegna and Stanley, 1998; Zdrozny, 2005); consequently many empirical studies test if financial data actually satisfy this assumption (Cont, 2001; Huisman, Koedijk, Kool, and Palm, 2001). For this particular dataset student's t and a Lambert $W \times$ Gaussian fit give different answers to the same question. Since previous empirical studies often use student's t as a baseline (Wong et al., 2009), it might be worthwhile to re-examine their findings in light of heavy tail Lambert $W \times$ Gaussian distributions.

521 6.2.3 “Gaussianizing” Returns

A typical parameter inference study would conclude here. Using Lambert’s W function we can analyze the back-transformed $\mathbf{x}_{\hat{\tau}_{MLE}}$ to test if a Lambert W $\times$ Gaussian distribution is indeed appropriate. Figure 7b shows that $\mathbf{x}_{\hat{\tau}_{MLE}}$ is indistinguishable from a Gaussian sample. Not even one Normality test can reject Gaussianity: p-values are 0.18, 0.18, 0.31, and 0.24, respectively (Anderson Darling, Cramer-von-Mises, Shapiro-Francia, Shapiro-Wilk; see Thode (2002)). Table 3b also shows that Lambert W “Gaussianization” was successful: $\hat{\gamma}_2(\mathbf{x}_{\hat{\tau}}) = 2.93$ and $\hat{\gamma}_2(\mathbf{x}_{\hat{\tau}}) = -0.039$ are within the typical variation for a Gaussian sample. Thus

$$Y = \left(U e^{\frac{0.172}{2} U^2} \right) 0.705 + 0.055, \quad U = \frac{X - 0.055}{0.705}, \quad U \sim \mathcal{N}(0, 1) \quad (37)$$

522 is an adequate (unconditional) Lambert W $\times$ Gaussian model for the S&P 500 log-returns
 523 $\mathbf{y}$. For trading, this means that the expected return is significantly larger than zero ($\hat{\mu}_x =$
 524 $0.055 > 0$), and thus replicating certificates should be bought.

525 6.2.4 Gaussian MLE for Gaussianized Data

526 For $\delta_l = \delta_r \equiv \delta < 1$, also $\mu_x \equiv \mu_y$. We can therefore replace testing $\mu_y = 0$ versus $\mu_y \neq 0$
 527 for a non-Gaussian $\mathbf{y}$, with the very well understood hypothesis test $\mu_x = 0$ versus $\mu_x \neq 0$
 528 for the Gaussian $\mathbf{x}_{\hat{\tau}_{MLE}}$. In particular, standard errors based on $\frac{\hat{\sigma}}{\sqrt{N}}$ - and thus t and p-
 529 values - should be closer to the “truth” (Table 4c and 4d) than a Gaussian MLE on the
 530 non-Gaussian $\mathbf{y}$ (Table 4e). Table 4f shows that standard errors for $\hat{\mu}_{\mathbf{x}}$ are even a bit too
 531 small compared to the heavy-tailed versions. Since the “Gaussianizing” transformation was
 532 estimated, treating $\mathbf{x}_{\hat{\tau}_{MLE}}$ as if it was original data is too optimistic regarding its Gaussian-
 533 ity (recall the penalty (31) in the total likelihood (29)).

534

535 This example confirms that if a model and its theoretical properties are based on Gaus-
 536 sianity, but the observed data is heavy-tailed, then Gaussianizing the data first gives more
 537 reliable inference than applying the Gaussian methods to the original, heavy-tailed data
 538 (Fig. 1). Clearly, a joint estimation of the model parameters based on Lambert W $\times$ Gaus-

sian errors (or any other heavy-tailed distribution) would be optimal. However, theoretical properties and estimation techniques may not have been developed and implemented yet, or are simply not known to researchers who are non-experts in heavy-tailed statistics. The Lambert Way to Gaussianize data thus is a pragmatic method to improve statistical inference on heavy-tailed data, while preserving the ease of usage and interpretation of Gaussian models.

6.3 Removing Power Law From Solar Flare Counts

The previous section focused on Lambert $W \times F_X$ distributions as a “true” model for the data $\mathbf{y}$. Here I consider it merely as a data transformation to remove heavy tails. In the same way as scaling $\mathbf{y}$ to zero-mean, unit-variance data, $(\mathbf{y} - \bar{\mathbf{y}})/\hat{\sigma}_y$, does not necessarily mean we believe the underlying process is Gaussian, we can also convert $\mathbf{y}$ to $\mathbf{x}_\tau = W_\tau(\mathbf{y})$ without assuming that $\mathbf{y}$ is actually Lambert $W \times$ Gaussian. While $\mathbf{x}_\tau$ might lose the interpretability of the observed data (e.g. units become distorted), it can be helpful for exploratory data analysis (EDA), as the eye is a bad judgment to detect regularities corrupted by heavy tails. Removing them can reveal hidden patterns and thus greatly improve the accuracy of statistical inference for $\mathbf{y}$.

Here I study solar flare gamma-ray count rates (Clauset, Shalizi, and Newman, 2009; Newman, 2005). The data¹⁰ were collected approximately four times a day from Feb. 1980 until Nov. 1989 giving $T = 12,773$ observations. See Dennis, Orwig, Kennard, Labow, Schwartz, Shaver, and Tolbert (1991) for details and scientific background.

The gamma-ray count rates exhibit a strong right heavy tail (Fig. 8a), which makes more detailed visual inspection as well as simple EDA difficult. A zoom to $y_i \leq 400$ in Fig. 8d shows that a lot of counts lie between 50 and 100 and this level drops off at the end of the observation cycle. This drop is not an intrinsic characteristic of solar flares but due to a decreasing sensitivity of the X-ray detectors over time (Dennis et al., 1991). For the sake of comparison with Clauset et al. (2009); Newman (2005) most estimates are based on all

¹⁰Dataset `SolarFlares` in the `LambertW` package.




Figure 8: Peak X-ray count rates of solar flares.

567 $T = 12,773$ observations. Figures 8d, 8e, and 8f also show separate density estimates for
 568 the first 4,000 and last 2,273 observations, and while the estimates change, the qualitative
 569 findings do not.

570

571 Clauset et al. (2009) find that a power-law ($\hat{a} = 1.79(0.02)$) with cut-off ($\hat{y}_{\min} = 323(89)$)
 572 gives the best fit amongst various alternatives. However, this first EDA might not be
 573 complete: not only visually heavy tails can obscure underlying non-trivial structure, but
 574 also estimates - such as the power law fit or non-parametric density estimates (Fig. 8d and
 575 8b) - are affected by the heavy right tail. Here I show that Gaussianizing this data reveals
 576 new insights for the data-generating process, with a new interpretation for the optimality
 577 of the cut-off.

578 A Lambert $W \times$ Gaussian MLE fit $\hat{\theta} = (\hat{\mu}_x, \hat{\sigma}_x, \hat{\delta}_\ell, \hat{\delta}_r) = (86.97, 26.80, 0, 2.37)$ confirms

that only the right tail (tail index $1/2.373 = 0.421$) needs a Gaussianizing transformation.¹¹

The last column of Fig. 8 shows EDA for the Gaussianized data. Removing the heavy right tail reveals a bimodal structure, which gives additional meaning to $\hat{y}_{\min} = 323$. The Gaussianized cut-off value equals $W_{\hat{\tau}}(323) = 121.16$ with the transformed standard deviation interval $[117.74, 123.37]$ (corresponding to 323 ± 89). Fitting a two component Gaussian mixture model to $\mathbf{x}_{\hat{\tau}}$ yields $\hat{\lambda}\mathcal{N}_1(67.10, 14.04^2) + (1 - \hat{\lambda})\mathcal{N}_2(113.12, 14.27^2)$ with $\hat{\lambda} = 0.52$ and optimal decision boundary between classes of 90.48. The mean of the larger component, 113.12, lies within one standard deviation of the optimal Gaussianized cut-off 121.16: for lower cut-offs the left-tail of the larger component – or for much lower cut-offs even the smaller component – would counteract the power-law decay of the upper gamma-ray count rates.

As mentioned above, this analysis is not intended to describe the underlying process of solar flare gamma rays; it should rather show new insights that can be gained by Gaussianizing. Future research based on these new findings might lead to new physical interpretations of the statistical properties gamma-ray count rates, see for example [Aschwanden \(2011\)](#).

7 Discussion and Outlook

I adapt the skewed Lambert W input / output framework to introduce heavy tails in continuous RVs $X \sim F_X(x)$. For Gaussian input this not only contributes to existing work on Tukey’s h distribution, but also gives convincing empirical results: unimodal data with heavy tails can be transformed to Gaussian data/RVs. Properties of a Gaussian model $\mathcal{M}_{\mathcal{N}}$ on the back-transformed data mimic the features of the “true” skewed, heavy-tailed model $\mathcal{M}_G$ very closely.

Since Gaussianity is the single most typical, and often required, assumption in many areas of statistics, machine learning, and signal processing, future research can take many directions. From a theoretical perspective properties of Lambert W $\times F_X$ distributions viewed as a generalization of already well-known distributions F_X can be studied. This area

¹¹For comparison Fig. 8e also shows the back-transformed data $\mathbf{x}_{\hat{\tau}_1}$ using the same δ on each tail ($\hat{\tau}_1 = (74.46, 26.32, 1.53)$). However, due to the clear right heavy tail I will continue with the (δ_l, δ_r) transformation.

606 will profit from existing literature on the Lambert W function, which has been discovered
607 only recently by the statistics community. Empirical work can focus on transforming the
608 data and compare performances of approximate Gaussian versus joint heavy-tail analysis.
609 The comparisons in this work showed that approximate inference for Gaussianized data is
610 comparable with the direct heavy tail modeling, and so provides an easy tool to improve
611 inference for heavy-tailed data in statistical practice.

612 I also provide the R package [LambertW](#), publicly available at [CRAN](#), to facilitate the use
613 of Lambert $W \times F_X$ distributions in practice.

614 Acknowledgments

615 I want to thank Andrew F. Siegel who brought Tukey's h distribution to my attention, and
616 Brian R. Dennis who gave detailed background information and suggestions on the solar
617 flares dataset.

References

- Achim, A., P. Tsakalides, and A. Bezerianos (2003). SAR image denoising via Bayesian wavelet shrinkage based on heavy-tailed modeling. *Geoscience and Remote Sensing, IEEE Transactions on* 41(8), 1773 – 1784.
- Aschwanden, M. J. (2011). The State of Self-Organized Criticality of the Sun During the Last Three Solar Cycles. II. Theoretical Model. *Solar Physics* 274, 119–129.
- Aysal, T. C. and K. E. Barner (2006). Second-order heavy-tailed distributions and tail analysis. *IEEE Transactions on Signal Processing* 54(7), 2827–2832.
- Azzalini, A. and A. Capitanio (2003). Distributions generated by perturbation of symmetry with emphasis on a multivariate skew t distribution. *Journal of the Royal Statistical Society ser B* 65, 367–389.
- Baek, C. and V. Pipiras (2010). Estimation of parameters in heavy-tailed distribution when its second order tail parameter is known. *Journal of Statistical Planning and Inference* 140(7), 1957 – 1967.
- Blaylock, J. R., L. E. Salathe, and R. D. Green (1980). A note on the Box-Cox transformation under heteroskedasticity. *Western Journal of Agricultural Economics* 05(02), 129–136.
- Bollerslev, T. (1986). Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics* 31, 307 – 327.
- Brockwell, P. J. and R. A. Davis (1998). *Time Series: Theory and Methods*. Springer Series in Statistics.
- Clauset, A., C. R. Shalizi, and M. E. J. Newman (2009). Power-law distributions in empirical data. *SIAM Review* 51, 661–703.
- Cont, R. (2001). Empirical properties of asset returns: stylized facts and statistical issues. *Quantitative Finance* 1, 223–236.

- 643 Corless, R. M., G. H. Gonnet, D. E. G. Hare, and D. J. Jeffrey (1996). On the Lambert W
644 function. *Advances in Computational Mathematics* 5, 329–359.
- 645 Dennis, B. R., L. E. Orwig, G. S. Kennard, G. J. Labow, R. A. Schwartz, A. R. Shaver, and
646 A. K. Tolbert (1991). The complete Hard X Ray Burst Spectrometer event list, 1980-
647 1989. Available <http://adsabs.harvard.edu/abs/1991chxb.book.....D> and <http://umbra.nascom.nasa.gov/smm/hxrbs.html>.
648
- 649 Deo, R., C. Hurvich, and Y. Lu (2006). Forecasting Realized Volatility Using a Long
650 Memory Stochastic Volatility Model: Estimation, Prediction and Seasonal Adjustment.
651 *Journal of Econometrics* 127, 29 – 58.
- 652 Dutta, K. K. and D. Babbel (2002). On Measuring Skewness and Kurtosis in Short Rate
653 Distributions: The Case of the US Dollar London Inter Bank Offer Rates. Technical
654 report, Wharton School Center for Financial Institutions, University of Pennsylvania.
- 655 Engle, R. (1982). Autoregressive conditional heteroskedasticity with estimates of the vari-
656 ance of U.K. inflation. *Econometrica* 50, 987 – 1008.
- 657 Fernandez, C. and M. F. J. Steel (1999). Multivariate Student-t Regression Models: Pitfalls
658 and Inference. *Biometrika* 86, 153–167.
- 659 Field, C. and M. G. Genton (2006). The Multivariate g-and-h Distribution. *Technomet-*
660 *rics* 48(1), 104–111.
- 661 Field, C. A. (2004). Using the gh distribution to model extreme wind speeds. *Journal of*
662 *Statistical Planning and Inference* 122(1-2), 15 – 22.
- 663 Fischer, M. (2010). Generalized Tukey-type distributions with application to financial and
664 teletraffic data. *Statistical Papers* 51, 41–56. 10.1007/s00362-007-0114-z.
- 665 Galassi, M., J. Davies, J. Theiler, B. Gough, G. Jungman, P. Alken, M. Booth, and F. Rossi
666 (2011). *GNU Scientific Library Reference Manual* (3rd ed.). ISBN 0954612078; www.gnu.org/software/gsl.
667
- 668 Gidlund, M. and N. Debernardi (2009). Scheduling performance of heavy-tailed data traffic
669 in wireless high-speed shared channels. In *Proceedings of the 2009 IEEE conference on*

670 *Wireless Communications & Networking Conference, WCNC'09*, Piscataway, NJ, USA,
671 pp. 1818–1823. IEEE Press.

672 Goerg, G. M. (2011). Lambert W Random Variables - A New Family of Generalized
673 Skewed Distributions with Applications to Risk Estimation. *The Annals of Applied Statis-*
674 *tics* 5(3), p. 2197 – 2230. arxiv.org/abs/0912.4554.

675 Goncalves, S. and N. Meddahi (2011). Box-Cox transforms for realized volatility. *Journal*
676 *of Econometrics* 160(1), 129–144.

677 Granger, C. W. J. and R. Joyeux (2001). An introduction to long-memory time series
678 models and fractional differencing. *Journal of Time Series Analysis* 1, 15 – 30.

679 Headrick, T. C., R. K. Kowalchuk, and Y. Sheng (2008). Parametric Probability Densities
680 and Distribution Functions for Tukey g-and-h Transformations and their Use for Fitting
681 Data. *Applied Mathematical Sciences* 2(9), 449 – 462.

682 Hoaglin, D. C. (2006). *Summarizing Shape Numerically: The g-and-h Distributions*, pp.
683 461–513. Hoboken, NJ, USA: John Wiley and Sons, Inc.

684 Huisman, R., K. G. Koedijk, C. J. M. Kool, and F. Palm (2001). Tail-index estimates in
685 small samples. *Journal of Business & Economic Statistics* 19(2), 208–16.

686 Hwang, J., S. Lay, and A. Lippman (1994). Nonparametric multivariate density estimation:
687 A comparative study. *IEEE Trans. Signal Processing* 42, 2795–2810.

688 Ilow, J. (2000). Forecasting network traffic using farima models with heavy tailed inno-
689 vations. *Acoustics, Speech, and Signal Processing, IEEE International Conference on* 6,
690 3814–3817.

691 Jodrá, P. (2009). A closed-form expression for the quantile function of the Gompertz-
692 Makeham distribution. *Math. Comput. Simul.* 79, 3069–3075.

693 Kim, T.-H. and H. White (2003). On More Robust Estimation of Skewness and Kurtosis:
694 Simulation and Application to the S&P500 Index.

- 695 Lawrance, A. J. (1987). A note on the variance of the Box-Cox regression transformation
696 estimate. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 36(2),
697 221–223.
- 698 Lehmann, E. L. and G. Casella (1998). *Theory of Point Estimation* (2 ed.). Springer Texts
699 in Statistics.
- 700 Liu, C. and D. B. Rubin (1995). ML Estimation of the t distribution using EM and its
701 extensions, ECM and ECME. *Statistica Sinica* 5, 19–39.
- 702 Liu, H., J. Lafferty, and L. Wasserman (2009). The Nonparanormal: Semiparametric Es-
703 timation of High Dimensional Undirected Graphs. *Journal of Machine Learning Re-*
704 *search* 10, 2295–2328.
- 705 Maiboroda, R. and N. Markovich (2004). Estimation of heavy-tailed probability den-
706 sity function with application to web data. *Computational Statistics* 19, 569–592.
707 10.1007/BF02753913.
- 708 Mantegna, R. N. and H. E. Stanley (1998). Modeling of financial data: Comparison of
709 the truncated Lévy flight and the ARCH(1) and GARCH(1,1) processes. *Physica A:*
710 *Statistical and Theoretical Physics* 254(1-2), 77 – 84.
- 711 Markovich, N. M. (2005). Accuracy of transformed kernel density estimates for a heavy-
712 tailed distribution. *Autom. Remote Control* 66, 217–232.
- 713 Melino, A. and S. M. Turnbull (1990). Pricing foreign currency options with stochastic
714 volatility. *Journal of Econometrics* 45(1-2), 239 – 265.
- 715 Morgenthaler, S. and J. W. Tukey (2000). Fitting quantiles: Doubling, hr, hq, and hhh
716 distributions. *Journal of Computational and Graphical Statistics* 9(1), pp. 180–195.
- 717 Newman, M. E. J. (2005). Power laws, Pareto distributions and Zipf’s law. *Contemporary*
718 *Physics* 46, 323–351.
- 719 Nowicka-Zagrajek, J. and R. Weron (2002). Modeling electricity loads in California: ARMA
720 models with hyperbolic noise. *Signal Process.* 82, 1903–1915.

- 721 Pakes, A. G. (2011). Lambert’s W, infinite divisibility and Poisson mixtures. *Journal of*
722 *Mathematical Analysis and Applications* 378, 480492.
- 723 Palma, W. and M. Zavallos (2011). Fitting non-gaussian persistent data. *Applied Stochastic*
724 *Models in Business and Industry* 27(1), 23–36.
- 725 R Development Core Team (2010). *R: A Language and Environment for Statistical Com-*
726 *puting*. Vienna, Austria: R Foundation for Statistical Computing. ISBN 3-900051-07-0.
- 727 Rathie, R. N. and P. Silva (2011). Applications of Lambert W Function. *International*
728 *Journal of Applied Mathematics & Statistics* 23(D11), –.
- 729 Rayner, G. D. and H. L. MacGillivray (2002). Numerical maximum likelihood estimation for
730 the g-and-k and generalized g-and-h distributions. *Statistics and Computing* 12, 57–75.
- 731 Rosenlicht, M. (1969). On the explicit solvability of certain transcendental equations. *Pub.*
732 *Math. Institut des Hautes Etudes Scientifiques* 36, 15 – 22.
- 733 Sakia, R. M. (1992). The Box-Cox transformation technique: A review. *Journal of the*
734 *Royal Statistical Society. Series D (The Statistician)* 41(2), 169–178.
- 735 Smith, V. K. (1973). Least Squares Regression with Cauchy Errors. *Oxford Bulletin of*
736 *Economics and Statistics* 35(3), 223–31.
- 737 Thode, Jr., H. C. (2002). *Testing for Normality*. CRC Press.
- 738 Tsiotas, G. (2007). On the use of the Box-Cox transformation on conditional variance
739 models. *Finance Research Letters* 4(1), 28–32.
- 740 Valluri, S. R., D. J. Jeffrey, and R. M. Corless (2000). Some Applications of the Lambert
741 W Function to Physics. *Canadian Journal of Physics* 78, 823 – 831.
- 742 Vázquez, A., J. G. Oliveira, Z. Dezső, K.-I. Goh, I. Kondor, and A.-L. Barabási (2006).
743 Modeling bursts and heavy tails in human dynamics. *Phys. Rev. E* 73(3), 036127.
- 744 Wasserman, L. (2007). *All of Nonparametric Statistics*. Springer Texts in Statistics.

- 745 Wong, C. S., W. S. Chan, and P. L. Kam (2009). A Student t-mixture autoregressive model
746 with applications to heavy-tailed financial data. *Biometrika* 96(3), 751–760.
- 747 Yan, J. (2005). Asymmetry, fat-tail, and autoregressive conditional density in finan-
748 cial return data with systems of frequency curves. [citeseerx.ist.psu.edu/viewdoc/
749 summary?doi=10.1.1.76.2741](https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.76.2741).
- 750 Zadrozny, P. A. (2005). *Necessary and Sufficient Restrictions for Existence of a Unique*
751 *Fourth Moment of a Univariate GARCH(p, q) Process*, Volume 20 of *Advances in Econo-*
752 *metrics*, Chapter –, pp. 365–379. Emerald Group Publishing Limited. [ideas.repec.
753 org/p/ces/ceswps/_1505.html](https://ideas.repec.org/p/ces/ceswps/_1505.html).

755 A Auxiliary Results and Properties

756 A.1 Inverse Transformation $W_\delta(z)$

757 The function $W_\delta(z)$ is the building block of Lambert $W \times F_X$ distributions. This section
758 lists useful properties of $W_\delta(z)$ as a function of z as well as a function of δ .

Properties A.1. For $\delta = 0$,

$$W_\delta(z_i) |_{\delta=0} = z_i, \quad W'(\delta z_i^2) |_{\delta=0} = z_i^2, \quad \text{and } W(\delta z_i^2) |_{\delta=0} = 0. \quad (38)$$

By definition $\frac{W_\delta(z)}{z} = e^{-\frac{\delta}{2}W_\delta(z)^2}$ and therefore

$$\log \frac{W_\delta(z)}{z} = -\frac{\delta}{2}W_\delta(z)^2 = -\frac{W(\delta z^2)}{2}. \quad (39)$$

Lemma A.2 (Derivative of $W_\delta(z)$ with respect to z). *It holds*

$$\frac{d}{dz}W_\delta(z) = -\frac{W_\delta(z)}{z(1 + \delta W_\delta(z)^2)} = e^{-\frac{1}{2}W(\delta z^2)} \frac{1}{1 + W(\delta z^2)} \quad (40)$$

Proof. One of the many interesting properties of the Lambert W function relates to its derivative which satisfies

$$W'(z) = \frac{W(z)}{z(1 + W(z))} = \frac{1}{e^{W(z)}(1 + W(z))}, \quad z \neq 0, -1/e. \quad (41)$$

759 Hence,

$$\frac{d}{dz} \frac{W(\delta z^2)}{\delta} = W'(\delta z^2) \cdot 2z = \frac{W(\delta z^2)}{\delta z^2(1 + W(\delta z^2))} \cdot 2z = \frac{2W(\delta z^2)}{\delta z(1 + W(\delta z^2))} \quad (42)$$

Therefore,

$$\frac{d}{dz} W_\delta(z) = \frac{1}{2} \left(\frac{1}{\delta} W(\delta z^2) \right)^{-1/2} \cdot \frac{d}{dz} \frac{W(\delta z^2)}{\delta} \quad (43)$$

$$= \frac{1}{2} \left(\frac{1}{\delta} W(\delta z^2) \right)^{-1/2} \cdot \frac{2W(\delta z^2)}{\delta z (1 + W(\delta z^2))} \quad (44)$$

$$= \frac{1}{\delta^{1/2}} (W(\delta z^2))^{-1/2} \cdot \frac{W(\delta z^2)}{z (1 + W(\delta z^2))} \quad (45)$$

As $W(\delta z^2) = \delta u^2$ the last line simplifies to

$$\frac{1}{\delta^{1/2}} \frac{1}{\delta^{1/2} u} \cdot \frac{\delta u^2}{z (1 + \delta u^2)} = \frac{u}{z (1 + \delta u^2)}. \quad (46)$$

Now use again $u = W_\delta(z)$. □

Lemma A.3 (Derivative of $W_\delta(z)^2$ with respect to δ). *For all $z \in \mathbb{R}$*

$$\frac{\partial}{\partial \delta} [W_\delta(z)]^2 = -\frac{1}{1 + W(\delta z^2)} W_\delta(z)^4 \leq 0. \quad (47)$$

Proof. By definition $[W_\delta(z)]^2 = \frac{W(\delta z^2)}{\delta}$. Thus

$$\frac{\partial}{\partial \delta} \frac{W(\delta z^2)}{\delta} = \frac{\delta \frac{\partial}{\partial \delta} W(\delta z^2) - W(\delta z^2) \cdot 1}{\delta^2} \quad (48)$$

$$= \frac{\delta W'(\delta z^2) z^2 - W(\delta z^2)}{\delta^2} \quad (49)$$

$$= \frac{\delta \frac{W(\delta z^2)}{\delta z^2 (1 + W(\delta z^2))} z^2 - W(\delta z^2)}{\delta^2} \quad (50)$$

$$= \frac{\frac{W(\delta z^2)}{1 + W(\delta z^2)} - W(\delta z^2)}{\delta^2} \quad (51)$$

$$= \frac{\frac{-W(\delta z^2)^2}{1 + W(\delta z^2)}}{\delta^2} \quad (52)$$

$$= -\frac{1}{1 + W(\delta z^2)} [W_\delta(z)]^4. \quad (53)$$

Since both terms are non-negative for all $z \in \mathbb{R}$, the result follows. □

That is $W_\delta(z)^2$ is a decreasing function in δ for every $z \in \mathbb{R}$, i.e. the more we remove heavy tails the more z gets shrinked (non-linearly) towards $0 = \lim_{\delta \rightarrow \infty} W_\delta(z)$. In particular,

$$[W_\delta(z)]^2 < z^2 \Leftrightarrow \frac{W_\delta(z)}{z} < 1 \text{ and } \frac{W_{\delta+\varepsilon}(z)}{z} < \frac{W_\delta(z)}{z} \text{ for } \delta \geq 0 \text{ and } \varepsilon > 0.$$

Lemma A.4 (Derivative of $W_\delta(z)$ with respect to δ). *It holds*

$$\frac{\partial}{\partial \delta} W_\delta(z) = -\frac{1}{2} \frac{1}{1 + W(\delta z^2)} W_\delta(z)^3 \quad (54)$$

Proof.

$$\frac{\partial}{\partial \delta} W_\delta(z) = \text{sgn}(z) \frac{\partial}{\partial \delta} \left(\frac{W(\delta z^2)}{\delta} \right)^{1/2} \quad (55)$$

$$= \text{sgn}(z) \frac{1}{2} \left(\frac{W(\delta z^2)}{\delta} \right)^{-1/2} \frac{\partial}{\partial \delta} \frac{W(\delta z^2)}{\delta} \quad (56)$$

$$= \frac{1}{2} \frac{1}{W_\delta(z)} \frac{\partial}{\partial \delta} [W_\delta(z)]^2 \quad (57)$$

$$= -\frac{1}{2} \frac{1}{1 + W(\delta z^2)} W_\delta(z)^3, \quad (58)$$

where the last line follows by Lemma A.3. □

A.2 Penalty $\log R(\delta \mid z_i)$ for Standard Gaussian Input

For $\mu_x = 0$ and $\sigma_x = 1$ the penalty equals ($y_i = z_i$)

$$R(\delta \mid z_i) = \frac{W_\delta(z_i)}{z_i [1 + \delta (W_\delta(z_i))^2]} = \frac{W_\delta(z_i)}{z_i [1 + W(\delta z_i^2)]} \quad (59)$$

and thus

$$\log R(\delta \mid z_i) = \log \frac{W_\delta(z_i)}{z_i} - \log [1 + W(\delta z_i^2)] \quad (60)$$

$$= -\frac{W(\delta z_i^2)}{2} - \log [1 + W(\delta z_i^2)] \quad (61)$$

Lemma A.5 (Derivative of $\log R(\delta \mid z)$ with respect to δ). *For all $\delta \geq 0$ and all $z \in \mathbb{R}$*

$$\frac{\partial \log R(\delta \mid z)}{\partial \delta} = -z^2 W'(\delta z^2) \left(\frac{1}{2} + \frac{1}{1 + W(\delta z^2)} \right) \leq 0. \quad (62)$$

Proof. We have

$$\frac{\partial \log R(\delta | z)}{\partial \delta} = \frac{1}{W_\delta(z)} \frac{\partial W_\delta(z)}{\partial \delta} - \frac{1}{1 + W(\delta z^2)} W'(\delta z^2) z^2 \quad (63)$$

$$\stackrel{\text{Lemma A.4}}{=} \frac{1}{W_\delta(z)} \left(-\frac{1}{2} \frac{1}{1 + W(\delta z^2)} W_\delta(z)^3 \right) - \frac{1}{1 + W(\delta z^2)} W'(\delta z^2) z^2 \quad (64)$$

$$= -\frac{1}{1 + W(\delta z^2)} \left(\frac{1}{2} W_\delta(z)^2 + W'(\delta z^2) z^2 \right) \quad (65)$$

Using $W'(\delta z^2) = \frac{W(\delta z^2)}{\delta z^2(1+W(\delta z^2))}$ and re-factorizing gives (62).

□

A.3 Gaussian log-Likelihood at $W_\delta(z)$

Lemma A.6 (Derivative of the Gaussian log-likelihood at $W_\delta(z)$). *For all $z \in \mathbb{R}$ and for $\delta \geq 0$*

$$\frac{\partial}{\partial \delta} \ell(\mu_x = 0, \sigma_x = 1 | W_\delta(z)) = \frac{1}{2} \frac{1}{1 + W(\delta z^2)} [W_\delta(z)]^4 \geq 0. \quad (66)$$

Proof. The log of the standard Gaussian pdf evaluated at $W_\delta(z)$ simplifies to

$$\log \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2} [W_\delta(z)]^2} = \log \frac{1}{\sqrt{2\pi}} - \frac{1}{2} [W_\delta(z)]^2. \quad (67)$$

The rest follows by Lemma A.3. □

Lemma A.6 shows that increasing δ always increases the input log-likelihood $\ell(\delta | \mathbf{u}_\delta = W_\delta(\mathbf{z}))$ - see also Fig. 6b. For $\delta \rightarrow \infty$ the Gaussianized $\mathbf{u}_\delta$ goes to $\mathbf{0}$, which clearly maximizes the Gaussian likelihood if $\mu = 0$.

B Proofs

B.1 Inverse transformation

Proof of Lemma 2.5. Without loss of generality assume that $\mu_x = 0$ and $\sigma_x = 1$. Squaring (2) and multiplying by δ yields

$$\delta Z^2 = \delta U^2 \exp(\delta U^2) \quad (68)$$

The inverse of (68) is by definition Lambert's $W(z)$ function (Rosenlicht, 1969)

$$W(z) \exp W(z) = z, \quad z \in \mathbb{C}. \quad (69)$$

779 $W(z)$ is bijective for $z \geq 0$. Since $\delta U^2 \geq 0$ for all $\delta \geq 0$, applying $W(\cdot)$ to (68), dividing by
780 δ , and taking the square root gives

$$U = \pm \sqrt{\frac{W(\delta Z^2)}{\delta}} \quad (70)$$

781 Since $\exp(\frac{\delta}{2}U^2) > 0$ for all $\delta \in \mathbb{R}$ and all U , it follows that $Z = U \exp(\delta/2U^2)$ and U must
782 have the same sign, which concludes the proof. $\square$

783 B.2 Cdf and pdf

Proof of Theorem 2.8. By definition,

$$G_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}\left(\left\{U \exp\left(\frac{\delta}{2}U^2\right)\right\} \sigma_x + \mu_x \leq y\right) \quad (71)$$

$$= \mathbb{P}\left(U \exp\left(\frac{\delta}{2}U^2\right) \leq z\right) = \mathbb{P}(U \leq W_\delta(z)) \quad (72)$$

$$= F_U(U \leq W_\delta(z)). \quad (73)$$

784 Taking the derivative with respect to y gives

$$\frac{d}{dy}G_Y(y \mid \beta, \delta) = f_X(W_\delta(z)\sigma_x + \mu_x \mid \beta) \cdot \sigma_x \frac{d}{dy}W_\delta\left(\frac{y - \mu_x}{\sigma_x}\right) \quad (74)$$

$$= f_U(W_\delta(z) \mid \beta) \cdot \sigma_x \frac{1}{\sigma_x} \frac{d}{dz}W_\delta\left(\frac{y - \mu_x}{\sigma_x}\right) \quad (75)$$

$$= f_U(W_\delta(z) \mid \beta) \cdot \frac{d}{dz}W_\delta(z). \quad (76)$$

785 Using Lemma A.2 yields (15). $\square$

786 B.3 MLE for δ

Lemma B.1 (Derivative of the Lambert $W \times$ Gaussian log-likelihood). *We have*

$$D(\delta \mid \mathbf{z}) := \frac{\partial}{\partial \delta} \ell(\delta \mid \mathbf{z}) = \sum_{i=1}^N z_i^2 W'(\delta z_i^2) \left(\frac{1}{2} W_\delta(z_i)^2 - \left(\frac{1}{2} + \frac{1}{1 + W(\delta z_i^2)} \right) \right) \quad (77)$$

$$= \frac{1}{2} \sum_{i=1}^N \frac{W_\delta(z_i)^4}{1 + \delta W_\delta(z_i)^2} - \sum_{i=1}^N \frac{W_\delta(z_i)^2}{1 + \delta W_\delta(z_i)^2} \left(\frac{1}{2} + \frac{1}{1 + \delta W_\delta(z_i)^2} \right) \quad (78)$$

$$= \frac{1}{2} \sum_{i=1}^N \frac{W_\delta(z_i)^4}{1 + W(\delta z_i^2)} - \sum_{i=1}^N \frac{W_\delta(z_i)^2}{1 + W(\delta z_i^2)} \left(\frac{1}{2} + \frac{1}{1 + W(\delta z_i^2)} \right). \quad (79)$$

787 *Proof.* Apply Lemmas A.5 and A.6 to $\frac{\partial}{\partial \delta} \ell(\delta \mid \mathbf{z}) = \frac{\partial}{\partial \delta} \log R(\delta \mid \mathbf{z}) + \frac{\partial}{\partial \delta} \ell(\mu_x = 0, \sigma_x = 1 \mid$
788 $W_\delta(z))$. □

789 *Proof sketch of Theorem 4.1.* a) If condition (35) holds, then $D(\delta \mid \mathbf{z}) < 0$ at $\delta = 0$ and
790 stays negative for all $\delta > 0$. Hence the maximizer occurs at the boundary $\delta = 0$.

791 b) If (35) does not hold, then $D(\delta = 0 \mid \mathbf{z}) > 0$, decreases in δ and crosses the zero line
792 (one candidate for $\hat{\delta}_{MLE}$ occurs here).

793 c) As δ gets larger, $D(\delta \mid \mathbf{z})$ reaches a minimum (negative value) and starts increasing.
794 However, for $\delta \rightarrow \infty$ the derivative approaches zero from below and never equals zero
795 again; thus $\hat{\delta}_{MLE}$ is unique. □

796

Proof of Theorem 4.1. a) The log-likelihood is increasing at $\delta = 0$ if and only if (set $\delta = 0$
in (79) and use Property A.1)

$$\sum_{i=1}^N z_i^4 > 3 \sum_{i=1}^N z_i^2. \quad (80)$$

797 Eq. (80) means that transforming the data (choosing $\hat{\delta} > 0$) increases the overall like-
798 lihood only if the data is heavy-tailed enough. Note that the sum of squares is not
799 squared again. Hence condition (80) is not equivalent for the data having empirical
800 kurtosis larger than 3.

801 b) If (80) does not hold, then $\hat{\delta}_{MLE}$ must satisfy $D(\delta \mid \mathbf{z})|_{\delta=\hat{\delta}_{MLE}} = 0$ from (77) in Lemma
 802 **B.1**. It remains to be shown that this equation has (at least) one positive solution.

803 i) Since $\lim_{\delta \rightarrow \infty} W_\delta(z) = 0$ for all $z \in \mathbb{R}$, (79) is also true in the limit; however, we
 804 can ignore this solution as we require $\hat{\delta}_{MLE} \in \mathbb{R}$.

ii) By continuity and $\lim_{\delta \rightarrow \infty} W_\delta(z) = 0$, for sufficiently large δ_M , $W_{\delta_M}(z_i) < 1$ for all
 $z_i \in \mathbb{R}$. Hence $W_{\delta_M}(z_i)^4 < W_{\delta_M}(z_i)^2$ and therefore

$$\frac{1}{2} \sum_{i=1}^N \frac{W_\delta(z_i)^4}{1 + \delta W_\delta(z_i)^2} < \frac{1}{2} \sum_{i=1}^N \frac{W_\delta(z_i)^2}{1 + \delta W_\delta(z_i)^2} \quad (81)$$

$$< \sum_{i=1}^N \frac{W_\delta(z_i)^2}{1 + \delta W_\delta(z_i)^2} \left(\frac{1}{2} + \frac{1}{1 + \delta W_\delta(z_i)^2} \right) \text{ for } \delta \geq \delta_M, \quad (82)$$

805 showing that $D(\delta \mid \mathbf{z})|_{\delta \geq \delta_M} < 0$. That is, $D(\delta \mid \mathbf{z})$ approaches 0 from below for
 806 $\delta \rightarrow \infty$.

807 iii) By continuity and $D(\delta \mid \mathbf{z})|_{\delta=0} > 0$ (if (80) does not hold), it must cross the
 808 $D(\delta \mid \mathbf{z}) = 0$ line at least once in the interval $(0, \delta_M)$, proving the existence of
 809 $\hat{\delta}_{MLE}$.

c) The log-likelihood can be decomposed in

$$\ell(\delta \mid \mathbf{z}) \propto \underbrace{-\frac{1}{2} \sum_{i=1}^N [W_\delta(z_i)]^2}_{\ell(\mu_x=0, \sigma_x=1 \mid W_\delta(\mathbf{z}))} + \underbrace{\sum_{i=1}^N \log \frac{W_\delta(z_i)}{z_i} - \log [1 + W(\delta z_i^2)]}_{\mathcal{R}(\delta \mid \mathbf{z})}. \quad (83)$$

810 Lemmas **A.5** and **A.6** show that $\mathcal{R}(\delta \mid \mathbf{z})$ is monotonically decreasing and $\ell(\mu_x = 0, \sigma_x =$
 811 $1 \mid W_\delta(\mathbf{z}))$ is monotonically increasing in δ .

812 Furthermore, $\lim_{\delta \rightarrow \infty} \ell(\mu_x = 0, \sigma_x = 1 \mid W_\delta(\mathbf{z})) = 0$, that is the input likelihood is
 813 monotonically increasing but bounded from above (by $0 = \log 1$). On the other hand the
 814 penalty is decreasing without bounds, $\lim_{\delta \rightarrow \infty} \mathcal{R}(\delta \mid \mathbf{z}) = -\infty$. Thus their sum attains
 815 a global maximum either at the unique mode of $\ell(\delta \mid \mathbf{z})$ or at the boundary $\delta = 0$ - see
 816 also Fig. **6b**.

817 □

Algorithm 1 Find optimal δ : function `delta_GMM(·)` in the `LambertW` package.

Input: standardized data vector $\mathbf{z}$; theoretical kurtosis $\gamma_2(X)$

Output: $\hat{\delta}_{GMM}$ as in (84)

- 1: $\hat{\delta}_{GMM} = \arg \min_{\delta} \|\hat{\gamma}_2(\mathbf{u}_{\delta}) - \gamma_2(X)\|$, where $\mathbf{u}_{\delta} = W_{\delta}(\mathbf{z})$ subject to $\delta \geq 0$
 - 2: **return** $\hat{\delta}_{GMM}$
-

C Details on IGMM

Here I present an iterative method to obtain $\hat{\tau}$, which builds on the input/output aspect and theoretical properties of the input X . For example, if a random variable should be exponentially distributed (e.g. waiting times), but the observed data shows heavier tails then it is natural to estimate $\sigma_x = \lambda^{-1}$ and δ such that the back-transformed data has skewness 2, as this is a particular property of exponential RVs - independent of the rate parameter λ ; to remove heavy tails in $\mathbf{y}$ we should choose τ such that the back-transformed data $\mathbf{x}_{\tau}$ has sample kurtosis 3; or for uniform input, we can try to find a τ such that $\mathbf{x}_{\tau}$ has a flat density estimate.

Here I describe the estimator for τ to remove heavy-tails in location-scale data, in the sense that the kurtosis of the input equals 3. It can be easily adapted to match other properties of the input as outlined above.

For a moment assume that $\mu_x = \mu_x^{(0)}$ and $\sigma_x = \sigma_x^{(0)}$ are known and fixed; only δ has to be estimated. A natural choice for δ is the one that results in back transformed data $\mathbf{x}_{\tau}$ ($\tau = (\mu_x^{(0)}, \sigma_x^{(0)}, \delta)$) with sample kurtosis $\hat{\gamma}_2(\mathbf{x}_{\tau})$ equal to the theoretical kurtosis $\gamma_2(X)$. Formally,

$$\hat{\delta}_{GMM} = \arg \min_{\delta} \|\gamma_2(X) - \hat{\gamma}_2(\mathbf{x}_{\tau})\|, \quad (84)$$

where $\|\cdot\|$ is a proper norm in $\mathbb{R}$.

While the concept of this estimator is identical to its skewed version (Goerg, 2011), it has one important advantage: the inverse transformation is bijective. Thus here we do not have to consider “lost” data points when applying the inverse transformation.

Discussion of Algorithm 1: The kurtosis of Y as a function of δ is continuous and monotonically increasing (see (22)). Also $u = W_{\delta}(z)$ has a smaller slope than the identity

Algorithm 2 Iterative Generalized Method of Moments (IGMM) : function IGMM($\cdot$) in the LambertW package.

Input: data vector $\mathbf{y}$; tolerance level tol ; theoretical kurtosis $\gamma_2(X)$

Output: IGMM parameter estimate $\hat{\tau}_{IGMM} = (\hat{\mu}_x, \hat{\sigma}_x, \hat{\delta})$

```

1: Set  $\tau^{(-1)} = (0, 0, 0)$ 
2: Starting values:  $\tau^{(0)} = (\mu_x^{(0)}, \sigma_x^{(0)}, \delta^{(0)})$ , where  $\mu_x^{(0)} = \tilde{\mathbf{y}}$  and  $\sigma_x^{(0)} = \bar{\sigma}_y \cdot \left( \frac{1}{\sqrt{(1-2\delta^{(0)})^{3/2}}} \right)^{-1}$ 
   are the sample median and standard deviation of  $\mathbf{y}$  divided by the standard deviation
   factor (see also (21)), respectively.  $\delta^{(0)} = \frac{1}{66} \left( \sqrt{66\hat{\gamma}_2(\mathbf{y}) - 162} - 6 \right) \rightarrow$  see (??) for
   details.
3:  $k = 0$ 
4: while  $\|\tau^{(k)} - \tau^{(k-1)}\| > tol$  do
5:    $\mathbf{z}^{(k)} = (\mathbf{y} - \mu_x^{(k)})/\sigma_x^{(k)}$ 
6:   Pass  $\mathbf{z}^{(k)}$  to Algorithm 1  $\rightarrow \delta^{(k+1)}$ 
7:   back-transform  $\mathbf{z}^{(k)}$  to  $\mathbf{u}^{(k+1)} = W_{\delta^{(k+1)}}(\mathbf{z}^{(k)})$ ; compute  $\mathbf{x}^{(k+1)} = \mathbf{u}^{(k+1)}\sigma_x^{(k)} + \mu_x^{(k)}$ 
8:   Update parameters:  $\mu_x^{(k+1)} = \bar{\mathbf{x}}_{k+1}$  and  $\sigma_x^{(k+1)} = \hat{\sigma}_{x_{k+1}}$ 
9:    $\tau^{(k+1)} = (\mu_x^{(k+1)}, \sigma_x^{(k+1)}, \delta^{(k+1)})$ 
10:   $k = k + 1$ 
11: return  $\tau_{IGMM} = \tau^{(k)}$ 

```

837 $u = z$, and the slope is decreasing as δ is increasing. Thus if the kurtosis of the original data
838 is larger than the target kurtosis of the back-transformed data, $\hat{\gamma}_2(\mathbf{y}) > \gamma_2(X)$, then there
839 always exists a $\delta^{(*)}$ that achieves $\hat{\gamma}_2(\mathbf{x}_{\tau^*}) \equiv \gamma_2(X)$. By the re-parametrization $\tilde{\delta} = \log \delta$
840 the bounded optimization problem can be solved by standard (unbounded) optimization
841 algorithms.

842

843 In practice, μ_x and σ_x are rarely known but also have to be estimated from the data.
844 As $\mathbf{y}$ is shifted and scaled *ahead of* the back-transformation $W_\delta(\cdot)$, the initial choice of μ_x
845 and σ_x affects the optimal choice of δ . Therefore the optimal triple $\hat{\tau} = (\hat{\mu}_x, \hat{\sigma}_x, \hat{\delta})$ must be
846 obtained iteratively.

847 **Discussion of Algorithm 2:** Algorithm 2 first computes $\mathbf{z}^{(k)} = (\mathbf{y} - \mu_x^{(k)})/\sigma_x^{(k)}$ using $\mu_x^{(k)}$
848 and $\sigma_x^{(k)}$ from the previous step. This normalized output can then be passed to Algorithm
849 1 to obtain an updated $\delta^{(k+1)} = \hat{\delta}_{GMM}$. Using this new $\delta^{(k+1)}$ one can back-transform $\mathbf{z}^{(k)}$
850 to $\mathbf{u}^{(k+1)} = W_{\delta^{(k+1)}}(\mathbf{z}^{(k)})$, and consequently obtain a better approximation to the “true”
851 latent $\mathbf{x}$ by $\mathbf{x}^{(k+1)} = \mathbf{u}^{(k+1)}\sigma_x^{(k)} + \mu_x^{(k)}$. However, $\delta^{(k+1)}$ - and therefore $\mathbf{x}^{(k+1)}$ - has been

852 obtained using $\mu_x^{(k)}$ and $\sigma_x^{(k)}$, which are not necessarily the most accurate estimates in light
853 of the updated approximation $\widehat{\mathbf{x}}_{(\mu_x^{(k)}, \sigma_x^{(k)}, \delta^{(k+1)})}$. Thus Algorithm 2 computes new estimates
854 $\mu_x^{(k+1)}$ and $\sigma_x^{(k+1)}$ by the sample mean and standard deviation of $\widehat{\mathbf{x}}_{(\mu_x^{(k)}, \sigma_x^{(k)}, \delta^{(k+1)})}$, and starts
855 another iteration by passing the updated normalized output $\mathbf{z}^{(k+1)} = \frac{\mathbf{y} - \mu_x^{(k+1)}}{\sigma_x^{(k+1)}}$ to Algorithm
856 1 to obtain a new $\delta^{(k+2)}$.

857 It returns the optimal $\widehat{\tau}_{\text{IGMM}}$ once convergence has been reached, i.e., if $||\tau^{(k)} - \tau^{(k+1)}|| <$
858 tol .

859

Remark C.1 (IGMM for double-tail Lambert $W \times F_X$). *For a double-tail fit the one-dimensional optimization in Algorithm 1 has to be replaced with a two-dimensional optimization*

$$\left(\widehat{\delta}_\ell, \widehat{\delta}_r\right)_{\text{GMM}} = \arg \min_{\delta_\ell, \delta_r} h\left(\gamma_2(X) - \widehat{\gamma}_2(\mathbf{x}_{(\mu_x^*, \sigma_x^*, \delta_\ell, \delta_r)})\right). \quad (85)$$

860 Algorithm 2 remains unchanged.

Algorithm 3 Random sample generation : function `rLambertW(·)` in `LambertW` package.

Input: number of observations n ; parameter vector θ ; specification of the input distribution $F_X(x)$

Output: random sample $(y_1, \dots, y_n)$ of a Lambert $W \times F_X$ RV.

- 1: Simulate n samples $\mathbf{x} = (x_1, \dots, x_n) \sim F_X(x)$.
 - 2: Compute $\mu_x = \mu_x(\boldsymbol{\beta})$ and $\sigma_x = \sigma_x(\boldsymbol{\beta})$ (for scale family set $\mu_x = 0$, for non-central, non-scaled also set $\sigma_x = 1$)
 - 3: Compute normalized $\mathbf{u} = (\mathbf{x} - \mu_x)/\sigma_x$.
 - 4: $\mathbf{z} = \mathbf{u} \exp\left(\frac{\delta}{2} \mathbf{u}^2\right)$
 - 5: **return** $\mathbf{y} = \mathbf{z}\sigma_x + \mu_x$
-

D Simulation Details

Slightly heavy-tailed: $\delta = 1/10$. Here the RV Y has slight excess kurtosis ($3 + 2.51$) and $\sigma_y(\delta, \sigma_x = 1) = 1.18$. The Lambert W estimates of $\hat{\tau}$ are unbiased, and have smaller empirical standard deviation for $\hat{\mu}_x$ than the Gaussian MLE or the median. Also using Lambert W estimators does not give worse estimates for σ_y .

$\delta = 1/10$	median	Gaussian MLE		IGMM				Lambert W MLE				NA
N		μ_y	σ_y	μ_x	σ_x	δ	σ_y	μ_x	σ_x	δ	σ_y	ratio
50	-0.02	-0.02	1.15	-0.02	1.02	0.08	1.18	-0.02	0.99	0.09	∞	0
100	0.00	0.00	1.17	0.00	1.02	0.09	1.18	0.00	1.00	0.09	1.18	0
250	0.00	0.00	1.18	0.00	1.01	0.09	1.18	0.00	1.00	0.10	1.18	0
1000	0.00	0.00	1.18	0.00	1.00	0.10	1.18	0.00	1.00	0.10	1.18	0
50	0.56	0.53	0.61	0.54	0.48	0.64	0.55	0.53	0.55	0.58	0.56	0
100	0.50	0.49	0.57	0.50	0.45	0.61	0.54	0.49	0.51	0.56	0.54	0
250	0.50	0.48	0.56	0.47	0.46	0.56	0.53	0.48	0.51	0.54	0.53	0
1000	0.48	0.49	0.53	0.48	0.50	0.54	0.51	0.48	0.52	0.51	0.52	0
50	1.27	1.22	1.13	1.18	1.03	0.52	1.27	1.16	1.07	0.62	NA	0
100	1.28	1.19	1.21	1.15	1.07	0.60	1.26	1.12	1.09	0.64	1.28	0
250	1.26	1.19	1.20	1.12	1.09	0.63	1.22	1.09	1.09	0.65	1.23	0
1000	1.23	1.17	1.26	1.11	1.14	0.66	1.26	1.08	1.11	0.63	1.23	0

(a) Slightly heavy-tailed data: $\delta = 1/10$

Table 5: Based on $n = 1,000$ replications. In each sub-table: (first rows) average, (middle rows) proportion of estimates below true value, (bottom rows) empirical standard deviation times $\sqrt{N}$.