

A BAYESIAN STATISTICAL APPROACH FOR INFERENCE ON STATIC ORIGIN-DESTINATION MATRICES

BY LUIS CARVALHO*

Boston University

We address the problem of static OD matrix estimation from a formal statistical viewpoint. We adopt a novel Bayesian framework to develop a class of models that explicitly cast trip configurations in the study region as random variables. As a consequence, classical solutions from growth factor, gravity, and maximum entropy models are identified to specific estimators under the proposed models. We show that each of these solutions usually account for only a small fraction of the posterior probability mass in the ensemble and we then contend that the uncertainty in the inference should be propagated to later analyses or next-stage models. We also propose alternative, more robust estimators and devise Markov chain Monte Carlo sampling schemes to obtain them and perform other types of inference. We present several examples showcasing the proposed models and approach, and highlight how other sources of data can be incorporated in the model and inference in a principled, non-heuristic way.

1. Introduction. Consider a study region divided into n zones where trips can occur between any pair of zones. During a certain time period we observe the number of trips *originated* at zone i , O_i , and the number of trips *destined* to zone j , D_j , for $i, j = 1, \dots, n$. Our objective is to estimate the number of trips T_{ij} from each zone i to each zone j —including intrazonal trips T_{ii} —conditional on the $\mathcal{O} = \{O_i\}_{i=1}^n$ and $\mathcal{D} = \{D_j\}_{j=1}^n$. Since the trips $\mathcal{T} = \{T_{ij}\}_{i,j=1,\dots,n}$ can be represented by the matrix

$$(1.1) \quad M = \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1n} \\ T_{21} & T_{22} & \cdots & T_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ T_{n1} & T_{n2} & \cdots & T_{nn} \end{bmatrix},$$

and we are fixing a time window for the trip realizations, our problem is usually referred to as *static OD matrix estimation*. We note that the OD

*Supported by NSF grant DMS-1107067.

Keywords and phrases: static OD matrix estimation, random matrix, constrained sampling

matrix M has restrictions on its row and column margins,

$$(1.2) \quad \begin{aligned} \sum_{j=1}^n T_{ij} &= O_i, \quad i = 1, \dots, n, \\ \sum_{i=1}^n T_{ij} &= D_j, \quad j = 1, \dots, n. \end{aligned}$$

and thus the estimation is constrained. We also require that $\sum_{i=1}^n O_i = \sum_{j=1}^n D_j \doteq T$ for consistency. It should be immediate from this formulation that static OD matrix estimation is a *contingency table* problem[2]; our goal here is to provide a broader treatment from a more applied perspective.

This problem has been studied for many decades in the transportation literature. The first contributions to its solution adopted a physical interpretation and assumed \mathcal{T} could be described by a gravitational law [1]: $T_{ij} \propto O_i D_j d_{ij}^{-2}$, where d_{ij} is the distance between zones i and j . This functional relation was later generalized to include decreasing functions of traveling costs c_{ij} between zones i and j , called “deterrence” functions:

$$(1.3) \quad T_{ij} \propto O_i D_j d(c_{ij}).$$

Common choices for d include exponential linear functions of costs, such as $d(c_{ij}) = \exp(-\beta c_{ij})$ or $d(c_{ij}) = \exp(-\beta c_{ij} - \alpha \log c_{ij})$. These gravity models are *synthetic* models since they do not incorporate previously observed trip patterns. In contrast, *growth factor* models regard \mathcal{T} as possible future trip patterns and incorporate previous observations in a doubly constrained formulation. Let the “seed” matrix $\mathcal{T}_0 = \{t_{ij}\}_{i,j=1,\dots,n}$ be previous observations from the same or similar study region. Based on the method proposed by Furness [4], we assume

$$(1.4) \quad T_{ij} = A_i O_i B_j D_j t_{ij},$$

where A_i and B_j are “balancing factors” that are known up to a proportionality constant. Furness method defines \mathcal{T} by iteratively solving for the balancing factors to respect constraints (1.2) until convergence.

Both gravity and growth factor models provide estimates for \mathcal{T} based on heuristic, functional arguments. Wilson [13, 14] defined a formulation based on entropy maximization that would unify both previous approaches. If

$$W(\mathcal{T}) = \frac{T!}{\prod_{i,j} T_{ij}!}$$

is the number of “micro” states associated with “meso” state \mathcal{T} , then the trip configuration that maximizes W , or equivalently

$$\log W(\mathcal{T}) - \log T! \approx - \sum_{i,j} \left(T_{ij} \log T_{ij} - T_{ij} \right),$$

subject to constraints (1.2) is a maximum entropy solution. If instead of $\log W$ we maximize

$$\log W'(\mathcal{T}, \mathcal{T}_0) = - \sum_{i,j} \left(T_{ij} \log \frac{T_{ij}}{t_{ij}} - T_{ij} \right)$$

the solution would coincide with the one provided by the Furness model. By adding an additional cost constraint, such as

$$(1.5) \quad \sum_{i,j} c_{ij} T_{ij} = C_T$$

we obtain the same estimates from the gravity model with $d(c_{ij}) = \exp(-\beta c_{ij})$.

We can make two important observations from the maximum entropy approach. First, we note that the functional expressions for T_{ij} from the gravity and Furness models can actually be regarded as closed form expressions that can be used to iteratively obtain solutions to a mathematical program that maximizes $\log W$ or $\log W'$ subject to certain constraints. Second, since there are many feasible configurations for \mathcal{T} , we can define weights—in Wilson’s case given by W —to help us find the best trip configuration; it is, however, implicit from this formulation that any other trip pattern but the “optimal” is also possible, or even likely, to occur.

In this paper we propose a formulation for the OD matrix estimation problem where \mathcal{T} is explicitly *random*. As we will show, this formulation corresponds to a Bayesian statistical approach, e.g. [5]. Even though our focus will be on exploring the randomness associated with the trip patterns instead of simply extracting a single trip pattern through optimization, we show that the maximum entropy solutions, including the classical gravity and growth model solutions, are identified with maximum *a posteriori* (MAP) estimates under our setup. Besides this unifying consequence, Bayesian methods also provide other types of estimators and, more generally, are able to quantify the uncertainty in estimation and to propagate it to posterior analyses in a principled, integrated framework.

2. Proposed Model. Let us say that the trips \mathcal{T} are $(\mathcal{O}, \mathcal{D})$ -consistent, denoted by $\mathcal{T} \in C(\mathcal{O}, \mathcal{D})$, if \mathcal{T} satisfies equations (1.2). That is, we define

$$C(\mathcal{O}, \mathcal{D}) = \left\{ \tilde{\mathcal{T}} = \{\tilde{T}_{ij}\} : \sum_{j=1}^n \tilde{T}_{ij} = O_i \text{ and } \sum_{i=1}^n \tilde{T}_{ij} = D_j \right\}.$$

As stated before, we regard \mathcal{T} as *random* while margin trips \mathcal{O} and \mathcal{D} are observed data. As usual in the fully Bayesian approach we pursue next, all inferences are driven by the posterior distribution on \mathcal{T} conditional on data \mathcal{O} and \mathcal{D} ,

$$P(\mathcal{T} | \mathcal{O}, \mathcal{D}) = \frac{P(\mathcal{O}, \mathcal{D} | \mathcal{T})P(\mathcal{T})}{\sum_{\tilde{\mathcal{T}}} P(\mathcal{O}, \mathcal{D} | \tilde{\mathcal{T}})P(\tilde{\mathcal{T}})}.$$

Let us then consider the simple likelihood

$$(2.1) \quad P(\mathcal{O}, \mathcal{D} | \mathcal{T}) = I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})]$$

where $I(\cdot)$ is the indicator function: $I(A) = 1$ if and only if A is true. By the definition of OD consistency, the likelihood in equation (2.1) just states that the margin trips satisfy equations (1.2), that is, it is a simple indicator for $(\mathcal{O}, \mathcal{D})$ -consistency.

The randomness in trips \mathcal{T} comes initially from our belief, before observing any data in the margins, of how the trips are distributed. This belief is hardly subjective, but often arises from experience on similar regions and zones; in the next section we discuss how to incorporate knowledge gathered from small scale studies in the same region. To establish a parallel to the maximum entropy approach of the previous section, we assume that \mathcal{T} has a conditional multinomial prior distribution given by $\mathcal{T} | T \sim \text{MN}(T, \mathbf{p})$, that is,

$$P(\mathcal{T} | T) = \frac{T!}{\prod_{i,j} T_{ij}!} \prod_{i,j} p_{ij}^{T_{ij}},$$

where T is the total number of trips in the region and $\mathbf{p} = \{p_{ij}\}_{i,j=1,\dots,n}$ with p_{ij} being the proportion of trips between zones i and j . Of course, we require that $\sum_{i,j} p_{ij} = 1$ and p_{ij} are nonnegative. The hyper-prior parameter T has an improper non-informative distribution $P(T) \propto 1$, and so the prior

becomes

$$\begin{aligned}
 P(\mathcal{T}) &= \sum_{T=0}^{\infty} P(\mathcal{T} | T) P(T) \\
 (2.2) \quad &= \sum_{T=0}^{\infty} \frac{T!}{\prod_{i,j} T_{ij}!} \prod_{i,j} p_{ij}^{T_{ij}} I\left(\sum_{i,j} T_{ij} = T\right) \\
 &= \frac{(\sum_{i,j} T_{ij})!}{\prod_{i,j} T_{ij}!} \prod_{i,j} p_{ij}^{T_{ij}}.
 \end{aligned}$$

The prior on \mathcal{T} resembles the number of micro states W defined by Wilson, but with the proportions as extra parameters. The proportions \mathbf{p} have the important role of conveying prior information on the *structure* of trip distribution in the study area. From a behavioral perspective, p_{ij} corresponds to the probability of a trip in the system, out of the total T available, occurring between zones i and j ; we could, for example, borrowing from random decision theory, define a multinomial logit model on each p_{ij} that depends on a set of covariates \mathbf{x}_{ij} for each OD pair such as transport costs, time, and user preferences:

$$p_{ij} = \frac{\exp(\mathbf{x}_{ij}^T \boldsymbol{\beta})}{\sum_{k,l=1,\dots,n} \exp(\mathbf{x}_{kl}^T \boldsymbol{\beta})},$$

where $\boldsymbol{\beta}$ are known coefficients.

While we are now assuming that \mathbf{p} is known and thus fully specifies $P(\mathcal{T})$ above, we can further incorporate uncertainty by adding another level of randomness to the prior parameters to form a hierarchical model; we postpone such considerations to Section 3.

2.1. Estimation. The inference we wish to carry out is driven by our updated belief in \mathcal{T} after observing \mathcal{O} and \mathcal{D} as summarized by the posterior distribution

$$\begin{aligned}
 P(\mathcal{T} | \mathcal{O}, \mathcal{D}) &= \frac{P(\mathcal{O}, \mathcal{D} | \mathcal{T}) P(\mathcal{T})}{\sum_{\tilde{\mathcal{T}}} P(\mathcal{O}, \mathcal{D} | \tilde{\mathcal{T}}) P(\tilde{\mathcal{T}})} \\
 (2.3) \quad &= \frac{I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})] P(\mathcal{T})}{\sum_{\tilde{\mathcal{T}} \in C(\mathcal{O}, \mathcal{D})} P(\tilde{\mathcal{T}})} \\
 &\propto \frac{T!}{\prod_{i,j} T_{ij}!} \prod_{i,j} p_{ij}^{T_{ij}} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})].
 \end{aligned}$$

One important consequence of $\mathcal{T} \in C(\mathcal{O}, \mathcal{D})$ in the posterior above is that the prior parameter T implicitly satisfies

$$(2.4) \quad T = \sum_{i,j} T_{ij} = \sum_{i=1}^n O_i = \sum_{j=1}^n D_j,$$

that is, \mathcal{O} and \mathcal{D} are self-consistent through \mathcal{T} .

A common estimator for \mathcal{T} is the maximum *a posteriori* (MAP) estimator, the posterior mode:

$$\begin{aligned} \hat{\mathcal{T}} &= \arg \max_{\mathcal{T}} \left\{ \log P(\mathcal{T} | \mathcal{O}, \mathcal{D}) \right\} \\ &= \arg \max_{\mathcal{T} \in C(\mathcal{O}, \mathcal{D})} \left\{ \sum_{i,j} T_{ij} \log p_{ij} - \log T_{ij}! \right\} \\ &\approx \arg \max_{\mathcal{T} \in C(\mathcal{O}, \mathcal{D})} \left\{ \sum_{i,j} T_{ij} \log p_{ij} - (T_{ij} \log T_{ij} - T_{ij}) \right\} \\ &= \arg \max_{\mathcal{T} \in C(\mathcal{O}, \mathcal{D})} \left\{ - \sum_{i,j} \left(T_{ij} \log \frac{T_{ij}}{p_{ij}} - T_{ij} \right) \right\}. \end{aligned}$$

Note the similarity between the maximand and $\log W'$. It is now straightforward to show that

$$\hat{T}_{ij} = A_i O_i B_j D_j p_{ij},$$

where A_i and B_j are balancing factors. Thus, the MAP estimator is equivalent to the solution obtained from the Furness method for the maximum entropy formulation. In fact, if we use a prior seed matrix $\mathcal{T}_0 = \{t_{ij}\}$ to set $p_{ij} = t_{ij} / \sum_{i,j} t_{ij}$, the prior proportions, we recover the growth factor solution.

To obtain gravity model solutions we just have to define \mathbf{p} based on an entropy maximizing principle: we want \mathbf{p} that maximizes the entropy $\mathcal{H}(\mathbf{p}) = - \sum_{i,j} p_{ij} \log p_{ij}$ possibly subject to additional constraints on \mathbf{p} other than $\sum_{i,j} p_{ij} = 1$. Since entropy uniquely measures the amount of uncertainty in a probability distribution, a maximum entropy assignment is justified as the only unbiased assumption we can attain under a state of partial knowledge of the system. As Wilson [13, pg. 10] points out, “the probability distribution which maximizes entropy makes the weakest assumption which is consistent with what is known”. If we then constraint on trip costs by requiring a fixed mean cost in the region

$$(2.5) \quad \sum_{i,j} c_{ij} p_{ij} = C_p,$$

we obtain $p_{ij} \propto \exp(-\beta c_{ij})$, and hence a gravity model with a familiar exponential deterrence function.

Even though setting \mathbf{p} as above provides the same solution, there is a subtle but important difference to the original maximum entropy formulation: in Wilson’s model we constraint the trip patterns using (1.5), effectively reducing the number of feasible trip configurations, while in our proposed model we only restrict the proportions using (2.5) to redefine the weights on trip patterns. In other words, our feasible space is still only constrained by (1.2), but we set the proportions as a structural guide for estimation since the shape of the posterior distribution on \mathcal{T} depends on \mathbf{p} . In this sense, we can think of (2.5) as a “soft” constraint. We can argue that such a formulation is more natural since we can certainly have prior knowledge of overall transport expenditures in the system while it seems artificial to establish a rigid cost constraint on the whole study region.

Another good estimator is the posterior mean, defined as

$$\bar{\mathcal{T}} = \mathbb{E}[\mathcal{T} | \mathcal{O}, \mathcal{D}] = \sum_{\tilde{\mathcal{T}}} \tilde{\mathcal{T}} \cdot \mathbb{P}(\tilde{\mathcal{T}} | \mathcal{O}, \mathcal{D}).$$

The posterior mean is more “robust” than the posterior mode since it averages the uncertainty on trip patterns across all possible \mathcal{T} —weighted by their respective posterior probability mass—as opposed to simply picking the trip pattern with highest posterior probability. Moreover, since the posterior mean is a linear combination of feasible trip patterns, it also satisfies the linear constraints in (1.2). There is, however, one major difficulty in this venue: we need to know $\mathbb{P}(\mathcal{T} | \mathcal{O}, \mathcal{D})$ for each \mathcal{T} .

The main hurdle in evaluating the posterior on \mathcal{T} in (2.3) is the normalizing factor $Z(\mathcal{O}, \mathcal{D}) \doteq \sum_{\tilde{\mathcal{T}} \in \mathcal{C}(\mathcal{O}, \mathcal{D})} \mathbb{P}(\tilde{\mathcal{T}})$. Computing $Z(\mathcal{O}, \mathcal{D})$ requires summing over all possible pairwise trip assignments that are $(\mathcal{O}, \mathcal{D})$ -consistent, a daunting task. Before addressing this central issue, we offer some motivation in the next subsection.

2.2. A simple example. Suppose that, for $n = 2$ zones, we observe O_1 , O_2 , D_1 , D_2 , and wish to estimate the entries \mathcal{T} in the OD matrix

$$\begin{array}{cc|c} T_{11} & T_{12} & O_1 \\ T_{21} & T_{22} & O_2 \\ \hline D_1 & D_2 & T \end{array}$$

with margins and total number of trips T displayed.

Since \mathcal{T} is consistent, we know that $T_{12} = O_1 - T_{11}$, $T_{21} = D_1 - T_{11}$ and $T_{22} = O_2 - T_{11} = T_{11} - (T - O_2 - D_2) = T_{11} - \Delta$, where we set

$\Delta \doteq T - O_2 - D_2$. The posterior on \mathcal{T} is then a posterior on T_{11} due to these linear constraints:

$$\begin{aligned}
 (2.6) \quad P(T_{11} | \mathcal{O}, \mathcal{D}) &\propto \frac{T!}{T_{11}!T_{12}!T_{21}!T_{22}!} p_{11}^{T_{11}} p_{12}^{T_{12}} p_{21}^{T_{21}} p_{22}^{T_{22}} \\
 &\propto \frac{p_{11}^{T_{11}} p_{12}^{O_1 - T_{11}} p_{21}^{D_1 - T_{11}} p_{22}^{T_{11} - \Delta}}{T_{11}!(O_1 - T_{11})!(D_1 - T_{11})!(T_{11} - \Delta)!} \\
 &\propto \binom{O_1}{T_{11}} \binom{D_1 - \Delta}{D_1 - T_{11}} \psi^{T_{11}} \\
 &\doteq H(T_{11}; O_1, D_1, \Delta, \psi),
 \end{aligned}$$

where $\psi = (p_{11}p_{22})/(p_{12}p_{21})$ can be interpreted as a intra-interzonal odds ratio. Since $D_1 - \Delta = T - O_1$, we can see that T_{11} follows a *non-central hypergeometric distribution*[11]:

$$T_{11} | \mathcal{O}, \mathcal{D} \sim \text{HG}(O_1, D_1, \Delta; \psi).$$

Note that $\mathcal{T} \in C(\mathcal{O}, \mathcal{D})$ is equivalent to requiring that $\max\{0, \Delta\} \leq T_{11} \leq \min\{O_1, D_1\}$, and so the normalizing constant for (2.6) is the sum of its right-hand side over the values of T_{11} above. In practice, however, it is simpler to obtain posterior samples of T_{11} using a *Metropolis-Hastings* algorithm [9, 7, 8].

As proposal we adopt a random walk: given our actual position $T_{11}^{(t-1)}$ at iteration $t-1$, we set our candidate T_{11}^* a step to the left, $T_{11}^* = T_{11}^{(t-1)} - 1$ with probability 0.5 or a step to the right, $T_{11}^* = T_{11}^{(t-1)} + 1$ with probability 0.5. If $T_{11}^* < \max\{0, \Delta\}$ or $T_{11}^* > \min\{O_1, D_1\}$ we immediately reject T_{11}^* —and set $T_{11}^{(t)} = T_{11}^{(t-1)}$ —as it is out of bounds. Otherwise we accept T_{11}^* —and thus set $T_{11}^{(t)} = T_{11}^*$ —with probability $\min\{R(T_{11}^{(t-1)}, T_{11}^*), 1\}$, where $R(T_{11}^{(t-1)}, T_{11}^*)$ is the acceptance ratio

$$R(T_{11}^{(t-1)}, T_{11}^*) = \frac{H(T_{11}^*; O_1, D_1, \Delta, \psi)}{H(T_{11}^{(t-1)}; O_1, D_1, \Delta, \psi)}.$$

We denote this Metropolis step by

$$T_{11}^{(t)} = MS(T_{11}^{(t-1)}; O_1, D_1, \Delta, \psi).$$

To summarize, we can obtain samples from T_{11} by doing:

Step 1. Start at some arbitrary initial $T_{11}^{(0)}$.

Step 2. For $t = 1, 2, \dots$ do (until convergence): execute a Metropolis step,

$$T_{11}^{(t)} = MS(T_{11}^{(t-1)}; O_1, D_1, \Delta, \psi),$$

that is,

Step 2.1. Sample candidate T_{11}^* : sample $U \sim U(0, 1)$; if $U < 0.5$ set $T_{11}^* = T_{11}^{(t-1)} - 1$, otherwise set $T_{11}^* = T_{11}^{(t-1)} + 1$.

Step 2.2. If $T_{11}^* < \max\{0, \Delta\}$ or $T_{11}^* > \min\{O_1, D_1\}$ set $T_{11}^{(t)} = T_{11}^{(t-1)}$ (reject). Otherwise, sample $U \sim U(0, 1)$: if $U < \min\{R(T_{11}^{(t-1)}, T_{11}^*), 1\}$ then set $T_{11}^{(t)} = T_{11}^*$ (accept), else set $T_{11}^{(t)} = T_{11}^{(t-1)}$ (reject).

A numerical example should help us further gain intuition on the problem.

EXAMPLE 1. Let $O_1 = 40$, $O_2 = 40$, $D_1 = 60$, $D_2 = 20$, $p_{11} = 0.1$, $p_{12} = 0.2$, $p_{21} = 0.3$, and $p_{22} = 0.4$. It follows that $T = O_1 + O_2 = D_1 + D_2 = 80$, $\Delta = T - O_2 - D_2 = 20$, and $\psi = (p_{11}p_{22})/(p_{12}p_{21}) = (0.1 \cdot 0.4)/(0.2 \cdot 0.3) = 2/3$, and so $T_{11} \sim \text{HG}(40, 60, 20; 2/3)$.

Using random walk Metropolis samples $T_{11}^{(1)}, \dots, T_{11}^{(G)}$ we can produce point estimates for T_{11} if desired: the posterior mean,

$$\bar{T}_{11} = E[T_{11} \mid \mathcal{O}, \mathcal{D}] \approx \frac{1}{G} \sum_{g=1}^G T_{11}^{(g)},$$

and the posterior mode,

$$\hat{T}_{11} = \arg \max_{x=\max\{0, \Delta\}, \dots, \min\{O_1, D_1\}} P(T_{11} = x \mid \mathcal{O}, \mathcal{D}).$$

\hat{T}_{11} can be obtained from estimates for $P(T_{11} \mid \mathcal{O}, \mathcal{D})$, by Monte Carlo simulation,

$$(2.7) \quad P(T_{11} = x \mid \mathcal{O}, \mathcal{D}) \approx \frac{1}{G} \sum_{g=1}^G I(T_{11}^{(g)} = x),$$

or from the Furness method. Using $G = 10,000$, we obtain $\bar{T}_{11} = 28.43$ and $\hat{T}_{11} = 28.49$, and so both the posterior mean and posterior mode, estimated from our samples and rounded to the nearest feasible integer, are ≈ 28 . It is not uncommon for both estimates to coincide, especially when the distribution is unimodal and close to symmetric, as in this case.

Interestingly, $P(T_{11} = 28 | \mathcal{O}, \mathcal{D}) \approx 0.20$; even for this simple example with a small number of trips we can see that the probability of the most probable trip configuration corresponds to a small fraction of possible configurations. This effect should not come as a surprise: as the number of zones and margins grow, so do the number of possible consistent configurations, and so the probability of any single trip configuration becomes even smaller.

We have previously remarked on the structural role of the proportions \mathbf{p} , serving as a guide when searching for a representative trip pattern among the many possible feasible configurations. We note, however, that there is no principled reason to expect a close relation between \mathbf{p} and actual proportions \mathcal{T}/T since the latter is constrained by origin and destination margins. As an example, consider Figure 1, where we show the marginal posterior distributions of T_{11} , T_{12} , T_{21} , and T_{22} , along with expected “structural” number of trips given by $T\mathbf{p}$. The discrepancies are clear once we observe that $Tp_{11} + Tp_{12} = 24 < 40 = O_1$ and similarly for the other margins; equivalently, $(T_{11} + T_{12})/T = 0.5 > 0.3 = p_{11} + p_{12}$ for any (feasible) trip pattern \mathcal{T} .




FIG 1. *Estimated posterior distributions of \mathcal{T} from 10,000 samples. Squares mark expected structural trips.*

2.3. Posterior sampler. Let us now extend the results from the last section to our problem. In general, for n zones we have the following OD matrix

with margins displayed:

$$\begin{array}{cccc|c}
 T_{11} & T_{12} & \cdots & T_{1n} & O_1 \\
 T_{21} & T_{22} & \cdots & T_{2n} & O_2 \\
 \vdots & \vdots & \ddots & \vdots & \vdots \\
 T_{n1} & T_{n2} & \cdots & T_{nn} & O_n \\
 \hline
 D_1 & D_2 & \cdots & D_n & T
 \end{array}$$

We now proceed to eliminate the first $n - 1$ entries in the last row and column by means of the linear constraints in the margins:

$$\begin{aligned}
 (2.8) \quad T_{nj} &= D_j - \sum_{i=1}^{n-1} T_{ij}, \quad j = 1, \dots, n-1, \\
 T_{in} &= O_i - \sum_{j=1}^{n-1} T_{ij}, \quad i = 1, \dots, n-1.
 \end{aligned}$$

The corner entry T_{nn} requires special handling:

$$\begin{aligned}
 (2.9) \quad T_{nn} &= O_n - \sum_{j=1}^{n-1} T_{nj} \\
 &= O_n - \sum_{j=1}^{n-1} \left(D_j - \sum_{i=1}^{n-1} T_{ij} \right) \\
 &= \sum_{i,j=1}^{n-1} T_{ij} - \left(\sum_{j=1}^{n-1} D_j - O_n \right) \\
 &= \sum_{i,j=1}^{n-1} T_{ij} - \underbrace{(T - O_n - D_n)}_{\Delta}.
 \end{aligned}$$

Ultimately, T_{nn} stems from the symmetry in equation (2.4).

To sample from the entries in the $(n - 1)$ -by- $(n - 1)$ upper submatrix S we adopt a *Gibbs sampler* [6]; see also [7, 8]. The conditional posterior distributions are $P(T_{ij} | T_{[ij]}, \mathcal{O}, \mathcal{D})$, for $i, j = 1, \dots, n - 1$, where $T_{[ij]}$ denotes all the entries in \mathcal{T} but T_{ij} , that is, $T_{[ij]} \doteq \{T_{kl}\}_{k,l=1,\dots,n-1, k \neq i, l \neq j}$. The only terms in $P(\mathcal{T} | \mathcal{O}, \mathcal{D})$ that depend on T_{ij} are now related to T_{in} and T_{nj}

through equations (2.8) and to T_{nn} through equation (2.9). Namely,

$$\begin{aligned}
 (2.10) \quad P(T_{ij} | T_{[ij]}, \mathcal{O}, \mathcal{D}) &\propto \frac{p_{ij}^{T_{ij}} p_{in}^{T_{in}} p_{nj}^{T_{nj}} p_{nn}^{T_{nn}}}{T_{ij}! T_{in}! T_{nj}! T_{nn}!} \\
 &\doteq \frac{p_{ij}^{T_{ij}} p_{in}^{O_{ij}-T_{ij}} p_{nj}^{D_{ij}-T_{ij}} p_{nn}^{T_{ij}-\Delta_{ij}}}{T_{ij}! (O_{ij}-T_{ij})! (D_{ij}-T_{ij})! (T_{ij}-\Delta_{ij})!} \\
 &\propto \binom{O_{ij}}{T_{ij}} \binom{D_{ij}-\Delta_{ij}}{D_{ij}-T_{ij}} \psi_{ij}^{T_{ij}},
 \end{aligned}$$

where we define $O_{ij} \doteq O_i - \sum_{l=1, \dots, n-1, l \neq j} T_{il}$, $D_{ij} \doteq D_j - \sum_{k=1, \dots, n-1, k \neq i} T_{kj}$, $\Delta_{ij} \doteq \Delta - \sum_{k,l=1, \dots, n-1, k \neq i, l \neq j} T_{kl}$, and $\psi_{ij} \doteq (p_{ij} p_{nn}) / (p_{in} p_{nj})$ —a “within-between” odds trip ratio—to simplify the expressions. Thus,

$$(2.11) \quad T_{ij} | T_{[ij]}, \mathcal{O}, \mathcal{D} \sim \text{HG}(O_{ij}, D_{ij}, \Delta_{ij}; \psi_{ij}).$$

It is now straightforward to sample from the posterior for \mathcal{T} using a hybrid Metropolis-within-Gibbs sampling scheme since we know how to sample from the non-central hypergeometric:

Step 1. Start at some arbitrary initial configuration $\mathcal{T}^{(0)}$.

Step 2. For $t = 1, 2, \dots$ do (until convergence):

Step 2.1. For $i, j = 1, \dots, n-1$ do: sample $T_{ij}^{(t)} \sim T_{ij} | T_{[ij]}^{(t-1)}, \mathcal{O}, \mathcal{D}$ in (2.11) using a Metropolis step,

$$T_{ij}^{(t)} = MS(T_{ij}^{(t-1)}; O_{ij}^{(t-1)}, D_{ij}^{(t-1)}, \Delta_{ij}^{(t-1)}, \psi_{ij}),$$

with $O_{ij}^{(t-1)}$, $D_{ij}^{(t-1)}$, $\Delta_{ij}^{(t-1)}$, and ψ_{ij} defined as above. Note that all the parameters but ψ_{ij} depend on $T_{[ij]}^{(t-1)}$ and so carry an iteration index.

It should be noted that this sampling scheme is similar to the more general scheme from algebraic statistics and based on Markov basis [3].


EXAMPLE 2. We end this section with an example taken from [12, pg. 179]. The costs $\{c_{ij}\}$ between four zones are listed in Table 1, along with observed origin and destination margins.

Let us now assume that $p_{ij} \propto \exp(-\beta c_{ij})$ with $\beta = 0.10$. After running our Gibbs sampler until assumed convergence, we take $G = 10,000$ samples to perform posterior inference; the marginal posterior distributions for T_{ij} in the upper 3-by-3 matrix are summarized in Figure 2.

TABLE 1

Trip costs between four zones with observed origin and destination margins. Reproduced from [12, table 5.8].

Zone	1	2	3	4	O_i
1	3	11	18	22	400
2	12	3	13	19	460
3	15.5	13	5	7	400
4	24	18	8	5	702
D_j	260	400	500	802	1962

FIG 2. *Estimated posterior distributions of \mathcal{T} from 10,000 samples.*

The posterior mean $\overline{\mathcal{T}}$, estimated from our samples by

$$(2.12) \quad \overline{\mathcal{T}} = \mathbb{E}[\mathcal{T} \mid \mathcal{O}, \mathcal{D}] \approx \frac{1}{G} \sum_{g=1}^G \mathcal{T}^{(g)}$$

is very similar to the Furness solution reported in [12]. We list $\overline{\mathcal{T}}$ along with 95% credible intervals for each T_{ij} in Table 2. The credible intervals are wider than in our previous simple example due to the much higher number of feasible configurations in $\mathcal{C}(\mathcal{O}, \mathcal{D})$. In fact, we estimate from the posterior samples that $\mathbb{P}(\mathcal{T} = \overline{\mathcal{T}} \mid \mathcal{O}, \mathcal{D}) \approx \mathbb{P}(\mathcal{T} = \hat{\mathcal{T}} \mid \mathcal{O}, \mathcal{D}) \approx 2 \cdot 10^{-3}$. Since the most probable trip pattern accounts for only 0.2% of the posterior probability mass, we can conclude that even the Furness solution has little support from the data. Interval estimators now become more attractive representatives of the posterior space of trip configurations given a desired credibility level.

TABLE 2
Posterior mean and 95% credible intervals.

Zone	1		2		3		4	
1	157.14	[147, 169]	97.37	[85, 110]	68.73	[56, 81]	76.75	[64, 91]
2	58.70	[48, 68]	206.35	[190, 221]	101.27	[84, 116]	93.69	[79, 91]
3	24.16	[16, 33]	44.91	[33, 56]	138.32	[125, 151]	192.61	[177, 207]
4	20.00	[12, 29]	51.37	[40, 64]	191.68	[172, 211]	438.95	[418, 460]

An even better alternative is to use the whole posterior distribution to propagate the randomness in \mathcal{T} in our subsequent analyses. Consider, for instance, the mean regional cost

$$c(\mathcal{T}) = \sum_{i,j} c_{ij} T_{ij} / T,$$

and let us compare its posterior distribution, as induced by \mathcal{T} , to the fixed value C_p —the mean prior regional cost—we set as a restriction in (2.5) to define β . Since $\beta = 0.1$, $C_p = 8.51$. We can now use our samples $\mathcal{T}^{(1)}, \dots, \mathcal{T}^{(G)}$ from the Gibbs sampler to generate realizations

$$(2.13) \quad c(\mathcal{T}^{(g)}) = \sum_{i,j} c_{ij} T_{ij}^{(g)} / T$$

and estimate $P(c(\mathcal{T}) \mid \mathcal{O}, \mathcal{D})$. Figure 3 shows a histogram based on $\{c(\mathcal{T}^{(g)})\}$. The estimated posterior mean cost is $E[c(\mathcal{T}) \mid \mathcal{O}, \mathcal{D}] = c(\bar{\mathcal{T}}) = 8.67$, the posterior mode cost—the Furness solution cost—is $c(\hat{\mathcal{T}}) = 8.70$, both higher than C_p , while a 95% credible interval for $c(\mathcal{T})$ is $[8.46, 8.88]$, barely covering C_p ; moreover,

$$P(c(\mathcal{T}) \geq C_p \mid \mathcal{O}, \mathcal{D}) \approx \frac{1}{G} \sum_{g=1}^G I[c(\mathcal{T}^{(g)}) \geq C_p] = 0.93.$$

That a great proportion of possible trip patterns is spending more than previously expected strongly suggests that a lower value for β would be more realistic given the restrictions on \mathcal{T} by \mathcal{O} and \mathcal{D} .

We might also want to analyse the trip length distribution (TLD) of the system: given a set of K cost ranges $(c_0, c_1], \dots, (c_{K-1}, c_K]$, where $0 \leq c_0 < c_1 < \dots < c_K < \infty$, we bin the proportion of trips T_k/T with costs in the k -th range $(c_{k-1}, c_k]$ for each $k = 1, \dots, K$. We again use our samples to generate an estimate for each T_k :

$$(2.14) \quad T_k^{(g)} = \sum_{i,j} T_{ij}^{(g)} I\{c_{ij} \in (c_{k-1}, c_k]\}.$$




FIG 3. *Estimated posterior distribution of mean regional cost from 10,000 samples. Solid line indicates posterior mean, dashed line marks prior mean, and dash-dotted line marks posterior mode cost.*

Table 3 compares the mean posterior TLD with the prior TLD using aggregated range proportions $\{p_k\}_{k=1,\dots,K}$, where $p_k = \sum_{i,j} p_{ij} I\{c_{ij} \in (c_{k-1}, c_k]\}$. Figure 4 represents both TLD with additional 95% credible intervals for each range. The discrepancy between prior proportions \mathbf{p} and posterior proportions T_{ij}/T is now more evident due to the structure in the TLD. In the next section we will propose a principled way to narrow the gap between these two regional features.

TABLE 3
Mean posterior TLD and prior TLD from proportions \mathbf{p} .

Range	(0, 4]	(4, 8]	(8, 12]	(12, 16]	(16, 20]	(20, 24]
$E[T_k/T \mathcal{O}, \mathcal{D}]$	0.18	0.49	0.08	0.09	0.11	0.05
p_k	0.26	0.38	0.11	0.13	0.08	0.04

3. Extensions to the Proposed Model. As we have seen in the last example in the previous section, prior beliefs might be deceptively outdated or based on regions that are not similar to the current study region. As a consequence, the related posterior distribution might be wrongly biased and scaled, affecting the estimation. In addition, it is possible that during the process of eliciting the prior proportions we realize that the trip structure in the region is uncertain as it might change during the study time frame




FIG 4. Mean posterior TLD (bars) with 95% credible intervals (whiskers), and prior TLD (squares).

due to, for example, seasonal effects.

A natural approach is then to adopt our same viewpoint with respect to trip patterns and to explicitly quantify the uncertainty by regarding the proportions themselves as random, yielding a *hierarchical* model. Under this updated model the proportions \mathbf{p} are now random and our samples from the last section are now conditional on \mathbf{p} , that is, $P(\mathcal{T} | \mathcal{O}, \mathcal{D})$ becomes $P(\mathcal{T} | \mathbf{p}, \mathcal{O}, \mathcal{D})$. Nevertheless, we can still proceed in the same way we have done before if we integrate out the uncertainty in the nuisance parameters, the proportions, to obtain the marginal posterior distribution on the trips \mathcal{T} ,

$$(3.1) \quad P(\mathcal{T} | \mathcal{O}, \mathcal{D}) = \int P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}) d\mathbf{p}.$$

It is noteworthy that similarly to the previous posterior derivations,

$$P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}) \propto P(\mathcal{O}, \mathcal{D} | \mathcal{T}, \mathbf{p}) P(\mathcal{T}, \mathbf{p}) = P(\mathcal{O}, \mathcal{D} | \mathcal{T}) P(\mathcal{T} | \mathbf{p}) P(\mathbf{p}),$$

that is, we now simply condition \mathcal{T} on \mathbf{p} (compare with the numerator in (2.3)). The integral in (3.1) can be hard to evaluate directly, but we can again resort to Monte Carlo methods to sample from $P(\mathcal{T} | \mathcal{O}, \mathcal{D})$ and conduct the inference, as we will see shortly.

Even though a hierarchical model increases complexity, it has two main advantages. First, we can now explain the uncertainty in trip pattern struc-

ture by specifying a suitable probability distribution for \mathbf{p} . This way, lack of information about trip pattern behaviors in the study region is reflected by more variability in the proportions, which, in turn, results in more dispersed trip pattern posterior distributions.

Secondly, we can better incorporate additional data that are related to the trip pattern structure. For instance, if there is available preliminary data \mathcal{T}_0 —usually from a small scale study in the same region or from a region with very similar structure—we can seamlessly incorporate it in the inference through the posterior $P(\mathcal{T} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0)$. This last posterior distribution can be obtained by adding the extra conditional on \mathcal{T}_0 in (3.1) and defining the likelihood $P(\mathcal{T}_0 | \mathbf{p})$ to derive

$$(3.2) \quad \begin{aligned} P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0) &\propto P(\mathcal{O}, \mathcal{D}, \mathcal{T}_0 | \mathcal{T}, \mathbf{p}) P(\mathcal{T}, \mathbf{p}) \\ &= P(\mathcal{O}, \mathcal{D} | \mathcal{T}) P(\mathcal{T}_0 | \mathbf{p}) P(\mathcal{T} | \mathbf{p}) P(\mathbf{p}). \end{aligned}$$

Note that we make the reasonable assumption that \mathcal{T} and \mathcal{T}_0 are conditionally independent given \mathbf{p} .

An alternative, common approach is to assume that the proportions \mathbf{p} are unknown, use \mathcal{T}_0 to estimate them, and then adopt the obtained estimate as if it were the “true” value of \mathbf{p} ; this approach is called *empirical Bayes* in the statistical literature, but is traditionally referred to as *calibration* in OD matrix estimation. Albeit being computationally simpler, this treatment has the drawback of underestimating variance, that is, it does not fully reflect the total uncertainty in the inference [10].

To better elucidate the proposed hierarchical models we present two applications next.

3.1. Incorporating seed matrices. A good candidate for the hyperprior distribution on \mathbf{p} is the multinomial conjugate distribution, the Dirichlet distribution, $\mathbf{p} \sim \text{Dir}(\boldsymbol{\pi})$, with mass function

$$P(\mathbf{p}) \propto \prod_{i,j} p_{ij}^{\pi_{ij}-1}.$$

We then have

$$\begin{aligned} P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}) &\propto \prod_{i,j} \frac{p_{ij}^{T_{ij}}}{T_{ij}!} \prod_{i,j} p_{ij}^{\pi_{ij}-1} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})] \\ &= \prod_{i,j} \frac{p_{ij}^{T_{ij}+\pi_{ij}-1}}{T_{ij}!} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})]. \end{aligned}$$

A non-informative prior on \mathbf{p} is attained by setting $\boldsymbol{\pi} = (1, \dots, 1)$ which is equivalent to \mathbf{p} having a uniform distribution over all $\{p_{ij}\} \in [0, 1]^{n^2}$ such that $\sum_{i,j} p_{ij} = 1$. In this case, the expression for $P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0)$ above is exactly the same as (2.3), but with the important distinction of now being a joint distribution since \mathbf{p} is random.

Suppose now that we have preliminary data $\mathcal{T}_0 = \{t_{ij}\}_{i,j=1,\dots,n}$ in the form of a seed matrix of trip counts. In the classical approach discussed in the introduction, \mathcal{T}_0 is commonly used to estimate the proportions as $\hat{p}_{ij} = t_{ij}/T_0$, where $T_0 = \sum_{k,l} t_{kl}$, or to simply kick-start an estimation procedure. This approach, however, effectively ignores the sample size T_0 since \hat{p}_{ij} remains the same if we observe κ times more counts, κT_0 , even for κ arbitrarily large; furthermore, similarly to empirical Bayes, it yields lower posterior variances for \mathcal{T} .

Following our discussion, here we offer a more principled way to incorporate the seed matrix \mathcal{T}_0 by performing posterior inference on \mathcal{T} through the distribution in (3.2). We assume that, similar to \mathcal{T} , the seed counts follow a conditional multinomial distribution, $\mathcal{T}_0 \sim \text{MN}(T_0, \mathbf{p})$ with flat prior $P(T_0) \propto 1$. Adopting the same Dirichlet distribution for \mathbf{p} we have

$$(3.3) \quad \begin{aligned} P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0) &\propto \prod_{i,j} \frac{p_{ij}^{T_{ij}}}{T_{ij}!} \prod_{i,j} \frac{p_{ij}^{t_{ij}}}{t_{ij}!} \prod_{i,j} p_{ij}^{\pi_{ij}-1} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})] \\ &\propto \prod_{i,j} \frac{p_{ij}^{T_{ij}+t_{ij}+\pi_{ij}-1}}{T_{ij}!} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})], \end{aligned}$$

and thus $\mathbf{p} | \mathcal{T}, \mathcal{T}_0 \sim \text{Dir}(\boldsymbol{\pi} + \mathcal{T} + \mathcal{T}_0)$.

To sample from $P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0)$ we adopt an extended Gibbs sampler with an extra step that accommodates the new hierarchical level: we iteratively sample from $P(\mathcal{T} | \mathbf{p}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0) = P(\mathcal{T} | \mathbf{p}, \mathcal{O}, \mathcal{D})$ exactly how we were doing in the previous section, and sample from the conditional Dirichlet $P(\mathbf{p} | \mathcal{T}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0) = P(\mathbf{p} | \mathcal{T}, \mathcal{T}_0)$. If a seed matrix is not available, the second step becomes simply sampling from $P(\mathbf{p} | \mathcal{T})$, still a Dirichlet distribution. The updated Gibbs sampler is listed below.

Step 1. Start at some arbitrary initial configuration $\mathcal{T}^{(0)}$ and initial proportions $\mathbf{p}^{(0)}$.

Step 2. For $t = 1, 2, \dots$ do (until convergence):

Step 2.1. For $i, j = 1, \dots, n-1$ do: sample $T_{ij}^{(t)} \sim T_{ij} | T_{[ij]}^{(t-1)}, \mathbf{p}^{(t-1)}, \mathcal{O}, \mathcal{D}$ from a non-central hypergeometric using a Metropolis step,

$$T_{ij}^{(t)} = MS(T_{ij}^{(t-1)}; O_{ij}^{(t-1)}, D_{ij}^{(t-1)}, \Delta_{ij}^{(t-1)}, \psi_{ij}^{(t-1)}),$$

with $O_{ij}^{(t-1)}$, $D_{ij}^{(t-1)}$, and $\Delta_{ij}^{(t-1)}$ as before, and

$$\psi_{ij}^{(t-1)} = (p_{ij}^{(t-1)} p_{nn}^{(t-1)}) / (p_{in}^{(t-1)} p_{nj}^{(t-1)}).$$

Step 2.2. Sample $\mathbf{p}^{(t)} \sim \text{Dir}(\mathcal{T}^{(t)} + \mathcal{T}_0 + \boldsymbol{\pi})$ or $\mathbf{p}^{(t)} \sim \text{Dir}(\mathcal{T}^{(t)} + \boldsymbol{\pi})$ if \mathcal{T}_0 is not available.

To perform inference on the marginal posterior $P(\mathcal{T} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0)$ we just need to use the realizations from the Gibbs sampler; the posterior mean, for instance, is readily available from (2.12). MAP estimates, however, are harder to obtain since we need to compute the integral in (3.1). One alternative is to use the joint posterior mode,

$$\tilde{\mathcal{T}} = \arg \max_{\mathcal{T} \in C(\mathcal{O}, \mathcal{D})} \left\{ \max_{\mathbf{p} \in [0,1]^{n^2} : \sum_{i,j} p_{ij} = 1} P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0) \right\},$$

but then the estimate might be biased since it is conditional on the optimal value of \mathbf{p} . In the same vein, we could first “calibrate” by setting some specific \mathbf{p} , say the marginal posterior mean

$$\bar{\mathbf{p}} = E[\mathbf{p} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0] \approx \frac{1}{G} \sum_{g=1}^G \mathbf{p}^{(g)},$$

and then produce

$$(3.4) \quad \hat{\mathcal{T}} = \arg \max_{\mathcal{T} \in C(\mathcal{O}, \mathcal{D})} P(\mathcal{T} | \bar{\mathbf{p}}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0).$$

It can be shown that the first estimator, $\tilde{\mathcal{T}}$, can be obtained by an extended Furness method that iteratively solves for \mathbf{p} while fitting the balancing factors by setting

$$\tilde{p}_{ij} = \frac{\tilde{T}_{ij} + t_{ij} + \pi_{ij} - 1}{\sum_{k,l=1,\dots,n} \tilde{T}_{kl} + t_{kl} + \pi_{kl} - 1},$$

but we will not pursue it further here.

3.2. Incorporating prior trip length distributions. Seed matrices provide information on each OD pair in the system and thus derive more accurate trip pattern inferences. More often than not, however, we do not have preliminary data \mathcal{T}_0 at this level of detail at our disposal. In some cases \mathcal{T}_0 contains censored observations; we might observe trips in a survey, but these trips are known only to have come from a certain origin, or to a destination, or

to have had some specific travel cost. For instance, recalling the trip length distribution (TLD) from Example 2, we might only discriminate a trip in our survey by specifying its cost “bin”, that is, within which range its cost falls.

Assume that we know the OD trip costs $\{c_{ij}\}$ and consider, as before, the K cost ranges $(c_0, c_1], \dots, (c_{K-1}, c_K]$. Our preliminary counts now fall into K possible strata, $\mathcal{T}_0 = \{t_1, \dots, t_K\}$, depending on their transport costs: we observe t_1 trips with costs between c_0 and c_1 , t_2 trips spending between c_1 and c_2 , and so on. If we again define range proportions aggregated by cost $\mathbf{p}_0 = \{p_k\}_{k=1, \dots, K}$, where $p_k = \sum_{i,j} p_{ij} I\{c_{ij} \in (c_{k-1}, c_k]\}$, we can then analogously set $\mathcal{T}_0 | \mathbf{p} \sim \text{MN}(\mathcal{T}_0, \mathbf{p}_0)$ with $P(\mathcal{T}_0) \propto 1$ as the preliminary data likelihood. We note that \mathbf{p}_0 is a function of \mathbf{p} .

We can assume the same Dirichlet distribution for the proportions, $\mathbf{p} \sim \text{Dir}(\boldsymbol{\pi})$, but since

$$P(\mathcal{T}, \mathbf{p} | \mathcal{O}, \mathcal{D}, \mathcal{T}_0) \propto \prod_{i,j} \frac{p_{ij}^{T_{ij}}}{T_{ij}!} \prod_k \frac{p_k^{t_k}}{t_k!} \prod_{i,j} p_{ij}^{\pi_{ij}-1} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})]$$

and each p_k is a sum of p_{ij} for all pairs i and j with cost in the k -th bin, we lose the conjugacy. Another approach, in case we are more informed about the censored proportions, is to opt for a Dirichlet prior on \mathbf{p}_0 ; but then we again lack conjugacy. Regardless, we can still obtain a Gibbs sampler that is very similar to the scheme shown in the previous subsection; we just need to substitute the direct Dirichlet sampling step, Step 2.2, by another Metropolis step. Next, we provide an updated sampling scheme in a simpler context.

Suppose that the proportions follow a gravity model with $p_{ij} \propto \exp(-\beta c_{ij})$, as in the previous section, but now we make β random to drive the uncertainty in \mathbf{p} . Moreover, we settle on a Dirichlet prior on \mathbf{p}_0 , $\mathbf{p}_0(\beta) \sim \text{Dir}(\boldsymbol{\pi})$, where $\boldsymbol{\pi} = \{\pi_1, \dots, \pi_K\}$. In what follows we explicitly represent the dependency of the proportions on β for clarity; we also note that now

$$p_k(\beta) \propto \sum_{i,j} \exp(-\beta c_{ij}) I\{c_{ij} \in (c_{k-1}, c_k]\}.$$

The joint posterior is thus given by

$$\begin{aligned}
 (3.5) \quad P(\mathcal{T}, \beta \mid \mathcal{O}, \mathcal{D}, \mathcal{T}_0) &\propto \prod_{i,j} \frac{p_{ij}(\beta)^{T_{ij}}}{T_{ij}!} \prod_k \frac{p_k(\beta)^{t_k}}{t_k!} \prod_k p_k(\beta)^{\pi_k-1} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})] \\
 &\propto \underbrace{\prod_{i,j} p_{ij}(\beta)^{T_{ij}} \prod_k p_k(\beta)^{t_k + \pi_k - 1}}_{\Phi(\beta; \mathcal{T}, \mathcal{T}_0)} I[\mathcal{T} \in C(\mathcal{O}, \mathcal{D})].
 \end{aligned}$$

From (3.5) we deduce that setting $\boldsymbol{\pi} = \{1, \dots, 1\}$ for a non-informative Dirichlet prior is equivalent to having a flat improper prior for the cost deterrence, $P(\beta) \propto 1$.

The Gibbs sampler has two iterative steps: we alternate between sampling from \mathcal{T} conditional on the impedance β and all the data, $P(\mathcal{T} \mid \beta, \mathcal{O}, \mathcal{D}, \mathcal{T}_0)$, and sampling from β conditional on trip patterns \mathcal{T} and margins and preliminary data, $P(\beta \mid \mathcal{T}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0)$. We already know, since Section 2, how to sample from $P(\mathcal{T} \mid \beta, \mathcal{O}, \mathcal{D}, \mathcal{T}_0) = P(\mathcal{T} \mid \mathbf{p}(\beta), \mathcal{O}, \mathcal{D})$ using random walk Metropolis steps for the non-central hypergeometric. To sample from $P(\beta \mid \mathcal{T}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0)$ we construct another random walk Metropolis step.

First, let us define the normalizing factors $Z_k(\beta) = \sum_{i,j} \exp(-\beta c_{ij}) I\{c_{ij} \in (c_{k-1}, c_k]\}$ and $Z(\beta) = \sum_{i,j} \exp(-\beta c_{ij}) = \sum_k Z_k(\beta)$, so that $p_{ij} = \exp(-\beta c_{ij})/Z(\beta)$ and $p_k = Z_k(\beta)/Z(\beta)$. Also, recall that $T = \sum_{i,j} T_{ij}$, $T_0 = \sum_k t_k$, and define $T_0^* = \sum_k (t_k + \pi_k - 1) = T_0 + \sum_k \pi_k - K$. The function $\Phi(\beta; \mathcal{T}, \mathcal{T}_0)$ in the joint posterior (3.5) then simplifies to

$$\begin{aligned}
 \Phi(\beta; \mathcal{T}, \mathcal{T}_0) &= \prod_{i,j} \left(\frac{\exp(-\beta c_{ij})}{Z(\beta)} \right)^{T_{ij}} \prod_k \left(\frac{Z_k(\beta)}{Z(\beta)} \right)^{t_k + \pi_k - 1} \\
 &= \exp \left\{ -\beta \sum_{i,j} c_{ij} T_{ij} + \sum_k (t_k + \pi_k - 1) \log Z_k(\beta) \right. \\
 &\quad \left. - (T + T_0^*) \log Z(\beta) \right\}.
 \end{aligned}$$

As proposal distribution, let us select a normal distribution centered at the current realization of β in the chain with small variance σ^2 . To get $\beta^{(t)}$ at the t -th iteration we then sample a candidate $\beta^* \sim N(\beta^{(t-1)}, \sigma^2)$ and accept or reject it based on the acceptance ratio

$$(3.6) \quad R(\beta^{(t-1)}, \beta^*) = \frac{P(\beta^* \mid \mathcal{T}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0)}{P(\beta^{(t)} \mid \mathcal{T}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0)} = \frac{\Phi(\beta^*; \mathcal{T}^{(t-1)}, \mathcal{T}_0^{(t-1)})}{\Phi(\beta^{(t-1)}; \mathcal{T}^{(t-1)}, \mathcal{T}_0^{(t-1)})}.$$

The final, updated Gibbs sampler is listed below.

Step 1. Start at some arbitrary initial configuration $\mathcal{T}^{(0)}$ and initial impedance $\beta^{(0)}$.

Step 2. For $t = 1, 2, \dots$ do (until convergence):

Step 2.1. For $i, j = 1, \dots, n-1$ do: sample $T_{ij}^{(t)} \sim T_{ij} | T_{[ij]}^{(t-1)}, \mathbf{p}(\beta^{(t-1)}), \mathcal{O}, \mathcal{D}$ from a non-central hypergeometric using a Metropolis step,

$$T_{ij}^{(t)} = MS(T_{ij}^{(t-1)}; O_{ij}^{(t-1)}, D_{ij}^{(t-1)}, \Delta_{ij}^{(t-1)}, \psi_{ij}(\beta^{(t-1)})).$$

with

$$\psi_{ij}(\beta^{(t-1)}) = \frac{p_{ij}(\beta^{(t-1)})p_{nn}(\beta^{(t-1)})}{p_{in}(\beta^{(t-1)})p_{nj}(\beta^{(t-1)})}.$$

Step 2.2. Sample candidate $\beta^* \sim N(\beta^{(t-1)}, \sigma^2)$ and set $\beta^{(t)} = \beta^*$ (accept) with probability $\min\{1, R(\beta^{(t-1)}, \beta^*)\}$ where $R(\cdot)$ is the ratio in (3.6); otherwise, set $\beta^{(t)} = \beta^{(t-1)}$ (reject.)

EXAMPLE 2, REVISITED. Under the same setting of Example 2, but now with β random, let us initially set $\boldsymbol{\pi} = \{1, \dots, 1\}$, that is, a non-informative prior on β . We run a Gibbs sampler with proposal variance $\sigma^2 = 10^{-4}$ until convergence and take $G = 10,000$ samples for posterior inference.

Our estimate for β , $\bar{\beta} = E[\beta | \mathcal{O}, \mathcal{D}] \approx \frac{1}{G} \sum_{g=1}^G \beta^{(g)} = 0.031$, is much lower than the assumed value in Example 2 ($\beta = 0.1$), which corroborates with our previous remark about a more realistic value for the cost impedance. Such lower values are expected since the inference is solely driven by the observed data and thus better represents the margin constraints. The estimated 95% credible interval for β is large, $[0.009, 0.056]$, reflecting the high degree of uncertainty that arises from trying to capture the structural trip proportions using a single parameter.

The effect of a random β in trip patterns can be appreciated in the estimated marginal posterior distributions for \mathcal{T} pictured in Figure 5. We draw attention to the increased spread when compared to the distributions in Figure 2. We also observe that the Furness solution, conditional on $\bar{\beta}$ and represented by squares, is similar to the posterior mean $E[\mathcal{T} | \mathcal{O}, \mathcal{D}]$.

The higher variability in \mathcal{T} is reproduced by wider credible intervals in the trip length distribution, as shown in Figure 6: each bar represents the estimated posterior mean of T_k/T for each cost range, the squares pinpoint the posterior mean of $p_k(\beta)$, while the dotted line corresponds to the prior mean $1/K$. As can be seen, the dependence of the proportions on a single parameter makes the distribution on \mathbf{p} not flexible enough to follow \mathcal{T} closely.




FIG 5. *Estimated marginal posterior distributions for T from hierarchical model with non-informative prior on β . Squares mark conditional Furness solution.*




FIG 6. *Mean posterior TLD (bars) with 95% credible intervals (whiskers), and mean posterior TLD proportions (squares). The dotted line marks the prior mean, $1/K$.*

We note again the higher variability in the posterior TLD as assessed by the wider 95% credible intervals (whiskers) when compared to Figure 4.

Suppose now that we observe preliminary data \mathcal{T}_0 from [12, pg. 186] in Table 4. Keeping the flat prior on β and $\sigma^2 = 10^{-4}$, we perform posterior inference from 10,000 samples taken from the Gibbs sampler after convergence.

TABLE 4
Preliminary TLD. Data reproduced from [12, table 5.14].

Range	(0, 4]	(4, 8]	(8, 12]	(12, 16]	(16, 20]	(20, 24]
t_k	365	962	160	150	230	95
t_k/T_0	0.19	0.49	0.08	0.08	0.12	0.05

The preliminary TLD counts are very informative, $T_0 = T = 1962$, and greatly affect the inference: our updated estimate for the cost deterrence is a higher $\bar{\beta} = E[\beta | \mathcal{O}, \mathcal{D}, \mathcal{T}_0] = 0.086$, closer to the original $\beta = 0.1$ in Example 2, and the 95% credible interval for β is much tighter, $[0.086, 0.093]$.

The posterior inference on trip patterns is summarized by Table 5, showing posterior mean $\bar{\mathcal{T}}$ and marginal 95% credible intervals, and Figure 7. The marginal distributions have increased variability when compared to Example 2 due to the randomness in the proportions, as expected. The variance is, however, not much higher since the preliminary TLD is very informative. The conditional Furness solution $\hat{\mathcal{T}}$, shown in square marks in Figure 7, is very similar to the posterior mean. The estimated posterior probabilities of these solutions are $P(\bar{\mathcal{T}} | \bar{\beta}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0) = 1.3 \cdot 10^{-3}$ and $P(\hat{\mathcal{T}} | \bar{\beta}, \mathcal{O}, \mathcal{D}, \mathcal{T}_0) = 1.5 \cdot 10^{-3}$, slightly smaller than in Example 2.

TABLE 5
Marginal posterior mean and 95% credible intervals.

Zone	1		2		3		4	
1	141.34	[128, 155]	101.49	[87, 118]	71.11	[57, 85]	86.07	[71, 103]
2	63.87	[52, 76]	184.96	[168, 204]	106.10	[89, 120]	105.07	[90, 122]
3	28.47	[20, 37]	51.32	[39, 63]	131.06	[116, 146]	189.14	[172, 205]
4	26.31	[17, 37]	62.23	[48, 77]	191.73	[174, 209]	421.72	[400, 444]

Since $\beta < 0.1$ with high posterior probability, we should expect the system to spend more when compared to the scenario in Example 2. Figure 8 displays the posterior distribution of trip costs $c(\mathcal{T})$, as estimated from (2.13). The posterior mean regional cost $c(\bar{\mathcal{T}}) = E[c(\mathcal{T}) | \mathcal{O}, \mathcal{D}, \mathcal{T}_0]$ is 9.12, with a 95% credible interval of $[8.81, 9.45]$, higher than before. The posterior mode cost $c(\hat{\mathcal{T}})$ is 9.09, close to $c(\bar{\mathcal{T}})$, as expected since the estimates are similar. The proportion cost $C_p(\beta) = \sum_{i,j} c_{ij} p_{ij}(\beta)$ in (2.5) inherits the randomness from β ; its estimated posterior mean, 8.95, is lower than $c(\bar{\mathcal{T}})$, which can




FIG 7. *Estimated marginal posterior distributions for \mathcal{T} from hierarchical model. Squares mark conditional Furness solution.*

also be attributed to the rigidity in \mathbf{p} .

Finally, we can also see the effect of \mathcal{T}_0 in reducing the inferential uncertainty in the posterior TLD at Figure 9, as illustrated by the tighter 95% credible intervals. We still see the discrepancy between the posterior TLD—whose mean $E[T_k/T \mid \mathcal{O}, \mathcal{D}, \mathcal{T}_0]$ is represented by bars—and the posterior proportion TLD—whose mean $E[p_k(\beta) \mid \mathcal{O}, \mathcal{D}, \mathcal{T}_0]$ is identified by squares. We note, however, that the posterior mean TLD is close to the prior mean TLD, t_k/T_0 , represented by diamonds and listed in Table 4, since \mathcal{T}_0 is highly informative and thus influential. The two mean posterior TLD are listed in Table 6.

TABLE 6
Posterior mean trip length distributions based on \mathcal{T} and \mathbf{p} .

Range	(0, 4]	(4, 8]	(8, 12]	(12, 16]	(16, 20]	(20, 24]
$E[T_k/T \mid \mathcal{O}, \mathcal{D}, \mathcal{T}_0]$	0.17	0.48	0.08	0.09	0.12	0.06
$E[p_k(\beta) \mid \mathcal{O}, \mathcal{D}, \mathcal{T}_0]$	0.24	0.36	0.12	0.14	0.10	0.04

4. Discussion. Static origin-destination matrix estimation has been traditionally regarded as an optimization problem. Here we draw from the contingency table literature and cast OD matrix estimation as a formal statistical inference problem and adopt a Bayesian approach where trip patterns are considered random. Furthermore, we make model assumptions on




FIG 8. *Estimated posterior distribution of mean regional cost. Solid line indicates posterior mean, dashed line marks posterior mean proportion cost, and dash-dotted line marks posterior mode cost.*




FIG 9. *Posterior mean TLD (bars) with 95% credible intervals (whiskers), posterior mean proportion TLD (squares), and prior mean TLD (diamonds).*

the parameters describing the probability distribution on trip patterns—trip proportions that govern the structure of trip distribution—as opposed to the classical assumptions on particular objective functions. The use of trip proportions frees us from requiring seemingly artificial constraints on trip configurations, provides more easily interpretable results, and allows us to better incorporate other sources of data in a principled way within a Bayesian framework.

By electing specific functional forms for the trip proportions—as based on the entropy maximizing principle, for example—we are able to recover classical solutions as MAP estimators and thus inherit the justifications and rich history behind traditional approaches. Yet, perhaps the main benefit of our proposed approach is to better characterize the uncertainty in the solutions and, in general, in trip distribution. As we have showed in many examples, it is common for any point estimate—such as the Furness solution or posterior mean—to capture only a small fraction of possible trip configurations given the large number of alternatives. Point estimators, when seen as ensemble summarizers, can be useful for preliminary planning purposes and gaining insight on the trip distribution in the study region; they can, however, be poor substitutes of the full posterior distribution in further analyses as they can dramatically underestimate the variability in trip patterns.

Preliminary data is traditionally used to calibrate specific parameters of the trip distribution model, such as cost deterrence. Nonetheless, fixing an optimal data fitting value for the parameter can further underestimate variance in the inference. In our fully Bayesian approach we explicitly acknowledge the uncertainty in the parameters by also making them random: we set a hyper-prior distribution on trip proportions to build a hierarchical model. As a consequence, and in contrast with a traditional approach, more informative preliminary data—for example, high counts in a seed matrix—yield more precise inference on trip configurations as we are able to more accurately characterize trip proportions.

The adoption of a Bayesian framework carries many other benefits not covered here: besides point and interval inference, we are also able to test hypotheses by explicitly comparing models through Bayes factors; moreover, Bayesian methods can be further explored to perform model validation through posterior predictive checks. In summary, the flexibility of Bayesian statistics is particularly helpful and really comes to bear when exploring high-dimensional spaces such as the ensemble of feasible trip configurations.

There is, however, a price to pay for such modeling power in higher computational costs, and thus the procedures discussed here still need to be more closely examined in this respect. Specifically, the increased complex-

ity in generating and analysing trip configuration samples instead of simply obtaining the most likely trip assignment needs to be assessed as the proposed routines are tried in real-world datasets comprising large systems. Future directions would also include the development of more efficient sampling schemes through improved algorithms—better proposal densities, for example—and faster implementations that would explore, for instance, parallel versions of the proposed procedures.

Finally, it should be noted that the models proposed here can serve as basis for an integrated higher level model that incorporates other traffic modeling steps; as an example, the effect of congested networks could be considered in OD matrix estimation if our model would jointly consider trip distribution and route assignment. As it is common in Bayesian modeling, we would then be able to propagate the uncertainty across steps while performing marginal inference on any aspect of the higher model conditional on data from all steps. Furthermore, other types of data could also be considered to obtain more refined models with, for instance, link count data and camera sensors or temporal variation for dynamic OD matrix estimation.

Acknowledgements. The author would like to thank Prof. Felipe Loureiro from Federal University of Ceará, Brazil, for many fruitful discussions and a constant source of motivation.

References.

- [1] Casey, H.J., 1955. Applications to traffic engineering of the law of retail gravitation. *Traffic Quarterly* IX, 23–35.
- [2] Diaconis, P., Gangolli, A., 1995. Rectangular arrays with fixed margins, in: Aldous, D., et al. (Eds.), *Discrete Probability and Algorithms*. Springer-Verlag, pp. 15–41.
- [3] Diaconis, P., Sturmfels, B., 1998. Algebraic algorithms for sampling from conditional distributions. *The Annals of Statistics* 26, 363–397.
- [4] Furness, K.P., 1965. Time function iteration. *Traffic Engineering and Control* 7, 458–460.
- [5] Gelman, A., Carlin, J.B., Stern, H.S., Rubin, D.B., 2003. *Bayesian Data Analysis*. Chapman and Hall. Second edition.
- [6] Geman, S., Geman, D., 1984. Stochastic relaxation, Gibbs distributions, and the Bayesian restoration of images. *IEEE Transactions on Pattern Analysis and Machine Intelligence* 6, 721–741.
- [7] Gilks, W., Richardson, S., Spiegelhalter, D.J., 1995. *Markov Chain Monte Carlo in Practice*. Chapman and Hall. First edition.
- [8] Givens, G.H., Hoeting, J.A., 2005. *Computational Statistics*. Wiley-Interscience. First edition.
- [9] Hastings, W., 1970. Monte carlo sampling methods using markov chains and their applications. *Biometrika* 57, 97–109.
- [10] Kass, R., Steffey, D., 1989. Approximate Bayesian inference in conditionally independent hierarchical models (parametric empirical Bayes models). *Journal of the American Statistical Association* 84, 717–726.

- [11] McCullagh, P., Nelder, J., 1989. Generalized Linear Models. Chapman and Hall. Second edition.
- [12] Ortúzar, J.D., Willusen, L.G., 2001. Modelling Transport. John Wiley & Sons, London. Third edition.
- [13] Wilson, A.G., 1970. Entropy in Urban and Regional Modelling. Pion, London.
- [14] Wilson, A.G., 1974. Urban and Regional Models in Geography and Planning. John Wiley & Sons, London.

DEPARTMENT OF MATHEMATICS AND STATISTICS
BOSTON UNIVERSITY
BOSTON, MASSACHUSETTS 02215
E-MAIL: lecarval@math.bu.edu