

VANISHING OF ALGEBRAIC BRAUER-MANIN OBSTRUCTIONS

MIKHAIL BOROVOI

ABSTRACT. Let X be a homogeneous space of a quasi-trivial k -group G , with geometric stabilizer \overline{H} , over a number field k . We prove that under certain conditions on the character group of \overline{H} , certain algebraic Brauer-Manin obstructions to the Hasse principle and weak approximation vanish, because the abelian groups where they take values vanish. When \overline{H} is connected or abelian, these algebraic Brauer-Manin obstructions to the Hasse principle and weak approximation are the only ones, so we prove the Hasse principle and weak approximation for X under certain conditions. As an application, we obtain new sufficient conditions for the Hasse principle and weak approximation for linear algebraic groups.

CONTENTS

1. Introduction: Sansuc's results	1
2. Introduction (continued): our main results	4
3. Preliminaries on Galois cohomology of finitely generated Galois modules	6
4. Preliminaries on quasi-trivial groups	8
5. Homogeneous spaces of quasi-trivial groups	9
6. Principal homogeneous spaces of connected groups	10
7. Connected groups as homogeneous spaces of quasi-trivial groups	10
8. Principal homogeneous spaces of connected groups again	12
References	13

1. INTRODUCTION: SANSUC'S RESULTS

We are inspired by Sansuc's paper [Sa]. In this section we state and discuss Sansuc's results on the Hasse principle and weak approximation for principal homogeneous spaces of connected linear algebraic groups admitting special coverings.

1.1. Let k be a field of characteristic 0 and \overline{k} be a fixed algebraic closure of k . If X is an algebraic variety over k , we write $\overline{X} = X \times_k \overline{k}$.

A k -torus T is called quasi-trivial if its character group $\hat{T} := \text{Hom}(\overline{T}, \mathbb{G}_{m, \overline{k}})$ is a permutation $\text{Gal}(\overline{k}/k)$ -module, i.e. \hat{T} has a \mathbb{Z} -basis which the Galois group permutes.

2000 *Mathematics Subject Classification.* Primary: 14F22, Secondary: 20G30, 14M17, 11G35, 14G25, 14C22.

Key words and phrases. Hasse principle, weak approximation, Brauer-Manin obstruction, Brauer group, linear algebraic groups, homogeneous spaces, algebraic fundamental group.

Partially supported by the Hermann Minkowski Center for Geometry and by the Israel Science Foundation (grant 807/07).

A *special covering* of a (connected) reductive k -group G is a short exact sequence

$$1 \rightarrow B \rightarrow G' \rightarrow G \rightarrow 1,$$

where G' is a product of a simply connected semisimple k -group and a quasi-trivial k -torus, and B is a finite central subgroup of G' .

Not all reductive groups admit special coverings. For example, a nonsplit one-dimensional k -torus does not admit such a covering.

A finite group Γ is called *metacyclic* if all its Sylow subgroups are cyclic. Any cyclic group is metacyclic. The group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ is not metacyclic. A finite Galois extension L/k is called metacyclic if $\text{Gal}(L/k)$ is a metacyclic group. Clearly any cyclic extension is metacyclic.

Let k be a field of characteristic 0 and M be a $\text{Gal}(\bar{k}/k)$ -module. We say that a field extension $K \subset \bar{k}$ of k *splits* M if $\text{Gal}(\bar{k}/K)$ acts trivially on M . If T is a k -torus, then K splits T if and only if K splits \hat{T} .

Now let k be a number field. We denote by \mathcal{V} the set of all places v of k , and by \mathcal{V}_∞ the set of infinite (archimedean) places. For $v \in \mathcal{V}$ we denote by k_v the completion of k at v .

1.2. Let G be a reductive k -group over a number field k admitting a special covering

$$1 \rightarrow B \rightarrow G' \rightarrow G \rightarrow 1.$$

Let \hat{B} denote the character group of B , i.e. $\hat{B} = \text{Hom}(B, \mathbb{G}_{m, \bar{k}})$. Let K be the smallest subfield of \bar{k} splitting \hat{B} (i.e. K is the subfield corresponding to the subgroup $\ker [\text{Gal}(\bar{k}/k) \rightarrow \text{Aut}(\hat{B})]$). Let X be a principal homogeneous space of G .

Sansuc [Sa] proved the following results:

Proposition 1.3 ([Sa], Cor. 5.2). *Let k , G , B , K , and X be as in 1.2. If K/k is a metacyclic extension, then X satisfies the Hasse principle and weak approximation, i.e. if $X(k_v) \neq \emptyset$ for any place v of k , then $X(k) \neq \emptyset$ and, moreover, $X(k)$ is dense in $\prod_{v \in \mathcal{V}} X(k_v)$.*

Remark 1.4. If the extension K/k is not metacyclic (e.g. $\text{Gal}(K/k) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$), then the Hasse principle or weak approximation may fail for X , see [Se, §III.4.7] and [Sa, Examples 5.6, 5.7, and 5.8].

Proposition 1.5 ([Sa], Thm. 3.3 and Lemma 1.6). *Let k , G , B , K , and X be as in 1.2. Assume that X has a k -point (hence we may identify X with G). Let $S \subset \mathcal{V}$ be a finite set formed by places of k with cyclic decomposition groups in K/k (for example, assume that K/k is unramified at all finite places in S). Then X has the weak approximation property in S , i.e. the set $X(k)$ is dense in $\prod_{v \in S} X(k_v)$.*

1.6. The results of Propositions 1.3 and 1.5 can be explained in terms of the Brauer group of X . Let X be a smooth k -variety. We write $\text{Br}(X)$ for the cohomological Brauer group of X , i.e. $\text{Br}(X) = H_{\text{ét}}^2(X, \mathbb{G}_m)$. We set $\text{Br}_1(X) = \ker[\text{Br}(X) \rightarrow \text{Br}(\bar{X})]$. We define the *algebraic Brauer group* $\text{Br}_a(X)$ by $\text{Br}_a(X) = \text{coker}[\text{Br}(k) \rightarrow \text{Br}_1(X)]$.

When k is a number field and $S \subset \mathcal{V}$ is a finite set of places of k , we set

$$\mathbb{B}_S(X) = \ker \left[\text{Br}_a(X) \rightarrow \prod_{v \notin S} \text{Br}_a(X_{k_v}) \right].$$

Set $\mathbb{B}_\omega(X) = \bigcup_S \mathbb{B}_S(X)$. We set $\mathbb{B}(X) := \mathbb{B}_\emptyset(X)$ and $\mathbb{B}_{S, \emptyset}(X) = \mathbb{B}_S(X) / \mathbb{B}_\emptyset(X) = \mathbb{B}_S(X) / \mathbb{B}(X)$.

Sansuc [Sa, (6.2.3)], following Manin [M], defined a natural pairing

$$\prod_{v \in \mathcal{V}} X(k_v) \times \mathbb{B}_\omega(X) \rightarrow \mathbb{Q}/\mathbb{Z}$$

(see also [Sk, (5.2)]), which is continuous in the first argument and is additive in the second one. This pairing induces a continuous map

$$(1) \quad m: \prod_{v \in \mathcal{V}} X(k_v) \rightarrow \text{Hom}(\mathbb{E}_\omega(X), \mathbb{Q}/\mathbb{Z}).$$

If $x_0 \in X(k) \subset \prod_{v \in \mathcal{V}} X(k_v)$, then $m(x_0) = 0$. If m is not identically 0, say, $m(x_{\mathcal{V}}) \neq 0$ for some $x_{\mathcal{V}} = (x_v)_{v \in \mathcal{V}} \in \prod_{v \in \mathcal{V}} X(k_v)$, then $x_{\mathcal{V}}$ is not contained in the closure of $X(k)$, weak approximation fails for X . We say that m is *the algebraic Brauer-Manin obstruction to the Hasse principle and weak approximation for X associated with \mathbb{E}_ω* .

Assume that X is a smooth k -variety having a k -point. Let $S \subset \mathcal{V}$ be a finite set of places of k . Inspired by [CTS] and [Sa], we defined in [B3, §1] a natural pairing

$$\prod_{v \in S} X(k_v) \times \mathbb{E}_{S, \emptyset}(X) \rightarrow \mathbb{Q}/\mathbb{Z}.$$

This pairing induces a continuous map

$$(2) \quad m_S: \prod_{v \in S} X(k_v) \rightarrow \text{Hom}(\mathbb{E}_{S, \emptyset}(X), \mathbb{Q}/\mathbb{Z}).$$

If $x_0 \in X(k) \subset \prod_{v \in S} X(k_v)$, then $m_S(x_0) = 0$. If m_S is not identically 0, say $m_S(x_S) \neq 0$ for some $x_S \in \prod_{v \in S} X(k_v)$, then x_S is not contained in the closure of $X(k)$, hence weak approximation in S fails for X . We say that m_S is *the algebraic Brauer-Manin obstruction to weak approximation in S for X associated with $\mathbb{E}_{S, \emptyset}(X)$* .

Using Sansuc's methods and results, one can show that under the assumptions of Proposition 1.3 we have $\mathbb{E}_\omega = 0$, hence $m = 0$, see Proposition 1.7(ii) below. Moreover, the Hasse principle and weak approximation hold for X because there is no Brauer-Manin obstruction. Similarly, under the assumptions of Proposition 1.5 we have $\mathbb{E}_{S, \emptyset} = 0$, hence $m_S = 0$, see Proposition 1.7(i) below. Again, weak approximation in S holds for X because there is no Brauer-Manin obstruction. We provide some details.

Proposition 1.7. *Let k , G , B , K , and X be as in 1.2. Let $S \subset \mathcal{V}$ be a finite set of places of k .*

- (i) *If any place $v \in S$ has a cyclic decomposition group in K/k , then $\mathbb{E}_{S, \emptyset}(X) = 0$.*
- (ii) *If K/k is a metacyclic extension, then $\mathbb{E}_\omega(X) = 0$.*

Proof. We use the notation of §3.1 below. By [Sa, Lemma 6.8] $\text{Br}_a(X) \cong \text{Br}_a(G)$, hence $\mathbb{E}_\omega(X) \cong \mathbb{E}_\omega(G)$ and $\mathbb{E}_{S, \emptyset}(X) \cong \mathbb{E}_{S, \emptyset}(G)$. By [Sa, Prop. 9.8] $\mathbb{E}_\omega(G) \cong \text{III}_\omega^1(k, \widehat{B})$ and $\mathbb{E}(G) \cong \text{III}^1(k, \widehat{B})$. One proves similarly that $\mathbb{E}_S(G) \cong \text{III}_S^1(k, \widehat{B})$, hence $\mathbb{E}_{S, \emptyset}(G) \cong \text{III}_{S, \emptyset}^1(k, \widehat{B}) := \text{III}_S^1(k, \widehat{B}) / \text{III}_\emptyset^1(k, \widehat{B})$. In case (i), since S is formed by places of k with cyclic decomposition groups in K/k , by [Sa, Lemma 1.1(iii)] (see also Lemma 3.2(iii) below) $\text{III}_{S, \emptyset}^1(k, \widehat{B}) = 0$, hence $\mathbb{E}_{S, \emptyset}(X) = 0$. In case (ii), since K/k is metacyclic, by [Sa, Lemma 1.3] (see also Lemma 3.4 below) $\text{III}_\omega^1(k, \widehat{B}) = 0$, hence $\mathbb{E}_\omega(X) = 0$. \square

An alternative proof of Proposition 1.3. By Proposition 1.7(ii) the Brauer-Manin obstruction m of formula (1) is identically zero in our case, i.e. there is no algebraic Brauer-Manin obstruction to the Hasse principle and weak approximation associated with \mathbb{E}_ω . Since by [Sa, Cor. 8.7 and Cor. 8.13] this obstruction is the only one, we conclude that the Hasse principle and weak approximation hold for X . \square

An alternative proof of Proposition 1.5. By Proposition 1.7(i) the Brauer-Manin obstruction m_S of formula (2) is identically zero in our case, i.e. there is no algebraic Brauer-Manin obstruction to weak approximation in S associated with $\mathbb{E}_{S, \emptyset}(X)$. Since by [Sa, Cor. 8.13] this obstruction is the only one, we conclude that weak approximation in S holds for X . \square

2. INTRODUCTION (CONTINUED): OUR MAIN RESULTS

In this section we state our generalizations of Sansuc's results in two cases: homogeneous spaces of quasi-trivial groups and principal homogeneous spaces of connected linear algebraic groups. Our main results are Theorems 2.1 and 2.7, generalizing Proposition 1.7 and proving that the groups $\mathcal{B}_{S,\emptyset}(X)$ and $\mathcal{B}_\omega(X)$ vanish under certain conditions on X .

In order to state our results we use the notion of a quasi-trivial group, introduced by Colliot-Thélène [CT, Definition 2.1], see also Definition 4.2 below.

Let X be a homogeneous space of a quasi-trivial k -group G over a number field k . Let $\overline{H} \subset \overline{G}$ be the stabilizer of a \overline{k} -point $\overline{x} \in X(\overline{k})$ (we do not assume that \overline{H} is connected or abelian). It is well known that the character group $\widehat{\overline{H}}$ of \overline{H} has a canonical structure of a Galois module, see [B3, 4.1] or [BvH2, Rem. 5.7(1)], see also §8.1 below.

Theorem 2.1. *Let X be a left homogeneous space of a quasi-trivial k -group G over a number field k . Let $\overline{H} \subset \overline{G}$ be the stabilizer of a \overline{k} -point $\overline{x} \in X(\overline{k})$. Let $S \subset \mathcal{V}$ be a finite subset. Let K/k be the smallest Galois extension in \overline{k} splitting the Galois module $\widehat{\overline{H}}$.*

- (i) *If any place $v \in S$ has a cyclic decomposition group in K/k , then $\mathcal{B}_{S,\emptyset}(X) = 0$.*
- (ii) *If K/k is a metacyclic extension, then $\mathcal{B}_\omega(X) = 0$.*

Theorem 2.1 will be proved in Section 5.

Corollary 2.2. *Let X be as in Theorem 2.1. Assume that X has a k -point, i.e. $X = G/H$, where G is a quasi-trivial k -group and $H \subset G$ is a k -subgroup. Assume that H is connected or abelian. Let $S \subset \mathcal{V}$ be a finite set of places of k formed by places with cyclic decomposition groups in K/k (for example assume that K/k is unramified at all finite places in S). Then X has weak approximation in S .*

Proof. Since H is connected or abelian, the algebraic Brauer-Manin obstruction m_S associated with $\mathcal{B}_{S,\emptyset}$ is the only obstruction to weak approximation in S for X , see [B3, Thm. 2.3]. By Theorem 2.1(i) $m_S = 0$, hence X has weak approximation in S . \square

Corollary 2.3. *Let X be as in Corollary 2.2, i.e. $X = G/H$, where G is a quasi-trivial k -group and $H \subset G$ is a k -subgroup. Assume that H is connected or abelian. Then X has the real approximation property, i.e. $X(k)$ is dense in $\prod_{v \in \mathcal{V}_\infty} X(k_v)$.*

Proof. For any $v \in \mathcal{V}_\infty$ the decomposition group of v in K/k is cyclic, and by Corollary 2.2 X has weak approximation in \mathcal{V}_∞ , i.e. real approximation. \square

Question 2.4. Does there exist a homogeneous space $X = G/H$, where G is a quasi-trivial k -group over a number field k , and $H \subset G$ is a nonconnected non-abelian k -subgroup, such that real approximation fails for X ?

Corollary 2.5. *Let X be a homogeneous space having a k -rational point with connected stabilizer, of a connected linear algebraic group (not necessarily quasi-trivial) over a number field k ; in other words, $X = G/H$, where G is a connected k -group and $H \subset G$ is a connected k -subgroup. Then X has the real approximation property.*

Proof. By Lemma 4.3 below we can write $X = G'/H'$, where G' is a quasi-trivial k -group and $H' \subset G'$ is a connected k -subgroup. Now by Corollary 2.3 X has real approximation. \square

Corollary 2.6. *Let X be as in Theorem 2.1. Assume that \overline{H} is connected or abelian. Assume that K/k is a metacyclic extension. Then X satisfies the Hasse principle and weak approximation.*

Proof. Since \overline{H} is connected or abelian, the algebraic Brauer-Manin obstruction m associated with \mathcal{B}_ω is the only obstruction to the Hasse principle and weak approximation, see [B3, Thms. 2.2

and 2.3]. By Theorem 2.1(ii) we have $m = 0$, hence X satisfies the Hasse principle and weak approximation. \square

Note that Propositions 1.3 and 1.5 (due to Sansuc) follow immediately from our Corollaries 2.6 and 2.2, respectively. Note also that the special case of Corollary 2.2 when G is simply connected and H is connected was earlier proved in [B1, Cor. 1.6] by a different method.

In order to state our results on principal homogeneous spaces of connected k -groups, we use the notion of the *algebraic fundamental group* $\pi_1(G)$ introduced in [B4, §1] (where we wrote $\pi_1(\overline{G})$ instead of $\pi_1(G)$), see also [CT, §6]. Note that $\pi_1(G)$ is a finitely generated Galois module.

Theorem 2.7. *Let G be a connected linear k -group over a number field k . Let X be a principal homogeneous space (torsor) of G over k . Let $S \subset \mathcal{V}$ be a finite set of places of k . Let K/k be the smallest Galois extension in \overline{k} splitting the Galois module $\pi_1(G)$.*

- (i) *If any place $v \in S$ has a cyclic decomposition group in K/k , then $\mathbb{B}_{S,\emptyset}(X) = 0$.*
- (ii) *If K/k is a metacyclic extension, then $\mathbb{B}_\omega(X) = 0$.*

Theorem 2.7 will be proved in Section 6.

Proposition 2.8. *Let G be a connected linear algebraic group over a field k of characteristic 0. Let K/k be the smallest Galois extension in \overline{k} splitting $\pi_1(G)$. Then there exists an exact sequence*

$$1 \rightarrow H \rightarrow G' \rightarrow G \rightarrow 1,$$

where G' is a quasi-trivial k -group and H is a central k -subgroup of multiplicative type, such that K splits both \widehat{H} and \widehat{G}' .

Proposition 2.8 will be proved in Section 7. In Section 8 we shall give an alternative proof of Theorem 2.7 based on Proposition 2.8 and Theorem 2.1.

Corollary 2.9. *Let k , G , and K be as in Theorem 2.7. Let $S \subset \mathcal{V}$ be a finite set of places of k formed by places with cyclic decomposition groups in K/k (for example assume that K/k is unramified at all finite places in S). Then G has weak approximation in S .*

Proof. Under our assumptions the algebraic Brauer-Manin obstruction m_S associated with $\mathbb{B}_{S,\emptyset}(X)$ is the only obstruction to weak approximation in S for G , see [Sa, Cor. 8.13]. By Theorem 2.7(i) $m_S = 0$, hence G has weak approximation in S . Alternatively, the corollary follows from Proposition 2.8 and Corollary 2.2. \square

Corollary 2.10. *Let k , G , X , and K be as in Theorem 2.7. Assume that K/k is a metacyclic extension. Then X satisfies the Hasse principle and weak approximation.*

Proof. Under our assumptions the algebraic Brauer-Manin obstruction m associated with \mathbb{B}_ω is the only obstruction to the Hasse principle and weak approximation for X , see [Sa, Cor. 8.7 and Cor. 8.13]. By Theorem 2.7(ii) we have $m = 0$, hence X satisfies the Hasse principle and weak approximation. Alternatively, the corollary follows from Proposition 2.8 and Corollary 2.6 (because K splits \widehat{H} , where \overline{H} is the stabilizer of \overline{x} in \overline{G}' , see §8.2 below). \square

Remark 2.11. Sansuc proved in [Sa, Cor. 3.5(iii)] that any connected k -group over a number field k has the real approximation property. This follows from our Corollary 2.9 (because infinite places have cyclic decomposition groups in $\text{Gal}(K/k)$) and from our Corollary 2.5 (because we may write $G = G/\{1\}$, and $\{1\}$ is a connected k -subgroup).

Note that Sansuc proved the following result:

Proposition 2.12 (Sansuc). *Let G be a connected linear k -group over a number field k . Let $S \subset \mathcal{V}$ be a finite subset. Assume that G splits over a finite Galois extension K/k .*

(i) *If any place $v \in S$ has a cyclic decomposition group in K/k , then G has weak approximation in S (cf. [Sa, Cor. 3.5(ii)]).*

(ii) *If K/k is a metacyclic extension, then $\mathbb{B}_\omega(X) = 0$ (cf. [Sa, Prop. 9.8]), hence any principal homogeneous space X of G over k satisfies the Hasse principle and weak approximation.*

Proposition 2.12 follows from our Theorem 2.7: if a finite Galois extension K/k splits G , then it splits $\pi_1(G)$. The following example shows that Theorem 2.7 is indeed stronger than Proposition 2.12.

Example 2.13. Let k be a number field, and let K_1 and K_2 be two different quadratic extensions of k in \bar{k} . Let K be the composite of K_1 and K_2 , then K/k is a Galois extension with non-metacyclic Galois group $\text{Gal}(K/k) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. Set $G_1 = \text{SU}_{2,K_1}$, it is a k -group. Set

$$G_2 = R_{K_2/k}^1 \mathbb{G}_m := \ker[N_{K_2/k} : R_{K_2/k} \mathbb{G}_{m,K_2} \rightarrow \mathbb{G}_{m,k}],$$

where $N_{K_2/k}$ is the norm map. Set $\mu = \{(-1, -1), (1, 1)\} \subset G_1 \times G_2$, and set $G = (G_1 \times G_2)/\mu$.

Clearly G does not admit a special covering. Let L/k be any finite Galois extension in \bar{k} splitting G . Then L splits both G_1 and G_2 , hence $L \supset K$, and therefore L/k is not metacyclic. We see that we cannot prove the Hasse principle and weak approximation for a principal homogeneous space X of G using Proposition 1.3 or Proposition 2.12.

On the other hand, $\pi_1(G) \cong \mathbb{Z}$ as an abelian group, and $\text{Gal}(\bar{k}/k)$ acts on $\pi_1(G)$ via $\text{Gal}(K_2/k)$. We see that the cyclic Galois extension K_2/k splits $\pi_1(G)$. By Theorem 2.7(ii) $\mathbb{B}_\omega(X) = 0$, and by Corollary 2.10 X satisfies the Hasse principle and weak approximation.

The plan of the rest of this paper is as follows. In Sections 3 we give preliminaries on Galois cohomology of finitely generated Galois modules. In Section 4 we give preliminaries on quasi-trivial groups and on the algebraic fundamental group of a connected linear algebraic group. In Sections 5 and 6 we prove Theorems 2.1 and 2.7, respectively. In Section 7 we prove Proposition 2.8, and in Section 8 we use this proposition in order to give an alternative proof of Theorem 2.7. Our proofs are based on the results of Section 3 and of our papers [BvH1] and [BvH2].

3. PRELIMINARIES ON GALOIS COHOMOLOGY OF FINITELY GENERATED GALOIS MODULES

3.1. In this section k denotes a number field, and B is a discrete $\text{Gal}(\bar{k}/k)$ -module which is finitely generated as an abelian group (we say just “a finitely generated Galois module”). By S we always denote a finite subset of \mathcal{V} . We write

$$\text{III}_S^i(k, B) = \ker \left[H^i(k, B) \rightarrow \prod_{v \notin S} H^i(k_v, B) \right].$$

We have $\text{III}_\emptyset^i(k, B) = \text{III}^i(k, B)$. We set $\text{III}_{S, \emptyset}^i(k, B) := \text{III}_S^i(k, B) / \text{III}_\emptyset^i(k, B)$ and $\text{III}_\omega^i(k, B) = \bigcup_S \text{III}_S^i(k, B)$.

Lemmas 3.2 and 3.4 below are immediate generalizations of Sansuc’s lemmas from [Sa, §1a]. The only difference is that Sansuc assumes that B is a *finite* Galois module, while we assume that B is a finitely generated Galois module. Since Sansuc’s proofs are concise, we give detailed proofs here.

Lemma 3.2. *Let K/k be a finite Galois extension with Galois group \mathfrak{g} and S be a finite set of places of k .*

(i) *If B is a constant $\text{Gal}(\bar{k}/k)$ -module (i.e. $\text{Gal}(\bar{k}/k)$ acts trivially), then $\text{III}_S^1(K/k, B) = 0$ and $\text{III}_S^1(k, B) = 0$.*

(ii) If the extension K/k trivializes B , there is a reduction

$$\text{III}_S^1(k, B) = \text{III}_S^1(K/k, B).$$

(iii) If S' is a finite subset of \mathcal{V} formed of places with cyclic decomposition groups in K/k , then

$$\text{III}_{S \cup S'}^1(K/k, B) = \text{III}_S^1(K/k, B).$$

Proof. We follow closely [Sa, Proof of Lemma 1.1].

We prove (i). We write $\mathfrak{g} = \text{Gal}(K/k)$. Since by assumption the finite group \mathfrak{g} acts on B trivially, the group $\text{III}_S^1(K/k, B)$ is the kernel of the homomorphism

$$(3) \quad \text{Hom}_{\text{gr}}(\mathfrak{g}, B) \longrightarrow \prod_{v \notin S} \text{Hom}_{\text{gr}}(\mathfrak{g}^v, B).$$

where Hom_{gr} denotes the group of group homomorphisms, and \mathfrak{g}^v denotes the decomposition group of a place w of K prolonging v . By Chebotarev's density theorem, any element γ of \mathfrak{g} is conjugate to an element of some \mathfrak{g}^v with $v \notin S$ (because the set S is finite). It follows that the homomorphism (3) is injective, and hence, $\text{III}_S^1(K/k, B) = 0$.

Now let $c \in H^1(k, B) = \text{Hom}_{\text{c.gr}}(\text{Gal}(\bar{k}/k), B)$, where $\text{Hom}_{\text{c.gr}}$ denotes the group of continuous homomorphisms of groups. Since B is discrete and $\text{Gal}(\bar{k}/k)$ is profinite, we see that $\ker c$ is an open normal subgroup of $\text{Gal}(\bar{k}/k)$, and we may write $\ker c = \text{Gal}(\bar{k}/K)$, where K is a finite Galois extension of k contained in \bar{k} . Let c_K denote the natural homomorphism $\text{Gal}(K/k) \rightarrow B$; then c is the inflation $\text{Inf } c_K$. Let $v \in \mathcal{V} \setminus S$ and let w be a prolongation of v to K . Let $c_{K,v}$ denote the image of c_K in $H^1(K_w/k_v, B) = \text{Hom}(\text{Gal}(K_w/k_v), B)$ and c_v denote the image of c in $H^1(k_v, B) = \text{Hom}(\text{Gal}(\bar{k}_v/k_v), B)$. By assumption $c_v = 0$. Since $c_v = \text{Inf } c_{K,v}$ and $\text{Gal}(\bar{k}_v/k_v)$ acts on B trivially, we see that $c_{K,v} = 0$. Thus $c_K \in \text{III}_S^1(K/k, B)$. Since $\text{III}_S^1(K/k, B) = 0$, see above, we conclude that $c_K = 0$, and hence, $c = 0$. Thus $\text{III}_S^1(k, B) = 0$, as required.

We prove (ii). We assume that the extension K/k trivializes B . Let S' denote the set of the prolongations to K of the places in S . Then by (i) we have $\text{III}_{S'}^1(K, B) = 0$. We consider the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^1(K/k, B) & \longrightarrow & H^1(k, B) & \longrightarrow & H^1(K, B) \\ & & \alpha \downarrow & & \beta \downarrow & & \downarrow \varphi \\ 0 & \longrightarrow & \prod_{v \notin S} H^1(K_w/k_v, B) & \longrightarrow & \prod_{v \notin S} H^1(k_v, B) & \longrightarrow & \prod_{w \notin S'} H^1(K_w, B) \end{array}$$

obtained from inflation-restriction exact sequences, after choosing, for each place v of k , a prolongation w of v from k to K . Since $\text{III}_{S'}^1(K, B) = 0$, the homomorphism φ in the diagram is injective. It follows that $\text{III}_S^1(K/k, B) = \ker \alpha \cong \ker \beta = \text{III}_S^1(k, B)$, as required.

Chebotarev's density theorem implies that for each place v with cyclic decompositions group \mathfrak{g}^v (defined up to conjugacy in \mathfrak{g}), there exists infinitely many other places v' with the same decomposition group (again, up to conjugacy), which proves (iii). \square

Consider the homomorphism $\text{Gal}(\bar{k}/k) \rightarrow \text{Aut}(B)$. The image \mathfrak{g} of this homomorphism is finite. Let K denote the subfield in \bar{k} corresponding to the kernel of this homomorphism, then K/k is a finite Galois extension with Galois group \mathfrak{g} . We say that K/k be the smallest Galois extension in \bar{k} splitting B .

Corollary 3.3. *Let K/k be the smallest Galois extension in \bar{k} splitting B , and let $S \subset \mathcal{V}$ be a finite set of places of k . If all places $v \in S$ have cyclic decomposition groups in $\text{Gal}(K/k)$, then $\text{III}_{S, \emptyset}^1(k, B) = 0$.* \square

Proof. By Lemma 3.2(ii), $\text{III}_S^1(k, B) = \text{III}_S^1(K/k, B)$ and $\text{III}_{\emptyset}^1(k, B) = \text{III}_{\emptyset}^1(K/k, B)$. By Lemma 3.2(iii) $\text{III}_S^1(K/k, B) = \text{III}_{\emptyset}^1(K/k, B)$. Thus $\text{III}_S^1(k, B) = \text{III}_{\emptyset}^1(k, B)$ and $\text{III}_{S, \emptyset}^1(k, B) = 0$. \square

Recall that the exponent of a finite group is the least common multiple of the orders of its elements. A finite group is metacyclic if and only if its exponent is equal to its order.

Lemma 3.4. *Let K/k be the smallest Galois extension in \bar{k} splitting B , and let n and e be the order and the exponent of $\mathfrak{g} = \text{Gal}(K/k)$, respectively. Then multiplication by n/e equals 0 in $\text{III}_\omega^1(k, B)$. In particular, if K/k is a metacyclic extension, then*

$$\text{III}_\omega^1(k, B) = 0, \text{ and hence, } \text{III}^1(k, B) = 0.$$

The same holds if B is finite and its order is relatively prime to n/e . \square

Proof. We follow closely [Sa, Proofs of Lemmas 1.2 and 1.3].

First, we show that the group $\text{III}_\omega^1(k, B)$ is a finite group, namely,

$$(4) \quad \text{III}_\omega^1(k, B) = \ker \left[H^1(\mathfrak{g}, B) \rightarrow \prod_{g \in \mathfrak{g}} H^1(\langle g \rangle, B) \right],$$

where \mathfrak{g} denotes the Galois group of the smallest Galois extension K/k in \bar{k} splitting B , and $\langle g \rangle$ denotes the subgroup of \mathfrak{g} generated by g . We can write

$$(5) \quad \text{III}_\omega^1(k, B) = \text{III}_{S_0}^1(K/k, B),$$

where S_0 denotes the finite subset of \mathcal{V} consisting of the places (ramified in K) whose decomposition groups in K/k are noncyclic.

Indeed, it follows from the definition of K and Lemma 3.2(ii), that $\text{III}_S^1(k, B) = \text{III}_S^1(K/k, B)$. Further, by Lemma 3.2(iii), $\text{III}_S^1(K/k, B) = \text{III}_{S \cap S_0}^1(K/k, B)$. This proves that the group $\text{III}_\omega^1(k, B)$ is finite, and gives (5). The formula (4) follows from (5), because by the construction of S_0 , for any $v \notin S_0$ we have $\mathfrak{g}^v = \langle g \rangle$ for some $g \in \mathfrak{g}$, and by Chebotarev's density theorem any cyclic subgroup $\langle g \rangle \subseteq \mathfrak{g}$ is conjugate to a decomposition group \mathfrak{g}^v .

Now let g be an element of \mathfrak{g} , and let n_g denote its order. The map

$$\text{Cor} \circ \text{Res}: H^1(\mathfrak{g}, B) \rightarrow H^1(\langle g \rangle, B) \rightarrow H^1(\mathfrak{g}, B)$$

is multiplication by n/n_g ; see [AW, Section 6, Proposition 8]. By (4), the restriction of this map to $\text{III}_\omega^1(k, B)$ is 0, which proves the first assertion of the lemma, from which the other assertions follow immediately. \square

4. PRELIMINARIES ON QUASI-TRIVIAL GROUPS

4.1. Let k be a field of characteristic 0, \bar{k} a fixed algebraic closure of k . Let G be a connected linear k -group. We set $\overline{G} = G \times_k \bar{k}$. We use the following notation:

G^u is the unipotent radical of G ;

$G^{\text{red}} = G/G^u$ (it is reductive);

G^{ss} is the derived group of G^{red} (it is semisimple);

G^{sc} is the universal cover of G^{ss} (it is simply connected);

$G^{\text{tor}} = G^{\text{red}}/G^{\text{ss}}$ (it is a torus);

Definition 4.2 ([CT], Prop. 2.2). A connected linear k -group G over a field k of characteristic 0 is called *quasi-trivial*, if G^{tor} is a quasi-trivial torus and G^{ss} is simply connected.

Lemma 4.3 (well known). *Let k be a field of characteristic 0 and let X be a left homogeneous space of a connected linear k -group G . Let $\overline{H} \subset \overline{G}$ be the stabilizer of a point $\bar{x} \in X(\bar{k})$; we assume that \overline{H} is connected. Then the variety X is a homogeneous space of some quasi-trivial k -group G' such that the stabilizer $\overline{H}' \subset \overline{G}'$ of \bar{x} in \overline{G}' is connected.*

Proof. The lemma follows e.g. from [CT, Prop.-Def. 3.1]. \square

5. HOMOGENEOUS SPACES OF QUASI-TRIVIAL GROUPS

In this section we prove Theorem 2.1.

5.1. Let k be a field of characteristic 0, and let $A \rightarrow B$ be a morphism of $\text{Gal}(\bar{k}/k)$ -modules. We write $\mathbb{H}^i(k, A \rightarrow B)$ for the Galois hypercohomology of the complex $A \rightarrow B$, where A is in degree 0 and B is in degree 1.

When k is a number field, we define $\mathbb{III}_S^i(k, A \rightarrow B)$, $\mathbb{III}_{S, \emptyset}^i(k, A \rightarrow B)$, and $\mathbb{III}_\omega^i(k, A \rightarrow B)$ as in §3.

The following lemma must be well known (see [B3, Proof of Lemma 4.4] and [BvH1, Proof of Cor. 2.15] for similar results) but we do not know a reference where it was stated, so we state and prove it here.

Lemma 5.2. *Let k be a number field and $P \rightarrow L$ a complex of $\text{Gal}(\bar{k}/k)$ -modules in degrees 0 and 1, where P is a permutation $\text{Gal}(\bar{k}/k)$ -module. Then for any finite set S of places of k we have a canonical isomorphism $\mathbb{III}_S^1(k, L) \xrightarrow{\sim} \mathbb{III}_S^2(k, P \rightarrow L)$.*

Proof. We have an exact sequence

$$0 = H^1(k, P) \rightarrow H^1(k, L) \rightarrow \mathbb{H}^2(k, P \rightarrow L) \rightarrow H^2(k, P),$$

and similar exact sequences for Galois cohomology over k_v for $v \notin S$. We obtain a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^1(k, L) & \longrightarrow & \mathbb{H}^2(k, P \rightarrow L) & \longrightarrow & H^2(k, P) \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \prod_{v \notin S} H^1(k_v, L) & \longrightarrow & \prod_{v \notin S} \mathbb{H}^2(k_v, P \rightarrow L) & \longrightarrow & \prod_{v \notin S} H^2(k_v, P). \end{array}$$

Since P is a permutation module, we have $\mathbb{III}_S^2(k, P) = 0$, cf. [Sa, (1.9.1)]. An easy diagram chase shows that the homomorphism $\mathbb{III}_S^1(k, L) \rightarrow \mathbb{III}_S^2(k, P \rightarrow L)$ induced by this diagram is an isomorphism. \square

Proposition 5.3. *Let X be a homogeneous space of a quasi-trivial k -group G over a number field k . Let $\bar{H} \subset \bar{G}$ be the stabilizer of a \bar{k} -point $\bar{x} \in X(\bar{k})$. Let $S \subset \mathcal{V}$ be a finite set of places of k . Then there is a canonical isomorphism $\mathbb{B}_S(X) \cong \mathbb{III}_S^1(k, \widehat{\bar{H}})$.*

Proof. By [BvH2, Thm. 7.2] we have a canonical, functorial in k isomorphism

$$\text{Br}_a(X) \xrightarrow{\sim} \mathbb{H}^2(k, \widehat{G} \rightarrow \widehat{\bar{H}}),$$

whence we obtain a canonical isomorphism

$$\mathbb{B}_S(X) \cong \mathbb{III}_S^2(k, \widehat{G} \rightarrow \widehat{\bar{H}}).$$

Since \widehat{G} is a permutation module, by Lemma 5.2 we have a canonical isomorphism

$$\mathbb{III}_S^1(k, \widehat{\bar{H}}) \xrightarrow{\sim} \mathbb{III}_S^2(k, \widehat{G} \rightarrow \widehat{\bar{H}}).$$

Thus we obtain a canonical isomorphism $\mathbb{B}_S(X) \cong \mathbb{III}_S^1(k, \widehat{\bar{H}})$. \square

5.4. Proof of Theorem 2.1. By Proposition 5.3 we have a canonical isomorphism $\mathbb{B}_S(X) \cong \mathbb{III}_S^1(k, \widehat{\bar{H}})$, hence we obtain a canonical isomorphism $\mathbb{B}_{S, \emptyset}(X) \cong \mathbb{III}_{S, \emptyset}^1(k, \widehat{\bar{H}})$ and a canonical isomorphism $\mathbb{B}_\omega(X) \cong \mathbb{III}_\omega^1(k, \widehat{\bar{H}})$. In case (i) by Corollary 3.3 we have $\mathbb{III}_{S, \emptyset}^1(k, \widehat{\bar{H}}) = 0$, hence $\mathbb{B}_{S, \emptyset}(X) = 0$. In case (ii) by Lemma 3.4 we have $\mathbb{III}_\omega^1(k, \widehat{\bar{H}}) = 0$, hence $\mathbb{B}_\omega(X) = 0$. \square

6. PRINCIPAL HOMOGENEOUS SPACES OF CONNECTED GROUPS

In this section we prove Theorem 2.7.

6.1. Let M be a $\text{Gal}(\bar{k}/k)$ -module, finitely generated over \mathbb{Z} . Choose a \mathbb{Z} -free resolution

$$(6) \quad 0 \rightarrow L \rightarrow P \rightarrow M \rightarrow 0,$$

where L and P are finitely generated \mathbb{Z} -free Galois modules. We write

$$\mathbb{H}^i(k, M^D) := \mathbb{H}^i(k, P^\vee \rightarrow L^\vee),$$

where $P^\vee := \text{Hom}_{\mathbb{Z}}(P, \mathbb{Z})$ is in degree 0 and $L^\vee := \text{Hom}_{\mathbb{Z}}(L, \mathbb{Z})$ is in degree 1. We regard $M^D := (P^\vee \rightarrow L^\vee)$ as a dual complex to M . Since the isomorphism class of M^D in the derived category does not depend on the choice of the resolution (6), the hypercohomology $\mathbb{H}^i(k, M^D)$ also does not depend on the resolution.

Lemma 6.2. *Let M be as in 6.1. Let K/k be the smallest Galois extension in \bar{k} splitting M . Let $S \subset \mathcal{V}$ be finite set of places of k .*

(i) *If any place $v \in S$ has a cyclic decomposition group in $\text{Gal}(K/k)$, then $\mathbb{III}_{S, \emptyset}^2(k, M^D) = 0$.*

(ii) *If K/k is a metacyclic extension, then $\mathbb{III}_{\omega}^2(k, M^D) = 0$.*

Proof. Set $\mathfrak{g} = \text{Gal}(K/k)$, then M is a \mathfrak{g} -module. We can choose a resolution (6) such that P is a permutation \mathfrak{g} -module and L is a \mathbb{Z} -free \mathfrak{g} -module. Then P^\vee is a permutation module as well, and by Lemma 5.2 we have a canonical isomorphism

$$\mathbb{III}_S^1(k, L^\vee) \xrightarrow{\sim} \mathbb{III}_S^2(k, P^\vee \rightarrow L^\vee) = \mathbb{III}_S^2(k, M^D),$$

whence we obtain canonical isomorphisms

$$\begin{aligned} \mathbb{III}_{S, \emptyset}^1(k, L^\vee) &\xrightarrow{\sim} \mathbb{III}_{S, \emptyset}^2(k, M^D), \\ \mathbb{III}_{\omega}^1(k, L^\vee) &\xrightarrow{\sim} \mathbb{III}_{\omega}^2(k, M^D). \end{aligned}$$

Since K splits L^\vee , in case (i) by Corollary 3.3 we have $\mathbb{III}_{S, \emptyset}^1(k, L^\vee) = 0$, hence $\mathbb{III}_{S, \emptyset}^2(k, M^D) = 0$. In case (ii) by Lemma 3.4 we have $\mathbb{III}_{\omega}^1(k, L^\vee) = 0$, hence $\mathbb{III}_{\omega}^2(k, M^D) = 0$. \square

6.3. Proof of Theorem 2.7. By [Sa, Lemma 6.8] there is a canonical isomorphism $\text{Br}_a(X) \xrightarrow{\sim} \text{Br}_a(G)$. By [BvH1, Cor. 7] there is a canonical isomorphism $\text{Br}_a(G) \xrightarrow{\sim} \mathbb{H}^2(k, \pi_1(G)^D)$. Hence $\mathbb{B}_S(X) \cong \mathbb{III}_S^2(k, \pi(G)^D)$, whence $\mathbb{B}_{S, \emptyset}(X) \cong \mathbb{III}_{S, \emptyset}^2(k, \pi_1(G)^D)$ and $\mathbb{B}_{\omega}(X) \cong \mathbb{III}_{\omega}^2(k, \pi_1(G)^D)$. In case (i) by Lemma 6.2(i) we have $\mathbb{III}_{S, \emptyset}^2(k, \pi_1(G)^D) = 0$, hence $\mathbb{B}_{S, \emptyset}(X) = 0$. In case (ii) by Lemma 6.2(ii) we have $\mathbb{III}_{\omega}^2(k, \pi_1(G)^D) = 0$, hence $\mathbb{B}_{\omega}(X) = 0$. \square

7. CONNECTED GROUPS AS HOMOGENEOUS SPACES OF QUASI-TRIVIAL GROUPS

In this section we prove Proposition 2.8.

7.1. Proof of Proposition 2.8. We may and shall assume that G is reductive, cf. proof of [CT, Prop.-Def. 3.1]. Consider the biggest quotient torus G^{tor} of G . Set $M = \pi_1(G)$, then $\mathbb{X}_*(G^{\text{tor}}) = M/M_{\text{tors}}$, where M_{tors} denotes the torsion subgroup of M . Since K splits M , we see that K splits $\mathbb{X}_*(G^{\text{tor}})$.

We follow the construction in the proof of [CT, Prop.-Def. 3.1]. Let Z^0 denote the radical (the identity component of the center) of our reductive group G . Since Z^0 is isogeneous to G^{tor} , we see that K splits $\mathbb{X}_*(Z^0)$. Set $\mathfrak{g} = \text{Gal}(K/k)$, then $\mathbb{X}_*(Z^0)$ is a \mathfrak{g} -module. Choose a surjective homomorphism of \mathfrak{g} -modules $P \twoheadrightarrow \mathbb{X}_*(Z^0)$, where P is a finitely generated permutation \mathfrak{g} -module. We regard P as a $\text{Gal}(\bar{k}/k)$ -module, then K splits P . Let Q be the quasi-trivial k -torus with $\mathbb{X}_*(Q) = P$. We have a surjective homomorphism $\theta: Q \twoheadrightarrow Z^0$.

Consider the canonical homomorphism

$$\rho: G^{\text{sc}} \rightarrow G^{\text{ss}} \hookrightarrow G.$$

Set $G' = G^{\text{sc}} \times_k Q$, then G' is a quasi-trivial group and K splits $\widehat{G}' = P^\vee$. We define a surjective homomorphism

$$\alpha: G' \rightarrow G, \quad \alpha(g, q) = \rho(g)\theta(q), \text{ where } g \in G^{\text{sc}}, q \in Q.$$

Set $H = \ker \alpha$. Note that H is a central subgroup of multiplicative type of G' . We have an exact sequence

$$(7) \quad 1 \rightarrow H \rightarrow G' \rightarrow G \rightarrow 1.$$

Set $M' := \pi_1(G') = P$, then K splits M and M' . By Lemma 7.2 below we have a canonical isomorphism of Galois modules $\widehat{H} \cong \text{Ext}_{\mathbb{Z}}^0(M' \rightarrow M, \mathbb{Z})$ (where M' is in degree 0). It follows that K splits \widehat{H} , which proves the proposition. \square

Lemma 7.2. *Assume we have a short exact sequence of Galois modules*

$$1 \rightarrow H \rightarrow G' \xrightarrow{\varphi} G \rightarrow 1,$$

where G and G' are connected reductive k -groups over a field k of characteristic 0, and $H \subset G'$ is a central k -subgroup of multiplicative type. Set $M := \pi_1(G)$, $M' := \pi_1(G')$. Then there is a canonical isomorphism

$$\mathbb{X}^*(H) \cong \text{Ext}_{\mathbb{Z}}^0(M' \rightarrow M, \mathbb{Z}),$$

where in the complex $M' \rightarrow M$, the Galois module M' is in degree 0 and M is in degree 1, and we write $\mathbb{X}^*(H) := \widehat{H}$.

Proof (C. Demarche). Note that the homomorphism $G' \rightarrow G$ induces an isomorphism $G'^{\text{sc}} \rightarrow G^{\text{sc}}$. Choose compatible maximal tori $T_G \subset G$, $T_{G'} \subset G'$, $T_{G^{\text{sc}}} \subset G^{\text{sc}}$ and $T_{G'^{\text{sc}}} \subset G'^{\text{sc}}$. It follows from the definition of M and M' that we have a commutative diagram with exact rows

$$(8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{X}_*(T_{G'^{\text{sc}}}) & \longrightarrow & \mathbb{X}_*(T_{G'}) & \longrightarrow & M' \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathbb{X}_*(T_{G^{\text{sc}}}) & \longrightarrow & \mathbb{X}_*(T_G) & \longrightarrow & M \longrightarrow 0. \end{array}$$

Since the homomorphism $\varphi_*: G'^{\text{sc}} \rightarrow G^{\text{sc}}$ is an isomorphism, the left-hand vertical arrow $\mathbb{X}_*(T_{G'^{\text{sc}}}) \rightarrow \mathbb{X}_*(T_{G^{\text{sc}}})$ in diagram (8) is an isomorphism, and the five lemma shows that the morphism of complexes of Galois modules

$$(\mathbb{X}_*(T_{G'}) \rightarrow \mathbb{X}_*(T_G)) \longrightarrow (M' \rightarrow M)$$

given by this diagram is a quasi-isomorphism.

The short exact sequence of complexes

$$0 \rightarrow (0 \rightarrow \mathbb{X}_*(T_G)) \rightarrow (\mathbb{X}_*(T_{G'}) \rightarrow \mathbb{X}_*(T_G)) \rightarrow (\mathbb{X}_*(T_{G'}) \rightarrow 0) \rightarrow 0$$

induces an exact sequence of Ext-groups

$$\text{Hom}_{\mathbb{Z}}(\mathbb{X}_*(T_G), \mathbb{Z}) \rightarrow \text{Hom}_{\mathbb{Z}}(\mathbb{X}_*(T_{G'}), \mathbb{Z}) \rightarrow \text{Ext}_{\mathbb{Z}}^0(\mathbb{X}_*(T_{G'}) \rightarrow \mathbb{X}_*(T_G), \mathbb{Z}) \rightarrow \text{Ext}_{\mathbb{Z}}^1(\mathbb{X}_*(T_G), \mathbb{Z}) = 0.$$

Since the complex $\mathbb{X}_*(T_{G'}) \rightarrow \mathbb{X}_*(T_G)$ is quasi-isomorphic to $M' \rightarrow M$, we obtain an exact sequence

$$\text{Hom}_{\mathbb{Z}}(\mathbb{X}_*(T_G), \mathbb{Z}) \rightarrow \text{Hom}_{\mathbb{Z}}(\mathbb{X}_*(T_{G'}), \mathbb{Z}) \rightarrow \text{Ext}_{\mathbb{Z}}^0(M' \rightarrow M, \mathbb{Z}) \rightarrow 0,$$

which we can write as

$$(9) \quad \mathbb{X}^*(T_G) \rightarrow \mathbb{X}^*(T_{G'}) \rightarrow \text{Ext}_{\mathbb{Z}}^0(M' \rightarrow M, \mathbb{Z}) \rightarrow 0,$$

where $\mathbb{X}^*(T_G) = \widehat{T}_G$ and $\mathbb{X}^*(T_{G'}) = \widehat{T}_{G'}$. On the other hand, the exact sequence of k -groups of multiplicative type

$$1 \rightarrow H \rightarrow T_{G'} \rightarrow T_G \rightarrow 1$$

gives an exact sequence

$$(10) \quad 0 \rightarrow \mathbb{X}^*(T_G) \rightarrow \mathbb{X}^*(T_{G'}) \rightarrow \mathbb{X}^*(H) \rightarrow 0.$$

Comparing exact sequences (9) and (10), we obtain a canonical isomorphism of Galois modules

$$\mathbb{X}^*(H) \cong \text{Ext}_{\mathbb{Z}}^0(M' \rightarrow M, \mathbb{Z}). \quad \square$$

Remark 7.3. Constructing and arguing as in the proof of [CT, Prop.-Def. 3.1], we can construct an exact sequence (7) with G' a quasi-trivial k -group and H a flasque k -torus (and not just some k -group of multiplicative type) such that the smallest Galois extension K/k in \bar{k} splitting $\pi_1(G)$ splits both tori G'^{tor} and H . This strengthens Remark 3.1.1 of [CT].

8. PRINCIPAL HOMOGENEOUS SPACES OF CONNECTED GROUPS AGAIN

In this section we give an alternative proof of Theorem 2.7 based on Proposition 2.8 and Theorem 2.1.

8.1. Let X be a homogeneous space of a quasi-trivial k -group G over a number field k . Let $\bar{H} \subset \bar{G}$ be the stabilizer of a \bar{k} -point $\bar{x} \in X(\bar{k})$. We describe the action of $\text{Gal}(\bar{k}/k)$ on $\widehat{\bar{H}}$ defined by the homogeneous space X .

Let $h \in \bar{H}(\bar{k})$, then $\bar{x} \cdot h = \bar{x}$. Let $\sigma \in \text{Gal}(\bar{k}/k)$, then $\sigma_x \cdot \sigma h = \sigma_x$. For any $\sigma \in \text{Gal}(\bar{k}/k)$ we choose $g_\sigma \in G(\bar{k})$ such that $\sigma_x = x \cdot g_\sigma$ and the function $\sigma \mapsto g_\sigma$ is locally constant, then

$$g_\sigma \cdot \sigma h \cdot g_\sigma^{-1} \in H(\bar{k}).$$

The map $h \mapsto g_\sigma \cdot \sigma h \cdot g_\sigma^{-1}$ comes from some σ -semialgebraic automorphism (see [B2, §1.1] for a definition) v_σ of \bar{H} , which induces an automorphism \widehat{v}_σ of $\widehat{\bar{H}}$ (namely, $\widehat{v}_\sigma(\chi)(h) = \chi(v_\sigma^{-1}(h))$ for $\chi \in \widehat{\bar{H}}$ and $h \in \bar{H}(\bar{k})$). If we choose another element $g'_\sigma \in G(\bar{k})$ such that $\sigma_x = x \cdot g'_\sigma$, then $g'_\sigma = h' g_\sigma$ for some $h' \in \bar{H}(\bar{k})$. Then we obtain $v'_\sigma = \text{Inn}(h') \circ v_\sigma$, where $\text{Inn}(h')$ is the inner automorphism of \bar{H} defined by h' . We have $\widehat{v}'_\sigma = \widehat{v}_\sigma$, because $\text{Inn}(h')$ acts trivially on $\widehat{\bar{H}}$. The well-defined map $\sigma \mapsto \widehat{v}_\sigma$ is a homomorphism defining an action of $\text{Gal}(\bar{k}/k)$ on $\widehat{\bar{H}}$.

8.2. Alternative proof of Theorem 2.7. We deduce Theorem 2.7 from Proposition 2.8 and Theorem 2.1. Since K splits $\pi_1(G)$, by Proposition 2.8 we can write $G = G'/H$, where G' is a quasi-trivial k -group and H is a central k -subgroup of multiplicative type in G' such that K splits \widehat{H} .

The group G' acts on X via G . Let $\bar{x} \in X(\bar{k})$, then the stabilizer of \bar{x} in \bar{G}' is $\bar{H} := H_{\bar{k}}$. Consider the action $\sigma \mapsto \widehat{v}_\sigma$ of $\text{Gal}(\bar{k}/k)$ on $\widehat{\bar{H}}$ defined in 8.1. Write $\sigma_x = x \cdot g'_\sigma$, where $g'_\sigma \in G'(\bar{k})$, then

$$v_\sigma(h) = g'_\sigma \cdot \sigma h \cdot (g'_\sigma)^{-1} = \sigma h$$

for $h \in H(\bar{k})$, because H is central in G' . It follows that the action of $\text{Gal}(\bar{k}/k)$ on $\widehat{\bar{H}}$ defined by the homogeneous space X coincides with the action on \widehat{H} defined by the k -structure of H .

Now, since K splits \widehat{H} , we see that K splits $\widehat{\bar{H}}$, and Theorem 2.7 follows from Theorem 2.1. \square

Acknowledgements. We thank J.-L. Colliot-Thélène for helpful e-mail correspondence, C. Demarche for proving Lemma 7.2, and the anonymous referee for a quick and thorough review. The author finished this paper while visiting the Max-Planck-Institut für Mathematik, Bonn; he thanks the Institute for hospitality, support, and excellent working conditions.

REFERENCES

- [AW] M. F. Atiyah and C. T. C. Wall, *Cohomology of groups*, Algebraic Number Theory (J.W.S. Cassels and A. Fröhlich, eds.), Academic Press/Thompson Book Co., Inc., London/Washington, D.C., 1967, pp. 94–115.
- [B1] M. Borovoi, *On weak approximation in homogeneous spaces of simply connected algebraic groups*, in: Automorphic functions and their applications (Khabarovsk, 1988), 64–81, Acad. Sci. USSR, Inst. Appl. Math., Khabarovsk, 1990.
- [B2] M. Borovoi, *Abelianization of the second nonabelian Galois cohomology*, Duke Math. J. **72** (1993), 217–239.
- [B3] M. Borovoi, *The Brauer-Manin obstructions for homogeneous spaces with connected or abelian stabilizer*, J. reine angew. Math. **473** (1996), 181–194.
- [B4] M. Borovoi, *Abelian Galois cohomology of reductive groups*, Mem. Amer. Math. Soc. **132** (1998), no. 626.
- [BvH1] M. Borovoi and J. van Hamel, *Extended Picard complexes and linear algebraic groups*, J. reine angew. Math. **627** (2009), 53–82.
- [BvH2] M. Borovoi and J. van Hamel, *Extended equivariant Picard complexes and homogeneous spaces*, Preprint, arXiv:1010.3414v2[math.AG].
- [CT] J.-L. Colliot-Thélène, *Résolutions flasques des groupes linéaires connexes*, J. reine angew. Math. **618** (2008), 77–133.
- [CTS] J.-L. Colliot-Thélène et J.-J. Sansuc, *La descente sur une variété rationnelle définie sur un corps de nombres*, C. R. A. S. Paris **284** (1977), A1215–1218.
- [M] Yu. I. Manin, *Le groupe de Brauer-Grothendieck en géométrie diophantienne*, in: Actes du Congrès Intern. Math. (Nice, 1970), Tome 1, Gauthier-Villars, Paris, 1971, pp. 401–411.
- [Sa] J.-J. Sansuc, *Groupe de Brauer et arithmétique des groupes algébriques linéaires sur un corps de nombres*, J. reine angew. Math. **327** (1981), 12–80.
- [Se] J.-P. Serre, *Cohomologie galoisienne*, Lecture Notes in Math. **5**, 5ème ed., Springer-Verlag, Berlin 1994.
- [Sk] A. N. Skorobogatov, *Torsors and Rational Points*, Cambridge Tracts in Mathematics **144**, Cambridge University Press, Cambridge 2001.

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, 69978
TEL AVIV, ISRAEL

E-mail address: borovoi@post.tau.ac.il