

Most attractive channel for self-breaking of grand unification group symmetry

Nobuhiro Uekusa

*Department of Physics, Osaka University
Toyonaka, Osaka 560-0043 Japan
E-mail: uekusa@het.phys.sci.osaka-u.ac.jp*

Abstract

Most attractive channel is studied for $SU(5)$ grand unified theory in five dimensions where $\mathbf{5}^*$ and $\mathbf{10}$ fermions and a $\mathbf{24}$ gauge boson propagate in the bulk. If there are bulk fermions with zero mode for the chirality opposite to quark and lepton multiplets, they can contribute to forming $\mathbf{5}$, $\mathbf{24}$, $\mathbf{50}$ and $\mathbf{75}$ scalar bound states. We find that the inclusion of a simple anomaly-free set for zero mode yields scalar bound states whose binding strengths are the largest for $\mathbf{24}$ and the second largest for $\mathbf{5}$. This corresponds to the breaking pattern of the $SU(5)$ to color and electromagnetism via the standard model gauge group.

1 Introduction

While elementary particles have mass and charge, the origin still remains unknown. When the generation of masses is related to symmetry breaking, the corresponding problem is how symmetry is broken. In the standard model, the Higgs mechanism describes physics of elementary particles very well up to the weak scale. However, it has been found in the own framework of the standard model that this picture is not valid at higher energy scales. In addition, the standard model does not have any explanation for quantization of charge. Hence, we need a deeper understanding for mass and charge, or symmetry breaking and charge beyond the standard model.

If extra dimensions are included in an effective theory, the strength of force would behave unlike four dimensions. The gauge bosons of the standard model propagating in extra dimensions can rapidly become strongly coupled and form scalar bound states of quarks and leptons. The self-breaking of the standard-model gauge symmetry was proposed in Ref. [1]. The authors proposed that the existence of a Higgs doublet is a consequence of the standard-model gauge symmetry and three generation of quarks and leptons provided the gauge bosons and fermions propagate in appropriate extra dimensions compactified at a TeV scale. It has been shown earlier that electroweak symmetry may be broken by fields propagating in extra dimensions in Ref. [2][3]. Also in the Randall-Sundrum warped space [4][5], transition of the strength of force and the type of bound states have been studied [6]-[11].

As for quantization of charge, grand unification has attracted much attention. The charges belong to subgroups of a unification group. Quantum numbers for quarks and leptons are fixed by the group structure. The $SU(3)$, $SU(2)$ and $U(1)$ couplings meet at a single point and the couplings are evaded from blowup of the values by renormalization group evolution. Beyond the unification scale, the energy dependence of the three couplings are shown graphically as one identical line due to contributions of twelve X and Y bosons. Thus, a grand unification with extra dimensions might contain the generation of masses and the quantization of charges automatically.

In this paper, we study most attractive channel for an $SU(5)$ grand unification in five dimensions. Our setup is that components in each $SU(5)$ multiplet obey the same boundary condition with respect to extra dimensions although gauge symmetry breaking by boundary conditions might be an interesting possibility [12] [13]. The energy dependence of couplings in gauge theories with extra dimensions has been found [14] [15] and has been examined in detail [16] [17]. In particular, the energy dependence of the three couplings should be shown as one identical line beyond the unification scale. It has been emphasized in Ref. [18] that this is not the case for gauge symmetry breaking by boundary conditions where the same boundary conditions do not span irreducible representations of the unification group. In order to keep gauge coupling unification above the unification scale, we consider only the self-breaking for the origin of symmetry breaking. The fundamental fields are a gauge boson and fermions. A gauge boson belongs to **24**. Quarks and leptons are included in **5*** and **10** Dirac fermions whose left-handed components have zero mode. If these fields are only objects propagating in the bulk, there are no scalar bound states composed of zero mode. When an anomaly-free set of fields with zero mode for the right-handed component is added, we find scalar bound states whose binding strengths are the largest for **24** and the second largest for **5**, correspondingly to the breaking pattern $SU(5) \rightarrow SU(3) \times SU(2) \times U(1) \rightarrow SU(3) \times U(1)$.

The paper is organized as follows. In Section 2, we start with the basics such as the scenario, framework, field content and group structure. With consideration for anomaly-free sets, scalar bound states and the binding strengths are given in Section 3. Here we find various composites. The simple model to form **24** and **5** for appropriate symmetry breaking is proposed. Furthermore, the composites **50** and **75** are discussed for the doublet-triplet splitting. We conclude in Section 4.

2 Basics of the model

We consider a gauge theory with two branes on fixed points of the orbifold S^1/Z_2 in five dimensions. Bulk fields can have strong coupling compared with the corresponding four-dimensional theory so that the ordinary four-dimensional massless gluon and quarks does not give rise to gauge symmetry breaking and that the five-dimensional effect may change potentials for composites. If non-vanishing vacuum expectation values are generated, symmetry is broken. The pattern of symmetry breaking depends on the attractive force of composites and the masses of the constituents. The analysis can be applied for both of flat spacetime and warped spacetime.

In the present model, there are no fundamental scalar fields. The gauge boson A_M has the four-dimensional vector component A_μ and the four-dimensional scalar component A_5 . The boundary conditions at the location of the branes are given by Neumann for all A_μ and Dirichlet for all A_5 . Quarks and leptons are included as zero mode for **5*** and **10** Dirac fermions, ψ_{5*} and ψ_{10} . For the left-handed component, the Neumann boundary condition is imposed. Since singlet neutrinos do not produce attractive force, they can be omitted in the present analysis. For these boundary conditions, fields which belong to irreducible representations of $SU(5)$ obey the same boundary conditions so that at higher energy scales above the scale of symmetry breaking by any condensation the theory behave as $SU(5)$. For simplicity, the number of the generation is chosen as one.

The source of symmetry breaking is scalar bound states represented by the form $\bar{\psi}_I \psi_J$ where I, J are denoted as the species of fermions. The attractive force is dominantly originated from exchange of a gauge boson between fermions and is determined by the gauge coupling and the group structure. As for the group structure, we take the notation $\text{tr}(t_r^a t_r^b) = C(r) \delta^{ab}$ where t_r^a is denoted as the representation matrices in the irreducible representation r and $C(r)$ is a constant for each representation r . The relation $d(r)C_2(r) = d(\text{Adj})C(r)$ is fulfilled where $C_2(r)$ is the quadratic Casimir operator for each representation and d is the dimension of each representation. The values C , C_2 and d are summarized for several irreducible representations of $SU(N)$ in Table 1. For $N = 5$, the diagonal generator $t^8 = \text{diag}(2, 2, 2, -3, -3)/(2\sqrt{15})$ plays the same role as the $U(1)$ hypercharge up to a overall factor. We call its charge Q_Y . For $SU(3) \times SU(2) \times U(1)$, ψ_{5*} and ψ_{10} are decomposed into $(\mathbf{3}^*, \mathbf{1})_{Q_Y=-\sqrt{15}/15} \oplus (\mathbf{1}, \mathbf{2})_{Q_Y=\sqrt{15}/10}$ and $(\mathbf{3}, \mathbf{2})_{Q_Y=-\sqrt{15}/30} \oplus (\mathbf{3}^*, \mathbf{1})_{Q_Y=2\sqrt{15}/15} \oplus (\mathbf{1}, \mathbf{1})_{Q_Y=-\sqrt{15}/5}$, respectively.

For the quark and lepton multiplets, possible combinations of the bound states are $\bar{\psi}_{5*} \psi_{5*}$, $\bar{\psi}_{10} \psi_{10}$ and $\bar{\psi}_{5*} \psi_{10}$. They are written in terms of four-dimensional fields dependent on the five-dimensional coordinates as chirality mixing operators $\bar{\psi}_L \psi_R + \bar{\psi}_R \psi_L$. Because ψ_R consists of Kaluza-Klein massive mode without zero mode, it is difficult to have potentials for bound states to yield nonzero vacuum expectation values. Hence, we need to add new fields with zero mode for the right-handed component. In order that ψ_{5*} and

Table 1: $SU(N)$ property. Each box in Young tableaux is symmetric for the horizontal direction and antisymmetric for the vertical direction.

	d	C	C_2
\square	N	$\frac{1}{2}$	$\frac{N^2-1}{2N}$
$\begin{array}{ c } \hline \square \\ \hline \end{array}$	$\frac{N(N-1)}{2}$	$\frac{N-2}{2}$	$\frac{(N+1)(N-2)}{N}$
$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$\frac{N(N+1)}{2}$	$\frac{N+2}{2}$	$\frac{(N-1)(N+2)}{N}$
$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$\frac{N(N-1)(N-2)}{6}$	$\frac{(N-2)(N-3)}{4}$	$\frac{3(N-3)(N+1)}{2N}$
$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$\frac{N(N^2-1)}{6}$	$\frac{(N^2-3)}{2}$	$\frac{3(N^2-3)}{2N}$
$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$\frac{N(N+1)(N+2)}{6}$	$\frac{(N+2)(N+3)}{4}$	$\frac{3(N-1)(N+3)}{2N}$
Adj	$N^2 - 1$	N	N

ψ_{10} do not form brane mass terms together with the new fields, candidates are except for **5*** and **10**. A singlet is also excluded because it does not contribute to attractive force.

For forming the fundamental **5** or the anti-fundamental **5*** scalar bound states together with quark and lepton multiplets, possible bulk fermions are **5**, **10*** and **24**. For forming an adjoint **24** scalar bound state together with quark and lepton multiplets, possible bulk fermions are **15**, **40*** and **45**. Added new fermions have zero mode only for the right-handed component. In general they are chiral in four dimensions. Therefore new fields need to be added in such a way that anomaly is canceled. An anomaly coefficient $A(r)$ is defined by

$$\text{tr} [t_r^a \{t_r^b, t_r^c\}] = \frac{1}{2} A(r) d^{abc} \quad (2.1)$$

where $\{t_N^a, t_N^b\} = \frac{1}{N} \delta^{ab} + d^{abc} t_N^c$ for $SU(N)$. For $N = 5$, the generators associated with Q_Y yield $d^{888} = -1/\sqrt{15}$. The anomaly coefficients for the above irreducible representations are given by

$$A(\mathbf{5}) = -\frac{1}{2}, \quad A(\mathbf{10}^*) = \frac{1}{2}, \quad A(\mathbf{15}) = \frac{9}{2}, \quad A(\mathbf{24}) = 0, \quad A(\mathbf{40}^*) = -8, \quad A(\mathbf{45}) = 3. \quad (2.2)$$

Thus the minimal anomaly-free set to obtain **5** or **5*** scalar bound states and a **24** scalar bound state is (**10***, **15**, **40***, **45**) Dirac fermions. In the next section, we will examine the binding strengths for bound states in the most attractive channel approximation.

3 Composites and binding strengths

In this section we derive possible bound states and their quantum numbers and binding strengths by adding bulk fields with zero mode for the right-handed component, χ . Here $\bar{\psi}\chi$ includes $\bar{\psi}_L^{(0)} \chi_R^{(0)}$ composed of only zero mode and strong attractive force can lead to potentials with phase transition. First we examine the minimal anomaly-free set. Taking

into account the pattern of symmetry breaking more, we study a simple anomaly-free set. In addition, the **50** and **75** bound states associated with the doublet-triplet splitting are discussed.

The minimal anomaly-free set

The quark and lepton multiplets are ψ_{5*} and ψ_{10} . The minimal set of addition fermions is given by χ_{10*} , χ_{15} , χ_{40*} and χ_{45} . It is necessary to examine the binding strengths for eight possible scalar bound states

$$\bar{\psi}_{5*}\chi_{10*}, \bar{\psi}_{5*}\chi_{15}, \bar{\psi}_{5*}\chi_{40*}, \bar{\psi}_{5*}\chi_{45}, \bar{\psi}_{10}\chi_{10*}, \bar{\psi}_{10}\chi_{15}, \bar{\psi}_{10}\chi_{40*}, \bar{\psi}_{10}\chi_{45}. \quad (3.1)$$

They are decomposed further in terms of irreducible representations into twenty four bound states

$$\mathbf{5} \otimes \mathbf{10*} = \mathbf{5*} \oplus \mathbf{45}, \quad (3.2)$$

$$\mathbf{5} \otimes \mathbf{15} = \mathbf{35} \oplus \mathbf{40}, \quad (3.3)$$

$$\mathbf{5} \otimes \mathbf{40*} = \mathbf{10*} \oplus \mathbf{15*} \oplus \mathbf{175_A^*}, \quad (3.4)$$

$$\mathbf{5} \otimes \mathbf{45} = \mathbf{24} \oplus \mathbf{75} \oplus \mathbf{126}, \quad (3.5)$$

$$\mathbf{10*} \otimes \mathbf{10*} = \mathbf{5} \oplus \mathbf{45*} \oplus \mathbf{50*}, \quad (3.6)$$

$$\mathbf{10*} \otimes \mathbf{15} = \mathbf{24} \oplus \mathbf{126}, \quad (3.7)$$

$$\mathbf{10*} \otimes \mathbf{40*} = \mathbf{24} \oplus \mathbf{75} \oplus \mathbf{126*} \oplus \mathbf{175_B^*}, \quad (3.8)$$

$$\mathbf{10*} \otimes \mathbf{45} = \mathbf{10} \oplus \mathbf{15} \oplus \mathbf{40*} \oplus \mathbf{175} \oplus \mathbf{210}, \quad (3.9)$$

The binding strength for $\bar{\psi}\chi$ is given by [19][1]

$$\frac{1}{2}g^2 [C_2(\bar{\psi}) + C_2(\chi) - C_2(\bar{\psi}\chi)]. \quad (3.10)$$

The coupling constant g is common for types of fields in the present model. For SU(5), the values of C_2 for the irreducible representations appearing here are summarized in Table 2. Using these values, we find the binding strengths for the twenty four bound

Table 2: The quadratic Casimir operator C_2 for each representation r .

r	5	10	15	24	35	40	45	50	75	126	175_A	175_B	210
C_2	$\frac{12}{5}$	$\frac{18}{5}$	$\frac{28}{5}$	5	$\frac{48}{5}$	$\frac{33}{5}$	$\frac{32}{5}$	$\frac{42}{5}$	8	10	$\frac{48}{5}$	12	$\frac{90}{7}$

states as shown in Table 3.

For the minimal field content, it is found that the binding strengths for **10** and **10*** are larger than that of the adjoint **24**. Usually SU(5) symmetry is broken by **24** or **75**. The results given in Table 3 means SU(5) symmetry would not be broken desirably unless there is some mechanism such that the binding for **10** and **10*** decreases effectively or the binding for **24** increases. In the following we consider a non-minimal but simple anomaly-free set where this problem does not arise.

Table 3: Binding strengths for $\bar{\psi}\chi$ for the minimal set in unit of g^2 .

Composite	Constituents	Binding strength	Composite	Constituents	Binding strength
10	$\bar{\psi}_{10}\chi_{45}$	16/5	75	$\bar{\psi}_{5*}\chi_{45}$	2/5
10*	$\bar{\psi}_{5*}\chi_{40*}$	27/10	45*	$\bar{\psi}_{10}\chi_{10*}$	2/5
24	$\bar{\psi}_{10}\chi_{40*}$	13/5	175_A	$\bar{\psi}_{10}\chi_{45}$	1/5
5	$\bar{\psi}_{10}\chi_{10*}$	12/5	126*	$\bar{\psi}_{10}\chi_{40*}$	1/10
15	$\bar{\psi}_{10}\chi_{45}$	11/5	45	$\bar{\psi}_{5*}\chi_{10*}$	-1/5
24	$\bar{\psi}_{10}\chi_{15}$	21/10	175_A*	$\bar{\psi}_{5*}\chi_{40*}$	-3/10
24	$\bar{\psi}_{5*}\chi_{45}$	19/10	126	$\bar{\psi}_{10}\chi_{15}$	-2/5
5*	$\bar{\psi}_{5*}\chi_{10*}$	9/5	126	$\bar{\psi}_{5*}\chi_{45}$	-3/5
15*	$\bar{\psi}_{5*}\chi_{40*}$	17/10	50*	$\bar{\psi}_{10}\chi_{10*}$	-3/5
40*	$\bar{\psi}_{10}\chi_{45}$	17/10	35	$\bar{\psi}_{5*}\chi_{15}$	-4/5
75	$\bar{\psi}_{10}\chi_{40*}$	11/10	175_B*	$\bar{\psi}_{10}\chi_{40*}$	-9/10
40	$\bar{\psi}_{5*}\chi_{15}$	7/10	210	$\bar{\psi}_{10}\chi_{45}$	-10/7

The simple anomaly-free set

As given in the previous section, a **24** scalar bound state can be made of **15**, **40*** and **45** Dirac fermions with quark and lepton multiplets. From Table 3, it is read that a composite **24** has the largest binding strength only for a **15** χ field and the other composites are largest for different quantum numbers: the composite **10*** for a **40*** χ field and the composite **10** for a **45** χ field. Therefore the fermion with zero mode for the right-handed component to form a **24** scalar bound state is uniquely fixed as a **15** χ field.

One of the simplest anomaly-free fermion content with **15** is 9 χ_5 and 1 χ_{15} . The composite are given by $\bar{\psi}_{5*}\chi_5$ $\bar{\psi}_{5*}\chi_{15}$ $\bar{\psi}_{10}\chi_5$ and $\bar{\psi}_{10}\chi_{15}$ as well as 8 copies of χ_5 . They are decomposed into eight irreducible representations,

$$\mathbf{5} \otimes \mathbf{5} = \mathbf{10} \oplus \mathbf{15}, \quad (3.11)$$

$$\mathbf{10*} \otimes \mathbf{5} = \mathbf{5*} \oplus \mathbf{45}, \quad (3.12)$$

with Eqs. (3.3) and (3.7). For this simple set, the binding strengths are given in Table 4.

Table 4: Binding strengths for $\bar{\psi}\chi$ for the simple set in unit of g^2 .

Composite	Constituents	Binding strength	Composite	Constituents	Binding strength
24	$\bar{\psi}_{10}\chi_{15}$	21/10	45	$\bar{\psi}_{10}\chi_5$	-1/5
5*	$\bar{\psi}_{10}\chi_5$	9/5	126	$\bar{\psi}_{10}\chi_{15}$	-2/5
40	$\bar{\psi}_{5*}\chi_{15}$	7/10	15	$\bar{\psi}_{5*}\chi_5$	-2/5
10	$\bar{\psi}_{5*}\chi_5$	3/5	35	$\bar{\psi}_{5*}\chi_{15}$	-4/5

We find scalar bound states whose binding strengths are the largest for **24** in $\bar{\psi}_{10}\chi_{15}$ and the second largest for **5** in $\bar{\chi}_5\psi_{10}$. The largest strength for **24** corresponds to the

breaking of the $SU(5)$ to $SU(3) \times SU(2) \times U(1)$. After the $SU(5)$ is broken, nine **5** and **5*** composite scalar fields contribute for the breaking of $SU(2) \times U(1)$ to $U(1)$.

Now we compare the binding strengths given in Table 4 with the results of the standard model gauge group. When quarks Q, U, D , leptons L, E, N and $SU(3) \times SU(2) \times U(1)$ gauge bosons propagate in five dimensions, the binding strengths for the composites with zero mode are given in Table 5. Here Q, L are $SU(2)$ doublets and U, D, E, N are $SU(2)$ singlets.

Table 5: Binding strength for the standard model gauge group [11].

Composite	Constituents	$SU(3) \times SU(2) \times U(1)$ representation	Binding strength	Binding for $\sqrt{\frac{3}{5}} g_1 = g_2 = g_3 \equiv g$
H_1	QU	$(\mathbf{1}, \mathbf{2}, \frac{1}{2})$	$\frac{4}{3}g_3^2 + \frac{1}{9}g_1^2$	$(7/5) g^2$
H_2	$\bar{Q}D$	$(\mathbf{1}, \mathbf{2}, -\frac{1}{2})$	$\frac{4}{3}g_3^2 - \frac{1}{18}g_1^2$	$(13/10) g^2$
S_{10}	EQ	$(\mathbf{3}, \mathbf{2}, \frac{7}{6})$	$-\frac{1}{6}g_1^2$	$(-1/10) g^2$
S_{11}	$\bar{L}U$	$(\mathbf{3}, \mathbf{2}, \frac{5}{6})$	0	0
S_{12}	$\bar{L}D$	$(\mathbf{3}, \mathbf{2}, \frac{1}{6})$	$\frac{1}{6}g_1^2$	$(1/10) g^2$
S_{13}	$\bar{L}E$	$(\mathbf{1}, \mathbf{2}, -\frac{1}{2})$	$\frac{1}{2}g_1^2$	$(3/10) g^2$
S_{14}	N or \bar{N} is included		0	0

From Tables 4 and 5, the binding strengths for **24** and **5*** are larger than that of the Higgs doublet H_1 . Because the self-breaking of the standard model gauge group has been claimed [1] [2] [3], the symmetry breaking associated with the larger binding strengths can occur. In other words, the symmetry breaking of $SU(5)$ to color and electromagnetism can be expected.

On the composites **50**, **50*** and **75**

A solution to the doublet-triplet splitting has been to break the $SU(5)$ symmetry by the real representation **75** instead of **24** and add **50** and **50*** [20] [21]. Now we consider the composites **50**, **50*** and **75**.

First we examine the composite **75**. For quark-lepton multiplets ψ_{5*} and ψ_{10} , fermions χ with the representations **10**, **40**, **45** and **50** can form **75** bound states as

$$\mathbf{5} \otimes \mathbf{45} = \mathbf{24} \oplus \mathbf{75} \oplus \mathbf{126}, \quad (3.13)$$

$$\mathbf{5} \otimes \mathbf{50} = \mathbf{75} \oplus \mathbf{175_B}, \quad (3.14)$$

$$\mathbf{10*} \otimes \mathbf{10} = \mathbf{1} \oplus \mathbf{24} \oplus \mathbf{75}, \quad (3.15)$$

$$\mathbf{10*} \otimes \mathbf{40*} = \mathbf{24} \oplus \mathbf{75} \oplus \mathbf{126*} \oplus \mathbf{175_B*}. \quad (3.16)$$

Among these constituents, only the **50** χ field yields a **75** bound state without the adjoint **24**. The binding strengths for the constituents $\bar{\psi}_{5*}\chi_{50}$ are $12/5$ for the composite **75** and $-3/5$ for the composite **175_B**. This point is favorable. However, in addition to the coupling with ψ_{5*} , $\bar{\psi}_{10}\chi_{50}$ needs to be taken into account where $\mathbf{10*} \otimes \mathbf{50} = \mathbf{10} \oplus \mathbf{175} \oplus \mathbf{315}$. The binding strength for the constituents $\bar{\psi}_{10}\chi_{50}$ are $21/5$ for the composite **10**. This means that the composite **75** with the quark-lepton multiplets for the constituents involves other composites bound with larger strengths.

Next we examine the composites **50** and **50***. When the **5*** and **10** quark-lepton multiplets and their conjugates coupled to fermions with right-handed zero mode form **50** or **50*** bound states, the composites would include **45** or **45*** bound states simultaneously. Since the quadratic Casimir operators have the relation $C_2(\mathbf{50}) < C_2(\mathbf{45})$, the composite **45** is bound stronger than the composite **50**. This gives rise to SU(2) doublet mass terms. It is because the representation **45** for SU(5) has a color singlet and weak doublet state for SU(3)×SU(2) and the composites **45** and **75** can become an SU(5) singlet with the composite **5** as seen in Eq. (3.13).

Therefore doublet-triplet splitting by the composites **50**, **50*** and **75** would not occur minimally. The model would need to be modified for application of the idea.

4 Conclusion

If extra dimensions are included in an effective theory, the strength of force would behave unlike four dimensions. We have studied most attractive channel for an SU(5) grand unification in five dimensions. We have assumed that components in each SU(5) multiplet obey the same boundary condition with respect to extra dimensions to keep the gauge coupling unification above the unification scale. In addition to quarks, leptons and gauge bosons, ψ_{5*} , ψ_{10} and A_M , anomaly-free sets of fields χ with zero mode for the right-handed component lead to various scalar bound states.

The minimal anomaly-free set to obtain **5** and **24** scalar bound states is $(\psi_{5*}, \psi_{10}, A_M)$ plus $(\chi_{10*}, \chi_{15}, \chi_{40*}, \chi_{45})$. For this field content, the binding strengths for **10** and **10*** have been found to be larger than that of **24**. Then, before SU(5) symmetry is broken to the standard model gauge group by **24**, symmetry breaking can occur by **10** and **10***.

A simple way to overcome this problem is to adopt $(\chi_5^{I=1,\dots,9}, \chi_{15})$ instead of the anomaly-free set $(\chi_{10*}, \chi_{15}, \chi_{40*}, \chi_{45})$. Here the **15** χ field is uniquely chosen because it is the only representation to form a **24** scalar bound state whose binding strength is large compared with other composites. The largest strength for the composite **24** corresponds to the breaking of the SU(5) to the standard model gauge group. After the SU(5) is broken, **5** in $\bar{\psi}_{10}\chi_5$ contribute for the breaking of SU(2)×U(1) to U(1). In addition, we have found that the binding strengths for **24** and **5** are larger than the known results for the composite Higgs doublet for the standard model gauge group.

We have shown that the symmetry breaking of the grand unification group to color and electromagnetism may occur by **24** and **5**. It would be more interesting to identify other fermion sets to assure the double-triplet splitting. However, we have found that the doublet-triplet splitting by the composites **50**, **50*** and **75** does not seem to occur minimally.

Finally, our analysis can be developed in various directions such as modification of field contents. It needs to be examined in more detail how masses and charges of elementary particles are derived appropriately.

Acknowledgments

This work is supported by Scientific Grants from the Ministry of Education and Science, Grant No. 20244028.

References

- [1] N. Arkani-Hamed, H. C. Cheng, B. A. Dobrescu and L. J. Hall, Phys. Rev. D **62**, 096006 (2000) [arXiv:hep-ph/0006238].
- [2] B. A. Dobrescu, Phys. Lett. B **461**, 99 (1999) [arXiv:hep-ph/9812349].
- [3] H. C. Cheng, B. A. Dobrescu and C. T. Hill, Nucl. Phys. B **589**, 249 (2000) [arXiv:hep-ph/9912343].
- [4] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
- [5] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064].
- [6] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Lett. B **473**, 43 (2000) [arXiv:hep-ph/9911262].
- [7] S. Chang, J. Hisano, H. Nakano, N. Okada and M. Yamaguchi, Phys. Rev. D **62**, 084025 (2000) [arXiv:hep-ph/9912498].
- [8] N. Rius and V. Sanz, Phys. Rev. D **64**, 075006 (2001) [arXiv:hep-ph/0103086].
- [9] T. Gherghetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000) [arXiv:hep-ph/0003129].
- [10] G. Burdman and L. Da Rold, JHEP **0712**, 086 (2007) [arXiv:0710.0623 [hep-ph]].
- [11] N. Uekusa, arXiv:1102.0607 [hep-ph].
- [12] Y. Kawamura, Prog. Theor. Phys. **105**, 999 (2001) [arXiv:hep-ph/0012125].
- [13] Y. Kawamura, Prog. Theor. Phys. **105**, 691 (2001) [arXiv:hep-ph/0012352].
- [14] K. R. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. B **436**, 55 (1998) [arXiv:hep-ph/9803466].
- [15] K. R. Dienes, E. Dudas and T. Gherghetta, Nucl. Phys. B **537**, 47 (1999) [arXiv:hep-ph/9806292].
- [16] G. Bhattacharyya, A. Datta, S. K. Majee and A. Raychaudhuri, Nucl. Phys. B **760**, 117 (2007) [arXiv:hep-ph/0608208].
- [17] N. Uekusa, Phys. Rev. D **75**, 064014 (2007) [arXiv:hep-th/0701159].

- [18] N. Uekusa, Int. J. Mod. Phys. A **23**, 3535 (2008) [arXiv:0803.1537 [hep-ph]].
- [19] S. Raby, S. Dimopoulos and L. Susskind, Nucl. Phys. B **169**, 373 (1980).
- [20] A. Masiero, D. V. Nanopoulos, K. Tamvakis and T. Yanagida, Phys. Lett. B **115**, 380 (1982).
- [21] B. Grinstein, Nucl. Phys. B **206**, 387 (1982).