

Inclusive production in a QCD and N=4 SYM motivated model for soft interactions

Errol Gotsman,¹ Eugene Levin,^{1,2} and Uri Maor¹

¹*Department of Particle Physics, Tel Aviv University, Tel Aviv 69978, Israel*

²*Departamento de Física, Universidad Técnica Federico Santa María,
Avda. España 1680, Casilla 110-V, Valparaíso, Chile*

(Dated: February 24, 2024)

The results presented in this paper differ from our previous unsuccessful attempt to predict the rapidity distribution at $W = 7\text{TeV}$. The original version of our model (GLMM) only summed a particular class of Pomeron diagrams (enhanced diagrams). We believe that this was the reason for our failure to describe the 7TeV inclusive LHC data. We have developed a new approach (GLM) that also includes the summation of the semi-enhanced diagrams. This contribution is essential for a successful description of the inclusive distributions, which is presented here.

Traditionally, inclusive hadron production at high energies has been considered a typical example of a soft process. Due to our lack of understanding of QCD at long distances, such processes are studied in the framework of high energy phenomenology based on soft Pomerons and their interactions. However, the first LHC data on hadron production [1–3] showed that the alternative approach, based on high density QCD[4–11], is able to predict[12] and describe the main features of the inclusive experimental data at the LHC [13–15]. On the other hand, our first attempt [16] to predict the inclusive rapidity spectra based on a model of soft interactions[17] failed to describe the experimental data, possibly giving the impression that soft models are unable to depict the LHC data. In this letter we show that this impression is incorrect, and that models based on soft Pomeron interactions are capable of reproducing the inclusive LHC data. In our two papers (see Refs. [17, 18]) we have built a model for soft interactions that is based on two theoretical precepts: high energy scattering in N=4 SYM, which stems from Pomeron interactions; and its matching with perturbative QCD, where we can use the BFKL Pomeron calculus to obtain the scattering amplitude. Our model includes:

- (i) Pomeron $\Delta_P \approx 0.2$ and $\alpha'_P \approx 0$. The large intercept of the Pomeron follows from N=4 SYM, and its value is compatible with N=4 SYM models for DIS data[19].
- (ii) A large Good-Walker component as in N=4 SYM, where the main contributions are due to elastic scattering and diffraction production [20, 21].
- (iii) Weak Pomeron interaction, which is of the order of $2/\sqrt{\lambda} \ll 1$ in N=4 SYM.
- (iv) Only the triple Pomeron vertex is necessary to provide a natural matching with perturbative QCD.

In our recent paper[18] we summed all essential Pomeron diagrams. Therefore, it is important for consistency, to check whether our postulates are necessary for the description of the inclusive hadron production at the LHC.

The inclusive cross section can be calculated in Pomeron calculus[22] (see also Refs.[23–26]) using Mueller diagrams[27] shown in Fig. 1. They lead to the following expression for the single inclusive cross section

$$\begin{aligned} \frac{1}{\sigma_{NSD}} \frac{d\sigma}{dy} &= \frac{1}{\sigma_{NSD}(Y)} \left\{ a_{PP} \left(\int d^2b \left(\alpha^2 G_1(b, Y/2 - y) + \beta^2 G_2(b, Y/2 - y) \right) \right. \right. \\ &\times \int d^2b \left(\alpha^2 G_1(b, Y/2 + y) + \beta^2 G_2(b, Y/2 + y) \right) \\ &- a_{PR} (\alpha^2 g_1^R + \beta^2 g_2^R) \left[\alpha^2 \int d^2b \left(\alpha^2 G_1(b, Y/2 - y) + \beta^2 G_2(b, Y/2 - y) \right) e^{\Delta_R(Y/2+y)} \right. \\ &\left. \left. + \int d^2b \left(\alpha^2 G_1(b, Y/2 + y) + \beta^2 G_2(b, Y/2 + y) \right) e^{\Delta_R(Y/2-y)} \right] \right\}. \end{aligned} \quad (1)$$

In Eq. (1) $G_i(b, Y)$ denotes the sum of 'fan' diagrams

$$G_i(b, Y) = (g_i(b)/\gamma) G_{enh}(y) / \left(1 + (G_{3P}/\gamma) g_i(b) G_{enh}(y) \right), \quad (2)$$

where the Green's function of the Pomeron, obtained by the summation of the enhanced diagrams[17], is equal to

$$G_{enh}(Y) = 1 - \exp\left(-\frac{1}{T(Y)}\right) \frac{1}{T(Y)} \Gamma\left(0, \frac{1}{T(Y)}\right). \quad (3)$$

In Eq. (2) and Eq. (3) we denote (see Refs.[17, 18] for details):

$$g_i(b) = g_i S_i(b) = \frac{g_i}{4\pi} m_i^3 b K_1(m_i b); \quad T(Y) = \gamma e^{\Delta_P Y}; \quad \gamma^2 = \int d^2k G_{3P}. \quad (4)$$




FIG. 1: The Mueller diagrams [27] for single inclusive cross section. The wavy bold line denotes the exact Pomeron Green's function of Eq. (3), which is the sum of the enhanced diagrams. The zig-zag line stands for the exchange of a Reggeon.

g_i denotes the Pomeron vertex of interaction with the i state at $b=0$; m_i is the parameter which determines the impact parameter dependence of this vertex; Δ_P and Δ_R are the Pomeron and Reggeon trajectory intercepts respectively. The triple Pomeron vertex G_{3P} and the parameters α and β , which determine the decomposition of the proton wave function into its GW components, $\Psi_{proton} = \alpha\Psi_1 + \beta\Psi_2$, have been discussed in Refs.[17, 18]. In our calculations we took the numerical values of these parameters from Ref.[18].

In Eq. (1) we introduce two new vertices: a_{PP} and a_{PR} , which describe the emission of hadrons from Pomeron and from the secondary Reggeon (see Fig. 1). In practice, we have to deal with two more dimensional parameters Q and Q_0 . Q is the average transverse momentum of the produced minijets, and $Q_0/2$ is the mass of the slowest hadron produced in the decay of the minijet. As we shall see below, the only parameter that determines the inclusive spectra is the ratio Q_0/Q .

We need these parameters to calculate the pseudo-rapidity η , which we use instead of the rapidity y . The relation between y and η is well known (see Ref.[28])

$$y(\eta, Q_0/Q) = \frac{1}{2} \ln \left\{ \frac{\sqrt{\frac{Q_0}{Q} + 1 + \sinh^2 \eta} + \sinh \eta}{\sqrt{\frac{Q_0}{Q} + 1 + \sinh^2 \eta} - \sinh \eta} \right\}, \quad (5)$$

with the Jacobian

$$h(\eta, Q_0/Q) = \frac{\cosh \eta}{\sqrt{\frac{Q_0}{Q} + 1 + \sinh^2 \eta}}. \quad (6)$$

Using Eq. (5) and Eq. (6) we can re-write Eq. (1) in the form

$$\begin{aligned} \frac{1}{\sigma_{NSD}} \frac{d\sigma}{d\eta} &= h(\eta, Q_0/Q) \frac{1}{\sigma_{NSD}(Y)} \left\{ a_{PP} \left(\int d^2b \left\{ \alpha^2 G_1(b, Y/2 - y(\eta, Q_0/Q)) + \beta^2 G_2(b, Y/2 - y(\eta, Q_0/Q)) \right\} \right) \right. \\ &\times \int d^2b \left\{ \alpha^2 G_1(b, Y/2 + y(\eta, Q_0/Q)) + \beta^2 G_1(b, Y/2 + y(\eta, Q_0/Q)) \right\} \Big) \\ &- a_{PR} (\alpha^2 g_1^R + \beta^2 g_2^R) \left[\alpha^2 \int d^2b \left\{ \alpha^2 G_1(b, Y/2 - y(\eta, Q_0/Q)) + \beta^2 G_2(b, Y/2 - y(\eta, Q_0/Q)) \right\} \right] e^{\Delta_R(Y/2+y)} \\ &+ \left. \int d^2b \left\{ \alpha^2 G_1(b, Y/2 + y(\eta, Q_0/Q)) + \beta^2 G_2(b, Y/2 + y(\eta, Q_0/Q)) \right\} e^{\Delta_R(Y/2-y)} \right] \Big\}. \end{aligned} \quad (7)$$

We extract the three new parameters: a_{PP} , a_{PR} and Q_0/Q from the experimental inclusive data. We made two separate fits: (a) fitting only the CMS data at different LHC energies (see Fig. 2-a); and (b) fitting all inclusive data for $W \geq 546 \text{ GeV}$ (see Fig. 2-b). We choose only data in the central region of rapidity, as we have not included energy conservation, and therefore our model is inadequate to describe the data behavior in the fragmentation region. The values of fitted parameters are presented in the table. As stated, all other parameters were taken from Ref.[18].

Fig. 2 shows that the soft model based on the Pomeron approach is able to describe the behavior and the value of the inclusive production observed experimentally. Our predictions are shown in the same figure. We note that the final version of our approach which includes the contributions of enhanced, semi-enhanced and net diagrams (see Ref.[18]) provides a much better description of the data than we obtained in our previous attempt[16], where only enhanced diagrams were summed.




FIG. 2: The single inclusive density versus energy. The data were taken from Refs.[1–3] and from Ref.[29]. The fit to the CMS data is plotted in Fig. 2-a, while Fig. 2-b presents the description of all inclusive spectra with $W \geq 546\text{GeV}$.

Data	$a_{\mathcal{P}\mathcal{P}}$	$a_{\mathcal{P}\mathcal{R}}$	Q_0/Q
CMS	0.39	0.186	0.427
All	0.413	0.194	0.356

TABLE I: Values of parameters for the fit of inclusive spectra.

We believe that our description of the inclusive production presented here will be efficacious in calculations of other observables at high energies, such as correlations and multiplicity dependences.

Unfortunately, up to now, we are the only group that has attempted to describe inclusive production in the framework of a soft model. We hope that this effort will provide a background for other microscopic approaches based on high density QCD.

Acknowledgement

We thank all participants of the conference Diffraction 2010 for fruitful discussion on the subject. The research of one of us (E.L.) was supported in part by the Fondecyt (Chile) grant # 1100648.

-
- [1] ALICE Collaboration, *Eur. Phys. J. C* **65** (2010) 111 [arXiv:0911.5430 [hep-ex]].
 - [2] CMS Collaboration, *JHEP* **1002** (2010) 041 [arXiv:1002.0621 [hep-ex]].
 - [3] ATLAS Collaboration, arXiv:1003.3124 [hep-ex].
 - [4] L. V. Gribov, E. M. Levin and M. G. Ryskin, *Phys. Rep.* **100** (1983) 1.
 - [5] A. H. Mueller and J. Qiu, *Nucl. Phys.* **B268** (1986) 427.
 - [6] L. McLerran and R. Venugopalan, *Phys. Rev.* **D49** (1994) 2233, 3352; **D50** (1994) 2225; **D53** (1996) 458; **D59** (1999) 09400.
 - [7] I. Balitsky, [arXiv:hep-ph/9509348]; *Phys. Rev.* **D60** (1999) 014020 [arXiv:hep-ph/9812311].
 - [8] Y. V. Kovchegov, *Phys. Rev.* **D60** (1999) 034008 [arXiv:hep-ph/9901281].
 - [9] J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, *Phys. Rev.* **D59** (1999) 014014 [arXiv:hep-ph/9706377]; *Nucl. Phys.* **B504** (1997) 415 [arXiv:hep-ph/9701284]; J. Jalilian-Marian, A. Kovner and H. Weigert, *Phys. Rev.* **D59** (1999) 014015 [arXiv:hep-ph/9709432]; A. Kovner, J. G. Milhano and H. Weigert, *Phys. Rev.* **D62** (2000) 114005 [arXiv:hep-ph/0004014]; E. Iancu, A. Leonidov and L. D. McLerran, *Phys. Lett.* **B510** (2001) 133 [arXiv:hep-ph/0102009]; *Nucl. Phys.* **A692** (2001) 583 [arXiv:hep-ph/0011241]; E. Ferreira, E. Iancu, A. Leonidov and L. McLerran, *Nucl. Phys.* **A703** (2002) 489 [arXiv:hep-ph/0109115]; H. Weigert, *Nucl. Phys.* **A703** (2002) 823 [arXiv:hep-ph/0004044].
 - [10] D. Kharzeev, E. Levin and M. Nardi, *Nucl. Phys.* **A730** (2004) 448 [Erratum-ibid. **A743** (2004) 329] [arXiv:hep-ph/0212316]; *Phys. Rev.* **C71** (2005) 054903 [arXiv:hep-ph/0111315]; D. Kharzeev and E. Levin, *Phys. Lett.* **B523** (2001) 79 [arXiv:nucl-th/0108006]; D. Kharzeev and M. Nardi, *Phys. Lett.* **B507** (2001) 121 [arXiv:nucl-th/0012025].
 - [11] D. Kharzeev, E. Levin and M. Nardi, *Nucl. Phys.* **A747** (2005) 609 [arXiv:hep-ph/0408050].
 - [12] E. Levin and A. H. Rezaeian, *Phys. Rev.* **D82** (2010) 014022. [arXiv:1005.0631 [hep-ph]].
 - [13] E. Levin, A. H. Rezaeian, [arXiv:1102.2385 [hep-ph]]; *Phys. Rev.* **D82** (2010) 054003. [arXiv:1007.2430 [hep-ph]].
 - [14] L. McLerran, M. Praszalowicz, *Acta Phys. Polon.* **B42** (2011) 99 [arXiv:1011.3403 [hep-ph]]; *Acta Phys. Polon.* **B41** (2010) 1917. [arXiv:1006.4293 [hep-ph]].
 - [15] M. Praszalowicz, [arXiv:1101.6012 [hep-ph]]; [arXiv:1101.0585 [hep-ph]].
 - [16] E. Gotsman, E. Levin, U. Maor, *Phys. Rev.* **D81** (2010) 051501 [arXiv:1001.5157 [hep-ph]].
 - [17] E. Gotsman, E. Levin, U. Maor and J. S. Miller, *Eur. Phys. J.* **C57** (2008) 689 [arXiv:0805.2799 [hep-ph]].
 - [18] E. Gotsman, E. Levin, U. Maor, *Eur. Phys. J.* **C71** (2011) 1553 [arXiv:1010.5323 [hep-ph]].
 - [19] R. C. Brower, M. Djuric, I. Sarcevic, C. I. Tan, *JHEP* **1011** (2010) 051 [arXiv:1007.2259 [hep-ph]]; E. Levin, I. Potashnikova, *JHEP* **1008** (2010) 112 [arXiv:1007.0306 [hep-ph]].
 - [20] R. C. Brower, J. Polchinski, M. J. Strassler and C. I. Tan, *JHEP* **0712** (2007) 005 [arXiv:hep-th/0603115]; R. C. Brower, M. J. Strassler and C. I. Tan, "On The Pomeron at Large 't Hooft Coupling" arXiv:0710.4378 [hep-th].
 - [21] Y. Hatta, E. Iancu and A. H. Mueller, *JHEP* **0801** (2008) 026 [arXiv:0710.2148 [hep-th]].
 - [22] V. N. Gribov, *Sov. Phys. JETP* **26** (1968) 414 *Zh. Eksp. Teor. Fiz.* **53** (1967) 654].
 - [23] P.D.B. Collins, "An introduction to Regge theory and high energy physics", Cambridge University Press 1977.
 - [24] L. Caneschi (editor), "Regge Theory of Low -p_T Hadronic Interaction", North-Holland 1989.
 - [25] A. B. Kaidalov, *Phys. Usp.* **46** (2003) 1121; E. Levin, "An introduction to Pomerons", arXiv:hep-ph/9808486; "Everything about Reggeons. I: Reggeons in *soft* interaction", arXiv:hep-ph/9710546.
 - [26] V. A. Abramovsky, V. N. Gribov and O. V. Kancheli, *Yad. Fiz.* **18** (1973) 595 [*Sov. J. Nucl. Phys.* **18** (1974) 308].
 - [27] A. H. Mueller, *Phys. Rev.* **D2** (1970) 2963.
 - [28] D. Kharzeev and E. Levin, *Phys. Lett.* **B523** (2001) 79 [arXiv:nucl-th/0108006].
 - [29] C. Amsler et al. (Particle Data Group), *Phys. Lett.* **B667** (2008) 1.