Sharp Convergence Rate and Support Consistency of Multiple Kernel Learning with Sparse and Dense Regularization

Taiji Suzuki, Ryota Tomioka Department of Mathematical Informatics, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo t-suzuki@mist.i.u-tokyo.ac.jp, tomioka@mist.i.u-tokyo.ac.jp Masashi Sugiyama Department of Computer Science, Tokyo Institute of Technology, 2-12-1 O-okayama, Meguro-ku, Tokyo sugi@cs.titech.ac.jp

Abstract

We theoretically investigate the convergence rate and support consistency (i.e., correctly identifying the subset of non-zero coefficients in the large sample limit) of multiple kernel learning (MKL). We focus on MKL with block- ℓ_1 regularization (inducing sparse kernel combination), block- ℓ_2 regularization (inducing uniform kernel combination), and elastic-net regularization (including both block- ℓ_1 and block- ℓ_2 regularization). For the case where the true kernel combination is sparse, we show a sharper convergence rate of the block- ℓ_1 and elastic-net MKL methods than the existing rate for block- ℓ_1 MKL. We further show that elastic-net MKL requires a milder condition for being consistent than block- ℓ_1 MKL. For the case where the optimal kernel combination is not exactly sparse, we prove that elastic-net MKL can achieve a faster convergence rate than the block- ℓ_1 and block- ℓ_2 MKL methods by carefully controlling the balance between the block- ℓ_1 and block- ℓ_2 regularizers. Thus, our theoretical results overall suggest the use of elastic-net regularization in MKL.

1 Introduction

The choice of kernel functions is a key issue for *kernel methods* such as support vector machines to work well (Vapnik, 1998). A traditional but very powerful approach to optimizing the kernel function is the use of *cross-validation* (CV) (Stone, 1974). Although the CV-based kernel choice often leads to better generalization, it is computationally expensive when the kernel contains multiple tuning parameters.

To overcome this limitation, the framework of *multiple kernel learning* (MKL) has been introduced, which tries to learn the optimal linear combination of prefixed base-kernels by convex optimization (Lanckriet et al., 2004, Micchelli and Pontil, 2005, Lin and Zhang, 2006, Sonnenburg et al., 2006, Rakotomamonjy et al., 2008, Suzuki and Tomioka, 2009). The seminal paper by Bach et al. (2004) showed that this MKL formulation can be interpreted as block- ℓ_1 regularization (i.e., ℓ_1 regularization across the kernels and ℓ_2 regularization within the same kernel). We refer to this MKL formulation as 'block- ℓ_1 MKL'. Based on this interpretation, block- ℓ_1 MKL was proved to be *support consistent* (i.e., correctly identifying the subset of non-zero coefficients with probability one in the large sample limit) when the true kernel combination is sparse (Bach, 2008). Furthermore, the convergence rate of block- ℓ_1 MKL has also been elucidated in Koltchinskii and Yuan (2008), which can be regarded as an extension of the theoretical analysis for ordinary (non-block) ℓ_1 regularization (Bickel et al., 2009, Zhang, 2009).

However, in many practical applications, the true kernel combination may not be exactly sparse. In such a non-sparse situation, $block-\ell_1$ MKL was shown to perform rather poorly—just the uniform combination of base kernels obtained by $block-\ell_2$ regularization (Micchelli and Pontil, 2005) (which we call 'block- ℓ_2 MKL') often works better in practice (Cortes, 2009). Furthermore, recent works showed that some 'intermediate' regularization between $block-\ell_1$ and $block-\ell_2$ regularization is more promising, e.g., $block-\ell_p$ regularization with $1 \le p \le 2$ (Cortes et al., 2009, Kloft et al., 2009), and *elastic-net* regularization (Zou and Hastie, 2005) which includes both $block-\ell_1$ and $block-\ell_2$ regularization (Tomioka and Suzuki, 2010) (we call this method 'elastic-net MKL'). Theoretically, the support consistency and the convergence rate for parametric elastic-nets have been elucidated in Yuan and Lin (2007) and Zou and Zhang (2009), respectively, and that for non-parametric cases has been investigated in Meier et al. (2009) focusing on the Sobolev space.

In this paper, we theoretically analyze the support consistency and convergence rate of MKL, and provide three new results.

- For the case where the true kernel combination is sparse, we show that elastic-net MKL achieves a faster convergence rate than the one shown for block-l₁ MKL (Koltchinskii and Yuan, 2008). More specifically, we show that the L₂ convergence error is given by O_p(min{dn^{-2/2+s} + dlog(M)/n, d^{1-s/1+s}n^{-1/1+s} + dlog(M)/n}), where d is the number of active components of the target function, s is the complexity of RKHSs, M is the number of candidate kernels, and n is the number of samples.
- For the case where the optimal kernel combination is not exactly sparse, we prove that elastic-net MKL achieves a faster convergence rate than the block-l₁ and block-l₂ MKL methods by carefully controlling the balance between block-l₁ and block-l₂ regularization. Our theoretical result well agrees with the experimental results reported in Tomioka and Suzuki (2010).
- For the case where the true kernel combination is sparse, we prove that the necessary and sufficient conditions of the support consistency for elastic-net MKL is milder than the conditions required for block- ℓ_1 MKL (Bach, 2008).

Overall, our theoretical results suggest the use of elastic-net regularization in MKL.

2 Preliminaries

In this section, we formulate the elastic-net MKL approach and summarize mathematical tools that are needed for the theoretical analysis.

2.1 Formulation

Suppose we are given n samples $(x_i, y_i)_{i=1}^n$ where x_i belongs to an input space \mathcal{X} and $y_i \in \mathbb{R}$. $(x_i, y_i)_{i=1}^n$ are independent and identically distributed from a probability measure P. We denote the marginal distribution of X by Π . We consider a MKL regression problem in which the unknown target function is represented as a form of $f(x) = \sum_{m=1}^M f_m(x)$, where each f_m belongs to different RKHSs $\mathcal{H}_m(m = 1, \ldots, M)$ corresponding to M different base kernels k_m over $\mathcal{X} \times \mathcal{X}$.

Elastic-net MKL learns a decision function f as¹

$$\hat{f} = \operatorname*{arg\,min}_{f_m \in \mathcal{H}_m \ (m=1,\dots,M)} \frac{1}{n} \sum_{i=1}^n \left(y_i - \sum_{m=1}^M f_m(x_i) \right)^2 + \lambda_1^{(n)} \sum_{m=1}^M \|f_m\|_{\mathcal{H}_m} + \lambda_2^{(n)} \sum_{m=1}^M \|f_m\|_{\mathcal{H}_m}^2, \quad (1)$$

where the first term is the squared-loss of function fitting and, the second and the third terms are block- ℓ_1 and block- ℓ_2 regularizers, respectively. It can be seen from (1) that elastic-net MKL is reduced to block- ℓ_1 MKL if $\lambda_2^{(n)} = 0$, which tends to induce sparse kernel combination (Lanckriet et al., 2004, Bach et al., 2004). On the other hand, it is reduced to block- ℓ_2 MKL if $\lambda_1^{(n)} = 0$, which results in uniform kernel combination (Micchelli and Pontil, 2005). It is worth noting that, elastic-net MKL allows us to obtain various levels of sparsity by controlling the ratio between $\lambda_1^{(n)}$ and $\lambda_2^{(n)}$.

2.2 Notations and Assumptions

Here, we prepare technical tools needed in the following sections.

Due to Mercer's theorem, there are an orthonormal system $\{\phi_{k,m}\}_{k,m}$ in $L_2(\Pi)$ and the spectrum $\{\mu_{k,m}\}_{k,m}$ such that k_m has the following spectral representation:

$$k_m(x, x') = \sum_{k=1}^{\infty} \mu_{k,m} \phi_{k,m}(x) \phi_{k,m}(x').$$
(2)

By this spectral representation, the inner-product of RKHS can be expressed as $\langle f_m, g_m \rangle_{\mathcal{H}_m} = \sum_{k=1}^{\infty} \mu_{k,m}^{-1} \langle f_m, \phi_{k,m} \rangle_{L_2(\Pi)} \langle \phi_{k,m}, g_m \rangle_{L_2(\Pi)}$.

Let $\mathcal{H} = \mathcal{H}_1 \oplus \cdots \oplus \mathcal{H}_M$. For $f = (f_1, \ldots, f_M) \in \mathcal{H}$ and a subset of indices $I \subseteq \{1, \ldots, M\}$, we denote by f_I the restriction of f to an index set I, i.e., $f_I = (f_m)_{m \in I}$.

We denote by I_0 the indices of truly active kernels, i.e.,

$$I_0 = \{ m \mid ||f_m^*||_{\mathcal{H}_m} > 0 \},\$$

and define the complement of I_0 as $J_0 = I_0^{c}$.

Throughout the paper, we assume the following technical conditions (see also Bach (2008)).

¹ For simplicity, we focus on the squared-loss function here. However, we note that it is straightforward to extend our convergence analysis and support consistency results given in Sections 3 and 4 to general loss functions that are strongly convex and Lipschitz continuous, by following the line of Koltchinskii and Yuan (2008).

Table 1: Summary of the constants we use in this article.

M	The number of candidate kernels.
d	The number of active kernels of the truth; i.e., $d = I_0 $.
R	The upper bound of $\sum_{m=1}^{M} (\ f_{m}^{*}\ _{\mathcal{H}_{m}} + \ f_{m}^{*}\ _{\mathcal{H}_{m}}^{2})$; see (A4).
s	The spectral decay coefficient; see (A5).
β	The approximate sparsity coefficient; see (A7).
b	The parameter that tunes the correlation between kernels; see (A8).

Assumption 1 (Basic Assumptions)

- (A1) There exists $f^* = (f_1^*, \ldots, f_M^*) \in \mathcal{H}$ such that $\mathbb{E}[Y|X] = \sum_{m=1}^M f_m^*(X)$, and the noise $\epsilon := Y f^*(X)$ has a strictly positive variance; there exists $\sigma > 0$ such that $\mathbb{E}[\epsilon^2|X] > \sigma^2$ for all $X \in \mathcal{X}$. We also assume that ϵ is bounded as $|\epsilon| \leq L$.
- (A2) For each m = 1, ..., M, \mathcal{H}_m is separable and $\sup_{X \in \mathcal{X}} |k_m(X, X)| < 1$.
- (A3) There exists $g_m^* \in \mathcal{H}_m$ such that

$$f_m^*(x) = \int_{\mathcal{X}} k_m^{(1/2)}(x, x') g_m^*(x') d\Pi(x') \qquad (\forall m = 1, \dots, M),$$
(3)

where
$$k_m^{(1/2)}(x, x') = \sum_{k=1}^{\infty} \mu_{k,m}^{1/2} \phi_{k,m}(x) \phi_{k,m}(x')$$
 is the operator square-root of k_m .

The first assumption in (A1) ensures the model \mathcal{H} is correctly specified, and the technical assumption $|\epsilon| < L$ allows ϵf to be Lipschitz continuous with respect to f.

It is known that the assumption (A2) gives the following relation:

$$\|f_m\|_{\infty} \leq \sup_x \langle k_m(x,\cdot), f_m \rangle_{\mathcal{H}_m} \leq \sup_x \|k_m(x,\cdot)\|_{\mathcal{H}_m} \|f_m\|_{\mathcal{H}_m} \leq \sup_x \sqrt{k_m(x,x)} \|f_m\|_{\mathcal{H}_m} \leq \|f_m\|_{\mathcal{H}_m}$$

The assumption (A3) was used in Caponnetto and de Vito (2007) and also in Bach (2008). It ensures the consistency of the least-squares estimates in terms of the RKHS norm. Using the spectral representation (2), the condition $g_m^* \in \mathcal{H}_m$ is expressed as

$$\|g_m^*\|_{\mathcal{H}_m}^2 = \sum_{k=1}^{\infty} \mu_{k,m}^{-2} \langle f_m^*, \phi_{k,m} \rangle_{L_2(\Pi)}^2 < \infty.$$
(4)

This condition was also assumed in Koltchinskii and Yuan (2008). Proposition 9 of Bach (2008) gave a sufficient condition to fulfill (3) for translation invariant kernels $k_m(x, x') = h_m(x - x')$.

Constants we use later are summarized in Table 1.

3 Convergence Rate of Elastic-net MKL

In this section, we derive the convergence rate of elastic-net MKL in two situations:

- (i) A sparse situation where the truth f^* is sparse (Section 3.1).
- (ii) A near sparse situation where the truth is not exactly sparse, but $||f_m||_{\mathcal{H}_m}$ decays polynomially as m increases (Section 3.2).

For (i), we show that elastic-net MKL (and block- ℓ_1 MKL) achieves a faster convergence rate than the rate shown for block- ℓ_1 MKL (Koltchinskii and Yuan, 2008). Furthermore, for (ii), we show that elastic-net MKL can outperform block- ℓ_1 MKL and block- ℓ_2 MKL depending on the sparsity of the truth and the condition of the problem. Throughout this section, we assume the following conditions.

Assumption 2 (Boundedness Assumption) There exists constants C_1 and R such that

(A4)
$$\max_{m \in I_0} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} \le C_1, \quad \sum_{m=1}^M (\|f_m^*\|_{\mathcal{H}_m} + \|f_m^*\|_{\mathcal{H}_m}^2) \le R.$$

Assumption 3 (Spectral Assumption) There exist 0 < s < 1 and C_2 such that

(A5)
$$\mu_{k,m} \le C_2 k^{-\frac{1}{s}}, \quad (1 \le \forall k, 1 \le \forall m \le M),$$

where $\{\mu_{k,m}\}_k$ is the spectrum of the kernel k_m (see Eq.(2)).

The first assumption in (A4) appeared in Theorem 2 of Koltchinskii and Yuan (2008). The second assumption in (A4) bounds the amplitude of f^* . It was shown that the spectral assumption (A5) is equivalent to the classical covering number assumption (Steinwart et al., 2009). Recall that the ϵ -covering number $\mathcal{N}(\epsilon, \mathcal{B}_{\mathcal{H}_m}, L_2(\Pi))$ with respect to $L_2(\Pi)$ is the minimal number of balls with radius ϵ needed to cover the unit ball $\mathcal{B}_{\mathcal{H}_m}$ in \mathcal{H}_m (van der Vaart and Wellner, 1996). If the spectral assumption (A5) holds, there exists a constant *c* that depends only on *s* such that

$$\mathcal{N}(\varepsilon, \mathcal{B}_{\mathcal{H}_m}, L_2(\Pi)) \le c\varepsilon^{-2s},$$
(5)

and the converse is also true (see Theorem 15 of Steinwart et al. (2009) and Steinwart (2008) for details). Therefore, if s is large, at least one RKHS is "complex", and if s is small, the RKHSs are regarded as "simple".

For a given set of indices $I \subseteq \{1, \ldots, M\}$, let $\kappa(I)$ be defined as follows:

$$\kappa(I) := \sup\left\{\kappa \ge 0 \mid \kappa \le \frac{\|\sum_{m \in I} f_m\|_{L_2(\Pi)}^2}{\sum_{m \in I} \|f_m\|_{L_2(\Pi)}^2}, \, \forall f_m \in \mathcal{H}_m \ (m \in I)\right\}.$$

 $\kappa(I)$ represents the correlation of RKHSs inside the indices I. Similarly, we define the correlations of RKHSs between I and I^c as follows:

$$\rho(I) := \sup \left\{ \frac{\langle f_I, g_{I^c} \rangle_{L_2(\Pi)}}{\|f_I\|_{L_2(\Pi)} \|g_{I^c}\|_{L_2(\Pi)}} \mid f_I \in \mathcal{H}_I, g_{I^c} \in \mathcal{H}_{I^c}, f_I \neq 0, g_{I^c} \neq 0 \right\}.$$

In Subsections 3.1 and 3.2, we will assume that the kernels have *no perfect canonical dependence*, implying that the kernels are not similar to each other (see (A6) and (A8) below).

Throughout this paper, we assume $\frac{\log(Mn)}{n} \leq 1$ and $\log(M)$ is slower than any polynomial order against the number of samples $n: \log(M) = o(n^{\epsilon})$ for all $\epsilon > 0$. With some abuse, we use C to denote constants that are independent of d and n; its value may be different.

3.1 Sparse Situation

Here we derive the convergence rate of the estimator \hat{f} when the truth f^* is sparse. Let $d = |I_0|$ and suppose that the number of kernels M and the number of active kernels d are increasing with respect to the number of samples n. We further assume the following condition in this subsection.

Assumption 4 (Incoherence Assumption) There exists a constant $C_3 > 0$ such that

(A6)
$$0 < C_3^{-1} < \kappa(I_0)(1 - \rho^2(I_0)).$$
 (6)

This condition is known as the *incoherence condition* (Koltchinskii and Yuan, 2008, Meier et al., 2009), i.e., kernels are not too dependent on each other and the problem is well conditioned. Then we have the following convergence rate.

Theorem 1 Under assumptions (A1-A6), there exist constants C, F and K depending only on $\kappa(I_0)$, $\rho(I_0)$, s, C_1 , C_2 , L, and R such that the $L_2(\Pi)$ -norm of the residual $\hat{f} - f^*$ can be bounded as follows: when $d^{3+s}n^{-1} \leq 1$, for $\lambda_1^{(n)} = \lambda_2^{(n)} = \max\{Kn^{-\frac{1}{2+s}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\}$,

$$\|\hat{f} - f^*\|_{L_2(\Pi)}^2 \le C \Big(dn^{-\frac{2}{2+s}} + \frac{dt}{n} \Big),\tag{7}$$

and, when $d^{3+s}n^{-1} > 1$, for $\lambda_1^{(n)} = \max\{K(1+\sqrt{t})n^{-\frac{1}{2}}, F\sqrt{\frac{\log(Mn)}{n}}\}$ and $\lambda_2^{(n)} \le \lambda_1^{(n)}$,

$$\|\hat{f} - f^*\|_{L_2(\Pi)}^2 \le C \Big(d^{\frac{1-s}{1+s}} n^{-\frac{1}{1+s}} + \frac{d(\log(Mn) + t)}{n} \Big), \tag{8}$$

where each inequality holds with probability at least $1 - e^{-t} - n^{-1}$ for all $t \ge \log \log(R\sqrt{n}) + \log M$.

The above theorem indicates that the learning rate depends on the complexity of RKHSs (the simpler, the faster) and the number of *active* kernels rather than the number of kernels M (the influence of M is at most $\frac{d\log(M)}{n}$). It is worth noting that the convergence rate in (7) and (8) is faster than or equal to the rate of block- ℓ_1 MKL shown by Koltchinskii and Yuan (2008) which established the learning rate $O_p\left(d^{\frac{1-s}{1+s}}n^{-\frac{1}{1+s}} + \frac{d\log(M)}{n}\right)$ under the same conditions as ours ².

²In our second bound (8), there is the additional $\frac{d \log(n)}{n}$ term. However this can be eliminated by replacing the probability $1 - e^{-t} - n^{-1}$ with $1 - e^{-t} - M^{-A}$ as in Koltchinskii and Yuan (2008). Moreover, if $\sqrt{n} \log(n)^{-\frac{1+s}{2s}} \ge d$, then the term $\frac{d \log(n)}{n}$ is dominated by the first term $d^{\frac{1-s}{1+s}} n^{-\frac{1}{1+s}}$.

3.2 Near-Sparse Situation

In this subsection, we analyze the convergence rate under a situation where f^* is not sparse but *near sparse*. We have shown a faster learning rate than existing bounds in the previous subsection. However, the assumptions we used might be too restrictive to capture the situation where MKL is used in practice. In fact, it was pointed out in Zou and Hastie (2005) in the context of (non-block) ℓ_1 regularization that ℓ_1 regularization could fail in the following situations:

- When the truth f^* is not sparse, the ℓ_1 regularization shrinks many small but non-zero components to zero.
- When there exist strong correlations between different kernels, the solution of block- ℓ_1 MKL becomes unstable.
- When the number of kernels M is not large, there is no need to impose the estimator to be sparse.

In order to analyze these situations in the MKL setting, we introduce three parameters β , b, and τ : β controls the level of sparsity (see (A7)), b controls the correlation between candidate kernels (see (A8)), and τ controls the growth of the number of kernels against the number of samples (see (A9)).

We show that naturally block- ℓ_2 MKL is preferable when there are only few candidate kernels or the truth is dense. Importantly, if the candidate kernels are correlated, the convergence of block- ℓ_1 MKL can be slow even when the truth is sparse. Our analysis shows that elastic-net MKL is most valuable in such an intermediate situation.

By permuting indices, we can assume without loss of generality that $||f_m^*||_{\mathcal{H}_m}$ is decreasing with respect to m, i.e., $||f_1^*||_{\mathcal{H}_1} \ge ||f_2^*||_{\mathcal{H}_2} \ge ||f_3^*||_{\mathcal{H}_3} \ge \cdots$. We further assume the following conditions in this subsection.

Assumption 5 (Approximate Sparsity) The truth is approximately sparse, i.e., $||f_m^*||_{\mathcal{H}_m} > 0$ for all m and thus $I_0 = \{1, \ldots, M\}$. However, $||f_m^*||_{\mathcal{H}_m}$ decays polynomially with respect to m as follows:

(A7)
$$\|f_m^*\|_{\mathcal{H}_m} \le C_3 m^{-\beta}$$

We call β (> 1) *the approximate sparsity coefficient.*

Assumption 6 (Generalized Incoherence) There exist b > 0 and C_4 such that for all $I \subseteq \{1, \ldots, M\}$,

(A8)
$$(1 - \rho^2(I))\kappa(I) \ge C_4|I|^{-b}$$

Assumption 7 (Kernel-Set Growth) The number of kernels M is increasing polynomially with respect to the number of samples n, i.e., $\exists \tau > 0$ such that

(A9)
$$M = \lceil n^{\tau} \rceil.$$

For notational convenience, let $\tau_1 = \frac{1}{(2\beta+b)(2+s)-1-s}$, $\tau_2 = \frac{(s-1)(2\beta-1)+bs}{(2\beta+b)(2+s)-1-s}$, $\tau_3 = \frac{s\{2(b+\beta)-1\}}{2(2+s)(b+\beta)-s}$, $\tau_4 = \frac{s}{2+s}$, $\tau_5 = \frac{b+1}{(\beta+b)\{b(2+s)+2\}}$ and $\tau_6 = \frac{1}{(1-s)(1+b)}$. In addition, we denote by K some sufficiently large constant.

Theorem 2 Suppose assumptions (A1-A5) and (A7-A9), $2\beta(1-s) < s(b-1)$, and $\tau_1 < \tau < \tau_4$ are satisfied. Then the estimator of elastic-net MKL possesses the following convergence rate each of which holds with probability at least $1 - e^{-t} - n^{-1}$ for all $t \ge \log \log(R\sqrt{n}) + \log M$: 1. When $\tau_1 < \tau < \tau_2$,

$$\|\hat{f} - f^*\|_{L_2(\Pi)}^2 \le C \Big\{ n^{-\gamma_1} + (n^{-\frac{(2\beta+b)(2+s)-3-s+2\beta}{2\{(2\beta+b)(2+s)-1-s\}}} + \lambda_2^{(n)})(\sqrt{t}+t) \Big\},$$

where $\gamma_1 = \frac{4\beta+b-2}{(2+s)(2\beta+b)-1-s},$ (9)

with $\lambda_1^{(n)} = \max\{Kn^{-\frac{3\beta+b-1}{(2\beta+b)(2+s)-1-s}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\}$ and $\lambda_2^{(n)} = Kn^{-\frac{2\beta+b-1}{(2\beta+b)(2+s)-1-s}}$. 2. When $\tau_2 \leq \tau < \tau_3$,

$$\begin{split} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 \leq & C \Big\{ n^{\tau \frac{(2+s)b+2}{2\{(2+s)(b+\beta)-s\}} - \gamma_2} + (n^{\frac{\tau(2+s)(1-\beta) - (4\beta+2b+sb-2)}{2\{(\beta+b)(2+s)-s\}}} + \lambda_2^{(n)^2})(\sqrt{t} + t) \Big\}, \\ where \ \gamma_2 = \frac{4\beta + b(2+s) - 2}{2\{(2+s)(b+\beta) - s\}}, \end{split}$$
(10)

with $\lambda_1^{(n)} = \max\{K\sqrt{\frac{M}{n}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\} \text{ and } \lambda_2^{(n)} = Kn^{\frac{\tau - \{2(b+\beta)-1\}}{2\{(2+s)(b+\beta)-s\}}}.$ 3. When $\tau_3 \leq \tau < \tau_4$,

$$\|\hat{f} - f^*\|_{L_2(\Pi)}^2 \le C \Big(n^{\tau \gamma_3 - \gamma_3} + (n^{\frac{\tau(\beta - 1) + 1 - 2\beta - b}{2(b + \beta)}} + \lambda_2^{(n)^2})(\sqrt{t} + t) \Big),$$

$$where \ \gamma_3 = \frac{b + 2\beta - 1}{2(b + \beta)},$$
(11)

with $\lambda_1^{(n)} = \max\{K\sqrt{\frac{M}{n}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\} \text{ and } \lambda_2^{(n)} = K(M/n)^{\frac{2(b+\beta)-1}{4(b+\beta)}}.$

Theorem 3 Under assumptions (A1-A5) and (A7-A9), if $\tau_5 < \tau$, the estimator \hat{f}_{ℓ_1} of block- ℓ_1 MKL has the following convergence rate with probability at least $1 - e^{-t} - n^{-1}$ for all $t \ge \log \log(R\sqrt{n}) + \log M$:

$$(block - \ell_1 MKL) \qquad \|\hat{f}_{\ell_1} - f^*\|_{L_2(\Pi)}^2 \le C \left(n^{-\gamma_4} + n^{-\frac{4\beta + 2b - 2 + s(b+\beta)}{2(2+s)(b+\beta)}} (\sqrt{t} + t) \right),$$

where $\gamma_4 = \frac{2\beta + b - 1}{(\beta + b)(2+s)},$ (12)

with $\lambda_1^{(n)} = \max\{Kn^{-\frac{1}{2+s}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\}$ and $\lambda_2^{(n)} = 0$. Moreover, if $\tau < \tau_6$, the estimator \hat{f}_{ℓ_2} of block- ℓ_2 MKL has the following convergence rate with probability at least $1 - e^{-t} - n^{-1}$ for all $t \ge \log \log(R\sqrt{n}) + \log M$:

$$(block-\ell_2 MKL) \qquad \|\hat{f}_{\ell_2} - f^*\|_{L_2(\Pi)}^2 \le C \left(n^{\tau(b+\frac{2}{2+s})-\gamma_5} + \left(\lambda_2^{(n)^2} + \frac{M^{1+b}}{n}\right) t \right),$$

where $\gamma_5 = \frac{2}{2+s},$ (13)

with
$$\lambda_2^{(n)} = \max\{K(\frac{M}{n})^{\frac{1}{2+s}}, F\sqrt{\frac{\log(Mn)}{n}}\} \text{ and } \lambda_1^{(n)} = 0.$$

In all convergence rates presented in Theorems 2 and 3, the leading terms are the terms that do not contain *t*. The convergence order of the terms containing *t* are faster than the leading terms, thus negligible.

By simple calculation, we can confirm that elastic-net MKL always converges faster than block- ℓ_1 MKL and block- ℓ_2 MKL if β and M satisfy the condition of Theorem 2. The convergence rate of elastic-net MKL becomes identical with block- ℓ_2 MKL and block- ℓ_1 MKL at the two extreme points of the interval $\tau = \tau_1$ and τ_4 , respectively. Outside the region, block- ℓ_1 MKL or block- ℓ_2 MKL has a faster convergence rate than elastic-net MKL. Moreover, at $\tau = \tau_2$, the convergence rates (9) and (10) of elastic-net MKL are identical, and at $\tau = \tau_3$, the convergence rates (10) and (11) are identical. The relation between the most preferred method and the growth rate τ of the number of kernels is illustrated in Figure 1.

The condition $\tau_1 < \tau < \tau_4$ in Theorem 2 indicates that when the number of kernels is not too small or too large, an 'intermediate' effect of elastic-net MKL becomes advantageous. Roughly speaking, if M is large, sparsity is needed to ensure the convergence and thus block- ℓ_1 MKL performs the best. On the other hand, if M is small, there is no need to make the solution sparse and thus block- ℓ_2 MKL becomes the best. For an intermediate M, elastic-net MKL becomes the best.

The condition $2\beta(1-s) < s(b-1)$ in Theorem 2 ensures the existence of M that satisfies the condition in the theorem, i.e., $\tau_1 < \tau_2 < \tau_3 < \tau_4$. It can be seen that as b becomes large (the condition of the problem becomes worse), the range of β and M in which elastic-net MKL performs better than block- ℓ_1 MKL and block- ℓ_2 MKL becomes large. This indicates that the worse the condition of the problem becomes, the more important to control the balance of $\lambda_1^{(n)}$ and $\lambda_2^{(n)}$ appropriately.

4 Support Consistency of Elastic-net MKL

In this section, we derive necessary and sufficient conditions for the statistical support consistency of the estimated sparsity pattern, i.e., the probability of $\{m \mid ||\hat{f}_m||_{\mathcal{H}_m} \neq 0\} = I_0$ goes to 1 as the number of samples n tends to infinity. Due to the additional squared regularization term, the necessary condition for the support consistency of elastic-net MKL is shown to be weaker than that for block- ℓ_1 MKL (Bach, 2008). In this section, we assume M and $d = |I_0|$ are fixed against the number of samples n.

Let \mathcal{H}_I be the restriction of $\mathcal{H}_1 \oplus \cdots \oplus \mathcal{H}_M$ to the index set *I*. Since $\mathbb{E}_X[k_m(X,X)] < \infty$ for all *m* (from assumption (A2)), we define the (non-centered) *cross covariance operator* $\Sigma_{I,J} : \mathcal{H}_I \to \mathcal{H}_J$ as a bounded



Figure 1: Relation between the convergence rate and the number of kernels. If the truth is intermediately sparse (the growth rate τ of the number of kernels is between τ_1 and τ_5), then elastic-net MKL performs best. At the edge of the interval, the convergence rate of elastic-net MKL coincides with that of block- ℓ_1 MKL or block- ℓ_2 MKL.

linear operator such that³

$$\langle f_I, \Sigma_{I,J}g_J \rangle_{\mathcal{H}_I} = \sum_{m \in I} \sum_{m' \in J} \langle f_m, \Sigma_{m,m'}g_{m'} \rangle_{\mathcal{H}_m} = \sum_{m \in I} \sum_{m' \in J} \mathcal{E}_X[f_m(X)g_{m'}(X)], \tag{14}$$

for all $f_I = (f_m)_{m \in I} \in \mathcal{H}_I$ and $g_J = (g_{m'})_{m' \in J} \in \mathcal{H}_J$. See Baker (1973) for the details of the cross covariance operator $(f, g) \mapsto \operatorname{cov}(f(X)g(X))$.

Moreover, we define the bounded (non-centered) cross-correlation operators⁴ $V_{l,m}$ by $\Sigma_{l,l}^{1/2} V_{l,m} \Sigma_{m,m}^{1/2} = \Sigma_{l,m}$. The joint cross-correlation operator $V_{I,J} : \mathcal{H}_J \to \mathcal{H}_I$ is defined analogously to $\Sigma_{I,J}$. In this section, we assume in addition to the basic assumptions (A1-A3) that

(A10) All $V_{l,m}$ are compact and the joint correlation operator V is invertible.

Let \hat{I} be the indices of *active kernels* for the estimated $\hat{f} \in \mathcal{H}$ by elastic-net MKL: $\hat{I} := \{m \mid \|\hat{f}_m\|_{\mathcal{H}_m} > 0\}$. Let $D := \text{Diag}(\|f_m^*\|_{\mathcal{H}_m}^{-1}) = \text{Diag}((\|f_m^*\|_{\mathcal{H}_m}^{-1})_{m \in I_0})$, where Diag is the $|I_0| \times |I_0|$ block-diagonal operator with operators $\|f_m^*\|_{\mathcal{H}_m}^{-1} \mathbf{I}_{\mathcal{H}_m}$ on diagonal blocks for $m \in I_0$. In this section, we assume that the true sparsity pattern I_0 and the number of kernels M are fixed independently of the number of samples n. The norm of $f \in \mathcal{H}$ is defined by $\|f\|_{\mathcal{H}} := \sqrt{\sum_{m=1}^M \|f_m\|_{\mathcal{H}_m}^2}$ and similarly that of $f_I \in \mathcal{H}_I$ is de-

The norm of $f \in \mathcal{H}$ is defined by $||f||_{\mathcal{H}} := \sqrt{\sum_{m=1}^{M} ||f_m||_{\mathcal{H}_m}^2}$ and similarly that of $f_I \in \mathcal{H}_I$ is defined by $||f_I||_{\mathcal{H}_I} := \sqrt{\sum_{m \in I} ||f_m||_{\mathcal{H}_m}^2}$. The following theorem gives a sufficient condition for the support consistency of sparsity patterns.

Theorem 4 Suppose
$$\lambda_2^{(n)} > 0$$
, $\lambda_1^{(n)} \to 0$, $\lambda_2^{(n)} \to 0$, $\lambda_1^{(n)} \sqrt{n} \to \infty$, and

$$\lim \sup_n \left\| \Sigma_{m,I_0} (\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1} \left(D + 2\frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) f_{I_0}^* \right\|_{\mathcal{H}_m} < 1, \quad (\forall m \in J = I_0^c).$$
(15)

Then⁵, under assumptions (A1-A3, A10), $\|\hat{f} - f^*\|_{\mathcal{H}} \xrightarrow{p} 0$ and $\hat{I} \xrightarrow{p} I_0$.

The condition $\lambda_2^{(n)} > 0$ is just for technical simplicity to let $\Sigma_{I_0,I_0} + \lambda_2^{(n)}$ invertible. The condition $\lambda_1^{(n)}\sqrt{n} \to \infty$ means that $\lambda_1^{(n)}$ does not decrease too quickly. The condition (15) corresponds to an infinitedimensional extension of the elastic-net 'irrepresentable' condition. In the paper of Zhao and Yu (2006), the irrepresentable condition was derived as a necessary and sufficient condition for the sign consistency of ℓ_1 regularization when the number of parameters is finite. Its elastic-net version was derived in Yuan and Lin (2007), and it was extended to a situation where the number of parameters diverges as n increases (Jia and Yu, 2010).

We also have a necessary condition for consistency.

³ If one fits a function with a constant offset (f(x) + b instead of f(x)) as in Bach (2008), then the centered version of cross covariance operator is required instead of the non-centered version, i.e., $\langle f_m, \Sigma_{m,m'}g_{m'} \rangle_{\mathcal{H}_m} = \mathbb{E}_X[(f_m(X) - \mathbb{E}_X[f_m])(g_{m'}(X) - \mathbb{E}_X[g_{m'}])]$. However, this difference is not essential because, without loss of generality, one can consider a situation where $\mathbb{E}_Y[Y] = 0$ and $\mathbb{E}_X[f_m(X)] = 0$ for all $f_m \in \mathcal{H}_M$ by centering all the functions.

⁴ Actually, such a bounded operator always exists (Baker, 1973).

⁵For random variables x_n and $y, x_n \xrightarrow{p} y$ means the convergence in probability, i.e., the probability $|x_n - y| > \epsilon$ goes to 0 for all ϵ as the number of samples n tends to infinity.

Theorem 5 If $\|\hat{f} - f^*\|_{\mathcal{H}} \xrightarrow{p} 0$ and $\hat{I} \xrightarrow{p} I_0$, then under assumptions (A1-A3, A10), there exist sequences $\lambda_1^{(n)}, \lambda_2^{(n)} \to 0$ such that

$$\limsup_{n} \left\| \Sigma_{m,I_{0}} (\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)})^{-1} \left(D + 2\frac{\lambda_{2}^{(n)}}{\lambda_{1}^{(n)}} \right) f_{I_{0}}^{*} \right\|_{\mathcal{H}_{m}} \leq 1, \quad (\forall m \in J = I_{0}^{c}).$$
(16)

Moreover, such $\lambda_1^{(n)}$ satisfies $\lambda_1^{(n)}\sqrt{n} \to \infty$.

The sufficient condition (15) contains the strict inequality ('<'), while similar conditions for ordinary (non-block) ℓ_1 regularization or ordinary (non-block) elastic-net regularization contain the weak inequality (' \leq '). The strict inequality appears because each block contains multiple variables in group lasso and MKL (Bach, 2008).

The condition $\lambda_1^{(n)}\sqrt{n} \to \infty$ is necessary to impose the RKHS-norm convergence $\|\hat{f} - f^*\|_{\mathcal{H}} \stackrel{p}{\to} 0$. Roughly speaking, this means that the block- ℓ_1 regularization term should be stronger than the noise level to suppress fluctuations by noise.

It is worth noting that the conditions (15) and (16) are weaker than the condition for block- ℓ_1 MKL presented in Bach (2008); the block- ℓ_1 MKL irrepresentable condition is⁶

$$\begin{cases} \text{(Sufficient condition)} & \left\| \Sigma_{m,m}^{1/2} V_{m,I_0} V_{I_0,I_0}^{-1} Dg_{I_0}^* \right\|_{\mathcal{H}_m} < 1, \ (\forall m \in J), \\ \text{(Necessary condition)} & \left\| \Sigma_{m,m}^{1/2} V_{m,I_0} V_{I_0,I_0}^{-1} Dg_{I_0}^* \right\|_{\mathcal{H}_m} \le 1, \ (\forall m \in J). \end{cases}$$

$$(17)$$

This is because the group- ℓ_2 regularization term eases the singularity of the problem. Examples that elasticnets successfully estimate the true sparsity pattern, while ℓ_1 regularization fails in parametric situations can be found in Jia and Yu (2010).

5 Conclusions

We provided three novel theoretical results on the support consistency and convergence rate of elastic-net MKL.

- (i) Elastic-net MKL was shown to be support consistent under a milder condition than block- ℓ_1 MKL.
- (ii) A tighter convergence rate than existing bounds was derived for the situation where the truth is sparse.
- (iii) The convergence rates of block- ℓ_1 MKL, elastic-net MKL, and block- ℓ_2 MKL when the truth is near sparse were elucidated, and elastic-net MKL was shown to perform better when the *decrease rate* β is not large, or the condition of the problem is bad.

Based on our theoretical findings, we conclude that the use of elastic-net regularization is recommended for MKL.

Elastic-net MKL can be regarded as 'intermediate' between block- ℓ_1 MKL and block- ℓ_2 MKL. Another popular intermediate variant is block- ℓ_p MKL for $1 \le p \le 2$ (Kloft et al., 2009, Cortes et al., 2009). Elastic-net MKL and block- ℓ_p MKL are conceptually similar, but they have a notable difference: elastic-net MKL with $\lambda_1^{(n)} > 0$ tends to produce sparse solutions, while block- ℓ_p MKL with $1 always produces dense solutions (i.e., all combination coefficients of kernels are non-zero). Sparsity of elastic-net MKL would be advantageous when the true kernel combination is sparse, as we proved in this paper. However, when the true kernel combination between elastic-net MKL and block-<math>\ell_p$ MKL is not clear yet. This needs to be further investigated in the future work.

A Proofs of the theorems

For a function f on $\mathcal{X} \times \mathbb{R}$, we define $P_n f := \frac{1}{n} \sum_{i=1}^n f(x_i, y_i)$ and $Pf := \mathbb{E}_{X,Y}[f(X, Y)]$. For a function $f_I \in \mathcal{H}_I$, we define $\|f_I\|_{\ell_1}$ as $\|f_I\|_{\ell_1} := \sum_{m \in I} \|f_m\|_{\mathcal{H}_m}$ and for $f \in \mathcal{H}$ we write $\|f\|_{\ell_1} := \sum_{m=1}^M \|f_m\|_{\mathcal{H}_m}$. Similarly we define $\|f_I\|_{\ell_2}$ as $\|f_I\|_{\ell_2}^2 := \sum_{m \in I} \|f_m\|_{\mathcal{H}_m}^2$ for $f_I \in \mathcal{H}_I$ and for $f \in \mathcal{H}$ we write $\|f\|_{\ell_2}^2 := \sum_{m=1}^M \|f_m\|_{\mathcal{H}_m}^2$. We write $\max\{a, b\}$ as $a \lor b$.

⁶ Note that in the original paper by Bach (2008), the RHS of (17) is $\sum_{m \in I_0} ||f_m^*||_{\mathcal{H}_m}$ because the squared group- ℓ_1 regularizer $(\sum_m ||f_m||_{\mathcal{H}_m})^2$ was used. We can show that the squared formulation is actually equivalent to the non-squared formulation in the sense that there exists one-to-one correspondence between the two formulations.

Lemma 6 For all $I \subseteq \{1, \ldots, M\}$, we have

$$\|f\|_{L_2(\Pi)}^2 \ge (1 - \rho(I)^2)\kappa(I)(\sum_{m \in I} \|f_m\|_{L_2(\Pi)}^2).$$
(18)

Proof: For $J = I^c$, we have

$$Pf^{2} = \|f_{I}\|_{L_{2}(\Pi)}^{2} + 2\langle f_{I}, f_{J} \rangle_{L_{2}(\Pi)} + \|f_{J}\|_{L_{2}(\Pi)}^{2} \ge \|f_{I}\|_{L_{2}(\Pi)}^{2} - 2\rho(I)\|f_{I}\|_{L_{2}(\Pi)}\|f_{J}\|_{L_{2}(\Pi)} + \|f_{J}\|_{L_{2}(\Pi)}^{2} \\ \ge (1 - \rho(I)^{2})\|f_{I}\|_{L_{2}(\Pi)}^{2} \ge (1 - \rho(I)^{2})\kappa(I)(\sum_{m \in I} \|f_{m}\|_{L_{2}(\Pi)}^{2}),$$
(19)

where we used Schwarz's inequality in the last line.

The following lemma gives an upper bound of $\sum_{m=1}^{M} \|\hat{f}\|_{\mathcal{H}_m}$ that hold with a high probability. This is an extension of Theorem 1 of Koltchinskii and Yuan (2008). The proof is given in Appendix B.

Lemma 7 There exists a constant F depending on only L in (A1) such that, if $\lambda_1^{(n)} \ge F\sqrt{\frac{\log(Mn)}{n}}$, we have, for $r = \frac{\lambda_1^{(n)}}{\lambda_1^{(n)} \lor \lambda_1^{(n)}}$, with probability $1 - n^{-1}$,

$$\sum_{m=1}^{M} \|\hat{f}_m\|_{\mathcal{H}_m} \le M^{\frac{1-r}{2-r}} \left(3\sum_{m=1}^{M} \|f_m^*\|_{\mathcal{H}_m} + 3\sum_{m=1}^{M} \|f_m^*\|_{\mathcal{H}_m}^2\right)^{\frac{1}{2-r}}$$

Moreover, if $\lambda_2^{(n)} \ge F\sqrt{\frac{\log(Mn)}{n}}$ and $\lambda_2^{(n)} \ge \lambda_1^{(n)}$, we have

$$\sum_{m=1}^{M} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m} \le M \left(3/2 + 2 \max_m \|f_m^*\|_{\mathcal{H}_m} \right).$$

The following lemma gives a basic inequality that is a start point for the following analyses. The proof is given in Appendix B.

Lemma 8 Suppose $\lambda_1^{(n)} \vee \lambda_2^{(n)} \ge F\sqrt{\frac{\log(Mn)}{n}}$ where F is the constant appeared in Lemma 7. Then there exist constants \tilde{K}_1 and \tilde{K}_2 depending only on L in (A1), R in (A4), s in (A6), C_2 in (A6) such that for all $I \subseteq \{1, \ldots, M\}$, and for all $t \ge \log \log(R\sqrt{n}) + \log M$, with probability at least $1 - e^{-t} - n^{-1}$,

$$\frac{1}{2} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_2^{(n)} \sum_{m \in I} \|\hat{f}_I - f_I^*\|_{\mathcal{H}_m}^2 + \lambda_2^{(n)} \sum_{m \in J} \|\hat{f}_m\|_{\mathcal{H}_m}^2 + \left(\lambda_1^{(n)} - \hat{\gamma}_n - \tilde{K}_2 \sqrt{\frac{t}{n}}\right) \sum_{m \in J} \|\hat{f}_m\|_{\mathcal{H}_m}^2 \\
\leq \tilde{K}_1 (1 + \|\hat{f} - f^*\|_{\ell_1}) \Big(\sum_{m \in I} \frac{\|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^s}{\sqrt{n}} \vee \frac{\|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}}{n^{\frac{1}{1+s}}} + \frac{t\|\hat{f} - f^*\|_{\ell_1}}{n} \Big) \\
+ \sum_{m \in I} \left(\lambda_1^{(n)} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} + 2\lambda_2^{(n)} \|g_m^*\|_{\mathcal{H}_m} + \tilde{K}_2 \sqrt{\frac{t}{n}} \right) \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} \\
+ \lambda_2^{(n)} \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m}^2 + \left(\lambda_1^{(n)} + \hat{\gamma}_n + \tilde{K}_2 \sqrt{\frac{t}{n}} \right) \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m},$$
(20)
where $J = I^c, \gamma_n := \frac{\tilde{K}_1}{\sqrt{n}} and \hat{\gamma}_n := \gamma_n (1 + \|\hat{f} - f^*\|_\infty).$

The above lemma is derived by *peeling device* or *localization method*. Details of those techniques can be found in, for example, Bartlett et al. (2005), Koltchinskii (2006), Mendelson (2002), van de Geer (2000). **Proof:** (Theorem 1) Since $\lambda_1^{(n)} \ge F \sqrt{\frac{\log(Mn)}{n}}$, we can assume that the inequality (20) is satisfied with $I = I_0$. For notational simplicity, we suppose I denotes I_0 in this proof. In addition, since $\lambda_1^{(n)} \ge \lambda_2^{(n)}$, $\|\hat{f}\|_{\infty} \le \sum_{m=1}^{M} \|f^*\|_{\mathcal{H}_m} \le 3R$ (with probability $1 - n^{-1}$) by Lemma 7. Note that $\|f^*_m\|_{\mathcal{H}_m} = 0$ for all $m \in J = I^c = I^c_0$, and $\hat{\gamma}_n + \tilde{K}_2 \sqrt{\frac{t}{n}} \leq \max\{Kn^{-\frac{1}{2+s}} + \tilde{K}_2 \sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\} = \lambda_1^{(n)}$ by taking K sufficiently large. Therefore by the inequality (20), we have

$$\frac{1}{2} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_2^{(n)} \|\hat{f}_I - f_I^*\|_{\ell_2}^2 \le K_1 \Big(\sum_{m \in I} \frac{\|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^s}{\sqrt{n}} + \frac{t}{n} \Big) \\
+ \sum_{m \in I} \left(\lambda_1^{(n)} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} + 2\lambda_2^{(n)} \|g_m^*\|_{\mathcal{H}_m} + \tilde{K}_2 \sqrt{\frac{t}{n}} \right) \|\hat{f}_m - f_m^*\|_{L_2(\Pi)}, \quad (21)$$

where K_1 is $\tilde{K}_1(1+3R)$ (here we omitted the term $\sum_{m \in I} n^{-\frac{1}{1+s}} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}$ for simplicity. One can show that term is negligible).

By Hölder's inequality, the first term in the RHS of the above inequality can be bounded as

$$K_{1} \sum_{m \in I} \frac{\|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)}^{1-s} \|\hat{f}_{m} - f_{m}^{*}\|_{\mathcal{H}_{m}}^{s}}{\sqrt{n}} \leq K_{1} \frac{(\sum_{m \in I} \|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)})^{1-s} (\|\hat{f}_{I} - f_{I}^{*}\|_{\ell_{1}})^{s}}{\sqrt{n}} \\ \leq \sqrt{d} K_{1} \frac{(\sum_{m \in I} \|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)}^{2})^{\frac{1-s}{2}} (\|\hat{f}_{I} - f_{I}^{*}\|_{\ell_{2}})^{\frac{s}{2}}}{\sqrt{n}}.$$

Applying Young's inequality, the last term can be bounded by

$$\frac{K_{1}(\lambda_{2}^{(n)}/2)^{-\frac{s}{2}}\sqrt{d}}{\sqrt{n}}\left(\sum_{m\in I}\|\hat{f}_{m}-f_{m}^{*}\|_{L_{2}(\Pi)}^{2}\right)^{\frac{1-s}{2}}\times\left(\lambda_{2}^{(n)}/2\right)^{\frac{s}{2}}\left(\|\hat{f}_{I}-f_{I}^{*}\|_{\ell_{2}}^{2}\right)^{\frac{s}{2}} \\
\leq C(n^{-\frac{1}{2}}\sqrt{d}\lambda_{2}^{(n)^{-\frac{s}{2}}})^{\frac{2}{2-s}}\left(\sum_{m\in I}\|\hat{f}_{m}-f_{m}^{*}\|_{L_{2}(\Pi)}^{2}\right)^{\frac{1-s}{2-s}}+\frac{\lambda_{2}^{(n)}}{2}\|\hat{f}_{I}-f_{I}^{*}\|_{\ell_{2}}^{2} \\
\leq C[(1-\rho^{2}(I))\kappa(I)]^{-1}n^{-1}d\lambda_{2}^{(n)^{-s}}+\frac{(1-\rho^{2}(I))\kappa(I)}{8}\sum_{m\in I}\|\hat{f}_{m}-f_{m}^{*}\|_{L_{2}(\Pi)}^{2}+\frac{\lambda_{2}^{(n)}}{2}\|\hat{f}_{I}-f_{I}^{*}\|_{\ell_{2}}^{2} \\
\leq Cn^{-1}d\lambda_{2}^{(n)^{-s}}+\frac{1}{8}\|\hat{f}-f^{*}\|_{L_{2}(\Pi)}^{2}+\frac{\lambda_{2}^{(n)}}{2}\|\hat{f}_{I}-f_{I}^{*}\|_{\ell_{2}}^{2}.$$
(22)

where C denotes a constant that is independent of d and n and changes by the contexts, and we used Lemma 6 in the last line. Similarly, by the inequality of arithmetic and geometric means, we obtain a bound as

$$\sum_{m \in I} 2 \left(\lambda_1^{(n)} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} + \lambda_2^{(n)} \|g_m^*\|_{\mathcal{H}_m} + \tilde{K}_2 \sqrt{\frac{t}{n}} \right) \|\hat{f}_m - f_m^*\|_{L_2(\Pi)}$$

$$\leq C[(1 - \rho^2(I))\kappa(I)]^{-1} \sum_{m \in I} \left\{ \left(\frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} \right)^2 \lambda_1^{(n)^2} + \|g_m^*\|_{\mathcal{H}_m}^2 \lambda_2^{(n)^2} + \frac{t}{n} \right\}$$

$$+ \frac{(1 - \rho^2(I))\kappa(I)}{8} \sum_{m \in I} \|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^2$$

$$\leq C(d\lambda_1^{(n)^2} + \lambda_2^{(n)^2} + dt/n) + \frac{1}{8} \|\hat{f} - f^*\|_{L_2(\Pi)}^2, \qquad (23)$$

where we used Lemma 6 in the last line. By substituting (22) and (23) to (21), we have

$$\frac{1}{4} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 \le C \left(dn^{-1} \lambda_2^{(n)^{-s}} + d\lambda_1^{(n)^2} + \lambda_2^{(n)^2} + \frac{(d+1)t}{n} \right).$$
(24)

The minimum of the RHS with respect to $\lambda_1^{(n)}, \lambda_2^{(n)}$ under the constraint $\lambda_1^{(n)} \ge \lambda_2^{(n)}$ is achieved by $\lambda_1^{(n)} = \max\{Kn^{-\frac{1}{2+s}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\}, \lambda_2^{(n)} = Kn^{-\frac{1}{2+s}}$ up to constants. Thus we have the first assertion (7).

Next we show the second assertion (8). By Hölder's inequality and Young's inequality, we have

$$K_{1} \sum_{m \in I} \frac{\|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)}^{1-s} \|\hat{f}_{m} - f_{m}^{*}\|_{\mathcal{H}_{m}}^{s}}{\sqrt{n}} \leq K_{1} \frac{(\sum_{m \in I} \|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)})^{1-s} (\|\hat{f}_{I} - f_{I}^{*}\|_{\ell_{1}})^{s}}{\sqrt{n}}$$
$$\leq C\tilde{\lambda}^{-\frac{s}{1-s}} n^{-\frac{1}{2(1-s)}} \sum_{m \in I} \|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)} + \frac{\tilde{\lambda}}{2} \|\hat{f}_{I} - f_{I}^{*}\|_{\ell_{1}}$$
$$\leq Cd\tilde{\lambda}^{-\frac{2s}{1-s}} n^{-\frac{1}{1-s}} + \frac{1}{8} \|\hat{f} - f^{*}\|_{L_{2}(\Pi)}^{2} + \frac{\tilde{\lambda}}{2} (\|\hat{f}_{I}\|_{\ell_{1}} + \|f_{I}^{*}\|_{\ell_{1}}), \tag{25}$$

where $\tilde{\lambda} > 0$ is an arbitrary positive real. By substituting (25) and (23) to (21), we have

$$\frac{1}{4} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 \le C \Big(d\tilde{\lambda}^{-\frac{2s}{1-s}} n^{-\frac{1}{1-s}} + \tilde{\lambda} + d\lambda_1^{(n)^2} + \lambda_2^{(n)^2} + \frac{(d+1)t}{n} \Big).$$

nized by $\tilde{\lambda} = C d^{\frac{1-s}{1+s}} n^{-\frac{1}{1+s}}, \lambda_1^{(n)} = (\frac{2\tilde{K}_1(1+3R)}{\sqrt{n}} + \tilde{K}_2\sqrt{\frac{t}{n}}) \lor F\sqrt{\frac{\log(Mn)}{n}} \ge (2\hat{\gamma}_n + \tilde{K}_2\sqrt{\frac{t}{n}}) \lor$

This is minimized by $\tilde{\lambda} = Cd^{\frac{1-s}{1+s}}n^{-\frac{1}{1+s}}, \lambda_1^{(n)} = (\frac{2K_1(1+3R)}{\sqrt{n}} + \tilde{K}_2\sqrt{\frac{t}{n}}) \vee F\sqrt{\frac{\log(Mn)}{n}} \ge (2\hat{\gamma}_n + \tilde{K}_2\sqrt{\frac{t}{n}}) \vee F\sqrt{\frac{\log(Mn)}{n}}, \text{ and } \lambda_2^{(n)} \le \lambda_1^{(n)}.$ Thus we obtain the assertion.

Proof: (Theorem 2) Let $I_d := \{1, \ldots, d\}$ and $J_d = I_d^c = \{d + 1, \ldots, M\}$. By the assumption (A7), we have $\sum_{m \in J_d} \|f_m^*\|_{\mathcal{H}_m}^2 \leq \frac{C_3}{2\beta - 1} d^{1-2\beta}$, $\sum_{m \in J_d} \|f_m^*\|_{\mathcal{H}_m} \leq \frac{C_3}{\beta - 1} d^{1-\beta}$. Therefore Lemma 8 gives

$$\begin{split} &\|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_2^{(n)} \|\hat{f}_{I_d} - f_{I_d}^*\|_{\ell_2}^2 + \lambda_2^{(n)} \|\hat{f}_{J_d}\|_{\ell_2}^2 \\ \leq & K_1 \Big(\sum_{m \in I_d} \frac{\|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^s}{\sqrt{n}} + \frac{t\|\hat{f} - f^*\|_{\ell_1}}{n}\Big) \\ &+ K_1 \left(\sum_{m=1}^M \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}\right) \Big(\sum_{m \in I_d} \frac{\|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^s}{\sqrt{n}} + \frac{t\|\hat{f} - f^*\|_{\ell_1}}{n}\Big) \\ &+ \sum_{m \in I_d} \left(\lambda_1^{(n)} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} + 2\lambda_2^{(n)} \|g_m^*\|_{\mathcal{H}_m} + \tilde{K}_2 \sqrt{\frac{t}{n}}\right) \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} \\ &+ C \left(\lambda_2^{(n)} d^{1-2\beta} + \left(\lambda_1^{(n)} + \hat{\gamma}_n + \sqrt{\frac{t}{n}}\right) d^{1-\beta}\right), \end{split}$$
(26)

$$\begin{split} & K_1 \left(\sum_{m=1}^M \| \hat{f}_m - f_m^* \|_{\mathcal{H}_m} \right) \left(\sum_{m \in I_d} \frac{\| f_m - f_m^* \|_{L_2(\Pi)}^{1-s} \| f_m - f_m^* \|_{\mathcal{H}_m}}{\sqrt{n}} + \frac{t \| f - f^* \|_{\ell_1}}{n} \right) \\ \stackrel{\text{Hölder}}{\leq} & K_1 \left(\sum_{m=1}^M \| \hat{f}_m - f_m^* \|_{\mathcal{H}_m} \right) \left\{ \frac{(\sum_{m \in I_d} \| \hat{f}_m - f_m^* \|_{L_2(\Pi)})^{1-s} (\sum_{m \in I_d} \| \hat{f}_m - f_m^* \|_{\mathcal{H}_m})^s}{\sqrt{n}} + \frac{t \| \hat{f} - f^* \|_{\ell_1}}{n} \right\} \\ &= & K_1 \frac{(\sum_{m \in I_d} \| \hat{f}_m - f_m^* \|_{L_2(\Pi)})^{1-s} \left(\sum_{m=1}^M \| \hat{f}_m - f_m^* \|_{\mathcal{H}_m} \right) (\sum_{m \in I_d} \| \hat{f}_m - f_m^* \|_{\mathcal{H}_m})^s}{\sqrt{n}} + \frac{t \| \hat{f} - f^* \|_{\ell_1}^2}{n} \\ \stackrel{\text{Jensen}}{\leq} & K_1 \frac{d^{\frac{1-s}{2}} (\sum_{m \in I_d} \| \hat{f}_m - f_m^* \|_{L_2(\Pi)}^2)^{\frac{1-s}{2}} M^{\frac{1}{2}} \left(\sum_{m=1}^M \| \hat{f}_m - f_m^* \|_{\mathcal{H}_m}^2 \right)^{\frac{1}{2}} d^{\frac{s}{2}} (\sum_{m \in I_d} \| \hat{f}_m - f_m^* \|_{\mathcal{H}_m}^2)^{\frac{s}{2}}}{\sqrt{n}} \end{split}$$

$$+ \frac{t \|f - f^*\|_{\ell_1}^2}{n}$$

$$\begin{split} & \underset{\leq}{\text{Lemma } 6} K_1 \frac{\{(1-\rho(I_d)^2)\kappa(I_d)\}^{-\frac{1-s}{2}} (\|\hat{f}-f^*\|_{L_2(\Pi)}^2)^{\frac{1-s}{2}} d^{\frac{1}{2}} M^{\frac{1}{2}} \|\hat{f}-f^*\|_{\ell_2}^1}{\sqrt{n}} + \frac{t\|\hat{f}-f^*\|_{\ell_1}^2}{n} \\ & \stackrel{\text{Young }}{\leq} \frac{\|\hat{f}-f^*\|_{L_2(\Pi)}^2}{2} + C \frac{\{(1-\rho(I_d)^2)\kappa(I_d)\}^{-\frac{1-s}{1+s}} d^{\frac{1}{1+s}} M^{\frac{1}{1+s}} \|\hat{f}-f^*\|_{\ell_2}^2}{n^{\frac{1}{1+s}}} + \frac{t\|\hat{f}-f^*\|_{\ell_1}^2}{n} \\ & \stackrel{\text{(A8)}}{\leq} \frac{\|\hat{f}-f^*\|_{L_2(\Pi)}^2}{2} + C \frac{d^{\frac{b(1-s)+1}{1+s}} M^{\frac{1}{1+s}}}{n^{\frac{1}{1+s}}} \|\hat{f}-f^*\|_{\ell_2}^2 + \frac{t\|\hat{f}-f^*\|_{\ell_1}^2}{n}. \end{split}$$

We will see that we may assume $C \frac{d^{\frac{b(1-s)+1}{1+s}}M^{\frac{1}{1+s}}}{n^{\frac{1}{1+s}}} \leq \frac{\lambda_2^{(n)}}{4}$. Thus the second term in the RHS of the above inequality can be upper bounded as

$$C\frac{d^{\frac{b(1-s)+1}{1+s}}M^{\frac{1}{1+s}}}{n^{\frac{1}{1+s}}}\|\hat{f}-f^*\|_{\ell_2}^2 \leq \frac{\lambda_2^{(n)}}{4}\|\hat{f}-f^*\|_{\ell_2}^2 \leq \frac{\lambda_2^{(n)}}{4}\left(\|\hat{f}_{I_d}-f^*_{I_d}\|_{\ell_2}^2+2\|\hat{f}_{J_d}\|_{\ell_2}^2+2\|f^*_{J_d}\|_{\ell_2}^2\right)$$
$$\leq \frac{\lambda_2^{(n)}}{2}\left(\|\hat{f}_{I_d}-f^*_{I_d}\|_{\ell_2}^2+\|\hat{f}_{J_d}\|_{\ell_2}^2+\|f^*_{J_d}\|_{\ell_2}^2\right). \quad (27)$$

Moreover Lemma 7 gives $\frac{\|\hat{f}-f^*\|_{\ell_1}}{n} \leq \frac{C\sqrt{RM}}{n} \leq C\lambda_2^{(n)^2}$ and $\frac{\|\hat{f}-f^*\|_{\ell_1}^2}{n} \leq \frac{CRM}{n} \leq CR\lambda_2^{(n)^2}$. Therefore (26) becomes

$$\frac{1}{2} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \frac{\lambda_2^{(n)}}{2} \|\hat{f}_{I_d} - f^*_{I_d}\|_{\ell_2}^2 + \frac{\lambda_2^{(n)}}{2} \|\hat{f}_{J_d}\|_{\ell_2}^2 \\
\leq C \Big(\sum_{m \in I_d} \frac{\|\hat{f}_m - f^*_m\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f^*_m\|_{\mathcal{H}_m}^s}{\sqrt{n}} + t\lambda_2^{(n)^2} \Big) \\
+ \sum_{m \in I_d} \left(C_1 \lambda_1^{(n)} + 2\lambda_2^{(n)} \|g^*_m\|_{\mathcal{H}_m} + \tilde{K}_2 \sqrt{\frac{t}{n}} \right) \|\hat{f}_m - f^*_m\|_{L_2(\Pi)} \\
+ C \left(\lambda_2^{(n)} d^{1-2\beta} + \left(\lambda_1^{(n)} + \hat{\gamma}_n + \sqrt{\frac{t}{n}} \right) d^{1-\beta} \right).$$

As in the proof of Theorem 1 (using the relations (23) and (22)), we have

$$\begin{aligned} &\frac{1}{2} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 \\ \leq & C \bigg\{ [(1 - \rho^2(I_d))\kappa(I_d))]^{-1} \left[dn^{-1}\lambda_2^{(n)^{-s}} + d\lambda_1^{(n)^2} + \lambda_2^{(n)^2} + \frac{t}{n} \right] \\ &+ \lambda_2^{(n)} d^{1-2\beta} + (\lambda_1^{(n)} + \hat{\gamma}_n + (t/n)^{\frac{1}{2}}) d^{1-\beta} + t\lambda_2^{(n)^2} \bigg\}. \end{aligned}$$

Now using the assumption $(1 - \rho^2(I_d))\kappa(I_d) \ge C_4 d^{-b}$, we have

$$\|\hat{f}_{I_d} - f_{I_d}^*\|_{L_2(\Pi)}^2 \le C \left[d^{1+b} n^{-1} \lambda_2^{(n)} + d^{1+b} \lambda_1^{(n)} + d^b \lambda_2^{(n)} + \lambda_2^{(n)} d^{1-2\beta} + (\lambda_1^{(n)} + \hat{\gamma}_n) d^{1-\beta} + t \lambda_2^{(n)} + d^{1-\beta} \sqrt{\frac{t}{n}} + \frac{d^{1+b} t}{n} \right].$$
(28)

Remind that $\hat{\gamma}_n = \tilde{K}_1(1 + \|\hat{f} - f^*\|_{\infty})/\sqrt{n}$. Since $\lambda_1^{(n)} \ge F\sqrt{\frac{\log(Mn)}{n}}$, Lemma 7 gives $\|\hat{f} - f^*\|_{\infty} \le \sqrt{M3R} + R \le c\sqrt{M}$ with probability $1 - n^{-1}$ for some constant c > 0. Therefore $\hat{\gamma}_n \le c\sqrt{M/n}$. The values of $\lambda_1^{(n)}$, $\lambda_2^{(n)}$ presented in the statement is achieved by minimizing the RHS of Eq. (28) under the constraint $\lambda_1^{(n)} \ge c\sqrt{M/n} + \tilde{K}_2\sqrt{\frac{t}{n}} \ge \hat{\gamma}_n + \tilde{K}_2\sqrt{\frac{t}{n}}$ and $C\frac{d^{\frac{b(1-s)+1}{1+s}}M^{\frac{1}{1+s}}}{n^{\frac{1}{1+s}}} \le \frac{\lambda_2^{(n)}}{4}$.

i) Suppose $n^{-\frac{b+3\beta-1}{(2\beta+b)(2+s)-1-s}} > c\sqrt{M/n}$, i.e., $\tau \leq \tau_2$. Then the RHS of the above inequality can be minimized by $d = n^{\frac{1}{(2\beta+b)(2+s)-1-s}}$, $\lambda_2^{(n)} = Kn^{-\frac{2\beta+b-1}{(2\beta+b)(2+s)-1-s}}$, and $\lambda_1^{(n)} = \max\{Kn^{-\frac{b+3\beta-1}{(2\beta+b)(2+s)-1-s}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\}$ up to constants independent of n, where the leading terms are $d^{1+b}n^{-1}\lambda_2^{(n)-s} + d^b\lambda_2^{(n)^2} + \lambda_2^{(n)}d^{1-2\beta} + \lambda_1^{(n)}d^{1-\beta}$. It should be noted that $\lambda_1^{(n)}$ is greater than $\hat{\gamma}_n + \tilde{K}_2\sqrt{\frac{t}{n}}$ because $n^{-\frac{b+3\beta-1}{(2\beta+b)(2+s)-1-s}} > c\sqrt{M/n} \geq \hat{\gamma}_n$, therefore (26) is valid. Using $\tau \leq \tau_2$, we can show that $Cd^{\frac{b(1-s)+1}{1+s}}(M/n)^{\frac{1}{1+s}} \leq \lambda_2^{(n)}/4$ by setting the constant K sufficiently large, hence (27) is valid. Moreover, since $M > n^{\frac{(2\beta+b)(2+s)-1-s}{(2\beta+b)(2+s)-1-s}} = n^{\tau_1}$, we can take d as $d = n^{\frac{(2\beta+b)(2+s)-1-s}{(2\beta+b)(2+s)-1-s}} \leq M$.

ii) Suppose $\tau_2 \leq \tau \leq \tau_3$. Then the RHS of the above inequality can be minimized by $d = (M^{2+s}n^{2-s})^{\frac{1}{2\{(2+s)(b+\beta)-s\}}}$, $\lambda_2^{(n)} = K(Mn^{-\{2(b+\beta)-1\}})^{\frac{1}{2\{(2+s)(b+\beta-1)+2\}}}$, and $\lambda_1^{(n)} = \max\left\{c\sqrt{M/n} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\right\} \geq \hat{\gamma}_n + \tilde{K}_2\sqrt{\frac{t}{n}}$ up to constants independent of n, where the leading terms are $d^{1+b}n^{-1}\lambda_2^{(n)-s} + d^b\lambda_2^{(n)}^2 + \lambda_1^{(n)}d^{1-\beta}$. Since $\lambda_1^{(n)} \geq \hat{\gamma}_n + \tilde{K}_2\sqrt{\frac{t}{n}}$, (26) is valid. Using $\tau \leq \tau_3$, we can show that $Cd^{\frac{b(1-s)+1}{1+s}}(M/n)^{\frac{1}{1+s}} \leq \lambda_2^{(n)}/4$ by setting the constant K sufficiently large, hence (27) is valid. Moreover, since $\beta \leq \frac{s(b-1)}{2(1-s)}$ and $\tau_2 \leq \tau$, we can show that $d \leq M$.

iii) Suppose $\tau_3 \leq \tau \leq \tau_4$. We take $\lambda_1^{(n)} = \max\left\{c\sqrt{M/n} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\right\}$. Then the RHS of the inequality (28) is minimized by $\lambda_2^{(n)} = K\sqrt{d}\lambda_1^{(n)} \sim \sqrt{dM/n}$ and $d = (\frac{n}{M})^{\frac{1}{2(b+\beta)}}$ up to constants, where the leading terms are $d^b\lambda_2^{(n)^2} + d^{1+b}\lambda_1^{(n)^2} + \lambda_1^{(n)}d^{1-\beta}$. Note that since $\lambda_1^{(n)} \geq \hat{\gamma}_n + \tilde{K}_2\sqrt{\frac{t}{n}}$, (26) is valid. Using $\tau \leq \tau_4$, we can show that $Cd^{\frac{b(1-s)+1}{1+s}}(M/n)^{\frac{1}{1+s}} \leq \lambda_2^{(n)}/4$ by setting the constant K sufficiently large, hence (27) is valid. Moreover, since $\beta \leq \frac{s(b-1)}{2(1-s)}$ and $n^{\tau_3} \leq M$, we have $d = (\frac{n}{M})^{\frac{1}{2(b+\beta)}} \leq M$.

In all settings i) to iii), we can show that $\frac{d^{1-\beta}}{\sqrt{n}} \gtrsim \frac{d^{1+b}}{n}$. Thus the terms regarding t is upper bounded as $d^{1-\beta}\sqrt{\frac{t}{n}} + \frac{d^{1+b}t}{n} + t\lambda_2^{(n)^2} \lesssim (\frac{d^{1-\beta}}{\sqrt{n}} + \lambda_2^{(n)^2})(\sqrt{t} + t)$. Through a simple calculation $\frac{d^{1-\beta}}{\sqrt{n}}$ is evaluated as i) $\frac{d^{1-\beta}}{\sqrt{n}} \simeq n^{-\frac{(2\beta+b)(2+s)-3-s+2\beta}{2\{(2\beta+b)(2+s)-1-s\}}}$, ii) $\frac{d^{1-\beta}}{\sqrt{n}} \simeq (M^{(2+s)(1-\beta)}n^{-(4\beta+2b+sb-2)})^{\frac{1}{2\{(\beta+b)(2+s)-s\}}}$, and iii) $\frac{d^{1-\beta}}{\sqrt{n}} \simeq (M^{\beta-1}n^{1-2\beta-b})^{\frac{1}{2(\beta+b)}}$ respectively. Thus we obtain the assertion.

Proof: (Theorem 3)

(Convergence rate of block- ℓ_1 MKL)

Note that since $\lambda_1^{(n)} > \lambda_2^{(n)} = 0$, we have $\frac{\lambda_1^{(n)}}{\lambda_1^{(n)} \vee \lambda_2^{(n)}} = 1$. Therefore Lemma 7 gives $\sum_{m=1}^M \|\hat{f}_m\|_{\mathcal{H}_m} \le 3R$ with probability $1 - n^{-1}$. Thus $\hat{\gamma}_n = \gamma_n (1 + \|\hat{f} - f^*\|_{\infty}) \le \gamma_n (1 + \sum_{m=1}^M \|\hat{f}_m\|_{\mathcal{H}_m} + \sum_{m=1}^M \|f_m^*\|_{\mathcal{H}_m}) \le \gamma_n (1 + 4R)$.

When $\lambda_2^{(n)} = 0$ and $\lambda_1^{(n)} > (1 + 4R)\gamma_n + \tilde{K}_2\sqrt{\frac{t}{n}}$, as in Lemma 8 we have with probability at least $1 - e^{-t} - n^{-1}$

$$\begin{aligned} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_1^{(n)} \sum_{m \in I} \|\hat{f}_m\|_{\mathcal{H}_m} \\ \leq K_1 \Big(\sum_{m \in I} \frac{\|\hat{f}_m - f^*_m\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f^*_m\|_{\mathcal{H}_m}^s}{\sqrt{n}} + \frac{t}{n} \Big) + \lambda_1^{(n)} \sum_{m \in I} \|f^*_m\|_{\mathcal{H}_m} + 2\lambda_1^{(n)} \sum_{m \in J} \|f^*_m\|_{\mathcal{H}_m} \\ + \tilde{K}_2 \sum_{m \in I} \sqrt{\frac{t}{n}} \|f^*_m - \hat{f}_m\|_{L_2(\Pi)}, \end{aligned}$$

$$(29)$$

for all $t \ge \log \log(R\sqrt{n}) + \log M$.

We lower bound the term $\lambda_1^{(n)} \sum_{m \in I} (\|\hat{f}_m\|_{\mathcal{H}_m} - \|f_m^*\|_{\mathcal{H}_m})$ in the LHS of the above inequality (21). There exists $c_1 > 0$ only depending R such that

$$\|f_m\|_{\mathcal{H}_m} = \sqrt{\|f_m - f_m^*\|_{\mathcal{H}_m}^2 - 2\langle f_m - f_m^*, f_m^* \rangle_{\mathcal{H}_m} + \|f_m^*\|_{\mathcal{H}_m}^2} \\ \ge c_1 \|f_m - f_m^*\|_{\mathcal{H}_m}^2 - 2\|f_m^*\|_{\mathcal{H}_m}^{-1}|\langle f_m - f_m^*, f_m^* \rangle_{\mathcal{H}_m}| + \|f_m^*\|_{\mathcal{H}_m}$$
(30)

for all $f_m \in \mathcal{H}_m$ such that $||f_m||_{\mathcal{H}_m} \leq 3R$ and $m \in I_0$. Remind that $f_m^* = T_m^{1/2} g_m^*$, then we have $||f_m||_{\mathcal{H}_m} \geq c_1 ||f_m - f_m^*||_{\mathcal{H}_m}^2 - 2 \frac{||g_m^*||_{\mathcal{H}_m}}{||f_m^*||_{\mathcal{H}_m}} ||f_m - f_m^*||_{L_2(\Pi)} + ||f_m^*||_{\mathcal{H}_m}$. Since $\max_m ||\hat{f}_m||_{\mathcal{H}_m} \leq 3R$ are met with probability $1 - n^{-1}$,

$$\|\hat{f}_m\|_{\mathcal{H}_m} \ge c_1 \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^2 - 2\frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} + \|f_m^*\|_{\mathcal{H}_m},$$

with probability $1 - n^{-1}$.

Therefore by the inequality (29), we have with probability at least $1 - e^{-t} - n^{-1}$

$$\begin{split} &\|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_1^{(n)} \sum_{m \in I} (c_1 \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^2 - 2 \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} + \|f_m^*\|_{\mathcal{H}_m}) \\ &\leq K_1 \Big(\sum_{m \in I} \frac{\|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^s}{\sqrt{n}} + \frac{t}{n} \Big) + \lambda_1^{(n)} \sum_{m \in I} \|f_m^*\|_{\mathcal{H}_m} + 2\lambda_1^{(n)} \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m} \\ &+ \tilde{K}_2 \sum_{m \in I} \sqrt{\frac{t}{n}} \|f_m^* - \hat{f}_m\|_{L_2(\Pi)}, \end{split}$$
(31)

for all $t \ge \log \log(R\sqrt{n}) + \log M$. Thus using Young's inequality

$$\|\hat{f} - f^*\|_{L_2(\Pi)}^2 \le C \left[d^{1+b} n^{-1} \lambda_1^{(n)^{-s}} + d^{1+b} \lambda_1^{(n)^2} + 2\lambda_1^{(n)} d^{1-\beta} + \frac{t(1+d^{1+b})}{n} \right].$$

The RHS is minimized by $d = n^{\frac{1}{(2+s)(\beta+b)}}$ and $\lambda_1^{(n)} = \max\left\{Kn^{-\frac{1}{2+s}} + \tilde{K}_2\sqrt{\frac{t}{n}}, F\sqrt{\frac{\log(Mn)}{n}}\right\}$ (up to constants independent of n). Note that since the optimal $\lambda_1^{(n)}$ obtained above satisfies $\lambda_1^{(n)} > (1+4R)\gamma_n + \tilde{K}_2\sqrt{\frac{t}{n}}$ by taking K sufficiently large, the inequality (31) is valid. Moreover the condition $M > n^{\tau_5} = n^{\frac{b+1}{(\beta+b)(b(2+s)+2)}}$ in the statement ensures d < M. Finally we evaluate the terms including t, that is, $\frac{t}{n}d^{1+b} + \sqrt{\frac{t}{n}}d^{1-\beta}$. We can check that $\frac{1}{n}d^{1+b} \lesssim \sqrt{\frac{1}{n}}d^{1-\beta}$. Therefore those terms are upper bounded as $\frac{t}{n}d^{1+b} + \sqrt{\frac{t}{n}}d^{1-\beta} \lesssim \sqrt{\frac{1}{n}}d^{1-\beta}(\sqrt{t}+t) \simeq n^{-\frac{4\beta+2b-2+s(b+\beta)}{2(2+s)(b+\beta)}}(\sqrt{t}+t)$. Thus we obtain the assertion.

(Convergence rate for block- ℓ_2 MKL)

When $\lambda_1^{(n)} = 0$, substituting I_M to I in Lemma 8, and using Young's inequality, as in the proof of Theorem 2, the convergence rate of block- ℓ_2 MKL can be evaluated as

$$\|\hat{f}_{I_d} - f_{I_d}^*\|_{L_2(\Pi)}^2 \le C \left[M^{1+b} n^{-1} \lambda_2^{(n)^{-s}} + M^b \lambda_2^{(n)^2} + t \lambda_2^{(n)^2} + \frac{t}{n} M^{1+b} \right],$$
(32)

with probability $1 - e^{-t} - n^{-1}$ (note that since $I = \{1, \ldots, M\}$ ($I^c = \emptyset$), we don't need the condition $\lambda_1^{(n)} \ge \hat{\gamma}_n + \tilde{K}_2 \sqrt{\frac{t}{n}}$). $\lambda_2^{(n)} = K(\frac{M}{n})^{\frac{1}{2+s}} \lor F \sqrt{\frac{\log(Mn)}{n}}$ gives the minimum of the RHS with respect to $\lambda_2^{(n)}$ up to constants. Using $\tau \le \tau_6$, we can show that $M^{\frac{b(1-s)+1}{1+s}}(M/n)^{\frac{1}{1+s}} = M^{\frac{b(1-s)+2}{1+s}}n^{-\frac{1}{1+s}} \lesssim \lambda_2^{(n)}$ by setting the constant K sufficiently large, hence (27) is valid.

B Proof of Lemmas 7 and 8

Proof: (Lemma 7) Since \hat{f} minimizes the empirical risk (1), we have

$$\frac{1}{n}\sum_{i=1}^{n}\left(\sum_{m=1}^{M}(\hat{f}_{m}(x_{i})-f_{m}^{*}(x_{i}))\right)^{2}+\lambda_{1}^{(n)}\|\hat{f}\|_{\ell_{1}}+\lambda_{2}^{(n)}\|\hat{f}\|_{\ell_{2}}^{2} \leq \frac{2}{n}\sum_{m=1}^{M}\sum_{i=1}^{n}\epsilon_{i}(\hat{f}_{m}(x_{i})-f_{m}^{*}(x_{i}))+\lambda_{1}^{(n)}\|f^{*}\|_{\ell_{1}}+\lambda_{2}^{(n)}\|f^{*}\|_{\ell_{2}}^{2}.$$
(33)

By Proposition 1 (Bernstein's inequality in Hilbert spaces, see also Theorem 6.14 of Steinwart (2008) for example), there exists a universal constant C such that we have

$$\frac{1}{n} \sum_{i=1}^{n} \epsilon_{i} (\hat{f}_{m}(x_{i}) - f_{m}^{*}(x_{i})) \leq \left| \frac{1}{n} \sum_{i=1}^{n} \epsilon_{i} k_{m}(x_{i}, \cdot) \right| \|\hat{f}_{m} - f_{m}^{*}\|_{\mathcal{H}_{m}} \\
\leq CL \sqrt{\frac{\log(Mn)}{n}} \|\hat{f}_{m} - f_{m}^{*}\|_{\mathcal{H}_{m}} \leq CL \sqrt{\frac{\log(Mn)}{n}} (\|\hat{f}_{m}\|_{\mathcal{H}_{m}} + \|f_{m}^{*}\|_{\mathcal{H}_{m}})$$
(34)

for all m with probability at least $1 - n^{-1}$, where we used the assumption $\frac{\log(Mn)}{n} \leq 1$. If $\lambda_1^{(n)} \geq 4CL\sqrt{\frac{\log(Mn)}{n}}$, then we have

$$\lambda_1^{(n)} \|\hat{f}\|_{\ell_1} + \lambda_2^{(n)} \|\hat{f}\|_{\ell_2}^2 \le 3(\lambda_1^{(n)} \lor \lambda_2^{(n)})(\|f^*\|_{\ell_1} + \|f^*\|_{\ell_2}^2), \tag{35}$$

with probability at least $1 - n^{-1}$. Set $r = \frac{\lambda_1^{(n)}}{\lambda_1^{(n)} \vee \lambda_2^{(n)}}$, then by Young's inequality and Jensen's inequality, the LHS of the above inequality (33) is lower bounded by

$$\lambda_{1}^{(n)} \|\hat{f}\|_{\ell_{1}} + \lambda_{2}^{(n)} \|\hat{f}\|_{\ell_{2}}^{2} \geq (\lambda_{1}^{(n)} \vee \lambda_{2}^{(n)}) (\sum_{m=1}^{M} \|\hat{f}_{m}\|_{\mathcal{H}_{m}}^{2-r})$$

$$\geq M(\lambda_{1}^{(n)} \vee \lambda_{2}^{(n)}) \left(\frac{1}{M} \sum_{m=1}^{M} \|\hat{f}_{m}\|_{\mathcal{H}_{m}}^{2-r}\right)$$

$$\geq M^{r-1} (\lambda_{1}^{(n)} \vee \lambda_{2}^{(n)}) \|\hat{f}\|_{\ell_{1}}^{2-r}.$$
(36)

Therefore we have the first assertion by setting F = 4CL.

The second assertion can be shown as follows: by the inequality (33) we have

$$M^{-1}\lambda_{2}^{(n)}\left(\|\hat{f}-f^{*}\|_{\ell_{1}}\right)^{2} \leq \lambda_{2}^{(n)}\|\hat{f}-f^{*}\|_{\ell_{2}}^{2}$$

$$\leq \frac{2}{n}\sum_{m=1}^{M}\sum_{i=1}^{n}\epsilon_{i}(\hat{f}_{m}(x_{i})-f^{*}_{m}(x_{i}))+\lambda_{1}^{(n)}\|\hat{f}-f^{*}\|_{\ell_{1}}+2\lambda_{2}^{(n)}\sum_{m=1}^{M}\langle f^{*}_{m},f^{*}_{m}-\hat{f}_{m}\rangle_{\mathcal{H}_{m}}$$

$$\leq \lambda_{2}^{(n)}\left(\frac{3}{2}+2\max_{m}\|f^{*}_{m}\|_{\mathcal{H}_{m}}\right)\|\hat{f}-f^{*}\|_{\ell_{1}}$$
(37)

with probability at least $1 - n^{-1}$, where we used (34), $\lambda_2^{(n)} \ge 4CL\sqrt{\frac{\log(Mn)}{n}}$ and $\lambda_2^{(n)} \ge \lambda_1^{(n)}$ in the last inequality.

Proof: (Lemma 8) In what follows, we assume $\|\hat{f} - f^*\|_{\ell_1} \leq \bar{R}$ where $\bar{R} = 4MR$ (the probability of this event is greater than $1 - n^{-1}$ by Lemma 7). Since \hat{f} minimizes the empirical risk we have

$$P_{n}(\hat{f}-Y)^{2} + \lambda_{1}^{(n)} \|\hat{f}\|_{\ell_{1}} + \lambda_{2}^{(n)} \|\hat{f}\|_{\ell_{2}}^{2} \leq P_{n}(f^{*}-Y)^{2} + \lambda_{1}^{(n)} \|f^{*}\|_{\ell_{1}} + \lambda_{2}^{(n)} \|f^{*}\|_{\ell_{2}}^{2}$$

$$\Rightarrow P(\hat{f}-f^{*})^{2} + \lambda_{1}^{(n)} \|\hat{f}_{J}\|_{\ell_{1}} + \lambda_{2}^{(n)} \|\hat{f}_{J}\|_{\ell_{2}}^{2} \leq (P-P_{n})((f^{*}-\hat{f})^{2} + 2(\hat{f}-f^{*})\epsilon) + \lambda_{1}^{(n)}(\|f^{*}_{I}\|_{\ell_{1}} - \|\hat{f}_{I}\|_{\ell_{1}}) + \lambda_{2}^{(n)}(\|f^{*}_{I}\|_{\ell_{2}}^{2} - \|\hat{f}_{I}\|_{\ell_{2}}^{2}) + \lambda_{1}^{(n)} \|f^{*}_{J}\|_{\ell_{1}} + \lambda_{2}^{(n)} \|f^{*}_{J}\|_{\ell_{2}}^{2}.$$
(38)

The second term in the RHS of the above inequality (38) can be bounded from above as

$$(\|f_{I}^{*}\|_{\ell_{1}} - \|\hat{f}_{I}\|_{\ell_{1}}) \leq \sum_{m \in I} \langle \nabla \|f_{m}^{*}\|_{\mathcal{H}_{m}}, \hat{f}_{m} - f_{m}^{*} \rangle_{\mathcal{H}_{m}}$$
$$= \sum_{m \in I} \frac{\langle g_{m}^{*}, T_{m}^{1/2}(\hat{f}_{m} - f_{m}^{*}) \rangle_{\mathcal{H}_{m}}}{\|f_{m}^{*}\|_{\mathcal{H}_{m}}} \leq \sum_{m \in I} \frac{\|g_{m}^{*}\|_{\mathcal{H}_{m}}}{\|f_{m}^{*}\|_{\mathcal{H}_{m}}} \|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)}, \qquad (39)$$

where we used $f_m^* = T_m^{1/2} g_m^*$ for $m \in I \subseteq I_0$. We also have

$$\lambda_{2}^{(n)}(\|f_{I}^{*}\|_{\ell_{2}}^{2} - \|\hat{f}_{I}\|_{\ell_{2}}^{2}) = \lambda_{2}^{(n)}(\sum_{m \in I} 2\langle f_{m}^{*}, f_{m}^{*} - \hat{f}_{m} \rangle_{\mathcal{H}_{m}} - \|\hat{f}_{I} - f_{I}^{*}\|_{\ell_{2}}^{2})$$

$$\leq \lambda_{2}^{(n)}(\sum_{m \in I} 2\|g_{m}^{*}\|_{\mathcal{H}_{m}}\|\hat{f}_{m} - f_{m}^{*}\|_{L_{2}(\Pi)} - \|\hat{f}_{I} - f_{I}^{*}\|_{\ell_{2}}^{2}).$$
(40)

Substituting (39) and (40) to (38), we obtain

$$\begin{aligned} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_2^{(n)} \|\hat{f}_I - f_I^*\|_{\ell_2}^2 + \lambda_1^{(n)} \|\hat{f}_J\|_{\ell_1} + \lambda_2^{(n)} \|\hat{f}_J\|_{\ell_2}^2 \\ \leq (P - P_n)((f^* - \hat{f})^2 + 2(\hat{f} - f^*)\epsilon) + \sum_{m \in I} (\lambda_1^{(n)} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} + 2\lambda_2^{(n)} \|g_m^*\|_{\mathcal{H}_m}) \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} \\ + \lambda_1^{(n)} \|f_J^*\|_{\ell_1} + \lambda_2^{(n)} \|f_J^*\|_{\ell_2}^2. \end{aligned}$$

$$(41)$$

Finally we evaluate the first term $(P-P_n)((f^*-\hat{f})^2+2(\hat{f}-f^*)\epsilon)$ in the RHS of the above inequality (41) by applying Talagrand's concentration inequality (Talagrand, 1996a,b, Bousquet, 2002). First we decompose $(P-P_n)((f^*-\hat{f})^2+2(\hat{f}-f^*)\epsilon)$ as

$$(P - P_n)((f^* - \hat{f})^2 + 2(\hat{f} - f^*)\epsilon) = \sum_{m=1}^M (P - P_n)((f^* - \hat{f})(f^*_m - \hat{f}_m) + 2(\hat{f}_m - f^*_m)\epsilon),$$

and bound each term $(P - P_n)((f^* - \hat{f})(f_m^* - \hat{f}_m) + 2(\hat{f}_m - f_m^*)\epsilon)$ in the summation. Here suppose $f \in \mathcal{H}$ satisfies $\|f\|_{\infty} \leq \|f\|_{\ell_1} \leq \hat{R}$ for a constant $\hat{R} (\leq \bar{R})$. Since $|\epsilon| \leq L$, we have

$$\begin{aligned} |ff_m + 2f_m\epsilon| &\leq 2(L+R)|f| \leq 2(L+R)||f_m||_{\mathcal{H}_m}, \\ \sqrt{P(ff_m + 2f_m\epsilon)^2} &= \sqrt{P(f^2f_m^2) + 4P(f_m^2\epsilon^2)} \leq \sqrt{||f||_{L_2(\Pi)}^2 ||f_m||_{L_2(\Pi)}^2 + 4L^2 ||f_m||_{L_2(\Pi)}} \\ &\leq ||f||_{L_2(\Pi)} ||f_m||_{L_2(\Pi)} + 2L ||f_m||_{L_2(\Pi)}, \end{aligned}$$
(42a)

for all $f \in \mathcal{H}$. Let $Q_n f := \frac{1}{n} \sum_{i=1}^n \varepsilon_i f(x_i, y_i)$ where $\{\varepsilon_i\}_{i=1}^n \in \{\pm 1\}^n$ is the Rademacher random variable, and $\Psi_m(\xi_m, \sigma_m)$ be

$$\Psi_m(\xi_m, \sigma_m) := \mathbb{E}[\sup\{Q_n(|f_m|) \mid f_m \in \mathcal{H}_m, \|f_m\|_{\mathcal{H}_m} \le \xi_m, \|f_m\|_{L_2(\Pi)} \le \sigma_m\}].$$

Then one can show that by the spectral assumptions (A5) (equivalently the covering number condition)

$$\Psi_m(\xi_m, \sigma_m) \le K_s \left(\frac{\sigma_m^{1-s}\xi_m^s}{\sqrt{n}} \vee n^{-\frac{1}{1+s}}\xi_m\right)$$

where K_s is a constant that depends on s and C_2 (Mendelson, 2002). Let $\Xi_m(\xi_m, \sigma_m) := \{f_m \in \mathcal{H}_m \mid \|f_m\|_{\mathcal{H}_m} \leq \xi_m, \|f_m\|_{L_2(\Pi)} \leq \sigma_m\}$. Now by Rademacher contraction inequality (Ledoux and Talagrand, 1991, Theorem 4.12), for given $\{\xi_m, \sigma_m\}_{m \in I}$ and \hat{R} we have

$$\mathbb{E}[\sup\{Q_n(ff_m + 2f_m\epsilon) \mid f \in \mathcal{H} \text{ such that } f_m \in \Xi_m(\xi_m, \sigma_m), \ \|f\|_{\ell_1} \le \hat{R}\}]$$

$$\le 2(L + \hat{R})\Psi_m(\xi_m, \sigma_m) \le 2K_s(L + \hat{R})\left(\frac{\sigma_m^{1-s}\xi_m^s}{\sqrt{n}} \lor n^{-\frac{1}{1+s}}\xi_m\right).$$

$$(43)$$

Therefore by the symmetrization argument (van der Vaart and Wellner, 1996), we have

$$\mathbb{E}[\sup\{(P_n - P)(ff_m + 2f_m\epsilon) \mid f \in \mathcal{H} \text{ such that } f_m \in \Xi_m(\xi_m, \sigma_m), \ \|f\|_{\ell_1} \le \hat{R}\}]$$

$$\le 4K_s(L + \hat{R}) \left(\frac{\sigma_m^{1-s}\xi_m^s}{\sqrt{n}} \lor n^{-\frac{1}{1+s}}\xi_m\right).$$

$$(44)$$

By Talagrand's concentration inequality with (42) and (44), for given $\hat{R}, \bar{\sigma}, \xi_m, \sigma_m$ with probability at least $1 - e^{-t}$ (t > 0), we have

$$\sup_{\substack{f \in \mathcal{H}:\\ \|f\|_{L_{2}(\Pi)} \leq \bar{\sigma}, \|f\|_{\infty} \leq \hat{\kappa}, f_{m} \in \Xi_{m}(\xi_{m}, \sigma_{m})}} (P_{n} - P)(ff_{m} + 2f_{m}\epsilon) \leq \sqrt{2} \left(4K_{s}(L + \hat{R}) \left(\frac{\sigma_{m}^{1-s}\xi_{m}^{s}}{\sqrt{n}} \lor \frac{\xi_{m}}{n^{\frac{1}{1+s}}} \right) + \sqrt{\frac{t}{n}}(\bar{\sigma}\sigma_{m} + 2L\sigma_{m}) + 2(L + \hat{R})\xi_{m}\frac{t}{n} \right).$$
(45)

where we used the relation (42). Our next goal is to derive an uniform version of the above inequality over

$$\frac{1}{\sqrt{n}} \leq \hat{R} \leq \bar{R}, \quad \frac{1}{\sqrt{n}} \leq \bar{\sigma} \leq \bar{R}, \quad \frac{1}{\sqrt{n}M} \leq \xi_m \leq \bar{R} \quad \text{and} \quad \frac{1}{\sqrt{n}M} \leq \sigma_m \leq \bar{R}.$$

By considering a grid $\{\hat{R}^{(k_1)}, \bar{\sigma}^{(k_2)}, \xi_m^{(k_3)}, \sigma_m^{(k_4)}\}_{k_i=0(i=1,...,4)}^{\log_2(M\bar{R}\sqrt{n})}$ such that $\hat{R}^{(k)} := \bar{R}2^{-k}$, $\bar{\sigma}^{(k)} := \bar{R}2^{-k}$, $\bar{\sigma}^{(k)} := \bar{R}2^{-k}$, $\bar{\sigma}^{(k)} := \bar{R}2^{-k}$, we have with probability at least $1 - (\log(M\bar{R}\sqrt{n}))^4 e^{-t} \ge 1 - (\log(4RM^2\sqrt{n}))^4 e^{-t}$

$$(P_n - P)(ff_m + 2f_m\epsilon) \leq K(1 + ||f||_{\ell_1}) \left(\frac{||f_m||_{L_2(\Pi)}^{1-s}||f_m||_{\mathcal{H}_m}}{\sqrt{n}} \vee \frac{||f_m||_{\mathcal{H}_m}}{n^{\frac{1}{1+s}}} + \frac{t||f_m||_{\mathcal{H}_m}}{n} \right) + \sqrt{\frac{2t}{n}} (||f||_{L_2(\Pi)} ||f_m||_{L_2(\Pi)} + 2L||f_m||_{L_2(\Pi)}),$$

for all $f \in \mathcal{H}$ such that $||f_m||_{\mathcal{H}_m} \leq \overline{R}$ and $||f||_{\ell_1} \leq \overline{R}$, and for all t > 1, where $K = 4(4K_sL \vee 4K_s \vee 2L \vee 2)$. Summing up this bound for m = 1, ..., M, then we obtain

$$(P_n - P)(f^2 + 2f\epsilon) \leq K(1 + ||f||_{\ell_1}) \left(\sum_{m=1}^M \frac{||f_m||_{L_2(\Pi)}^{1-s} ||f_m||_{\mathcal{H}_m}}{\sqrt{n}} \vee \frac{||f_m||_{\mathcal{H}_m}}{n^{\frac{1}{1+s}}} + \frac{t||f||_{\ell_1}}{n} \right) + \sqrt{\frac{2t}{n}} \left(||f||_{L_2(\Pi)} \sum_{m=1}^M ||f_m||_{L_2(\Pi)} + 2L \sum_{m=1}^M ||f_m||_{L_2(\Pi)} \right),$$

uniformly for all $f \in \mathcal{H}$ such that $||f_m||_{\mathcal{H}_m} \leq \bar{R} \; (\forall m)$ and $||f||_{\ell_1} \leq \bar{R}$ with probability at least $1 - M(\log(4RM^2\sqrt{n}))^4 e^{-t}$. Here set $\gamma_n = \frac{K}{\sqrt{n}}$ and note that $\sqrt{\frac{2t}{n}} ||f||_{L_2(\Pi)} \sum_{m=1}^M ||f_m||_{L_2(\Pi)} \leq \frac{1}{2} ||f||_{L_2(\Pi)}^2 + \frac{t}{n} (\sum_{m=1}^M ||f_m||_{L_2(\Pi)})^2 \leq \frac{1}{2} ||f||_{L_2(\Pi)}^2 + \frac{t}{n} (||f||_{\ell_1})^2$ then we have

$$(P_n - P)(f^2 + 2f\epsilon) \leq K(1 + ||f||_{\ell_1}) \left[\sum_{m \in I} \left(\frac{||f_m||_{L_2(\Pi)}^{1-s} ||f_m||_{\mathcal{H}_m}}{\sqrt{n}} \vee \frac{||f_m||_{\mathcal{H}_m}}{n^{\frac{1}{1+s}}} \right) + \frac{2t ||f||_{\ell_1}}{n} \right] + \gamma_n (1 + ||f||_{\ell_1}) ||f_J||_{\ell_1} + \frac{1}{2} ||f||_{L_2(\Pi)}^2 + 2\sqrt{2}L \sqrt{\frac{t}{n}} \sum_{m=1}^M ||f_m||_{L_2(\Pi)}.$$
(46)

for all $f \in \mathcal{H}$ such that $||f_m||_{\mathcal{H}_m} \leq \bar{R} \; (\forall m)$ and $||f||_{\ell_1} \leq \bar{R}$ with probability at least $1 - M(\log(4RM^2\sqrt{n}))^4e^{-t}$. We will replace t with $t + 5\log M + 4\log\log(R\sqrt{n})$, then the probability $1 - M(\log(4R\sqrt{n}M^2))^4e^{-t}$ can be replaced with $1 - e^{-t}$ and we have $t + 5\log M + 4\log\log(R\sqrt{n}) \leq 6t$ for all $t \geq \log M + \log\log(R\sqrt{n})$. On the event where $||\hat{f} - f^*||_{\ell_1} \leq \bar{R}$ holds, substituting $\hat{f} - f^*$ to f in (46) and replacing K appropriately, (41) yields

$$\frac{1}{2} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_2^{(n)} \sum_{m \in I} \|\hat{f}_I - f_I^*\|_{\mathcal{H}_m}^2 + \lambda_2^{(n)} \sum_{m \in J} \|\hat{f}_m\|_{\mathcal{H}_m}^2 + (\lambda_1^{(n)} - \hat{\gamma}_n) \sum_{m \in J} \|\hat{f}_m\|_{\mathcal{H}_m} \\
\leq \tilde{K}_1 (1 + \|\hat{f} - f^*\|_{\ell_1}) \Big(\sum_{m \in I} \frac{\|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^s}{\sqrt{n}} \vee \frac{\|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}}{n^{\frac{1}{1+s}}} + \frac{t\|\hat{f} - f^*\|_{\ell_1}}{n} \Big) \\
+ \sum_{m \in I} \Big(\lambda_1^{(n)} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} + 2\lambda_2^{(n)} \|g_m^*\|_{\mathcal{H}_m} \Big) \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} + \lambda_2^{(n)} \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m}^2 + (\lambda_1^{(n)} + \hat{\gamma}_n) \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m} \\
+ \tilde{K}_2 \sqrt{\frac{t}{n}} \sum_{m=1}^M \|\hat{f}_m - f_m^*\|_{L_2(\Pi)},$$
(47)

where \tilde{K}_1 and \tilde{K}_2 are constants and $\hat{\gamma}_n = \gamma_n (1 + \|f\|_{\ell_1})$. Finally since $\tilde{K}_2 \sqrt{\frac{t}{n}} \sum_{m=1}^M \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} = \tilde{K}_2 \sqrt{\frac{t}{n}} (\sum_{m \in I} \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} + \sum_{m \in J} \|\hat{f}_m\|_{L_2(\Pi)} + \sum_{m \in J} \|f_m^*\|_{L_2(\Pi)}) \le \tilde{K}_2 \sqrt{\frac{t}{n}} (\sum_{m \in I} \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} + \sum_{m \in J} \|\hat{f}_m\|_{\mathcal{H}_m} + \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m}),$ (47) becomes

$$\frac{1}{2} \|\hat{f} - f^*\|_{L_2(\Pi)}^2 + \lambda_2^{(n)} \sum_{m \in I} \|\hat{f}_I - f_I^*\|_{\mathcal{H}_m}^2 + \lambda_2^{(n)} \sum_{m \in J} \|\hat{f}_m\|_{\mathcal{H}_m}^2 + \left(\lambda_1^{(n)} - \hat{\gamma}_n - \tilde{K}_2 \sqrt{\frac{t}{n}}\right) \sum_{m \in J} \|\hat{f}_m\|_{\mathcal{H}_m}^2 \\
\leq \tilde{K}_1 (1 + \|\hat{f} - f^*\|_{\ell_1}) \Big(\sum_{m \in I} \frac{\|\hat{f}_m - f_m^*\|_{L_2(\Pi)}^{1-s} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}^s}{\sqrt{n}} \vee \frac{\|\hat{f}_m - f_m^*\|_{\mathcal{H}_m}}{n^{\frac{1}{1+s}}} + \frac{t\|\hat{f} - f^*\|_{\ell_1}}{n} \Big) \\
+ \sum_{m \in I} \left(\lambda_1^{(n)} \frac{\|g_m^*\|_{\mathcal{H}_m}}{\|f_m^*\|_{\mathcal{H}_m}} + 2\lambda_2^{(n)} \|g_m^*\|_{\mathcal{H}_m} + \tilde{K}_2 \sqrt{\frac{t}{n}} \right) \|\hat{f}_m - f_m^*\|_{L_2(\Pi)} \\
+ \lambda_2^{(n)} \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m}^2 + \left(\lambda_1^{(n)} + \hat{\gamma}_n + \tilde{K}_2 \sqrt{\frac{t}{n}} \right) \sum_{m \in J} \|f_m^*\|_{\mathcal{H}_m},$$
(48)

which yields the assertion.

C Proof of Theorems 4 and 5

We write the operator norm of $S_{I,J} : \mathcal{H}_J \to \mathcal{H}_I$ as $\|S_{I,J}\|_{\mathcal{H}_I,\mathcal{H}_J} := \sup_{g_J \in \mathcal{H}_J, g_J \neq 0} \frac{\|S_{I,J}g_J\|_{\mathcal{H}_I}}{\|g_J\|_{\mathcal{H}_J}}.$

Definition 9 For all $1 \le m, m' \le M$, we define the empirical (non centered) cross covariance operator $\hat{\Sigma}_{m,m'}$ as follows:

$$\langle f_m, \hat{\Sigma}_{m,m'} g_{m'} \rangle_{\mathcal{H}_m} := \frac{1}{n} \sum_{i=1}^n f_m(x_i) g_{m'}(x_i),$$
(49)

where $f_m \in \mathcal{H}_m, g_{m'} \in \mathcal{H}_{m'}$. Analogous to the joint covariance operator Σ , we define the joint empirical cross covariance operator $\hat{\Sigma} : \mathcal{H} \to \mathcal{H}$ as $(\hat{\Sigma}h)_m = \sum_{l=1}^M \hat{\Sigma}_{m,l}h_l$. We denote by $\hat{\Sigma}_{m,\epsilon}$ the element of \mathcal{H}_m such that

$$\langle f_m, \hat{\Sigma}_{m,\epsilon} \rangle_{\mathcal{H}_m} := \frac{1}{n} \sum_{i=1}^n \epsilon_i f_m(x_i).$$

Let \bar{R} be a constant such that $4(\sum_{m=1}^{M} \|f_m^*\|_{\mathcal{H}_m} + \sum_{m=1}^{M} \|f_m^*\|_{\mathcal{H}_m}) \leq \bar{R}$. We denote by F_n the objective function of elastic-net MKL

$$F_n(f) := \frac{1}{n} \sum_{i=1}^n (f(x_i) - y_i)^2 + \lambda_1^{(n)} \sum_{m=1}^M \|f_m\|_{\mathcal{H}_m} + \lambda_2^{(n)} \sum_{m=1}^M \|f_m\|_{\mathcal{H}_m}^2.$$

Proof: (Theorem 4) Let $\tilde{f} \in \bigoplus_{m \in I_0} \mathcal{H}_m$ be the minimizer of \tilde{F}_n :

$$\begin{split} \tilde{f} &:= \mathop{\arg\min}_{f \in \mathcal{H}_{I_0}} \tilde{F}_n(f), \\ \text{where } \tilde{F}_n(f) &:= \frac{1}{n} \sum_{i=1}^n (f(x_i) - y_i)^2 + \lambda_1^{(n)} \sum_{m \in I_0} \|f_m\|_{\mathcal{H}_m} + \lambda_2^{(n)} \sum_{m \in I_0} \|f_m\|_{\mathcal{H}_m}^2 \end{split}$$

(Step 1) We first show that $\tilde{f} \stackrel{p}{\to} f^*$ with respect to the RKHS norm. Since $\lambda_1^{(n)}\sqrt{n} \to \infty$, as in the proof of Lemma 7, the probability of $\sum_{m=1}^{M} \|\hat{f}_m - f_m^*\|_{\mathcal{H}_m} \leq \sqrt{M\bar{R}}$ goes to 1 (this can be checked as follows: by replacing $\sqrt{\frac{\log(Mn)}{n}}$ in Eq. (34) with $\log(M)\lambda_1^{(n)}$, then we see that Eq. (34) holds with probability $1 - \exp(-\lambda_1^{(n)^2}n)$). There exists c_1 only depending $\sqrt{M\bar{R}}$ such that

$$\|f_m\|_{\mathcal{H}_m} = \sqrt{\|f_m - f_m^*\|_{\mathcal{H}_m}^2 - 2\langle f_m - f_m^*, f_m^* \rangle_{\mathcal{H}_m} + \|f_m^*\|_{\mathcal{H}_m}^2}$$

$$\geq c_1 \|f_m - f_m^*\|_{\mathcal{H}_m}^2 - 2\|f_m^*\|_{\mathcal{H}_m}^{-1}|\langle f_m - f_m^*, f_m^* \rangle_{\mathcal{H}_m}| + \|f_m^*\|_{\mathcal{H}_m}$$
(50)

for all $m \in I_0$ and all $f_m \in \mathcal{H}_m$ such that $||f_m||_{\mathcal{H}_m} \leq \sqrt{M}\overline{R}$.

Since \tilde{f} minimizes \tilde{F}_n , if $\sum_{m=1}^M \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m} \le \sqrt{MR}$ (the probability of which event goes to 1) we have

$$\langle \tilde{f}_{I_0} - f_{I_0}^*, \hat{\Sigma}_{I_0, I_0} (\tilde{f}_{I_0} - f_{I_0}^*) \rangle_{\mathcal{H}_{I_0}} + c_1 \lambda_1^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 + \lambda_2^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2$$

$$\leq 2 \langle \hat{\Sigma}_{I_0, \epsilon}, \tilde{f} - f^* \rangle_{\mathcal{H}_{I_0}} + 2 \sum_{m \in I_0} \left(\frac{1}{\|f_m^*\|_{\mathcal{H}_m}} \lambda_1^{(n)} + \lambda_2^{(n)} \right) |\langle \tilde{f}_m - f_m^*, f_m^* \rangle_{\mathcal{H}_m} |,$$
(51)

where we used the relation (50). By the assumption $f_m^* = \Sigma_{m,m}^{1/2} g_m^*$, we have $|\langle \tilde{f}_m - f_m^*, f_m^* \rangle_{\mathcal{H}_m}| \le ||g_m^*||_{\mathcal{H}_m} ||\tilde{f}_m - f_m^*||_{L_2(\Pi)}$. By Lemma 10 and Lemma 11, we have

$$\|\Sigma_{m,m'} - \hat{\Sigma}_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} = O_p(1/\sqrt{n}), \quad \|\hat{\Sigma}_{I_0,\epsilon}\|_{\mathcal{H}_{I_0}} = O_p(1/\sqrt{n})$$

Substituting these inequalities to (51), we have

$$\|\tilde{f} - f^*\|_{L_2(\Pi)}^2 + c_1 \lambda_1^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 + \lambda_2^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2$$

$$\leq O_p \left(\frac{\sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}}{\sqrt{n}} + (\lambda_1^{(n)} + \lambda_2^{(n)}) \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{L_2(\Pi)} \right).$$
(52)

Remind that the (non centered) cross correlation operator is invertible. Thus there exists a constant c such that

$$\begin{split} \|\tilde{f} - f^*\|_{L_2(\Pi)}^2 &= \langle \tilde{f}_{I_0} - f_{I_0}^*, \Sigma_{I_0,I_0}(\tilde{f}_{I_0} - f_{I_0}^*) \rangle_{\mathcal{H}} = \langle \tilde{f}_{I_0} - f_{I_0}^*, \operatorname{Diag}(\Sigma_{m,m}^{1/2}) V_{I_0,I_0} \operatorname{Diag}(\Sigma_{m,m}^{1/2}) (\tilde{f}_{I_0} - f_{I_0}^*) \rangle_{\mathcal{H}_{I_0}} \\ \geq c \sum_{m \in I_0} \langle \tilde{f}_m - f_m^*, \Sigma_{m,m}(\tilde{f}_m - f_m^*) \rangle_{\mathcal{H}_m} = c \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{L_2(\Pi)}^2. \end{split}$$

This and Eq. (52) give that using $ab \leq (a^2 + b^2)/2$

$$\begin{split} \|\tilde{f} - f^*\|_{L_2(\Pi)}^2 + c_1 \lambda_1^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 + \lambda_2^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 \\ &\leq O_p \left(\frac{\sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}}{\sqrt{n}} + (\lambda_1^{(n)} + \lambda_2^{(n)}) \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{L_2(\Pi)} \right) \\ &\leq O_p \left(\frac{1}{n\lambda_1^{(n)}} + (\lambda_1^{(n)} + \lambda_2^{(n)})^2 \right) + \frac{c_1}{2}\lambda_1^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 + \frac{c}{2} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{L_2(\Pi)}^2 \\ &\leq O_p \left(\frac{1}{n\lambda_1^{(n)}} + (\lambda_1^{(n)} + \lambda_2^{(n)})^2 \right) + \frac{c_1}{2}\lambda_1^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 + \frac{1}{2} \|\tilde{f} - f^*\|_{L_2(\Pi)}^2. \end{split}$$

Therefore we have

$$\frac{1}{2} \|\tilde{f} - f^*\|_{L_2(\Pi)}^2 + \frac{c_1}{2} \lambda_1^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 + \lambda_2^{(n)} \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 \le O_p\left(\frac{1}{n\lambda_1^{(n)}} + (\lambda_1^{(n)} + \lambda_2^{(n)})^2\right) \\ \Rightarrow \sum_{m \in I_0} \|\tilde{f}_m - f_m^*\|_{\mathcal{H}_m}^2 \le O_p\left(\frac{1}{(c_1\lambda_1^{(n)} + \lambda_2^{(n)})n\lambda_1^{(n)}} + \frac{(\lambda_1^{(n)} + \lambda_2^{(n)})^2}{c_1\lambda_1^{(n)} + \lambda_2^{(n)}}\right) = O_p\left(\frac{1}{n\lambda_1^{(n)}} + (\lambda_1^{(n)} + \lambda_2^{(n)})^2\right).$$

This and $\lambda_1^{(n)}\sqrt{n} \to \infty$ gives $\|\tilde{f} - f_{I_0}^*\|_{\mathcal{H}_{I_0}} \to 0$ in probability.

(Step 2) Next we show that the probability of $\tilde{f} = \hat{f}$ goes to 1. Since $\|\tilde{f} - f_{I_0}^*\|_{\mathcal{H}_{I_0}} \to 0$, we can assume that $\|\tilde{f}_m\|_{\mathcal{H}_m} > 0$ ($m \in I_0$) without loss of generality. We identify \tilde{f} as an element of \mathcal{H} by setting $\tilde{f}_m = 0$ for $m \in J_0$. Now we show that \tilde{f} is also the minimizer of F_n , that is $\tilde{f} = \hat{f}$, with high probability, hence $\hat{I} = I_0$ with high probability. By the KKT condition, the necessary and sufficient condition that \tilde{f} also minimizes F_n is

$$\|2\hat{\Sigma}_{m,I_0}(\tilde{f}_{I_0} - f_{I_0}^*) - 2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m} \le \lambda_1^{(n)} \quad (\forall m \in J_0),$$
(53)

$$(2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D_n)(\tilde{f}_{I_0} - f_{I_0}^*) + \lambda_1^{(n)}D_nf_{I_0}^* + 2\lambda_2^{(n)}f_{I_0}^* - 2\hat{\Sigma}_{I_0,\epsilon} = 0,$$
(54)

where $D_n = \text{Diag}(\|\tilde{f}_m\|_{\mathcal{H}_m}^{-1})$. Note that (54) is satisfied (with high probability) because \tilde{f} is the minimizer of \tilde{F}_n and $\|\tilde{f}_m\|_{\mathcal{H}_m} > 0$ for all $m \in I_0$ (with high probability). Therefore if the condition (53) holds w.h.p., $\tilde{f} = \hat{f}$ w.h.p..

We will now show the condition (53) holds w.h.p.. Due to (54), we have

$$\tilde{f}_{I_0} - f_{I_0}^* = -(2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D_n)^{-1}[(\lambda_1^{(n)}D_n + 2\lambda_2^{(n)})f_{I_0}^* - 2\hat{\Sigma}_{I_0,\epsilon}].$$

Therefore the LHS of (53), $\|2\hat{\Sigma}_{m,I_0}(\tilde{f}_{I_0} - f^*_{I_0}) - 2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m}$, can be evaluated as

$$\begin{aligned} \| -2\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}[(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}-2\hat{\Sigma}_{I_{0},\epsilon}]-2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_{m}} \\ = \|2\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*} \\ -2\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}(2\hat{\Sigma}_{I_{0},\epsilon}+2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_{m}} \\ \leq \|2\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}\|_{\mathcal{H}_{m}} \\ + \|2\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}2\hat{\Sigma}_{I_{0},\epsilon}-2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_{m}}. \end{aligned}$$
(55)

We evaluate the probabilistic orders of the last two terms.

(i) (Bounding $B_{n,m} := \|2\hat{\Sigma}_{m,I_0}(2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D_n)^{-1}2\hat{\Sigma}_{I_0,\epsilon} - 2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m})$ We show that

$$\hat{\Sigma}_{m,I_0} (2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)} D_n)^{-1} \hat{\Sigma}_{I_0,\epsilon} = O_p \left(\frac{1}{\sqrt{n}}\right).$$

Since $O \preceq \begin{pmatrix} \hat{\Sigma}_{I_0,I_0} & \hat{\Sigma}_{I_0,m} \\ \hat{\Sigma}_{m,I_0} & \hat{\Sigma}_{m,m} \end{pmatrix}$, we have

$$O \preceq \begin{pmatrix} \hat{\Sigma}_{I_0,I_0} + \lambda_2^{(n)} + \lambda_1^{(n)} D_n/2 & \hat{\Sigma}_{I_0,m} \\ \hat{\Sigma}_{m,I_0} & \hat{\Sigma}_{m,m} + \lambda_2^{(n)} \end{pmatrix} \preceq \begin{pmatrix} 2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)} D_n & 0 \\ 0 & 2\hat{\Sigma}_{m,m} + 2\lambda_2^{(n)} \end{pmatrix}$$

The second inequality is due to the fact that for all $(f_{I_0}, f_m) \in \mathcal{H}_{I_0 \cup m}$ we have

$$\left\langle \begin{pmatrix} f_{I_0} \\ -f_m \end{pmatrix}, \begin{pmatrix} \hat{\Sigma}_{I_0,I_0} + \lambda_2^{(n)} + \lambda_1^{(n)} D_n/2 & -\hat{\Sigma}_{I_0,m} \\ -\hat{\Sigma}_{m,I_0} & \hat{\Sigma}_{m,m} + \lambda_2^{(n)} \end{pmatrix} \begin{pmatrix} f_{I_0} \\ -f_m \end{pmatrix} \right\rangle_{\mathcal{H}_{I_0 \cup m}} \ge 0$$

because of $O \preceq \begin{pmatrix} \hat{\Sigma}_{I_0,I_0} & \hat{\Sigma}_{I_0,m} \\ \hat{\Sigma}_{m,I_0} & \hat{\Sigma}_{m,m} \end{pmatrix}$. Thus we have

$$\left\| \begin{pmatrix} \hat{\Sigma}_{I_0,I_0} + \lambda_2^{(n)} + \frac{\lambda_1^{(n)}D_n}{2} & \hat{\Sigma}_{I_0,m} \\ \hat{\Sigma}_{m,I_0} & \hat{\Sigma}_{m,m} + \lambda_2^{(n)} \end{pmatrix} \begin{pmatrix} 2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D_n & 0 \\ 0 & 2\hat{\Sigma}_{m,m} + 2\lambda_2^{(n)} \end{pmatrix}^{-1} \begin{pmatrix} \hat{\Sigma}_{I_0,\epsilon} \\ \hat{\Sigma}_{m,\epsilon} \end{pmatrix} \right\|_{\mathcal{H}_{I_0\cup m}}$$

$$\leq \left\| \begin{pmatrix} \hat{\Sigma}_{I_0,\epsilon} \\ \hat{\Sigma}_{m,\epsilon} \end{pmatrix} \right\|_{\mathcal{H}_{I_0\cup m}} \leq O_p(1/\sqrt{n}).$$
(56)

Here the LHS of the above inequality is equivalent to

$$\left\| \begin{pmatrix} * \\ \hat{\Sigma}_{m,I_0} (2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)} D_n)^{-1} \hat{\Sigma}_{I_0,\epsilon} + (\hat{\Sigma}_{m,m} + \lambda_2^{(n)}) (2\hat{\Sigma}_{m,m} + 2\lambda_2^{(n)})^{-1} \hat{\Sigma}_{m,\epsilon} \end{pmatrix} \right\|_{\mathcal{H}_{I_0 \cup m}}.$$

Therefore we observe

$$\hat{\Sigma}_{m,I_0} (2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)} D_n)^{-1} \hat{\Sigma}_{I_0,\epsilon} + \frac{1}{2} \hat{\Sigma}_{m,\epsilon} \bigg\|_{\mathcal{H}_m} = O_p(1/\sqrt{n}).$$

Since $\|\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m} = O_p(1/\sqrt{n})$, we also have

$$\|\hat{\Sigma}_{m,I_0}(2\hat{\Sigma}_{I_0,I_0}+2\lambda_2^{(n)}+\lambda_1^{(n)}D_n)^{-1}\hat{\Sigma}_{I_0,\epsilon}\|_{\mathcal{H}_m}=O_p(1/\sqrt{n}).$$

This and $\|\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m} = O_p(1/\sqrt{n})$ yield

$$B_{n,m} = O_p(1/\sqrt{n}). \tag{57}$$

(ii) (Bounding $E_{n,m} := \|2\hat{\Sigma}_{m,I_0}(2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D_n)^{-1}(\lambda_1^{(n)}D_n + 2\lambda_2^{(n)})f_{I_0}^*\|_{\mathcal{H}_m})$ Note that, due to $\|\tilde{f} - f^*\|_{\mathcal{H}} \xrightarrow{p} 0$, we have $D_n \xrightarrow{p} D$, and we know that $\max_{m,m'} \|\hat{\Sigma}_{m,n'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} = O_p(\sqrt{\log(M)/n}) = O_p(\frac{1}{\sqrt{n}})$ by Lemma 10. Thus $S_n := (2\Sigma_{I_0,I_0} - 2\hat{\Sigma}_{I_0,I_0})/\lambda_1^{(n)} + D - D_n$ satisfies $S_n = o_p(1)$ and thus $D - S_n \succeq D/2$ with high probability. Hence

$$2\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}$$

$$=2\Sigma_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}+O_{p}\left(\frac{1}{\sqrt{n}}\right)$$

$$=2\Sigma_{m,I_{0}}(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D)^{-1}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}+$$

$$=2\Sigma_{m,I_{0}}(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D)^{-1}\lambda_{1}^{(n)}S_{n}(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}(D-S_{n}))^{-1}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}+O_{p}\left(\frac{1}{\sqrt{n}}\right).$$
(58)

Here we obtain

$$\begin{split} \|\Sigma_{m,I_0}(2\Sigma_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D)^{-\frac{1}{2}}\|_{\mathcal{H}_m,\mathcal{H}_{I_0}}^2 \\ = \|\Sigma_{m,I_0}(2\Sigma_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D)^{-1}\Sigma_{I_0,m}\|_{\mathcal{H}_m,\mathcal{H}_m} \\ \le \|\Sigma_{m,m}^{\frac{1}{2}}V_{m,I_0}(2V_{I_0,I_0})^{-1}V_{I_0,m}\Sigma_{m,m}^{\frac{1}{2}}\|_{\mathcal{H}_m,\mathcal{H}_m} = O_p(1), \end{split}$$
(59)

and due to the fact that $D-S_n \succeq D/2$ with high probability we have

$$\begin{split} \| (\Sigma_{I_0,I_0} + \lambda_2^{(n)} + \lambda_1^{(n)} (D - S_n))^{-\frac{1}{2}} (\lambda_1^{(n)} D_n + 2\lambda_2^{(n)}) f_{I_0}^* \|_{\mathcal{H}_{I_0}} \\ = \| (\Sigma_{I_0,I_0} + \lambda_2^{(n)} + \lambda_1^{(n)} (D - S_n))^{-\frac{1}{2}} \mathrm{Diag}(\Sigma_{m,m}^{\frac{1}{2}}) (\lambda_1^{(n)} D_n + 2\lambda_2^{(n)}) g_{I_0}^* \|_{\mathcal{H}_{I_0}} \\ \leq O_p(\| V_{I_0,I_0}^{-1} \|_{\mathcal{H}_{I_0},\mathcal{H}_{I_0}}^{-\frac{1}{2}} (\lambda_1^{(n)} + \lambda_2^{(n)})) = O_p(\lambda_1^{(n)} + \lambda_2^{(n)}). \end{split}$$

Therefore the second term in the RHS of Eq. (58) is evaluated as

$$\begin{split} \|\Sigma_{m,I_{0}}(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D)^{-1}\lambda_{1}^{(n)}S_{n}(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}(D-S_{n}))^{-1}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}\|_{\mathcal{H}_{m}} \\ \leq \|\Sigma_{m,I_{0}}(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D)^{-\frac{1}{2}}\|_{\mathcal{H}_{m},\mathcal{H}_{I_{0}}}\|(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D)^{-\frac{1}{2}}\|_{\mathcal{H}_{I_{0}},\mathcal{H}_{I_{0}}}\chi_{I_{0}} \\ & \|(2\Sigma_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}(D-S_{n}))^{-\frac{1}{2}}\|_{\mathcal{H}_{I_{0}},\mathcal{H}_{I_{0}}}\|(\Sigma_{I_{0},I_{0}}+\lambda_{2}^{(n)}+\lambda_{1}^{(n)}(D-S_{n}))^{-\frac{1}{2}}(\lambda_{1}^{(n)}D_{n}+2\lambda_{2}^{(n)})f_{I_{0}}^{*}\|_{\mathcal{H}_{I_{0}}} \\ \leq O_{p}(1\cdot(\lambda_{1}^{(n)}+\lambda_{2}^{(n)})^{-\frac{1}{2}}\cdot\lambda_{1}^{(n)}o_{p}(1)\cdot(\lambda_{1}^{(n)}+\lambda_{2}^{(n)})^{-\frac{1}{2}}\cdot(\lambda_{1}^{(n)}+\lambda_{2}^{(n)})) \\ = o_{p}(\lambda_{1}^{(n)}). \end{split}$$

Therefore this and Eq. (58) give

$$2\hat{\Sigma}_{m,I_0}(2\hat{\Sigma}_{I_0,I_0}+2\lambda_2^{(n)}+\lambda_1^{(n)}D_n)^{-1}(\lambda_1^{(n)}D_n+2\lambda_2^{(n)})f_{I_0}^*$$

=2\Sum_{m,I_0}(2\Sum_{I_0,I_0}+2\lambda_2^{(n)}+\lambda_1^{(n)}D)^{-1}(\lambda_1^{(n)}D_n+2\lambda_2^{(n)})f_{I_0}^*+o_p(\lambda_1^{(n)})+O_p\left(\frac{1}{\sqrt{n}}\right)
=2\Sum_{m,I_0}(2\Sum_{I_0,I_0}+2\lambda_2^{(n)}+\lambda_1^{(n)}D)^{-1}(\lambda_1^{(n)}D_n+2\lambda_2^{(n)})f_{I_0}^*+o_p(\lambda_1^{(n)}).

Define

$$A_{n} := \Sigma_{m,I_{0}} \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} + \frac{\lambda_{1}^{(n)}D}{2} \right)^{-1} \left(D_{n} + 2\frac{\lambda_{2}^{(n)}}{\lambda_{1}^{(n)}} \right) f_{I_{0}}^{*},$$
$$A := \Sigma_{m,I_{0}} (\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)})^{-1} \left(D + 2\frac{\lambda_{2}^{(n)}}{\lambda_{1}^{(n)}} \right) f_{I_{0}}^{*}.$$

We show $||A_n - A||_{\mathcal{H}_m} = o_p(1)$. By the definition, we have

$$A - A_n = \sum_{m,I_0} (\sum_{I_0,I_0} + \lambda_2^{(n)})^{-1} \frac{\lambda_1^{(n)} D}{2} \left(\sum_{I_0,I_0} + \lambda_2^{(n)} + \frac{\lambda_1^{(n)} D}{2} \right)^{-1} \left(D + 2\frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) f_{I_0}^* + \sum_{m,I_0} \left(\sum_{I_0,I_0} + \lambda_2^{(n)} + \frac{\lambda_1^{(n)} D}{2} \right)^{-1} (D - D_n) f_{I_0}^*.$$
(60)

On the other hand, as in Eq. (56), we observe that

$$2 \ge \left\| \begin{pmatrix} \Sigma_{I_0,I_0} & \Sigma_{I_0,m} \\ \Sigma_{m,I_0} & \Sigma_{m,m} \end{pmatrix} \begin{pmatrix} (\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1} & 0 \\ 0 & 0 \end{pmatrix} \right\|_{\mathcal{H}_{I_0 \cup m},\mathcal{H}_{I_0 \cup m}} \\ = \left\| \begin{pmatrix} * & * \\ \Sigma_{m,I_0} (\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1} & 0 \end{pmatrix} \right\|_{\mathcal{H}_{I_0 \cup m},\mathcal{H}_{I_0 \cup m}} \ge \|\Sigma_{m,I_0} (\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1} \|_{\mathcal{H}_m,\mathcal{H}_{I_0}}.$$
(61)

Moreover, since $f_m^* = \sum_{m,m}^{\frac{1}{2}} g_m^*$ ($\forall m$), we have

$$\begin{aligned} \left\| \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} + \frac{\lambda_{1}^{(n)}D}{2} \right)^{-1} \left(D + 2\frac{\lambda_{2}^{(n)}}{\lambda_{1}^{(n)}} \right) f_{I_{0}}^{*} \right\|_{\mathcal{H}_{I_{0}}} \\ &= \left\| \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} + \frac{\lambda_{1}^{(n)}D}{2} \right)^{-1} \operatorname{Diag}(\Sigma_{m,m}^{\frac{1}{2}}) \left(D + 2\frac{\lambda_{2}^{(n)}}{\lambda_{1}^{(n)}} \right) g_{I_{0}}^{*} \right\|_{\mathcal{H}_{I_{0}}} \\ &\leq \left\| \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} + \frac{\lambda_{1}^{(n)}D}{2} \right)^{-\frac{1}{2}} \right\|_{\mathcal{H}_{I_{0}},\mathcal{H}_{I_{0}}} \left\| \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} + \frac{\lambda_{1}^{(n)}D}{2} \right)^{-\frac{1}{2}} \operatorname{Diag}(\Sigma_{m,m}^{\frac{1}{2}}) \right\|_{\mathcal{H}_{I_{0}},\mathcal{H}_{I_{0}}} \\ &\times \left\| \left(D + 2\frac{\lambda_{2}^{(n)}}{\lambda_{1}^{(n)}} \right) g_{I_{0}}^{*} \right\|_{\mathcal{H}_{I_{0}}} \\ &\leq O_{p}((\lambda_{1}^{(n)} + \lambda_{2}^{(n)})^{-\frac{1}{2}} \left\| V_{I_{0},I_{0}}^{-\frac{1}{2}} \right\|_{\mathcal{H}_{I_{0}},\mathcal{H}_{I_{0}}} \right) \leq O_{p}(\lambda_{1}^{(n)^{-\frac{1}{2}}}). \end{aligned}$$

$$(62)$$

We can also bound the second term of (60) as

$$\begin{split} \left\| \Sigma_{m,I_{0}} \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} + \frac{\lambda_{1}^{(n)}D}{2} \right)^{-1} (D - D_{n}) f_{I_{0}}^{*} \right\|_{\mathcal{H}_{m}} \\ \leq \left\| \Sigma_{m,I_{0}} \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} + \frac{\lambda_{1}^{(n)}D}{2} \right)^{-1} \right\|_{\mathcal{H}_{m},\mathcal{H}_{I_{0}}} \left\| (D - D_{n}) f_{I_{0}}^{*} \right\|_{\mathcal{H}_{I_{0}}} \\ \leq \left\| \Sigma_{m,I_{0}} \left(\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)} \right)^{-1} \right\|_{\mathcal{H}_{m},\mathcal{H}_{I_{0}}} \left\| (D - D_{n}) f_{I_{0}}^{*} \right\|_{\mathcal{H}_{I_{0}}} \\ \leq 2 \left\| (D - D_{n}) f_{I_{0}}^{*} \right\|_{\mathcal{H}_{I_{0}}} \quad (\because \text{Eq. (61)}) \\ = o_{p}(1). \end{split}$$

Therefore applying the inequalities Eq. (61) and Eq. (62) to Eq. (60), we have

$$||A_n - A||_{\mathcal{H}_m} = O_p(\lambda_1^{(n)^{\frac{1}{2}}}) + o_p(1) = o_p(1).$$
(63)

Hence we have $E_{n,m} = \lambda_1^{(n)} ||A||_{\mathcal{H}_m} + o_p(\lambda_1^{(n)}).$

(iii) (Combining (i) and (ii)) Due to the above evaluations ((i) and (ii)), we have

$$\begin{split} & \max_{m \in J_0} \left\| 2\hat{\Sigma}_{m,I}(\tilde{f}_{I_0} - f_{I_0}^*) - 2\hat{\Sigma}_{m,\epsilon} \right\|_{\mathcal{H}_m} \\ &= \max_{m \in J} \lambda_1^{(n)} \left\| \Sigma_{m,I_0}(\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1} \left(D + 2\frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) f_{I_0}^* \right\|_{\mathcal{H}_m} + o_p(\lambda_1^{(n)}) < \lambda_1^{(n)}(1 - \eta) + o_p(\lambda_1^{(n)}). \end{split}$$

This yields

$$P\left(\|2\hat{\Sigma}_{m,I_0}(\tilde{f}_{I_0} - f_{I_0}^*) - 2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m} \ge \lambda_1^{(n)}, \forall m \in J_0\right) \to 0$$

Thus the probability of the condition (53) goes to 1.

Proof: (Theorem 5) First we prove that $\lambda_1^{(n)}\sqrt{n} \to \infty$ is a necessary condition for $\hat{I} \xrightarrow{p} I_0$. Assume that $\liminf \lambda_1^{(n)}\sqrt{n} < \infty$. Then we can take a sub-sequence that converges to a finite value, therefore by taking the sub-sequence, if necessary, we can assume $\lim \lambda_1^{(n)}\sqrt{n} \to \mu_1$ without loss of generality. We will derive a contradiction under the conditions of $\|\hat{f} - f^*\|_{\mathcal{H}} \xrightarrow{p} 0$ and $\hat{I} \xrightarrow{p} I_0$. Suppose $\hat{I} = I_0$.

By the KKT condition,

$$0 = 2(\hat{\Sigma}_{I_0,I_0}\hat{f}_{I_0} - \hat{\Sigma}_{I_0,\epsilon} - \hat{\Sigma}_{I_0,I_0}f_{I_0}^*) + \lambda_1^{(n)}D_n\hat{f}_{I_0} + 2\lambda_2^{(n)}\hat{f}_{I_0}$$

$$\Rightarrow 2(\hat{\Sigma}_{I_0,I_0} + \lambda_2^{(n)})(f_{I_0}^* - \hat{f}_{I_0}) = \lambda_1^{(n)}D_nf_{I_0}^* + 2\lambda_2^{(n)}f_{I_0}^* - 2\hat{\Sigma}_{I_0,\epsilon}$$

$$\Rightarrow 2\sqrt{n}(\Sigma_{I_0,I_0} + \lambda_2^{(n)})(f_{I_0}^* - \hat{f}_{I_0}) = \sqrt{n}\lambda_1^{(n)}Df_{I_0}^* + \sqrt{n}2\lambda_2^{(n)}f_{I_0}^* - 2\sqrt{n}\hat{\Sigma}_{I_0,\epsilon}$$

$$+ (2\sqrt{n}(\Sigma_{I_0,I_0} - \hat{\Sigma}_{I_0,I_0})(f_{I_0}^* - \hat{f}_{I_0}) + \sqrt{n}\lambda_1^{(n)}(D_n - D)f_{I_0}^*)$$

$$\Rightarrow 2\sqrt{n}(\Sigma_{I_0,I_0} + \lambda_2^{(n)})(f_{I_0}^* - \hat{f}_{I_0}) = \mu_1 Df_{I_0}^* + \sqrt{n}2\lambda_2^{(n)}f_{I_0}^* - 2\sqrt{n}\hat{\Sigma}_{I_0,\epsilon} + o_p(1),$$
(65)

where the last inequality is due to $\sqrt{n}\lambda_1^{(n)} \to \mu_1$, $\|D_n - D\|_{\mathcal{H}_{I_0},\mathcal{H}_{I_0}} = o_p(1)$, $\|\hat{f} - f^*\|_{\mathcal{H}} = o_p(1)$ and $\|\Sigma_{I_0,I_0} - \hat{\Sigma}_{I_0,I_0}\|_{\mathcal{H}_{I_0},\mathcal{H}_{I_0}} = o_p(1)$. Moreover since the second equality (64) indicates that $o_p(1) + o_p(\lambda_2^{(n)}) = \lambda_1^{(n)} Df_{I_0}^* + 2\lambda_2^{(n)}f_{I_0}^* + o_p(1)$, we have $\lambda_1^{(n)} = o_p(1)$ and $\lambda_2^{(n)} = o_p(1)$.

We now show that the KKT condition under which \hat{f} satisfying $\hat{I} = I_0$ is optimal with respect to F_n is violated with strictly positive probability:

$$\liminf P\left(\exists m \in J, \|2(\hat{\Sigma}_{m,I_0}\hat{f}_{I_0} - \hat{\Sigma}_{m,I_0}f^*_{I_0} - \hat{\Sigma}_{m,\epsilon})\|_{\mathcal{H}_m} > \lambda_1^{(n)}\right) > 0.$$
(66)

Obviously this indicates that the probability $\hat{I} = I_0$ does not converges to 1, which is a contradiction.

For all $v_m \in \mathcal{H}_m$ $(m \in J_0)$, there exists $w_{I_0} \in \mathcal{H}_{I_0}$ such that

$$\Sigma_{I_0,m} v_m = (\Sigma_{I_0,I_0} + \lambda_2^{(n)}) w_{I_0}.$$
(67)

Note that w_{I_0} is uniformly bounded for all $\lambda_2^{(n)} \ge 0$ because the range of $\Sigma_{I_0,m}$ is included in the range of Σ_{I_0,I_0} (Baker, 1973) and there exists \tilde{w}_{I_0} such that $\Sigma_{I_0,m}v_m = \Sigma_{I_0,I_0}\tilde{w}_{I_0}$ (\tilde{w}_{I_0} is independent of $\lambda_2^{(n)}$), hence $\Sigma_{I_0,I_0}\tilde{w}_{I_0} = (\Sigma_{I_0,I_0} + \lambda_2^{(n)})w_{I_0}$, and

$$\|w_{I_0}\|_{\mathcal{H}_{I_0}} \leq \sqrt{\langle \tilde{w}_{I_0}, \Sigma_{I_0, I_0}(\Sigma_{I_0, I_0} + \lambda_2^{(n)})^{-2}\Sigma_{I_0, I_0}\tilde{w}_{I_0}\rangle_{\mathcal{H}_{I_0}}} \leq \|\tilde{w}_{I_0}\|_{\mathcal{H}_{I_0}}$$

for $\lambda_2^{(n)} > 0$ and $\|w_{I_0}\|_{\mathcal{H}_{I_0}} = \|\tilde{w}_I\|_{\mathcal{H}_{I_0}}$ for $\lambda_2^{(n)} = 0$. Let $v_m \in \mathcal{H}_m$ be any non-zero element such that $\sum_{m,m}^{1/2} v_m \neq 0$ and w_{I_0} be satisfying the above equality (67), then

$$\begin{split} &\sqrt{n} \langle v_{m}, \hat{\Sigma}_{m,\epsilon} + \hat{\Sigma}_{m,I_{0}} f_{I_{0}}^{*} - \hat{\Sigma}_{m,I_{0}} \hat{f}_{I_{0}} \rangle_{\mathcal{H}_{m}} \\ = &\sqrt{n} \langle v_{m}, \hat{\Sigma}_{m,\epsilon} \rangle_{\mathcal{H}_{m}} + \langle v_{m}, \hat{\Sigma}_{m,I_{0}} \sqrt{n} (f_{I_{0}}^{*} - \hat{f}_{I_{0}}) \rangle_{\mathcal{H}_{m}} \\ = &\sqrt{n} \langle v_{m}, \hat{\Sigma}_{m,\epsilon} \rangle_{\mathcal{H}_{m}} + \langle v_{m}, \Sigma_{m,I} \sqrt{n} (f_{I_{0}}^{*} - \hat{f}_{I_{0}}) \rangle_{\mathcal{H}_{m}} + o_{p}(1) \\ = &\sqrt{n} \langle v_{m}, \hat{\Sigma}_{m,\epsilon} \rangle_{\mathcal{H}_{m}} + \langle w_{I_{0}}, (\Sigma_{I_{0},I_{0}} + \lambda_{2}^{(n)}) \sqrt{n} (f_{I_{0}}^{*} - \hat{f}_{I_{0}}) \rangle_{\mathcal{H}_{m}} + o_{p}(1) \\ = &\sqrt{n} \langle v_{m}, \hat{\Sigma}_{m,\epsilon} \rangle_{\mathcal{H}_{m}} - \sqrt{n} \langle w_{I_{0}}, \hat{\Sigma}_{I_{0},\epsilon} \rangle_{\mathcal{H}_{m}} + \left\langle w_{I_{0}}, \left(\frac{\mu_{1}}{2} D + \sqrt{n} \lambda_{2}^{(n)} \right) f_{I_{0}}^{*} \right\rangle_{\mathcal{H}_{m}} + o_{p}(1), \end{split}$$

where we used $\|\hat{\Sigma}_{m,I_0} - \Sigma_{m,I_0}\|_{\mathcal{H}_m,\mathcal{H}_{I_0}} = O_p(1/\sqrt{n})$ and $\|f^* - \hat{f}\|_{\mathcal{H}} \xrightarrow{p} 0$ in the second equality, and the relation (65) in the last equality. We can show that $Z_n := \sqrt{n} \langle v_m, \hat{\Sigma}_{m,\epsilon} \rangle - \sqrt{n} \langle w_{I_0}, \hat{\Sigma}_{I_0,\epsilon} \rangle$ has a positive variance as follows (see also Bach (2008)):

$$\begin{split} \mathbf{E}[Z_n] &= 0, \\ \mathbf{E}[Z_n^2] \geq \sigma^2 \left(\langle v_m, \Sigma_{m,m} v_m \rangle - 2 \langle v_m, \Sigma_{m,I_0} w_{I_0} \rangle + \langle w_{I_0}, \Sigma_{I_0,I_0} w_{I_0} \rangle \right) \\ &= \sigma^2 \left(\langle v_m, \Sigma_{m,m} v_m \rangle - \langle v_m, \Sigma_{m,I_0} w_{I_0} \rangle + o_p(1) \right) \quad (\because \lambda_2^{(n)} = o_p(1)) \\ &= \sigma^2 \langle \Sigma_{m,m}^{1/2} v_m, (I_{\mathcal{H}_m} - V_{m,I_0} \tilde{V}_{I_0,I_0}^{-1} V_{I_0,m}) \Sigma_{m,m}^{1/2} v_m \rangle + o_p(1), \end{split}$$

where $\tilde{V}_{I_0,I_0}^{-1} = \text{Diag}(\Sigma_{m,m}^{1/2})(\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1}\text{Diag}(\Sigma_{m,m}^{1/2})$ (note that \tilde{V}_{I_0,I_0} is invertible because $V_{I_0,I_0} \preceq \tilde{V}_{I_0,I_0}$ and V_{I_0,I_0} is invertible). Now since $V_{I_0,I_0} \preceq \tilde{V}_{I_0,I_0}$ and $I_{\mathcal{H}_m} - V_{m,I_0}V_{I_0,I_0}^{-1}V_{I_0,m} \succ O$ (this is because $V_{I_0\cup m,I_0\cup m} = \begin{pmatrix} V_{I_0,I_0} & V_{m,I_0} \\ V_{I_0,m} & I_{\mathcal{H}_m} \end{pmatrix}$ is invertible), we have $I_{\mathcal{H}_m} - V_{m,I_0}\tilde{V}_{I_0,I_0}^{-1}V_{I_0,m} \succ O$. Therefore by the central limit theorem Z_n converges Gaussian random variable with strictly positive variance in distribution. Thus the probability of

$$2|\langle v_m, \hat{\Sigma}_{m,\epsilon} + \hat{\Sigma}_{m,I_0} f_{I_0}^* - \hat{\Sigma}_{m,I_0} \hat{f}_{I_0} \rangle_m| > \lambda_1^{(n)} ||v_m||_{\mathcal{H}_m}$$

is asymptotically strictly positive because $\lambda_1^{(n)}\sqrt{n} \to \mu_1$ (Note that this is true whether $\sqrt{n}\lambda_2^{(n)}$ converges to finite value or not). This yields (66), i.e. \hat{f} does not satisfy $\hat{I} = I_0$ with asymptotically strictly positive probability.

We say Condition A as

Condition A :
$$\lambda_1^{(n)}\sqrt{n} \to \infty$$
.

Now that we have proven $\lambda_1^{(n)}\sqrt{n} \to \infty$, we are ready to prove the assertion (16). Suppose the condition (16) is not satisfied for any sequences $\lambda_1^{(n)}, \lambda_2^{(n)} \to 0$, that is, there exists a constant $\xi > 0$ such that

$$\lim_{n \to \infty} \left\| \Sigma_{m, I_0} (\Sigma_{I_0, I_0} + \lambda_2^{(n)})^{-1} \left(D + 2 \frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) g_{I_0}^* \right\|_{\mathcal{H}_m} > (1 + \xi), \quad (\exists m \in J_0), \tag{68}$$

for any sequences $\lambda_1^{(n)}, \lambda_2^{(n)} \to 0$ satisfying Condition A ($\lambda_1^{(n)}\sqrt{n} \to \infty$). Fix arbitrary sequences $\lambda_1^{(n)}, \lambda_2^{(n)} \to 0$ satisfying Condition A. If $\hat{I} = I_0$, the KKT condition

$$\|2\hat{\Sigma}_{m,I_0}(\hat{f}_{I_0} - f^*_{I_0}) - 2\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m} \le \lambda_1^{(n)} \quad (\forall m \in J_0),$$
(69)

$$(2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)}D_n)(\tilde{f}_{I_0} - f_{I_0}^*) + \lambda_1^{(n)}D_nf_{I_0}^* + 2\lambda_2^{(n)}f_{I_0}^* - 2\hat{\Sigma}_{I_0,\epsilon} = 0,$$
(70)

should be satisfied (see (53) and (54)). We prove that the first inequality (69) of the KKT condition is violated with strictly positive probability under the assumptions and the condition (70). We have shown that (see (55))

$$\lambda_{1}^{(n)^{-1}}(2\hat{\Sigma}_{m,I_{0}}(\hat{f}_{I_{0}}-f_{I_{0}}^{*})-2\hat{\Sigma}_{m,\epsilon})$$

$$=2\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}(D_{n}+2\frac{\lambda_{2}^{(n)}}{\lambda_{1}^{(n)}})f_{I_{0}}^{*}$$

$$-\frac{2}{\lambda_{1}^{(n)}}\hat{\Sigma}_{m,I_{0}}(2\hat{\Sigma}_{I_{0},I_{0}}+2\lambda_{2}^{(n)}+\lambda_{1}^{(n)}D_{n})^{-1}2\hat{\Sigma}_{I_{0},\epsilon}+\frac{2}{\lambda_{1}^{(n)}}\hat{\Sigma}_{m,\epsilon}.$$
(71)

As shown in the proof of Theorem 1, the first term can be approximated by $\Sigma_{m,I_0} \left(\Sigma_{I_0,I_0} + \lambda_2^{(n)} \right)^{-1} \left(D + 2 \frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) f_{I_0}^*$, more precisely Eq. (63) gives

$$\left\| \hat{\Sigma}_{m,I_0} \left(\hat{\Sigma}_{I_0,I_0} + \lambda_2^{(n)} + \frac{\lambda_1^{(n)} D_n}{2} \right)^{-1} \left(D_n + 2 \frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) f_{I_0}^* - \Sigma_{m,I_0} (\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1} \left(D + 2 \frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) g_I^* \right\|_{\mathcal{H}_m}$$

$$\xrightarrow{P} 0.$$

Since $\liminf_n \left\| \Sigma_{m,I_0} (\Sigma_{I_0,I_0} + \lambda_2^{(n)})^{-1} \left(D + 2 \frac{\lambda_2^{(n)}}{\lambda_1^{(n)}} \right) g_{I_0}^* \right\|_{\mathcal{H}_m} > (1+\xi)$ by the assumption, we observe that

$$P\left(\left\|2\hat{\Sigma}_{m,I_0}(2\hat{\Sigma}_{I_0,I_0}+2\lambda_2^{(n)}+\lambda_1^{(n)}D_n)^{-1}\left(D_n+2\frac{\lambda_2^{(n)}}{\lambda_1^{(n)}}\right)f_{I_0}^*\right\|_{\mathcal{H}_m} > (1+\xi)\right) \neq 0.$$
(72)

Now since $\lambda_1^{(n)}\sqrt{n} \to \infty$, we have proven that

$$\left\| -\frac{2}{\lambda_1^{(n)}} \hat{\Sigma}_{m,I_0} (2\hat{\Sigma}_{I_0,I_0} + 2\lambda_2^{(n)} + \lambda_1^{(n)} D_n)^{-1} 2\hat{\Sigma}_{I_0,\epsilon} + \frac{2}{\lambda_1^{(n)}} \hat{\Sigma}_{m,\epsilon} \right\|_{\mathcal{H}_m} = O_p(1/(\lambda_1^{(n)}\sqrt{n})) = o_p(1), \quad (73)$$

in the proof of Theorem 1 (Eq. (57)). Therefore, combining (71), (72) and (73), we have observed that the KKT condition (53) is violated with strictly positive probability if the condition (68) is satisfied. This yields the irrepresenter condition (16) is a necessary condition for the consistency of elastic-net MKL.

Lemma 10 If $\sup_X k_m(X, X) \leq 1$ and $\sup_X k_{m'}(X, X) \leq 1$, then

$$P(\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}'_m} \ge \mathbb{E}[\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}'_m}] + \varepsilon) \le \exp(-n\varepsilon^2/2).$$
(74)

In particular,

$$P\left(\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_m'} \ge \sqrt{\frac{1}{n}} + \epsilon\right) \le \exp(-n\varepsilon^2/2).$$
(75)

Proof: We use McDiarmid's inequality (Devroye et al., 1996). By definition

$$\langle g, \hat{\Sigma}_{mm'} f \rangle = \frac{1}{n} \sum_{i=1}^{n} \langle g, k_m(\cdot, x_i) \rangle_m \langle f, k_{m'}(\cdot, x_i) \rangle_{m'}.$$

We denote by $\tilde{\Sigma}_{m,m'}$ the empirical cross covariance operator with n samples $(x_1, \ldots, x_{j-1}, \tilde{x}_j, x_{j+1}, \ldots, x_n)$ where the *j*-th sample x_j is replaced by \tilde{x}_j independently distributed by the same distribution as x_j 's.

By the triangular inequality, we have

$$\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} - \|\tilde{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} \le \|\hat{\Sigma}_{m,m'} - \tilde{\Sigma}_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}}.$$

Now the RHS can be evaluated as follows:

$$\begin{aligned} \|\hat{\Sigma}_{m,m'} - \tilde{\Sigma}_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} \\ = \left\|\frac{1}{n}(k_m(\cdot,x_j)k_{m'}(x_j,\cdot) - k_m(\cdot,\tilde{x}_j)k_{m'}(\tilde{x}_j,\cdot))\right\|_{\mathcal{H}_m,\mathcal{H}_{m'}}. \end{aligned}$$
(76)

The RHS of (76) can be further evaluated as

$$\begin{aligned} \|\frac{1}{n}(k_{m}(\cdot,x_{j})k_{m'}(x_{j},\cdot)-k_{m}(\cdot,\tilde{x}_{j})k_{m'}(\tilde{x}_{j},\cdot))\|_{\mathcal{H}_{m},\mathcal{H}_{m'}} \\ &\leq \frac{1}{n}(\|k_{m}(\cdot,x_{j})k_{m'}(x_{j},\cdot)\|_{\mathcal{H}_{m},\mathcal{H}_{m'}}+\|k_{m}(\cdot,\tilde{x}_{j})k_{m'}(\tilde{x}_{j},\cdot))\|_{\mathcal{H}_{m},\mathcal{H}_{m'}}) \\ &\leq \frac{1}{n}(\|k_{m}(\cdot,x_{j})\|_{\mathcal{H}_{m}}\|k_{m'}(x_{j},\cdot)\|_{\mathcal{H}_{m'}}+\|k_{m}(\cdot,\tilde{x}_{j})\|_{\mathcal{H}_{m}}\|k_{m'}(\tilde{x}_{j},\cdot))\|_{\mathcal{H}_{m'}}) \\ &\leq \frac{1}{n}(\sqrt{k_{m}(x_{j},x_{j})k_{m'}(x_{j},x_{j})}+\sqrt{k_{m}(\tilde{x}_{j},\tilde{x}_{j})k_{m'}(\tilde{x}_{j},\tilde{x}_{j})}) \\ &\leq \frac{2}{n}, \end{aligned}$$
(77)

where we used $||k_m(\cdot, x_j)||_{\mathcal{H}_m} = \sqrt{\langle k_m(\cdot, x_j), k_m(\cdot, x_j) \rangle_{\mathcal{H}_m}} = \sqrt{k_m(x_j, x_j)}$. Bounding the norm of (76) by (77), we have

$$\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} - \|\tilde{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} \le \frac{2}{n}$$

By symmetry, changing $\hat{\Sigma}$ and $\tilde{\Sigma}$ gives

$$\left\| \hat{\Sigma}_{m,m'} - \Sigma_{m,m'} \right\|_{\mathcal{H}_m,\mathcal{H}_{m'}} - \left\| \tilde{\Sigma}_{m,m'} - \Sigma_{m,m'} \right\|_{\mathcal{H}_m,\mathcal{H}_{m'}} \le \frac{2}{n}.$$

Therefore by McDiarmid's inequality we obtain

$$P\left(\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}} - \mathbb{E}[\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}}] \ge \varepsilon\right)$$

$$\leq \exp\left(-\frac{2\varepsilon^2}{n(2/n)^2}\right) = \exp\left(-\frac{\varepsilon^2 n}{2}\right).$$

This gives the first assertion Eq. (74).

To show the second assertion (Eq. (75)), first we note that

$$E[\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}}] \leq \sqrt{E[\|\hat{\Sigma}_{m,m'} - \Sigma_{m,m'}\|_{\mathcal{H}_m,\mathcal{H}_{m'}}^2]}$$
$$= \sqrt{E[\|(\hat{\Sigma}_{m,m'} - \Sigma_{m,m'})(\hat{\Sigma}_{m',m} - \Sigma_{m',m})\|_{\mathcal{H}_m,\mathcal{H}_m}]}$$
$$\leq \sqrt{E[\|(\hat{\Sigma}_{m,m'} - \Sigma_{m,m'})(\hat{\Sigma}_{m',m} - \Sigma_{m',m})\|_{\mathrm{tr}}]}, \tag{78}$$

where $\|\cdot\|_{tr}$ is the trace norm and the last inequality. As in Lemma 1 of Gretton et al. (2005), we see that

$$\begin{aligned} &\|(\Sigma_{m,m'} - \Sigma_{m,m'})(\Sigma_{m',m} - \Sigma_{m',m})\|_{\mathrm{tr}} \\ &= \frac{1}{n^2} \sum_{i,j=1}^n \|k_m(\cdot, x_i)k_{m'}(x_i, x_j)k_m(x_j, \cdot)\|_{\mathrm{tr}} \\ &- \frac{2}{n} \sum_{i=1}^n \mathrm{E}_X[\|k_m(\cdot, x_i)k_{m'}(x_i, X)k_m(X, \cdot)\|_{\mathrm{tr}}] + \mathrm{E}_{X,X'}[\|k_m(\cdot, X)k_{m'}(X, X')k_m(X', \cdot)\|_{\mathrm{tr}}] \\ &= \frac{1}{n^2} \sum_{i,j=1}^n k_m(x_j, x_i)k_{m'}(x_i, x_j) - \frac{2}{n} \sum_{i=1}^n \mathrm{E}_X[k_m(X, x_i)k_{m'}(x_i, X)] + \mathrm{E}_{X,X'}[k_m(X', X)k_{m'}(X, X')], \end{aligned}$$

where X and X' are independent random variable distributed from Π . Thus

$$\begin{split} & \mathbf{E}[\|(\hat{\Sigma}_{m,m'} - \Sigma_{m,m'})(\hat{\Sigma}_{m',m} - \Sigma_{m',m})\|_{\mathrm{tr}}] \\ &= \frac{n}{n^2} \mathbf{E}_X[k_m(X,X)k_{m'}(X,X)] + \frac{n(n-1)}{n^2} \mathbf{E}_{X,X'}[k_m(X',X)k_{m'}(X,X')] \\ &- 2\mathbf{E}_{X,X'}[k_m(X',X)k_{m'}(X,X')] + \mathbf{E}_{X,X'}[k_m(X',X)k_{m'}(X,X')] \\ &= \frac{1}{n} \mathbf{E}_X[k_m(X,X)k_{m'}(X,X)] - \frac{1}{n} \mathbf{E}_{X,X'}[k_m(X',X)k_{m'}(X,X')] \leq \frac{1}{n}. \end{split}$$

This and Eq. (78) with the first assertion (Eq. (74)) gives the second assertion.

Lemma 11 If $E[\epsilon^2|X] \leq \sigma^2$ almost surely and $\sup_X k_m(X, X) \leq 1$, then we have

$$\|\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m} = O_p(\sigma/\sqrt{n}). \tag{79}$$

Proof: By definition, we have

$$E[\|\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m}] \leq \sqrt{E[\|\hat{\Sigma}_{m,\epsilon}\|_{\mathcal{H}_m}^2]}$$
$$= \sqrt{E\left[\frac{1}{n^2}\sum_{i,j=1}^n k_m(x_i, x_j)\epsilon_i\epsilon_j\right]}$$
$$\leq \sqrt{\frac{\sigma^2}{n}}.$$

Applying Markov's inequality we obtain the assertion.

Proposition 1 (Bernstein's inequality in Hilbert spaces) Let (Ω, \mathcal{A}, P) be a probability space, \mathcal{H} be a separable Hilbert space, $\mathcal{B} > 0$, and $\sigma > 0$. Furthermore, let $\xi_1, \ldots, \xi_n : \Omega \to \mathcal{H}$ be independent random variables satisfying $\mathbb{E}[\xi_i] = 0$, $\|\xi\|_{\mathcal{H}} \leq B$, and $\mathbb{E}[\|\xi_i\|_{\mathcal{H}}^2] \leq \sigma^2$ for all $i = 1, \ldots, n$. Then we have

$$P\left(\left\|\frac{1}{n}\sum_{i=1}^{n}\xi_{i}\right\|_{\mathcal{H}} \geq \sqrt{\frac{2\sigma^{2}\tau}{n}} + \sqrt{\frac{\sigma^{2}}{n}} + \frac{2B\tau}{3n}\right) \leq e^{-\tau}, \quad (\tau > 0).$$

Proof: See Theorem 6.14 of Steinwart (2008).

References

- F. Bach, G. Lanckriet, and M. Jordan. Multiple kernel learning, conic duality, and the SMO algorithm. In *the* 21st International Conference on Machine Learning, pages 41–48, 2004.
- F. R. Bach. Consistency of the group lasso and multiple kernel learning. *Journal of Machine Learning Research*, 9:1179–1225, 2008.
- C. R. Baker. Joint measures and cross-covariance operators. Transactions of the American Mathematical Society, 186:273–289, 1973.
- P. Bartlett, O. Bousquet, and S. Mendelson. Local Rademacher complexities. *The Annals of Statistics*, 33: 1487–1537, 2005.
- P. J. Bickel, Y. Ritov, and A. B. Tsybakov. Simultaneous analysis of Lasso and Dantzig selector. *The Annals of Statistics*, 37(4):1705–1732, 2009.
- O. Bousquet. A Bennett concentration inequality and its application to suprema of empirical process. C. R. Acad. Sci. Paris Ser. I Math., 334:495–500, 2002.
- A. Caponnetto and E. de Vito. Optimal rates for regularized least-squares algorithm. *Foundations of Computational Mathematics*, 7(3):331–368, 2007.
- C. Cortes. Can learning kernels help performance?, 2009. Invited talk at International Conference on Machine Learning (ICML 2009). Montréal, Canada, 2009.

- C. Cortes, M. Mohri, and A. Rostamizadeh. L₂ regularization for learning kernels. In *the 25th Conference* on Uncertainty in Artificial Intelligence (UAI 2009), 2009. Montréal, Canada.
- L. Devroye, L. Györfi, and G. Lugosi. A Probabilistic Theory of Pattern Recognition. Springer, 1996.
- A. Gretton, O. Bousquet, A. Smola, and B. Schölkopf. Measuring statistical dependence with Hilbert-Schmidt norms. In S. Jain, H. U. Simon, and E. Tomita, editors, *Algorithmic Learning Theory*, Lecture Notes in Artificial Intelligence, pages 63–77, Berlin, 2005. Springer-Verlag.
- J. Jia and B. Yu. On model selection consistency of the elastic net when $p \gg n$. *Statistica Sinica*, 20(2):to appear, 2010.
- M. Kloft, U. Brefeld, S. Sonnenburg, P. Laskov, K.-R. Müller, and A. Zien. Efficient and accurate ℓ_p -norm multiple kernel learning. In *Advances in Neural Information Processing Systems* 22, pages 997–1005, Cambridge, MA, 2009. MIT Press.
- V. Koltchinskii. Local Rademacher complexities and oracle inequalities in risk minimization. *The Annals of Statistics*, 34:2593–2656, 2006.
- V. Koltchinskii and M. Yuan. Sparse recovery in large ensembles of kernel machines. In Proceedings of the Annual Conference on Learning Theory, pages 229–238, 2008.
- G. Lanckriet, N. Cristianini, L. E. Ghaoui, P. Bartlett, and M. Jordan. Learning the kernel matrix with semidefinite programming. *Journal of Machine Learning Research*, 5:27–72, 2004.
- M. Ledoux and M. Talagrand. *Probability in Banach Spaces. Isoperimetry and Processes.* Springer, New York, 1991. MR1102015.
- Y. Lin and H. H. Zhang. Component selection and smoothing in multivariate nonparametric regression. *The Annals of Statistics*, 34(5):2272–2297, 2006.
- L. Meier, S. van de Geer, and P. Bühlmann. High-dimensional additive modeling. *The Annals of Statistics*, 37(6B):3779–3821, 2009.
- S. Mendelson. Improving the sample complexity using global data. *IEEE Transactions on Information Theory*, 48:1977–1991, 2002.
- C. A. Micchelli and M. Pontil. Learning the kernel function via regularization. *Journal of Machine Learning Research*, 6:1099–1125, 2005.
- A. Rakotomamonjy, F. Bach, S. Canu, and G. Y. SimpleMKL. *Journal of Machine Learning Research*, 9: 2491–2521, 2008.
- S. Sonnenburg, G. Rätsch, C. Schäfer, and B. Schölkopf. Large scale multiple kernel learning. Journal of Machine Learning Research, 7:1531–1565, 2006.
- I. Steinwart. Support Vector Machines. Springer, 2008.
- I. Steinwart, D. Hush, and C. Scovel. Optimal rates for regularized least squares regression. In *Proceedings* of the Annual Conference on Learning Theory, pages 79–93, 2009.
- M. Stone. Cross-validatory choice and assessment of statistical predictions. *Journal of the Royal Statistical Society, Series B*, 36:111–147, 1974.
- T. Suzuki and R. Tomioka. Spicymkl, 2009. arXiv:0909.5026.
- M. Talagrand. A new look at independence. The Annals of Statistics, 24:1–34, 1996a.
- M. Talagrand. New concentration inequalities in product spaces. *Inventiones Mathematicae*, 126:505–563, 1996b.
- R. Tomioka and T. Suzuki. Sparsity-accuracy trade-off in MKL, 2010. arXiv:1001.2615.
- S. van de Geer. Empirical Processes in M-Estimation. Cambridge University Press, 2000.
- A. W. van der Vaart and J. A. Wellner. *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer, New York, 1996.

- V. N. Vapnik. Statistical Learning Theory. Wiley, New York, 1998.
- M. Yuan and Y. Lin. On the nonnegative garrote estimator. *Journal of the Royal Statistical Society B*, 69(2): 143–161, 2007.
- T. Zhang. Some sharp performance bounds for least squares regression with l_1 regularization. *The Annals of Statistics*, 37(5):2109–2144, 2009.
- P. Zhao and B. Yu. On model selection consistency of lasso. *Journal of Machine Learning Research*, 7: 2541–2563, 2006.
- H. Zou and T. Hastie. Regularization and variable selection via the elastic net. *Journal of the Royal Statistical: Series B*, 67(2):301–320, 2005.
- H. Zou and H. H. Zhang. On the adaptive elastic-net with a diverging number of parameters. *The Annals of Statistics*, 37(4):1733–1751, 2009.