# **Fractional Modified Special Relativity**

Hosein Nasrolahpour\*

Department of Physics, Faculty of Basic Sciences, University of Mazandaran, P. O. Box 47416-95447, Babolsar, IRAN

Hadaf Institute of Higher Education, P.O. Box 48175 – 1666, Sari, Mazandaran, IRAN

#### **Abstract:**

Fractional calculus represents a natural tool for describing relativistic phenomena in pseudo-Euclidean space-time. In this study, Fractional modified special relativity is presented. We obtain fractional generalized relation for the time dilation.

Keywords: Special relativity; Time dilation; Fractional special relativity; Fractional calculus

#### 1- Introduction

The Theory of Special Relativity was first published as a paper in 1905[1]. Recently, some modifications on this theory have been proposed. For example, in ref. [3] some aspects of nonlocal special relativity have been studied. Dissipative Lagrangian for a relativistic particle also has been discussed in ref. [4].

In more recent years attention has been attracted to the application of Fractional Calculus (FC) to special (and general) relativity [9-13]. Fractional calculus can be used to model the nonlocal dissipative phenomena in space (-time) [7, 8]. In particular FC represents a natural tool for describing relativistic phenomena in pseudo-Euclidean space-time [9, 11].

The aim of this short paper is to introduce fractional modified special relativity as a new nonlocal dissipative model for the special theory of relativity. In the following, fractional calculus [5, 6] is briefly reviewed in Sec. 2. Fractional generalized relation for the time dilation is presented in Sec. 3. Finally, in Sec. 4, we will present some conclusions and discussion.

### 2-Fractional calculus: Riemann-Liouvill integral operator

Fractional calculus is the calculus of derivatives and integrals with arbitrary order, which unify and generalize the notions of integer order differentiation and n-fold integration. In contrast whit the definition of the ordinary derivative and integration, the definition of the fractional ones is not unique. In fact, there exists several definitions including, Grünwald - Letnikov, Riemann - Liouville, Weyl, Riesz and Caputo for fractional derivative and integral. For the purpose of this study we use the left-sided Riemann-Liouville fractional integral operator. The left-sided Riemann - Liouville fractional integral of order  $\alpha$  of a function f (t) is defined by [5, 6]

$${}_{0}^{RL}I_{t}^{\alpha}f(t) = \frac{1}{\Gamma(\alpha)}\int_{0}^{t}(t-\tau)^{\alpha-1}f(\tau)d\tau \qquad t > 0, \ \alpha \in \mathbb{R}^{+}$$

<sup>\*</sup> E-mail address: hnasrolahpour@gmail.com; h.nasrolah-pour@umz.ac.ir

Where,  $\mathbb{R}^+$  is the set of positive real numbers and  $\Gamma(\alpha)$  \*denotes the gamma function.

## 3-Time dilation for fractional special relativity

We start with the following expression for the fractional generalized position

$$x(t) = {}^{RL}_{0}I_{t}^{\alpha}(v_{\alpha}) \qquad 0 < \alpha \le 1$$

By use of the above equation for the fractional generalized position we can easily derive dilated time formula according to the famous light clock experiment [2] as below

$$t = \frac{\sqrt{\binom{RL}{0}I_{t}^{\alpha}(v_{\alpha})^{2} + (ct_{0})^{2}}}{C}$$
 (3)

Where c is the speed of light in vacuum and  $t_0$  is proper time. By use of Eq. (1), we can obtain the following relationship between  $t_0$  (proper time) and t (dilated time):

$$t^2 - \frac{\beta^2 t^{2\alpha}}{\alpha^2 \Gamma^2(\alpha)} = t_0^2 \tag{4}$$

We can write Eq. (4) as

$$t^{2} - \frac{\beta^{2} t^{2\alpha}}{\Gamma^{2}(\alpha + 1)} = t_{0}^{2}$$
 (5)

where 
$$\beta = \frac{v_{\alpha}}{c}$$
. Now we can see that as  $\alpha \to 1$ , Eq. (4) gives 
$$t^2 - \beta^2 t^2 = t_0^2$$
 (6)

this leads to well-known time dilation formula of ordinary special relativity:

$$t = \frac{t_0}{\sqrt{1-\beta^2}}. (7)$$

If we plot t (dilated time) with respect to  $t_0$  for the special case of v = 0.99c we will have

<sup>\*</sup> Definition of Gamma function is given by  $\Gamma(\alpha) = \int_0^\infty e^{-t} t^{\alpha-1} dt$ ,  $\alpha \in \mathbb{R}^+$ . For this function we have  $\Gamma(\alpha+1) = \alpha \Gamma(\alpha)$ .




Fig. 1. t with respect to  $t_0$  for the special case of v = 0.99c,  $\alpha = 1$ 

With the aid of expression (5) we can find the t (dilated time) versus  $t_0$  (proper time) for the arbitrary case of  $\alpha$ . In particular, for the special case of  $\alpha = \frac{1}{2}$  we can draw t versus  $t_0$ 




Fig. 2. t with respect to  $t_0$  for the special case of v = 0.99c,  $\alpha = \frac{1}{2}$ 

As shown in this figure, there is a major difference between Fig.1 and Fig.2 near the origin of coordinate system. We can interpret this anomalous behaviour as an immediate consequence of applications of fractional calculus in our model.

### 4- CONCLUSION AND DISCUSSION

Fractional calculus can be used to model the nonlocal dissipative phenomena in space (-time). In particular FC represents an effective tool for describing relativistic phenomena in pseudo-Euclidean space-time. With this motivation we obtain fractional generalized relation for the well-known time dilation formula, to show the possibilities for future studies on fractional modified special theory of relativity theory. As we shown in Fig. (2), there is a anomalous behaviour when we plot t with respect to  $t_0$  for the arbitrary case of  $\alpha$  ( $0 < \alpha < 1$ ) near the origin of coordinate system. We hope to study some other aspects of Fractional modified special relativity in future.

### **Rrefrences**

- [1] A. Einstein, Zur Elektrodynamik bewegeter Körper. Annalen der Physik 1905, **17**. (English translation: On the electrodynamics of moving bodies, reprinted in Lorentz et al., 1923.)
- [2] Robert Resnick, Introduction to Special Relativity (John Wiley and Sons, Inc. New York, London, Sydney 1968.)
  - A.P.French, Special Relativity (Nelson, London, 1968).
  - Miller, A.I., Albert Einstein's Special Theory of Relativity, Springer, 1998.
- [3] B. Mashhoon, Nonlocal special relativity, Ann. Phys. (Berlin) 17(2008)11705.
- [4] G. Gonzalez, Relativistic motion with linear dissipation. Int. J. Theor. Phys. 46 (2007) 417–423.
- [5] I. Podlubny, Fractional Differential Equations Academic Press, New York, 1999.
- [6] R. Herrmann, Fractional Calculus: An Introduction for Physicists, World Scientific Press, 2011.
  - K. S. Miller, B. Ross, An introduction to the fractional calculus and fractional differential equations, *John Wiley & Sons*, Inc., 1993.
  - K. B. Oldham, J. Spanier, the Fractional Calculus, Academic Press, NewYork, 1974.
  - S. G. Samko, A. A. Kilbas, O. I. Marichev, Fractional Integrals and derivatives, Theory and Applications, *Gordon and Breach, Yverdon*, 1993.
- [7] L. Vázquez, "A Fruitful Interplay: From Nonlocality to Fractional Calculus" en "Nonlinear Waves, Classical and Quantum Aspects", 129-133, Ed.: V.V. Konotop y F. Abdullaev, NATO Science Series, Kluwer Academic Publishers 2004.
- [8] V.E.Tarasov, Fractional Vector Calculus and Fractional Maxwell's Equations, Ann. Phys. **323** (2008) 2756–2778.
- [9] E. Goldfain, Fractional dynamics and the TeV regime of field theory, *Comm. Non. Sci. Num. Siml.* **13** (2008) 666–676.
- [10] E. Goldfain, Fractional dynamics, Cantorian space—time and the gauge hierarchy problem, *Chaos, Solitons & Fractals*, **22**(2004) 513-520.
- [11] V. E. Tarasov, Fractional Dynamics of Relativistic Particle, *Int. J. Theor. Phys.* **49**(2010)293-303.
- [12] R. A. EL-Nabulsi, Fractional Nottale's Scale Relativity and emergence of complexified gravity, Chaos, Solitons and Fractals **42** (2009) 2924–2933.

[13] R. A. EL-Nabulsi, Fractional illusion theory of space: Fractional gravitational field with fractional extra-dimensions, Chaos, Solitons and Fractals **42** (2009) 377–384.