

A novel strong coupling expansion of the QCD Hamiltonian

H.-P. Pavel^{1, 2, *}

¹*Institut für Kernphysik, Technische Universität Darmstadt, D-64289 Darmstadt, Germany*

²*Bogoliubov Laboratory of Theoretical Physics, JINR Dubna, 141980 Dubna, Russia*

Introducing an infinite spatial lattice with box length a , a systematic expansion of the physical QCD Hamiltonian in $\lambda = g^{-2/3}$ can be obtained. The free part is the sum of the Hamiltonians of the quantum mechanics of spatially constant fields for each box, and the interaction terms proportional to λ^n with contain n discretised spatial derivatives connecting different boxes. As an example, the energy of the vacuum and the lowest scalar glueball is calculated up to order λ^2 for the case of $SU(2)$ Yang-Mills theory.

I. INTRODUCTION

The quantum Hamiltonian of $SU(2)$ Yang-Mills theory, to which we limit ourselves here for simplicity, can be obtained [1] by exploiting the time-dependence of the gauge transformations to put $A_{a0}(x) = 0$, ($a = 1, 2, 3$), and quantizing the spatial fields in the Schrödinger representation, $\Pi_{ai}(\mathbf{x}) = -E_{ai}(\mathbf{x}) = -i\delta/\delta A_{ai}(\mathbf{x})$. The physical states Ψ have to satisfy the Schrödinger equ. and the three non-Abelian Gauss law constraints

$$H\Psi = E\Psi, \quad H = \int d^3\mathbf{x} \frac{1}{2} \sum_{a,i} \left[\left(\frac{\delta}{\delta A_{ai}(\mathbf{x})} \right)^2 + B_{ai}^2(A(\mathbf{x})) \right] \quad (1)$$

$$G_a(\mathbf{x})\Psi = 0, \quad G_a(\mathbf{x}) = -i(\delta_{ac}\partial_i + g\epsilon_{abc}A_{bi}(\mathbf{x})) \frac{\delta}{\delta A_{ci}(\mathbf{x})} \quad (2)$$

with the chromo-magnetic fields $B_{ai}(A) = \epsilon_{ijk} \left(\partial_j A_{ak} + \frac{1}{2} g\epsilon_{abc} A_{bj} A_{ck} \right)$ and the generators $G_a(\mathbf{x})$ of the residual time-independent gauge transformations, satisfying $[G_a(\mathbf{x}), H] = 0$, and $[G_a(\mathbf{x}), G_b(\mathbf{y})] = ig\delta^3(\mathbf{x} - \mathbf{y})\epsilon_{abc}G_c(\mathbf{x})$, The matrix elements have Cartesian measure

$$\langle \Phi_1 | \mathcal{O} | \Phi_2 \rangle = \int \prod_{\mathbf{x}} \prod_{ik} dA_{ik}(\mathbf{x}) \Phi_1^* \mathcal{O} \Phi_2. \quad (3)$$

In order to implement the Gauss laws (2) into (1), to obtain the physical Hamiltonian, it is very useful to Abelianise them by a suitable point transformation of the gauge fields.

*Electronic address: hans-peter.pavel@physik.tu-darmstadt.de

II. THE PHYSICAL HAMILTONIAN OF SU(2) YANG-MILLS THEORY AND ITS STRONG COUPLING EXPANSION

Point transformation to the new set of adapted coordinates [2], the 3 q_j ($j = 1, 2, 3$) and the 6 elements $S_{ik} = S_{ki}$ ($i, k = 1, 2, 3$) of the positive definite symmetric 3×3 matrix S ,

$$A_{ai}(q, S) = O_{ak}(q) S_{ki} - \frac{1}{2g} \epsilon_{abc} \left(O(q) \partial_i O^T(q) \right)_{bc}, \quad (4)$$

where $O(q)$ is an orthogonal 3×3 matrix parametrised by the q_i , leads to an Abelianisation of the Gauss law constraints

$$G_a \Phi = 0 \quad \Leftrightarrow \quad \frac{\delta}{\delta q_i} \Phi = 0 \quad (\text{Abelianisation})$$

Equ. (4) corresponds to the symmetric gauge $\chi_i(A) = \epsilon_{ijk} A_{jk} = 0$. It has been proven in [3] that, at least for strong coupling, the symmetric gauge exists, i.e. any time-independent gauge field can be carried over uniquely into the symmetric gauge, and in [2] that both indices of the tensor field S are spatial indices, leaving S a colorless local field.

According to the general scheme of [1], the correctly ordered physical quantum Hamiltonian in the symmetric gauge in terms of the colorless physical variables $S_{ik}(\mathbf{x})$ and the corresponding canonically conjugate momenta $P_{ik}(\mathbf{x}) \equiv -i\delta/\delta S_{ik}(\mathbf{x})$ reads [4]

$$H(S, P) = \frac{1}{2} \mathcal{J}^{-1} \int d^3 \mathbf{x} P_{ai} \mathcal{J} P_{ai} + \frac{1}{2} \int d^3 \mathbf{x} (B_{ai}(S))^2 - \mathcal{J}^{-1} \int d^3 \mathbf{x} \int d^3 \mathbf{y} \left\{ \left(D_i(S)_{ma} P_{im} \right)(\mathbf{x}) \mathcal{J} \langle \mathbf{x} a | {}^* D^{-2}(S) | \mathbf{y} b \rangle \left(D_j(S)_{bn} P_{nj} \right)(\mathbf{y}) \right\} \quad (5)$$

with the covariant derivative $D_i(S)_{kl} \equiv \delta_{kl} \partial_i - g \epsilon_{klm} S_{mi}$, the Faddeev-Popov (FP) operator

$${}^* D_{kl}(S) \equiv \epsilon_{kmi} D_i(S)_{ml} = \epsilon_{kli} \partial_i - g \gamma_{kl}(S), \quad \gamma_{kl}(S) \equiv S_{kl} - \delta_{kl} \text{tr} S \quad (6)$$

and the Jacobian $\mathcal{J} \equiv \det |{}^* D|$. The matrix element of a physical operator O is given by

$$\langle \Psi' | O | \Psi \rangle \propto \int \prod_{\mathbf{x}} [dS(\mathbf{x})] \mathcal{J} \Psi'^*[S] O \Psi[S]. \quad (7)$$

The inverse of the FP operator can be expanded in the number of spatial derivatives

$$\begin{aligned} \langle \mathbf{x} k | {}^* D^{-1}(S) | \mathbf{y} l \rangle &= -\frac{1}{g} \gamma_{kl}^{-1}(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}) + \frac{1}{g^2} \gamma_{ka}^{-1}(\mathbf{x}) \epsilon_{abc} \partial_c^{(\mathbf{x})} \left[\gamma_{bl}^{-1}(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}) \right] \\ &\quad - \frac{1}{g^3} \gamma_{ka}^{-1}(\mathbf{x}) \epsilon_{abc} \partial_c^{(\mathbf{x})} \left[\gamma_{bi}^{-1}(\mathbf{x}) \epsilon_{ijk} \partial_k^{(\mathbf{x})} \left[\gamma_{jl}^{-1}(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}) \right] \right] + \dots \end{aligned} \quad (8)$$

In order to perform a consistent expansion, also the non-locality in the Jacobian \mathcal{J} has to be taken into account [4]. The Jacobian \mathcal{J} factorizes $\mathcal{J} = \mathcal{J}_0 \tilde{\mathcal{J}}$ with the local

$$\mathcal{J}_0 \equiv \det |\gamma| = \prod_{\mathbf{x}} \prod_{i < j} (\phi_i(\mathbf{x}) + \phi_j(\mathbf{x})), \quad (\phi_i = \text{eigenvalues of } S) \quad (9)$$

and the non-local $\tilde{\mathcal{J}}$, which can be included into the wave functional $\tilde{\Psi}(S) := \tilde{\mathcal{J}}^{-1/2} \Psi(S)$ leading to the corresponding transformed Hamiltonian

$$\tilde{H}(S, P) := \tilde{\mathcal{J}}^{1/2} H(S, P) \tilde{\mathcal{J}}^{-1/2} = H(S, P) \Big|_{J \rightarrow J_0} + V_{\text{measure}}(S) . \quad (10)$$

It is Hermitean with respect to the local measure \mathcal{J}_0 on the cost of extra terms V_{measure} , and can be expanded in the number of spatial derivatives using (8).

Next, an ultraviolet cutoff a is put by introducing an infinite spatial lattice of granulas $G(\mathbf{n}, a)$, here cubes of length a , situated at sites $\mathbf{x} = a\mathbf{n}$ ($\mathbf{n} \in Z^3$), and considering the averaged variables

$$S(\mathbf{n}) := \frac{1}{a^3} \int_{G(\mathbf{n}, a)} d\mathbf{x} S(\mathbf{x}) \quad (11)$$

and discretised spatial derivatives relating the $S(\mathbf{n})$ of different granulas (see [4] for details).

After an appropriate rescaling of the dynamical fields a novel strong coupling expansion of the Hamiltonian in $\lambda = g^{-2/3}$ can be obtained [4]

$$\tilde{H} = \frac{g^{2/3}}{a} \left[\mathcal{H}_0 + \lambda \sum_{\alpha} \mathcal{V}_{\alpha}^{(\partial)} + \lambda^2 \left(\sum_{\beta} \mathcal{V}_{\beta}^{(\Delta)} + \sum_{\gamma} \mathcal{V}_{\gamma}^{(\partial\partial \neq \Delta)} \right) + \mathcal{O}(\lambda^3) \right] . \quad (12)$$

as an alternative to existing strong coupling expansions [5] and [6] based on Wilsonian lattice QCD. The "free part" in (12) is just the sum of Hamiltonians $\mathcal{H}_0 = \sum_{\mathbf{n}} \mathcal{H}_0^{QM}(\mathbf{n})$ of Yang-Mills quantum mechanics of spatially constant fields [7]-[10] at each site, and the $\mathcal{V}_{\alpha}^{(\partial)}$ and $\mathcal{V}_{\beta}^{(\Delta)}$ are interaction parts, relating different sites. The local measure $\mathcal{J}_0 = \prod_{\mathbf{n}} \mathcal{J}_0^{QM}(\mathbf{n})$ is correspondingly the product of the quantum mechanical measures at each site. In terms of the principal-axes variables of the positive definite symmetric 3×3 matrix field S

$$S = R^T(\alpha, \beta, \gamma) \text{diag}(\phi_1, \phi_2, \phi_3) R(\alpha, \beta, \gamma) , \quad (13)$$

with the $SO(3)$ matrix R parametrized by the three Euler angles $\chi = (\alpha, \beta, \gamma)$, we find

$$\mathcal{J}_0^{QM} \rightarrow \sin \beta \prod_{i < j} (\phi_i^2 - \phi_j^2) \rightarrow 0 < \phi_1 < \phi_2 < \phi_3 \quad (\text{principle orbits}) . \quad (14)$$

and (with the intrinsic spin angular momenta ξ_i)

$$\mathcal{H}_0^{QM} = \frac{1}{2} \sum_{ijk}^{\text{cyclic}} \left[\pi_i^2 - \frac{2i}{\phi_j^2 - \phi_k^2} (\phi_j \pi_j - \phi_k \pi_k) + \xi_i^2 \frac{\phi_j^2 + \phi_k^2}{(\phi_j^2 - \phi_k^2)^2} + \phi_j^2 \phi_k^2 \right] . \quad (15)$$

Its low energy spectrum and eigenstates at any site \mathbf{n}

$$\mathcal{H}_0^{QM}(\mathbf{n}) |\Phi_{i,M}^{(S)\pm}\rangle_{\mathbf{n}} = \epsilon_i^{(S)\pm}(\mathbf{n}) |\Phi_{i,M}^{(S)\pm}\rangle_{\mathbf{n}} , \quad (16)$$

characterised by the quantum numbers of spin S, M and parity P , are known with high accuracy [10]. Hence the eigenstates of \mathcal{H}_0 in (12) are free glueball excitations of the lattice. The interactions \mathcal{V} (12) can be included using perturbation theory in λ .

III. CALCULATION OF THE GLUEBALL SPECTRUM UP TO ORDER λ^2

Using 1st and 2nd order perturbation theory in λ give the results [4]

$$E_{\text{vac}}^+ = \mathcal{N} \frac{g^{2/3}}{a} \left[4.1167 + 29.894\lambda^2 + \mathcal{O}(\lambda^3) \right], \quad (17)$$

for the energy of the interacting glueball vacuum and

$$E_1^{(0)+}(k) - E_{\text{vac}}^+ = \left[2.270 + 13.511\lambda^2 + \mathcal{O}(\lambda^3) \right] \frac{g^{2/3}}{a} + 0.488 \frac{a}{g^{2/3}} k^2 + \mathcal{O}((a^2 k^2)^2), \quad (18)$$

for the energy spectrum of the interacting spin-0 glueball, up to λ^2 for the (+) b.c. and similar results for the (-) b.c. The first, zeroth order numbers, correspond to the result of Yang-Mills quantum mechanics. Note that Lorentz invariance asks for energy momentum relation $E = \sqrt{M^2 + k^2} \simeq M + (2M)^{-1} k^2$. The result, shown here for the scalar glueball, which limits itself to the terms in the Hamiltonian containing the Laplace-operator Δ as a first step, violates this condition by about a factor of two. Including all spin-orbit coupling terms in the Hamiltonian dropped in this first approach and considering all possible $J = L + S = 0$ states is expected restore Lorentz invariance.

To study the coupling constant renormalisation in the IR, consider the physical glueball mass

$$M = \frac{g_0^{2/3}}{a} \left[\mu + c g_0^{-4/3} \right]. \quad (19)$$

Independence of the box size a is given for the two cases, $g_0 = 0$ or $g_0^{4/3} = -c/\mu$. The first solution corresponds to the perturbative fixed point, and the second, if it exists ($c < 0$), to an infrared fixed point. My result for the lowest spin-0 glueball $c_1^{(0)}/\mu_1^{(0)} = 5.95$ suggests, that no infrared fixed points exist, in accordance with the corresponding result of Wilsonian lattice QCD [11]. Solving the above equation (19) for positive ($c > 0$) one obtains

$$g_0^{2/3}(Ma) = \frac{Ma}{2\mu} + \sqrt{\left(\frac{Ma}{2\mu}\right)^2 - \frac{c}{\mu}}, \quad a > a_c := 2\sqrt{c\mu}/M \quad (20)$$

with the physical glueball mass M . For a typical [12] $M \sim 1.6$ GeV we find $a_c \sim 1.4$ fm. Comparing the behaviour of the bare coupling constant (20), obtained for boxes of large size a , with those obtained for small boxes in [7, 8], should lead to information about the intermediate region, including the possibility of the existence of phase transitions.

The generalisation of the approach to include quarks is possible [13] and under current investigation.

Acknowledgments

I would like to thank A. Dorokhov and J. Wambach for their interest. Financial support by the LOEWE-Program HIC for FAIR is gratefully acknowledged.

-
- [1] N.H. Christ and T.D. Lee, Phys. Rev. D **22**, 939 (1980).
 - [2] A.M. Khvedelidze and H.-P. Pavel, Phys. Rev. D **59**, 105017 (1999).
 - [3] A.M. Khvedelidze, D. M. Mladenov, H.-P. Pavel, and G. Röpke, Phys. Rev. D **67**, 105013 (2003).
 - [4] H.-P. Pavel, Phys. Lett. B **685**, 353 (2010).
 - [5] J. Kogut, D.K. Sinclair, L. Susskind, Nucl. Phys. B **114**, 199 (1976).
 - [6] G. Münster, Nucl. Phys. B **190** [FS3], 439 (1981) (E: B **200** [FS4], 536 (1982) and E: B **205**, 648 (1982)).
 - [7] M. Lüscher, Nucl. Phys. B **219**, 233 (1983).
 - [8] M. Lüscher and G. Münster, Nucl. Phys. B **232**, 445 (1984).
 - [9] J. Koller and P. van Baal, Nucl. Phys. B **273**, 387 (1986); Nucl. Phys. B **302**, 1 (1988); P. van Baal and J. Koller, Ann. of Phys. (N.Y.) **174**, 299 (1987).
 - [10] H.-P. Pavel, Phys. Lett. B **648**, 97 (2007).
 - [11] M. Creutz, *Quarks, Gluons and Lattices*, (Cambridge Univ. Press, Cambridge, 1983).
 - [12] B. Lucini, M. Teper and U. Wenger, JHEP **0406**, 012 (2004).
 - [13] H.-P. Pavel, Phys. Lett. B **700**, 265 (2011).