

What is the Real Structure of Fundamental Forces?

H.P. Morsch

HOFF, Brockmüllerstr. 11, D-52428 Jülich, Germany

E-mail: h.p.morsch@gmx.de

Abstract

Based on general considerations, the Standard Model of particle physics with its extensions (SM) can be ruled out as a valid theory of fundamental forces: it requires far too many parameters, which are not determined from first principles. The only way to uncover the real structure of elementary forces is to resort to a complete and fundamental theory **without** external parameters.

This requirement is fulfilled in a finite theory, based on an extension of the QED Lagrangian by boson-boson coupling, with massless elementary fields, charged and neutral elementary fermions (quantons) and gauge bosons. In this description the observed bound states of nature, hadrons, leptons, atoms and gravitational objects are understood as stationary systems of quantons bound by electric or magnetic forces. Other fields do not exist.

PACS/ keywords: 11.15.-q, 12.10.-g/ Description of elementary forces in a complete (fundamental) theory, based on a Lagrangian with Maxwell term, boson-boson coupling and massless elementary fermions (quantons). Bound states of nature described by systems bound by electric or magnetic forces.

The study of the fundamental forces of nature allows to gain insight into the early development of the universe. But this requires a close theoretical description, in which all model assumptions can be verified and the needed parameters are derived from first principles. Up to now the Standard Model of particle physics [1] (SM), established about 40 years ago, represents the state of our knowledge. It is a heuristic model constructed from relativistic first order quantum field theories, different for each fundamental force (gravitation is not included in the SM, since a quantum theory of gravity is not well established), with a number of fields of different symmetry. Further, the understanding of the masses of elementary fermions requires background (Higgs) fields, but for neutrino masses

in addition the postulation of heavy Majorana neutrinos is needed. The flavour degree of freedom observed in hadrons and leptons is understood in supersymmetric extensions of the SM. In particular, the minimal supersymmetric SM (MSSM) has been favoured, since it allows to understand the hierarchy problem between the weak and gravitational forces. Further, in this model the running couplings of the electric, weak and strong forces meet on the grand unification scale $\Lambda_{GUT} = 10^{16}$ GeV. However, in the MSSM supersymmetric particles have been predicted in the few hundred GeV region, which have not been found experimentally. These negative results cast serious doubts on the validity of the SM including its proposed extensions. Since it is not very probable that other grand unified theories can resolve these problems, it is time to examine what model assumptions may be unrealistic or even wrong.

A serious point of criticism of the SM is that very different theories are assumed for each fundamental interaction. This gives rise to a large complexity¹ in the weak and strong interaction sectors with fields of different symmetry, 12 elementary fermions and quite a few massless but also massive gauge bosons. The consequence of this construction is that a large number of parameters have to be adjusted (the masses of elementary particles, mixing parameters, couplings between fields, etc.). An understanding of most of these parameters (those, which cannot be determined from first principles) requires further theoretical explanations² with still other parameters, which again have to be understood. This leads to an endless chain of related parameters without clearly defined solution, proving that the assumptions in the SM are erroneous or unrealistic.

Examples of such parameters are mass and flavour of elementary particles. Their explanation requires additional background (Higgs) and supersymmetric fields. The couplings of the Higgs-fields to massless fermions have to be considered as additional parameters, which could be understood only by a further theory with additional rather uncertain parameters. Even worse, supersymmetry gives rise to many combinations of fermion and boson fields with mixing parameters of large uncertainties, which cannot be resolved from first principles. This indicates clearly that these explanations cannot be correct, but also that a realistic theory should have a much simpler structure with very few parameters.

¹It should have been questioned, whether nature is not more economic and efficient to organise the creation of hadrons and leptons – the constituents of matter – in a much simpler way.

²SM parameters should not just be adjusted to experimental data, they have to be understood!

The only way to avoid the above problems is to resort to a complete and fundamental theory, in which all parameters are determined from first principles. Such a theory should have only a minimum of fields and thus significantly less degrees of freedom than the SM. Also in such a theory relativistic bound states have to be described explicitly.

If we inspect the structure of the SM, even quantum electrodynamics (QED), the best established part of the SM, cannot be regarded as a closed and fundamental theory, since the coupling constant $\alpha_{QED} \sim 1/137$ is an external parameter determined from experiment. However, the precise prediction of spin properties suggests that QED is close to a fundamental theory. Only the divergencies for $r \rightarrow 0$ and ∞ appear to be in conflict with nature, which is known to develop in a smooth and finite way. Because of this, a complete and fundamental theory is expected to be similar to QED, but with a structure more complex than a first order gauge theory.

During the last years a theoretical framework has been developed [2, 3, 4], based on a second order extension of QED, which shows all features expected of a fundamental theory: all needed parameters can be determined by self-consistency conditions, thus showing completeness. With massless elementary bosons and fermions (quantons) this model shows a coupling to the vacuum with average energy $E_{vac} = 0$. As a crucial test, by applying this model to the binding of light atoms [4], the self-consistently deduced coupling constant α was found to be consistent with α_{QED} . This is needed, since the Coulomb potential³ yields eigenvalues in agreement with the experimental spectrum.

It has to be noted that in the past many different Lagrangians have been studied, with the general conclusion that higher order theories should be discarded, because they could lead to unphysical solutions [5]. However, the important difference to the higher order theories discussed in ref. [5] is that in the present formalism the Lagrangian is gauge invariant and non-physical solutions can be eliminated by strict geometric, mass-radius and energy-momentum relations. Further, only in a second order formalism relativistic bound states can be generated, which are stable, see the discussion below.

The Lagrangian has been used in the form

$$\mathcal{L} = \frac{1}{\tilde{m}^2} \bar{\Psi} i \gamma_\mu D^\mu D_\nu D^\nu \Psi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} , \quad (1)$$

³The Coulomb potential has to be considered as effective potential, since the restoring force is taken over by the potential itself.

where \tilde{m} is a mass parameter and Ψ a two-component massless fermion field (with charge and neutral part), $\Psi = (\Psi^+, \Psi^o)$ and $\bar{\Psi} = (\Psi^-, \bar{\Psi}^o)$. Vector boson fields A_μ with coupling g to fermions are contained in the covariant derivatives $D_\mu = \partial_\mu - igA_\mu$. The second term of the Lagrangian represents the Maxwell term with Abelian field strength tensors $F^{\mu\nu}$ given by $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$, which gives rise to both electric and magnetic effects.

By inserting $D^\mu = \partial^\mu - igA^\mu$ and $D_\nu D^\nu = \partial_\nu \partial^\nu - ig(A_\nu \partial^\nu + \partial_\nu A^\nu) - g^2 A_\nu A^\nu$ in eq. (1), the first term of \mathcal{L} gives rise to a number of different terms, which contain boson and fermion fields and/or their derivatives, see ref. [3]. All terms, which contain the derivative of the fermion field $\partial^\nu \Psi$, are related to a rather complex dynamics of the system. For stationary solutions only two terms of the Lagrangian contribute

$$\mathcal{L}_{2g} = \frac{-ig^2}{\tilde{m}^2} \bar{\Psi} \gamma_\mu [A^\mu \partial_\nu A^\nu] \Psi \quad (2)$$

and

$$\mathcal{L}_{3g} = \frac{-g^3}{\tilde{m}^2} \bar{\Psi} \gamma_\mu [A^\mu A_\nu A^\nu] \Psi . \quad (3)$$

The gauge condition $\partial_\mu A^\mu = 0$ used for first order Lagrangians is replaced in the present case by $\partial(\partial_\nu A^\nu) = 0$.

From the Lagrangians (2) and (3) fermion matrix elements have been evaluated. This has been found to be a reliable method to generate bound state potentials, see refs. [3, 4]. Using $\alpha = g^2/4\pi$ and fermion wave functions $\psi(p) = \frac{1}{\tilde{m}^{3/2}} \Psi(p_i) \Psi(k)$, matrix elements are obtained of the form

$$\mathcal{M}_{2g} = \frac{\alpha^2}{\tilde{m}^8} \bar{\psi}(p') \gamma_\mu A^\mu(q) (\partial_\nu A^\nu(q)) (\partial_\sigma A^\sigma(q)) \gamma_\rho A^\rho(q) \psi(p) \quad (4)$$

and

$$\mathcal{M}_{3g} = \frac{-\alpha^3}{\tilde{m}^8} \bar{\psi}(p') \gamma_\mu A^\mu(q) A_\nu(q) A^\nu(q) A_\sigma(q) A^\sigma(q) \gamma_\rho A^\rho(q) \psi(p) , \quad (5)$$

These matrix elements have a structure more complex than obtained in a first order theory [6]. But this is necessary to describe relativistic bound states in a fundamental approach: only by interactions between fermions a stable system cannot be generated; the additional boson fields on the right and left of \mathcal{M}_{2g} and \mathcal{M}_{3g} have to provide the restoring force needed to stabilise the system.

From these matrix elements bound state potentials can be deduced. Following the derivations in refs. [3, 4] \mathcal{M}_{2g} and \mathcal{M}_{3g} can be simplified, using (analogue to the fermion wave

functions) normalised boson (quasi) wave functions $W_\mu^\nu(q) = \frac{1}{\tilde{m}} A_\mu(q) A^\nu(q)$ and a boson-exchange interaction $V_\mu^\nu(q) = W_\mu^\nu(q)$ ($\mu \neq \nu$). Further, by an equal time requirement the fermion and boson vectors can be reduced by one dimension, yielding boson wave functions⁴ of scalar and vector structure $w_s(q)$ and $w_v(q)$ and an interaction potential $v_v(q) = w_v(q)$. Going to r-space the fermion matrix elements (4) and (5) can be written by

$$\mathcal{M}_{2g}^f = \bar{\psi}(r) V_{2g}(r) \psi(r) \quad (6)$$

and

$$\mathcal{M}_{3g}^f = \bar{\psi}(r) V_{3g}^{s,v}(r) \psi(r) , \quad (7)$$

where the bosonic potentials $V_{2g}(r)$ and $V_{3g}^{s,v}(r)$ are given by

$$V_{2g}(r) = \frac{\alpha^2 (\hbar c)^2 F_{2g}}{4\tilde{m}} \left(\frac{d^2 w_s(r)}{dr^2} + \frac{2}{r} \frac{dw_s(r)}{dr} \right) \frac{1}{w_s(r)} + E_o \quad (8)$$

and

$$V_{3g}^{s,v}(r) = -\frac{\alpha^3 \hbar c}{\tilde{m}} \int dr' w_{s,v}(r') v_v(r-r') w_{s,v}(r') . \quad (9)$$

The factor F_{2g} in eq. (8) is due to the Fourier transformation of the boson kinetic energy and E_o the energy of the lowest eigenstate. A connection to the vacuum is made by assuming $E_o = E_{vac} = 0$. $V_{2g}(r)$ shows a quite linear rise towards larger radii, leading to confinement of the system; therefore it can be identified with the confinement potential required in hadron potential models [7]. The potential $V_{3g}^{s,v}(r)$ can be considered as boson matrix element, in which the wave functions $w_{s,v}(r)$ are bound in the potential $v_v(r)$.

From the structure of the fermion matrix element in eqs. (7) and (9) one can see that there are two $q\bar{q}$ states (with quantum numbers $J^\pi = 1^-$) with scalar and vector boson wave functions $w_{s,v}(r)$ and corresponding fermion wave functions $\psi_{s,v}(r) \sim w_{s,v}(r)$. Further, there are two $q\bar{q}$ p-states (with quantum numbers $J^\pi = 0^+$) with similar wave functions, see ref. [4, 8].

The fermion wave functions have to be orthogonal, leading to the constraint

$$\int r^2 dr \psi_s(r) \psi_v(r) = \int r^2 dr w_s(r) w_v(r) = \langle r_{w_s, w_v} \rangle = 0 . \quad (10)$$

To satisfy this condition, $w_v(r)$ can be written in the form of a p-wave function

$$w_v(r) = w_{v,o} \left[w_s(r) + \beta R \frac{dw_s(r)}{dr} \right] , \quad (11)$$

⁴with dimension [GeV].

where $w_{v,o}$ is obtained from the normalisation $2\pi \int r dr w_v^2(r) = 1$ and βR determined by $\beta R = - \int r^2 dr w_s(r) / \int r^2 dr [dw_s(r)/dr]$. Interestingly, orthogonality gives rise to another quite natural condition for the deepest bound state, requiring that the interaction takes place inside the bound state volume of $w_s^2(r)$. This leads to

$$|V_{3g}^v(r)| \sim c w_s^2(r) . \quad (12)$$

The conditions (10)-(12) lead to a boson wave function $w_s(r)$ of the form

$$w_s(r) = w_{s,o} \exp\{-(r/b)^\kappa\} , \quad (13)$$

where $w_{s,o}$ is fixed by the normalisation $2\pi \int r dr w_s^2(r) = 1$. The parameters b and κ have to be determined by boundary conditions as discussed below. Different flavour states are obtained by solutions with different slope parameter b , which are constrained by a vacuum potential sum rule, see e.g. ref. [4]. The interaction $v_v(r)$ is given by $v_v(r) = \hbar c w_v(r)$.

Binding energies have been calculated within the Hamiltonian formalism by using the virial theorem in the form $4\pi[\int r^2 dr \psi^2(r)V_{ng}(r) - \frac{1}{2} \int r^3 dr \psi^2(r)\frac{d}{dr}V_{ng}(r)] = E_f^{ng}$, where $\psi(r)$ are fermion wave functions with a form similar to the boson wave functions in eqs. (11) and (13). In addition, $V_{3g}^s(r)$ can be interpreted as bound state of bosons. Its binding energy E_g has been calculated by the corresponding form $2\pi[\int r dr w_s^2(r)v_v(r) - \frac{1}{2} \int r^2 dr w_s^2(r)\frac{d}{dr}v_v(r)] = E_g$. For massless fermions the mass of the system $M^{s,v}$ is given by the absolute binding energies in $V_{2g}(r)$ and $V_{3g}^{s,v}(r)$, yielding $M^{s,v} = -E_{fs,v}^{3g} + E_f^{2g}$, while the reduced mass is given by $\tilde{m} = M^s/2$.

In order to make a detailed evaluation of the potentials other constraints are needed, which connect the coupling constant α to the shape parameters κ and b and the mass of the system. If the mass is adjusted to that known experimentally, only two conditions are needed for a complete determination of all parameters. This has been done in most cases. However, by applying in addition a vacuum potential sum rule, also the masses of different solutions are constrained, thus leading to a fully complete description.

The first condition is energy-momentum conservation, important for relativistic bound states. For $(q\bar{q})$ as well as $(p e^-)$ and (e^+e^-) systems bound by electric interactions, the negative fermion and boson binding energies E_f^s and E_g have to be compensated by their root mean square momenta $\langle q_v^2 \rangle = \int dq q^3 v(q) / \int dq q v(q)$ in the potentials $V_{3g}(r)$ and

$v_v(r)$, respectively

$$\langle q_{V_{3g}}^2 \rangle + \langle q_{v_v}^2 \rangle = (E_f^s + E_g)^2 . \quad (14)$$

For more complex $(q\bar{q})^n$ and $(q\bar{q})^n q$ systems somewhat different energy-momentum relations are needed. Of particular interest are $(q\bar{q})^n q$ states with opposite fermion momenta $\sum_i \langle q_{\psi_i}^2 \rangle = 0$. In this case the motion of fermions give rise to magnetic interactions and thus to magnetic binding. This leads to leptonic bound states [9] with an energy-momentum relation of the form

$$\langle q_{v_v}^2 \rangle (v/c)^2 = (E_f^s + E_g)^2 , \quad (15)$$

where $(v/c)^2$ is the relative motion of fermions, $(v/c)^2 < 6 \cdot 10^{-19}$ for leptons, see ref. [9].

A mass-radius condition can be derived from the confinement potential (8), see ref. [3], which leads to

$$Rat_{conf} = \frac{(\hbar c)^2}{\tilde{m}^2 \langle r_{ws}^2 \rangle} = 1 . \quad (16)$$

For magnetic binding eq. (16) is multiplied by a factor of $(v/c)^2$. In this case p-wave states are not stable, leading only to one lepton bound state (for each flavour system).

The last ambiguity between α , κ , b and the mass of the bound state can be removed by a vacuum potential sum rule. Here a vacuum potential $v_{vac}(q)$ is assumed, which should be equal to the sum of all individual interactions $v_v^i(q)$. This is expected to have a simple form $\sim 1/q^n$ with a cut-off function $f_{cut}(q)$ to make $v_{vac}(q)$ finite. This leads to

$$v_{vac}(q) = f_{cut}(q) 1/q^n = \sum_i v_v^i(q) \quad (17)$$

and allows to determine $v_{vac}(q)$ by the sum of all individual bound state solutions. In turn, $v_{vac}(q)$ can be used to check the mass assumptions of the individual bound states. Further, hypothetical ground state wave functions can be defined, which allows to check the consistency of this approach. For atomic states this procedure yields very consistent results, see ref. [4].

The general structure of the bound state solutions is shown in fig. 1. In the upper part the radial dependence of the interaction $v_v(r)$ is compared to the $1/r$ dependence of the Coulomb potential. In the middle part the radial dependence of boson density $w_s^2(r)$ and potentials $V_{3g}^{s,v}(r)$ is shown, which indicates that relation (12) is well fulfilled. In the lower part the confinement potential $V_{2g}(r)$ is given, which has the typical linearly

increasing behaviour for larger radii, as needed in empirical hadron structure models [7]. The same behaviour of boson densities and potentials has been found for all systems studied, therefore the horizontal and vertical scales are given only in relative units.

Of large importance, by using boundary conditions for electric and magnetic binding, systems of very different radii are obtained. Whereas for hadrons (bound electrically) root mean square radii between $4 \cdot 10^{-3}$ and 0.5 fm have been extracted [3, 8], for leptons (bound magnetically) root mean square radii smaller than 10^{-8} fm have been found [9]. This indicates that the decay of hadrons to leptons is very weak, as found experimentally. The decay width of the systems in question is related to the dynamical structure of the potentials, given by kinetic energy distributions $T_{2g}(q)$ and $T_{3g}(q)$. $T_{2g}(q)$ is given by the Fourier transformed potential $V_{2g}(r)$, whereas $T_{3g}(q)$ is given by the Fourier transformed kinetic energy $T_{3g}(r)$ given by

$$T_{3g}(r) = \frac{1}{2} \langle r^2 \rangle (d^2 V_{3g}(r)/dr^2 + \frac{2}{r} dV_{3g}(r)/dr) . \quad (18)$$

Results on $T_{2g}(q)$ and $T_{3g}(q)$ are given in fig. 2. Although in many cases the confinement potential $V_{2g}(r)$ is much weaker than $V_{3g}^{s,v}(r)$, this potential gives rise to a very pronounced energy distribution. From these distributions all decay properties and branching ratios can be deduced by calculating the overlap of $T_{2g}(r)$ and $T_{3g}(r)$ for all systems involved. Interestingly, for mesonic $q\bar{q}$ systems the confinement potential gives rise to a sharp peak consistent with the experimental width, see the discussion in ref. [8], whereas the boson-exchange potentials $V_{3g}(r)$ yield very wide structures, in the case of $V_{3g}^s(r)$ a peak of ~ 2 GeV width for J/ψ and ~ 70 GeV width for the top system.

Applications of the present formalism have shown that hadrons are well described as states bound by electric forces, see ref. [3, 8], whereas leptons represent systems bound by magnetic forces, see ref. [9]. This underlines the inherent symmetry of electric and magnetic forces in Maxwell's theory. These results confirm the general conclusion drawn above that in the SM both the strong and weak forces are incorrectly assumed with too many parameters. Concerning the strong interaction, confinement is a basic property of all relativistic bound states, see fig. 1; therefore 'colour confinement' in a theory of the strong interaction is redundant, making a non-Abelian colour theory as quantum chromodynamics (QCD) dispensable. With respect to the weak interaction, magnetic

forces have been misinterpreted in the SM as heavy-boson exchange. Both, magnetic and heavy-boson exchange forces give rise to similar properties, extremely small radii and strength, but the latter requires further Higgs fields to make the theory gauge invariant. Finally, there is no basis for supersymmetry: the flavour degree of freedom is characterised in the present formalism by systems of different slope parameter b .

The implications are clear: apart from elementary fermion and boson fields all other fields assumed in the SM are unrealistic. This excludes colour, heavy boson, Higgs, supersymmetric and all other exotic fields. Consequently, the experimental states W^\pm (80.4 GeV) and Z (91.2 GeV) have to be reinterpreted as $(q\bar{q})^n q$ and $(q\bar{q})^n$ states, respectively. Actually, Z (91.2 GeV) has to be regarded [8] as low mass top $q\bar{q}$ -state, with a width in agreement with experiment. Further, the new state⁵ with a mass of 126 GeV, discovered [10] recently at LHC, has to be identified with a scalar $q\bar{q}$ -state (of p-wave structure), found with exactly this mass, see ref. [8]. A second scalar $q\bar{q}$ -state is predicted with a mass of 41 GeV, which should be found in high energy experiments.

A further strict consequence of the present formalism is that gravitation has to be understood also by electromagnetic forces. Strong support for this conclusion comes from the analysis of neutrinos, see ref. [9], indicating that a hierarchy problem between 'weak' and gravitational forces does not exist, if we assume that gravitation is due to magnetic interactions of particles in matter. The present formalism excludes definitely dark matter in form of exotic particles (the rotation of galactic systems, considered as the best evidence for dark matter, has been well described in the present approach, see ref. [2]). Finally, the present formalism has to be considered as the proper theory of quantum gravity. For more detailed studies other solutions of the Lagrangian (1) may be needed, which go beyond the evaluation of matrix elements.

In conclusion, by using general arguments the SM with its complex structure cannot be considered as a valid theory of fundamental forces. The only method to find a correct solution is to develop a complete and fundamental theory, which has no external parameters. Such a theory has been constructed, based on a second order extension of QED, with the result that **all elementary forces** have to be described by electromagnetic interactions only. Apart from basic fermion and boson fields, the existence of all other SM

⁵misinterpreted as SM Higgs-boson.

fields can be ruled out. In particular, supersymmetric particles and dark matter in form of exotic particles do not exist.

For cosmology a new scenario arises: we have to conclude that the universe emerged out of the absolute vacuum of fluctuating boson fields with average energy $E_{vac} = 0$. During overlap of two boson fields $q\bar{q}$ -pairs could be created, which have been confined to form hadrons and leptons (for this process $V_{2g}(r)$ is of prime importance). By electric forces neutral matter was created, which bound and compressed by gravitation to a dense cosmic bound state. Above a critical density the largest part of matter annihilated (the detailed mechanism have still to be worked out) and gave a big radial impact (Big Bang) to the remaining matter, mainly in form of bound $(p\ e^-)$ -pairs, resulting in an expansion of the system. By perpetual decrease of the binding energy of the cosmic bound state the expanding matter gained and still gains extra momentum, leading in this way to an (ever) increasing expansion of the universe, without dark matter in form of exotic particles.

After one century of modern physics with the development of very complex theories for fundamental forces, it may be surprising that all basic bound states of nature with a strikingly different phenomenology can be understood as stationary states bound by electric or magnetic forces. However, this is in perfect agreement with the general rules of nature, effectiveness and formal logic, by which complex systems should evolve from the simplest structure (the vacuum) by the simplest possible mechanisms.

For fruitful discussions, direct help in the derivation of the formalism and general support the author is indepted to many colleagues, with a list of their names given in a final publication.

References

- [1] Review of particle properties, K. Nakamura et al., J. Phys. G 37, 075021 (2010); <http://pdg.lbl.gov/> and refs. therein
- [2] H.P. Morsch, EPJ Web of Conferences 28, 12068 (2012), Hadron Collider Physics Symposium, Paris 2011 (open access)
- [3] H.P. Morsch, Universal Journal of Physics and App. 1, 252 (2013), arXiv:1112.6345 [gen-ph]
- [4] H.P. Morsch, arXiv:1104.2574 [hep-ph]
- [5] J.Z. Simon, Phys. Rev. D 41, 3720 (1990); A. Foussats, E. Manavella, C. Repetto, O.P. Zandron, and O.S. Zandron, Int. J. theor. Phys. 34, 1 (1995); V.V. Nesterenko, J. Phys. A: Math. Gen. 22, 1673 (1989); and refs. therein
- [6] see e.g. I.J.R. Aitchison and A.J.G. Hey, “Gauge theories in particle physics”, Adam Hilger Ltd, Bristol, 1982; or M.E. Peskin and D.V. Schroeder, “An introduction to quantum field theory”, Addison-Wesley Publ. 1995
- [7] R. Barbieri, R. Kögerler, Z. Kunszt, and R. Gatto, Nucl. Phys. B 105, 125 (1976); E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane, and T.M. Yan, Phys. Rev. D 17, 3090 (1978); S. Godfrey and N. Isgur, Phys. Rev. D 32, 189 (1985); D. Ebert, R.N. Faustov, and V.O. Galkin, Phys. Rev. D 67, 014027 (2003); and refs. therein
- [8] H.P. Morsch, arXiv:1210.0244 [gen-ph]
- [9] H.P. Morsch, arXiv:1104.2576 [hep-ph]
- [10] The Atlas collaboration, arXiv:1207.7214; CMS collaboration, arXiv:1207.7235




Figure 1: General structure of boson density and potentials. Upper part: Relative interaction $v_v(r)$ in comparison with the Coulomb potential ($\sim 1/r$) given by dot-dashed line. Middle part: Boson density $w_s^2(r)$ (dot-dashed line) and boson-exchange potentials $|V_{3g}^{s,v}(r)|$ given by dashed and solid lines, respectively. Lower part: Deduced confinement potential $V_{2g}(r)$.




Figure 2: Relative distributions of widths deduced for the confinement potential (upper part) and the boson-exchange potential $V_{3g}^s(r)$ (lower part).