

A Gaussian Process Approximation for a two-color Randomly Reinforced Urns

Li-Xin Zhang*
Zhejiang University

Abstract

We prove a Gaussian process approximation for the sequence of random compositions of a two-color randomly reinforced urn for both the cases with the equal and unequal reinforcement means. By using the Gaussian approximation, the law of the iterated logarithm and the functional limit central limit theorem in both the stable convergence sense and the almost-sure conditional convergence sense are established. Also as a consequence, we are able to prove that the distribution of the urn composition has no points masses both when the reinforcement means are equal and unequal under the assumption of only finite $(2+\epsilon)$ -th moments.

Keywords: Reinforced urn model; Gaussian process; strong approximation; functional central limit theorem; Pólya urn; law of the iterated logarithm

AMS 2000 subject classifications: Primary 60F15; secondary 62G10; 60F05; 60F10

1 Introduction

Asymptotic properties, including the strong consistency and asymptotic normality, of urn models and their applications are widely studied in recent years under various assumptions concerning the updating rules, for example, one may refer to Chauvin, Ponnyanne and Sahnoun (2009), Bai, Hu and Rosenberger (2002), Bai and Hu (2005), Janson (2004,2006), Zhang, Chueng and Hu (2006) etc. In this paper, we consider a kind of two-color urn model, called the randomly reinforced urn (RRU) model, which is a generalization of the original Pólya urn (c.f., Eggenberger and Polya (1923), Pólya (1931)). The main issue of this model different from most urn models in literature is that, as shown, the proportions of balls in the urn will not converge to a non-extreme constant and the numbers of different type balls may increase in different speeds. This issue makes its asymptotic properties quite different from the those of other urn models and difficult to study.

*Department of Mathematics, Zhejiang University, 310027, P.R. China

The RRU model is described as follows. Consider a two-color urn with the initial urn components $\mathbf{Y}_0 = (Y_{0,1}, Y_{0,2})$, where $Y_{0,k} > 0$ is the number of type k balls. The urn is sampled sequentially. Suppose the urn components are $\mathbf{Y}_m = (Y_{m,1}, Y_{m,2})$ after m samplings. At the $(m + 1)$ -th sampling, a ball of type k is drawn with a probability

$$p_{m+1,k} = \frac{Y_{m,k}}{|\mathbf{Y}_m|}, \quad \text{where } |\mathbf{Y}_m| = Y_{m,1} + Y_{m,2}.$$

And the sampled ball is replaced in the urn together with a nonnegative random number $U_{m+1,k}$ of balls of the same type k , generated from a distribution μ_k with mean $m_k > 0$. This is the model introduced and formally named the randomly reinforced urn in Mulier, Paganoni and Secchi (2006a). But it would appear in earlier literatures in different versions. For example, Durham and Yu (1990) proposed a similar model for sequential sampling in clinical trails. In our RRU setting, the numbers of balls take positive real values, not necessary integers. When $U_{m+1,1} = U_{m+1,2} = \alpha$ is a constant and a positive integer, a RRU is the original Pólya urn (c.f., Eggenberger and Polya (1923), Pólya (1931)) which is very popular in literatures. The RRU model is of fundamental importance in many areas of applications, for instance in economics (c.f., Erev and Roth (1998), Beggs (2005), Hopkins and Posch (2005)), in information science (c.f., Martin and Ho (2002)), in resampling theory etc. In clinical trial studies, the RRU model is utilized to define a response-adaptive design focusing to reduce the expected number of patients receiving inferior treatments (c.f., Durham, Flournory, Li (1998), Li, Durham and Flournory (1996), Melifer, Paganoni and Secchi (2006a,b), Paganoni and Secchi (2007), May and Flournory (2009) etc).

Suppose the reinforcement distributions μ_1 and μ_2 have bounded supports. In Melifer, Paganoni and Secchi (2006a), it is showed that the sequence $\{Z_n = Y_{n,1}/|\mathbf{Y}_n|\}$ of the random sample proportions in the urn converges to almost surely to a random limit $Z_\infty \in [0, 1]$. When $\mu_1 = \mu_2$, Crimaldi (2009) proved a central limit theorem by showing almost-sure conditional convergence to a Gaussian kernel of the sequence $\{\sqrt{n}(Z_n - Z_\infty)\}$. Aletti, May and Secchi (2009) extended Crimaldi's result to a general case that reinforcement means m_1 and m_2 are equal and proved that Z_∞ has no point masses in $[0, 1]$ by using this kind of conditional central limit theorem. When the means m_1 and m_2 are different, the limit proportion Z_∞ of a RRU is showed to be a point mass either 1 and 0 by Beggs (2005), Hopkins and Posch (2005) and Melifer, Paganoni and Secchi (2006a) under the assumption that the supports of μ_1 and μ_2 are bounded from 0, and by Aletti, May and Secchi (2009) only under the assumption that μ_1 and μ_2 have bounded supports. May and Flournory (2009) proved that the sequence $\{Y_{n,1}/Y_{n,2}^{m_1/m_2}\}$ converges to almost surely to a random limit $\psi_\infty \in (0, \infty)$ both when $m_1 = m_2$ and $m_1 \neq m_2$.

The purpose of this paper is to establish the Gaussian process approximation of the sequence $\{Z_n\}$ when $m_1 = m_2$ as well as the sequence $\{Y_{n,1}/Y_{n,2}^{m_1/m_2}\}$ when $m_1 \neq m_2$, under the assumption that μ_1 and μ_2 have only finite $(2 + \epsilon)$ -th

moments. This assumption is much weaker than that μ_1 and μ_2 have bounded supports. We will show that both these sequences can be approximated by a tail stochastic integral with respect to a Brownian motion mixed with a random variable. It is interesting that, as we will find, the mixed Gaussian process for approximating is nearly independent of the urn composition to be approximated. Our Gaussian process approximation enables us (i) to establish the law of the iterated logarithm; (ii) to establish the functional limit central limit theorem in both the stable convergence sense and the almost-sure conditional convergence sense; (iii) to prove that the limit ψ_∞ (resp. Z_∞) has no point masses in $[0, \infty]$ (resp. in $[0, 1]$) when $m_1 \neq m_2$ (resp. when $m_1 = m_2$) under the assumption that μ_1 and μ_2 have only finite $(2 + \epsilon)$ -th moments. Another implication of our Gaussian approximation is that we are able to establish the central limit theorem in a simple way for the random number $N_{n,k}$ of draws, where $N_{n,k}$ is the number of type k balls being drawn in the first n samplings. In a response-adaptive design in clinical trials driven by a RRU model, $N_{n,k}$ is the number of patients allocated to treatment k , and its asymptotic behaviors are of particular interest.

For the generalized Friedman urn models, Bai, Hu and Zhang (2002) and Zhang and Hu (2009) established the Gaussian approximation for both the urn proportions $Y_{n,k}/n$ and the sampling proportion $N_{n,k}/n$. But the RRU which we consider here is not covered by their assumptions. The main reason is that the mean replacement matrix $\text{diag}(m_1, m_2)$ of a RRU is not irreducible and hence the limit of $Y_{n,k}/n$ and $N_{n,k}/n$ is not a constant in $(0, 1)$.

The paper is organized as follows. The main approximation theorems with applications for equal and unequal reinforcement mean case are stated in Section 2 and Section 3, respectively, and the proofs of the approximations appear in the last section. Some remarks on unsolved problems are discussed in Section 4.

In the sequel of this paper if having not been specially mentioned, $(U_{l,1}, U_{l,2})$, $l = 1, 2, \dots$ are assumed to be independent identically distributed random vectors with finite second moments. Let $X_{m,k}$ be the result of the m -th drawing, i.e., $X_{m,k} = 1$ if the m -th drawn ball is of type k , and 0 otherwise. It is obvious that $N_{m,k} = \sum_{j=1}^m X_{j,k}$ and $X_{m,1} + X_{m,2} = 1$. Denote $\mathcal{F}_n = \sigma(U_{l,k}, X_{l,k}, Y_{l,k} : k = 1, 2; l = 1, \dots, n)$ be the history σ -field generated by all the observations up to stage n , and $\mathcal{F}_\infty = \bigvee_n \mathcal{F}_n$. Further, for two positive sequences $\{a_n\}$ and $\{b_n\}$, we write $a_n = O(b_n)$ if there is a constant C such that $a_n \leq Cb_n$, $a_n \sim b_n$ if $a_n/b_n \rightarrow 1$, and $a_n \approx b_n$ if $a_n = O(b_n)$ and $b_n = O(a_n)$.

2 Equal reinforcement mean case

In this section, we consider the case of $m_1 = m_2 > 0$. Let $\sigma_k^2 = \mathbf{E}[(U_{1,k}/m_k)^2]$, $k = 1, 2$,

$$Z_n = \frac{Y_{n,1}}{Y_{n,1} + Y_{n,2}}, \quad Z_\infty(\omega) = \lim Z_n(\omega), \quad H(\omega) = \frac{\sigma_1^2}{Z_\infty} + \frac{\sigma_2^2}{1 - Z_\infty}.$$

To start, we shall assume $P(Z_\infty = 0) = P(Z_\infty = 1) = 0$, for otherwise H may have no definition. This result is proved by May and Flournoy (2009) under the condition that the reinforcement distributions μ_1 and μ_2 have bounded supports. The next theorem tells us that May and Flournoy's condition can be relaxed at least to the assumption of finite second moments.

Theorem 2.1 *Suppose $m_1 = m_2 > 0$, $EU_{1,k}^q < \infty$ for some $q > 1$, $k = 1, 2$. Then the limit Z_∞ exists almost surely and $P(0 < Z_\infty < 1) = 1$.*

The following theorem is the main result on the Gaussian approximation.

Theorem 2.2 *Suppose $m_1 = m_2 > 0$, $EU_{1,k}^p < \infty$, $k = 1, 2$, where $2 \leq p < 4$. Then (possibly in an enlarged probability space) there is standard Brownian motion $B(y)$ such that*

$$Z_\infty - Z_n = Z_\infty(1 - Z_\infty)H \int_{nH}^{\infty} \frac{dB(y)}{y} + o(\lambda_n) \text{ a.s.} \quad (2.1)$$

where

$$\lambda_n = \begin{cases} n^{-1/2}(\log \log n)^{1/2}, & \text{if } p = 2 \\ n^{1/p-1}(\log n)^{1/2}, & \text{if } 2 < p < 4. \end{cases}$$

Furthermore, the Brownian motion $B(y)$ can be constructed with a filtration of σ -fields $\{\mathcal{G}_n\}$ and a non-decreasing sequence of stopping times $\{T_n\}$ satisfying the following properties:

Property (a) $\mathcal{F}_n \subset \mathcal{G}_n$, T_n is \mathcal{G}_n measurable;

Property (b) $T_n = nH + o(n^{2/p})$ a.s.;

Property (c) Conditional on \mathcal{F}_n , $B(T_n + y) - B(T_n)$, $y \geq 0$, is also a standard Brownian motion.

REMARK 2.1 Denote $W(t) = -t \int_t^\infty y^{-2} dB(y)$. By checking the covariance function, it is easily seen that $W(t)$, $t > 0$ is also a standard Brownian motion.

REMARK 2.2 The process in (2.1) for approximating is a tail stochastic integral respective to the Brownian motion. It looks like to be independent of Z_n . Actually, according to Property (b) nH can be replaced by T_n , and $\sqrt{T_n} \int_{T_n}^\infty y^{-1} dB(y)$ is indeed a normal random variable which is independent of Z_n . We will illustrate this interesting property in Corollary 2.3 in more details.

We will prove Theorem 2.2 by first approximating $Z_n - Z_\infty$ to an infinite summation of a weighted martingale sequence and then approximating the martingale to a Brownian motion by applying the Skorokhod embedding method. The detail proofs of Theorems 2.1 and 2.2 will be stated in Section A. In the sequel of this section, we give several corollaries as applications. Define

$$\tilde{\sigma}(\omega) = \sqrt{Z_\infty(1 - Z_\infty)} \sqrt{(1 - Z_\infty)\sigma_1^2 + Z_\infty\sigma_2^2},$$

$$\tilde{\sigma}_n(\omega) = \sqrt{Z_n(1-Z_n)}\sqrt{(1-Z_n)\sigma_1^2 + Z_n\sigma_2^2}.$$

The first corollary is the following law of the iterated logarithm.

Corollary 2.1 *Suppose $m_1 = m_2 > 0$, $EU_{1,k}^2 < \infty$, $k = 1, 2$. Then*

$$\limsup_{n \rightarrow \infty} \frac{\sqrt{n}(Z_n - Z_\infty)}{\sqrt{2 \log \log n}} = \tilde{\sigma} \text{ a.s..}$$

Proof. Write $\gamma(x) = \sqrt{x/(2 \log \log x)}$, $G(x) = -\int_x^\infty y^{-1} dB(y)$. By (2.1), we need to show that

$$\limsup_{n \rightarrow \infty} \gamma(nH)G(nH) = \limsup_{T \rightarrow \infty} \gamma(T)|G(T)| = 1 \text{ a.s.} \quad (2.2)$$

Note that $\gamma(x)G(x) = xG(x)/\sqrt{2x \log \log x}$, and that $xG(x)$ is also a standard Brownian motion. (2.2) follows from the law of the iterated logarithm of the Brownian motion. \square .

The next corollary is on the functional central limit theorem.

Corollary 2.2 *Suppose $m_1 = m_2 > 0$, $EU_{1,k}^p < \infty$ for some $p > 2$, $k = 1, 2$. Define*

$$W_n(t) = t\sqrt{n}(Z_{[nt]} - Z_\infty), t > 0.$$

Then

$$W_n(\cdot) \xrightarrow{d} \tilde{\sigma}B'(\cdot) \text{ stably,} \quad (2.3)$$

in the Skorokhod Topological space $D(0, \infty)$, where $B'(t)$ is a standard Brownian motion which is independent of \mathcal{F}_∞ . In particular,

$$\lim_{n \rightarrow \infty} P\left(\frac{\max_{0 \leq l \leq n} l(Z_l - Z_n)}{\tilde{\sigma}_n \sqrt{n}} \geq x\right) = e^{-2x^2}, \quad x > 0. \quad (2.4)$$

Here the stable convergence in (2.3) means that for any bounded and (uniformly) continuous function $f : D(0, \infty) \rightarrow (-\infty, \infty)$,

$$E[f(W_n(\cdot))I_E] \rightarrow E[f(\tilde{\sigma}B'(\cdot))I_E] \text{ for any event } E.$$

REMARK 2.3 *The convergence (2.4) does not depend on the unknown value of Z_∞ .*

Proof. (2.4) is due to fact that

$$\max_{0 \leq l \leq n} \frac{l(Z_l - Z_n)}{\tilde{\sigma}_n \sqrt{n}} = \sup_{0 < t \leq 1} \frac{W_n(t) - tW_n(1)}{\tilde{\sigma}} \xrightarrow{d} \sup_{0 < t \leq 1} (B'(t) - tB'(1))$$

and that $B'(t) - tB'(1)$ is a Brownian bridge. For (2.3), note that $W(x) = -t \int_t^\infty y^{-1} dB(y)$ is also a standard Brownian motion. By (2.1),

$$n(Z_n - Z_\infty) - \frac{\tilde{\sigma}}{\sqrt{H}}W(nH) = o(n^{1/2}) \text{ a.s.,}$$

which implies that for any $T > 0$,

$$\sup_{0 < t \leq T} \left| W_n(t) - \tilde{\sigma} \frac{W(nHt)}{\sqrt{nH}} \right| \rightarrow o(1) \text{ a.s.}$$

For the Brownian motion $W(\cdot)$, we have

$$W(n\cdot)/\sqrt{n} \xrightarrow{d} B'(\cdot) \text{ mixing,}$$

i.e., for any given event E with $\mathbb{P}(E) > 0$, the conditional distribution of $W(n\cdot)/\sqrt{n}$ converges to a Brownian motion. It follows that

$$(\tilde{\sigma}/\sqrt{H}, H, W(n\cdot)/\sqrt{n}) \xrightarrow{d} (\tilde{\sigma}/\sqrt{H}, H, B'(\cdot)) \text{ stably.}$$

Note that $\tilde{\sigma} \frac{W(nH\cdot)}{\sqrt{nH}}$ is a continuous function of $(\tilde{\sigma}/\sqrt{H}, H, W(n\cdot)/\sqrt{n})$ of the form $f(r, h, x(\cdot)) = rx(\cdot/h)$. It follows that

$$\tilde{\sigma} \frac{W(nH\cdot)}{\sqrt{nH}} \xrightarrow{d} \tilde{\sigma} \frac{B'(\cdot/H)}{\sqrt{H}} \stackrel{d}{=} \tilde{\sigma} B'(\cdot) \text{ stably.}$$

The proof is now completed. \square

Corollary 2.2 implies the central limit theorem for $\sqrt{n}(Z_n - Z_\infty)$. Aletti, May and Secchi (2009) proved a strong version of the central limit theorem. For every Borel set B , every ω , and $n = 1, 2, \dots$, define

$$K_n(\omega, B) = \mathbb{P}(\sqrt{n}(Z_n - Z_\infty) \in B | \mathcal{F}_n)(\omega),$$

i.e., K_n is a version of the condition distribution of $\sqrt{n}(Z_n - Z_\infty)$ given \mathcal{F}_n . Aletti, May and Secchi (2009) showed that, if $m_1 = m_2$ and the distributions of μ_1 and μ_2 have bounded supports, then for almost every ω , the sequence of probability distributions $K_n(\omega, \cdot)$ converges weakly to the normal distribution

$$N(0, \tilde{\sigma}^2(\omega)).$$

We denote this kind of convergence by

$$\sqrt{n}(Z_n - Z_\infty) \Big|_{\mathcal{F}_n} \xrightarrow{d} N(0, \tilde{\sigma}^2(\omega)) \text{ a.s.} \quad (2.5)$$

This kind of conditional central limit theorem was first established by Grimadi (2008) for a special case that $\mu_1 = \mu_2$. Aletti, May and Secchi (2009) also showed that (2.5) implies that Z_∞ has no point masses in $(0, 1)$. Our next corollary tells that (2.5) and a type of conditional functional central limit theorem are conclusions of the Gaussian approximation.

Corollary 2.3 *Suppose $m_1 = m_2 > 0$, $EU_{1,k}^p < \infty$ for some $p > 2$, $k = 1, 2$. Then*

$$\sup_{t \geq 1} \left| \sqrt{n}(Z_\infty - Z_{[nt]}) - \tilde{\sigma}_n \sqrt{T_n} \int_{T_n t}^{\infty} \frac{dB(y)}{y} \right| = o(n^{-\epsilon}) \text{ a.s.} \quad (2.6)$$

for some $\epsilon > 0$, and further, $\bar{B}_n(t) = -t\sqrt{T_n} \int_{T_n t}^{\infty} y^{-1} dB(y)$, $t \geq 1$, is also a standard Brownian on $[1, \infty)$ which is independent of \mathcal{F}_n .

As a consequence,

$$W_n(\cdot)|_{\mathcal{F}_n} \xrightarrow{d} \tilde{\sigma}(\omega)B'(\cdot) \text{ a.s.}, \quad (2.7)$$

in the Skorokhod Topological space $D[1, \infty)$, where $W_n(t)$ is defined as in Corollary 2.2, $B'(t)$ is a standard Brownian motion which is independent of \mathcal{F}_∞ .

In particular, (2.5) holds, Z_∞ has no point masses in $(0, 1)$, and there exists a sequence of standard normal random variables for which ζ_n is independent of \mathcal{F}_n and

$$\sqrt{n}(Z_n - Z_\infty) = \tilde{\sigma}_n \zeta_n + o(n^{-\epsilon}) \text{ a.s.}, \quad (2.8)$$

Proof. We first prove (2.6). By (2.1),

$$\sup_{t \geq 1} \left| \sqrt{n}(Z_\infty - Z_{[nt]}) - Z_\infty(1 - Z_\infty)H\sqrt{n} \int_{ntH}^{\infty} \frac{dB(y)}{y} \right| = o(n^{-\epsilon}) \text{ a.s.}$$

Let $H_n = \sigma_1^2/Z_n + \sigma_2^2/(1 - Z_n)$. Note that $Z_n - Z_\infty = O(\sqrt{n^{-1} \log \log n})$ a.s. by Corollary 2.1, and $T_n/n = H + o(n^{2/p-1})$ a.s.. It follows that $\sqrt{n}Z_\infty(1 - Z_\infty)H - \tilde{\sigma}_n\sqrt{T_n} = o(n^{2/p-1/2})$ a.s.. Further,

$$n^{2/p-1/2} \sup_{t \geq 1} \left| \int_{ntH}^{\infty} \frac{dB(y)}{y} \right| = n^{2/p-1/2} O(\sqrt{n^{-1} \log \log n}) = o(n^{-\epsilon}) \text{ a.s.}$$

It remains to show that

$$\sup_{t \geq 1} \left| \int_{ntH}^{T_n t} \frac{dB(y)}{y} \right| = o(n^{-1/2-\epsilon}) \text{ a.s.}$$

Write $a_n = T_n - nH$. Note that $W(x) = -x \int_x^{\infty} y^{-1} dB(y)$ is a standard Brownian motion and

$$\int_{ntH}^{T_n t} \frac{dB(y)}{y} = \frac{W(nHt + a_n t) - W(nHt)}{T_n t} + \frac{W(ntH) nH - T_n}{ntH T_n}.$$

The second term on the right hand of the above equality does not exceed

$$O(\sqrt{n^{-1} \log \log n})o(n^{2/p-1})$$

uniformly in $t \geq 1$ almost surely, by the law of the iterated logarithm. The first term does not exceed

$$O\left(\frac{\sqrt{a_n t (\log(nH) + \log \log(nHt))}}{T_n t}\right) = o(n^{1/p-1}(\log n)^{1/2})$$

uniformly in $t \geq 1$ almost surely, by the path properties of a Brownian motion (c.f. Hanson and Russo (1983)). The proof of (2.6) is now proved.

Let $B_n(y) = B(T_n + y) - B(T_n)$. Then conditional on \mathcal{G}_n , $B_n(y)$ is a standard Brownian motion. It is obvious that

$$\overline{B}_n(t) = -t\sqrt{T_n} \int_{T_n(t-1)}^{\infty} \frac{dB_n(y)}{T_n + y}.$$

Hence, conditional on \mathcal{G}_n , $\overline{B}_n(t)$, $t \geq 1$, is a mean zero Gaussian process with covariance function

$$T_n t s \int_{T_n(t-1)}^{\infty} \frac{dy}{(T_n + y)^2} = s \quad \text{for } t \geq s \geq 1.$$

It follows that $\overline{B}_n(t)$, $t \geq 1$, is a standard Brownian motion and is independent of \mathcal{G}_n . So, it is independent of \mathcal{F}_n because $\mathcal{F}_n \subset \mathcal{G}_n$. The proof of the main part of the corollary is now completed.

Now for (2.7), from (2.6) it follows that

$$\text{dist}(W_n(\cdot), \tilde{\sigma}_n B_n(\cdot)) \rightarrow 0 \quad \text{a.s. in } D[1, \infty),$$

where $\text{dist}(\cdot, \cdot)$ is a metric in $D[1, \infty)$. Note that $\tilde{\sigma}_n$ is \mathcal{F}_n -measurable and $\tilde{\sigma}_n \rightarrow \tilde{\sigma}$ a.s., and, conditional on \mathcal{F}_n , $\tilde{\sigma}_n B_n(\cdot)$ and $\tilde{\sigma}_n B'(\cdot)$ has the same distribution. Hence, (2.7) follows from (2.8) by noting the following fact that:

$$\begin{aligned} \xi_n(\cdot) \Big|_{\mathcal{F}_n} &\xrightarrow{d} \tilde{\sigma} B'(\cdot) \quad \text{a.s. and } \text{dist}(\xi_n(\cdot), \eta_n(\cdot)) \rightarrow 0 \quad \text{a.s. in } D[1, \infty) \\ \implies \eta_n(\cdot) \Big|_{\mathcal{F}_n} &\xrightarrow{d} \tilde{\sigma} B'(\cdot) \quad \text{a.s. in } D[1, \infty). \end{aligned}$$

This fact follows from that for any bounded and uniformly continuous function $f : D[1, \infty) \rightarrow (-\infty, \infty)$,

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \left| \mathbb{E}[f(\eta_n) - f(\xi_n) | \mathcal{F}_n] \right| \\ &\leq \mathbb{E} \left[\limsup_{n \rightarrow \infty} |f(\eta_n) - f(\xi_n)| \Big| \bigvee_n \mathcal{F}_n \right] = 0 \quad \text{a.s.}, \end{aligned}$$

due to Lemma A.2 of Crimaldi (2009).

Finally, (2.8) follows from (2.6) by letting $\zeta_n = \overline{B}_n(1)$, and (2.5) is a conclusion of (2.8) or (2.7). Aletti, May and Secchi (2009) showed that (2.5) implies Z_∞ having no point masses in $(0, 1)$ by utilizing a metric of the weak convergence of probability measures with the limit distribution being absolutely continuous. Here we give a straightforward proof. Let $f(t) = e^{-t^2/2}$ be the characteristic function of a standard normal distribution. Firstly, note that (2.5) implies that for every $\bigvee_n \mathcal{F}_n$ -measurable event E ,

$$\lim_{n \rightarrow \infty} \mathbb{E}[e^{it\sqrt{n}(Z_n - Z_\infty)} I_E | \mathcal{F}_n] = f(\tilde{\sigma}t) I_E \quad \text{a.s.}$$

In fact, if let $I_n = \mathbb{E}[I_E | \mathcal{F}_n]$, then $I_n \rightarrow I_E$ a.s.. And hence

$$\begin{aligned} &\lim_{n \rightarrow \infty} \mathbb{E}[e^{it\sqrt{n}(Z_n - Z_\infty)} I_E | \mathcal{F}_n] = \lim_{n \rightarrow \infty} \mathbb{E}[e^{it\sqrt{n}(Z_n - Z_\infty)} I_n | \mathcal{F}_n] \\ &= \lim_{n \rightarrow \infty} \mathbb{E}[e^{it\sqrt{n}(Z_n - Z_\infty)} | \mathcal{F}_n] I_n = f(\tilde{\sigma}t) I_E \quad \text{a.s.}, \end{aligned}$$

where in the first equality we use the fact that

$$\eta_n \rightarrow 0 \text{ a.s. and } |\eta_n| \leq M \text{ a.s.} \implies \mathbf{E}[\eta_n | \mathcal{F}_n] \rightarrow 0 \text{ a.s..}$$

This fact is due to Lemma A.2 of Crimaldi (2009). Next, choosing $E = \{Z_\infty = p\}$, $p \in (0, 1)$, yields

$$\begin{aligned} f(\tilde{\sigma}t)I_E &= \lim_{n \rightarrow \infty} \mathbf{E}[e^{it\sqrt{n}(Z_n - p)} I_E | \mathcal{F}_n] \\ &= \lim_{n \rightarrow \infty} e^{it\sqrt{n}(Z_n - p)} \mathbf{E}[I_E | \mathcal{F}_n] = \lim_{n \rightarrow \infty} e^{it\sqrt{n}(Z_n - p)} I_E \text{ a.s..} \end{aligned}$$

Hence, $|f(\tilde{\sigma}t)I_E| = I_E$ a.s.. So $I_E = 0$ a.s. because $|f(\tilde{\sigma}t)| < 1$ on E . The proof is now completed. \square .

From the above proof, we obtain the following corollary.

Corollary 2.4 *Suppose $\{Y_\infty, Y_n, n \geq 1\}$ is a sequence of random variables, $\{a_n\}$ is a sequence of constants with $a_n \rightarrow \infty$, and \mathcal{G}_n is a filtration of σ -fields such that Y_n is \mathcal{G}_n -measurable. If for almost every ω , the distribution of $a_n(Y_n - Y_\infty)$ conditional on \mathcal{G}_n converges to a non-degenerate distribution, then Y_∞ has no point masses.*

REMARK 2.4 *By (2.5), it can also be shown that for every event E ,*

$$\begin{aligned} & \limsup_{|t| \rightarrow \infty} \limsup_{n \rightarrow \infty} \left| \mathbf{E} \left[\exp \{ -itn^{1/2} Z_\infty \} I_E \right] \right| \\ & \leq \limsup_{|t| \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbf{E} \left| \mathbf{E} \left[\exp \{ itn^{1/2} (Z_n - Z_\infty) \} I_E | \mathcal{F}_n \right] \right| \\ & = \limsup_{|t| \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbf{E} \left| \mathbf{E} [f(t\tilde{\sigma}_n) I_E | \mathcal{F}_n] \right| \\ & = \limsup_{|t| \rightarrow \infty} \mathbf{E} [|f(t\tilde{\sigma}_\infty)| I_E] = 0. \end{aligned}$$

If we denote $f_E(t)$ be the characteristic function of the conditional distribution of Z_∞ given E with $P(E) > 0$. Then the above equality means that

$$\lim_{|t| \rightarrow \infty} \limsup_{n \rightarrow \infty} \left| f_E(tn^{1/2}) \right| = 0.$$

Note that for any $t_0 \geq 1$, $n_0 \geq 1$ and $|s| \geq 2t_0n_0$, there exist a real number t with $|t| \geq t_0$ and an integer $n \geq n_0$ such that $s = tn^{1/2}$. We conclude that

$$\lim_{|s| \rightarrow \infty} f_E(s) = 0.$$

This is related to the Cramér condition. Obviously, if $E = \{Z_\infty = p\}$, $p \in (0, 1)$, and $P(E) > 0$, then $|f_E(t)| \equiv 1$ which is a contradiction.

The next corollary is the central limit theorem for the random number of draws.

Corollary 2.5 Suppose $m_1 = m_2 > 0$, $EU_{1,k}^p < \infty$ for some $p > 2$, $k = 1, 2$. Then

$$\sqrt{n} \left(\frac{N_{n,1}}{n} - Z_\infty \right) \xrightarrow{d} h(\omega) \cdot N(0, 1) \text{ stably,} \quad (2.9)$$

where $h(\omega) = \sqrt{Z_\infty(1 - Z_\infty)} \sqrt{(1 - Z_\infty)(2\sigma_1^2 - 1) + Z_\infty(2\sigma_2^2 - 1)}$, and $N(0, 1)$ is a standard normal random variable which is independent of \mathcal{F}_∞ .

Proof. We need to prove

$$\begin{aligned} & \sqrt{n} \left(\frac{N_{n,1}}{n} - Z_\infty \right) \\ & \xrightarrow{d} -\sqrt{Z_\infty(1 - Z_\infty)} N_1(0, \sigma_1^2 - 1) \\ & \quad + \sqrt{1 - Z_\infty} Z_\infty N_2(0, \sigma_2^2 - 1) + \tilde{\sigma} \cdot N_3(0, 1) \text{ stably,} \end{aligned}$$

where $N_k(0, \sigma_k^2 - 1)$, $k = 1, 2$, $N_3(0, 1)$ are three independent normal random variables which are independent of \mathcal{F}_∞ . Write

$$A_{n,k} = \frac{\sum_{l=1}^n X_{l,k}(U_{l,k}/m_k - 1)}{N_{n,k}}, \quad k = 1, 2. \quad (2.10)$$

Then

$$A_{n,k} = O(\sqrt{n^{-1} \log \log n}) \text{ a.s. and } \frac{Y_{n,k}}{m_k N_{n,k}} = 1 + A_{n,k}.$$

By the Taylor expansion and (2.8), we have that

$$\begin{aligned} \frac{N_{n,1}}{n} - Z_\infty &= \frac{N_{n,1}}{N_{n,1} + N_{n,2}} - Z_\infty = \frac{\frac{Z_n}{1+A_{n,1}}}{\frac{Z_n}{1+A_{n,1}} + \frac{1-Z_n}{1+A_{n,2}}} - Z_\infty \\ &= -Z_n(1 - Z_n)(A_{n,1} - A_{n,2}) + (Z_n - Z_\infty) + O\left(\frac{\log \log n}{n}\right) \\ &= -\frac{Z_n(1 - Z_n)}{\sqrt{N_{n,1}}}(\sqrt{N_{n,1}}A_{n,1}) + \frac{Z_n(1 - Z_n)}{\sqrt{N_{n,2}}}(\sqrt{N_{n,2}}A_{n,2}) \\ & \quad + \tilde{\sigma}_n \zeta_n + o(n^{-1/2}) \text{ a.s..} \end{aligned} \quad (2.11)$$

Note that ζ_n is a standard normal random variable which is independent of $\tilde{\sigma}_n$, Z_n , $N_{n,k}$ and $A_{n,k}$, $k = 1, 2$. Also, $\tilde{\sigma}_n \rightarrow \tilde{\sigma}$ a.s., $Z_n(1 - Z_n)\sqrt{n}/\sqrt{N_{n,1}} \rightarrow \sqrt{Z_\infty(1 - Z_\infty)}$ a.s. and $Z_n(1 - Z_n)\sqrt{n}/\sqrt{N_{n,2}} \rightarrow Z_\infty\sqrt{1 - Z_\infty}$. The proof is completed if we have shown that

$$(\sqrt{N_{n,1}}A_{n,1}, \sqrt{N_{n,2}}A_{n,2}) \xrightarrow{d} (N_1(0, \sigma_1^2 - 1), N_2(0, \sigma_2^2 - 1)) \text{ mixing.} \quad (2.12)$$

Note that $\sigma_k^2 - 1 = \text{Var}(U_{1,k}/m_k)$, $k = 1, 2$. The above convergence follows from Theorem 4.1 of May and Fournoy (2009). \square

The next corollary tells us that conditional on \mathcal{F}_n , the conditional distribution of $\sqrt{n}(N_{n,1}/n - Z_\infty)$ does not converge.

Corollary 2.6 *Suppose $m_1 = m_2 > 0$, $EU_{1,k}^p < \infty$ for some $p > 2$, $k = 1, 2$. Let E be an event that for $\omega \in E$ there is a distribution F_ω for which*

$$\sqrt{n}\left(\frac{N_{n,1}}{n} - Z_\infty\right)\Big|_{\mathcal{F}_n} \xrightarrow{d} F_\omega. \quad (2.13)$$

Then $P(E) = 0$.

Proof. Recall (2.11). Let $\eta_n = \sqrt{nZ_n(1-Z_n)}(A_{n,1} - A_{n,2})$. Note that $Z_n \rightarrow Z_\infty$ a.s.,

$$\tilde{\sigma}_n \xi_n \Big|_{\mathcal{F}_n} \xrightarrow{d} \tilde{\sigma} N(0, 1) \text{ a.s.}$$

and that Z_n , $A_{n,1}$ and $A_{n,2}$ are \mathcal{F}_n -measurable. By (2.11) and (2.13), there exists an event Ω_0 with $P(\Omega_0) = 1$ such that $\sqrt{n}Z_n(1-Z_n)(A_{n,1} - A_{n,2})$ converges on $E \cap \Omega_0$. So there exists a random variable η such that

$$\eta_n(\omega) \rightarrow \eta(\omega) \quad \forall \omega \in E \cap \Omega_0.$$

Suppose $P(E) > 0$. Choose x such that $P(\eta > x, E) > 0$. Then it follows that

$$P(\eta_n \leq x, \eta > x, E) \rightarrow P(\eta \leq x, \eta > x, E) = 0.$$

So $P(\xi_n \leq x | \xi > x, E) \rightarrow 0$. On the other hand, according to (2.12) we have

$$\eta_n = \frac{\sqrt{nZ_n(1-Z_n)}}{\sqrt{N_{n,1}}} \sqrt{N_{n,1}} A_{n,1} - \frac{\sqrt{nZ_n(1-Z_n)}}{\sqrt{N_{n,2}}} \sqrt{N_{n,2}} A_{n,2} \xrightarrow{d} N(0, 1) \text{ mixing,}$$

where $N(0, 1)$ is independent of \mathcal{F}_∞ . It follows that

$$\lim_n P(\eta_n \leq x | \eta > x, E) = \Phi(x) > 0.$$

We get a contradiction. The proof is completed. \square

3 Unequal reinforcement mean case

In this section, we consider the case of $m_1 \neq m_2$. Without loss of generality, we assume that $0 < m_1 < m_2$. Denote $\rho = m_1/m_2$, and

$$\psi_n = \frac{Y_{n,1}}{Y_{n,2}^\rho}, \quad \psi_\infty = \lim_{n \rightarrow \infty} \psi_n.$$

May and Flournoy (2009) proved that the limit ψ_∞ exists almost surely with $P(0 < \psi_\infty < \infty) = 1$ when the reinforcement distributions μ_1 and μ_2 have bounded supports. Durham and Yu (1990) proved a similar result as that

$$\frac{N_{n,1}}{N_{n,2}^\rho} \text{ converges almost surely to a finite limit } \eta_\infty.$$

It is easily seen that

$$\eta_\infty = \frac{m_2^\rho}{m_1} \psi_\infty \text{ a.s.}$$

and

$$\lim_{n \rightarrow \infty} \frac{N_{n,1}}{n^\rho} = \frac{m_2^\rho}{m_1} \psi_\infty \text{ a.s.}, \quad \lim_{n \rightarrow \infty} \frac{Y_{n,1}}{n^\rho} = m_2^\rho \psi_\infty \text{ a.s.}$$

In a recent manuscript of Zhang et al (2010), it is proved that the weakest condition for $P(0 < \psi_\infty < \infty) = 1$ is that $E[U_{1,k} \log^+ U_{1,K}] < \infty$, $k = 1, 2$, and a general multi-color RRU is consider. For the completeness of this paper, we will give a simple proof under the assumption of finite $(1 + \epsilon)$ -th moments for the two-color case. The following is the result.

Theorem 3.1 *Suppose $EU_{1,k}^q < \infty$ for some $q > 1$, $m_k > 0$, $k = 1, 2$. Then the limit ψ_∞ exists almost surely and $P(0 < \psi_\infty < \infty) = 1$ both when $m_1 = m_2$ and $m_1 \neq m_2$.*

The following theorem is our main result on the Gaussian process approximation for ψ_n . From the Gaussian approximation we are able to show that ψ_∞ has no point masses in $(0, \infty)$. And accordingly, all the limits of the sequences $\{Y_{n,1}/Y_{n,2}^\rho\}$, $\{Y_{n,1}/n^\rho\}$, $\{Y_{n,1}/|\mathbf{Y}_n|^\rho\}$, $\{N_{n,1}/N_{n,2}^\rho\}$ and $\{N_{n,1}/n^\rho\}$ have no point masses in $[0, \infty]$.

Theorem 3.2 *Suppose $m_2 > m_1 > 0$, $EU_{1,k}^p < \infty$, $k = 1, 2$, for some $p > 2$. Denote $\sigma_k^2 = E[(U_{1,k}/m_k)^2]$, $k = 1, 2$. Let $\delta_0 = \min\{(1 - \rho)/\rho, 1/2 - 1/p\}$. Then (possibly in an enlarged probability space) there is standard Brownian motion $B(y)$ such that for any $0 < \delta < \delta_0$,*

$$\psi_\infty - \psi_n = \frac{\sigma_1 \sqrt{m_1}}{m_2^{\rho/2}} \int_{n^\rho/\psi_\infty}^{\infty} \frac{dB(y)}{y} + o(n^{-\rho(1+\delta)/2}) \quad (3.1)$$

$$= -n^{-\rho/2} \frac{\sigma_1 \sqrt{m_1 \psi_\infty}}{m_2^{\rho/2}} \frac{W(n^\rho/\psi_\infty)}{\sqrt{n^\rho/\psi_\infty}} + o(n^{-\rho(1+\delta)/2}) \text{ a.s.}, \quad (3.2)$$

where $W(x) = -x \int_x^\infty y^{-1} dB(y)$ is also a standard Brownian motion.

Furthermore, the Brownian motion $B(y)$ can be constructed with a filtration of σ -fields $\{\mathcal{G}_n\}$ and a non-decreasing sequence of stopping times $\{T_n\}$ satisfying Properties (a) and (c) in Theorem 2.2, and

Property (b') $T_n = n^\rho/\psi_\infty + o(n^{\rho(1-\delta)})$ a.s. for $0 < \delta < \delta_0$.

The proof of this theorem will be given in the last section. Next, we state several corollaries. The first one is on the law of iterated logarithm and the central limit theorem for ψ_n .

Corollary 3.1 *Under the conditions in Theorem 3.2,*

$$\limsup_{n \rightarrow \infty} \frac{n^{\rho/2}(\psi_n - \psi_\infty)}{\sqrt{2 \log \log n}} = \frac{\sigma_1 \sqrt{m_1 \psi_\infty}}{m_2^{\rho/2}} \text{ a.s.} \quad (3.3)$$

and there exists a sequence $\{\zeta_n\}$ of standard normal random variables for which ζ_n is independent of \mathcal{F}_n and

$$n^{\rho/2}(\psi_n - \psi_\infty) = \frac{\sigma_1 \sqrt{m_1 \psi_n}}{m_2^{\rho/2}} \zeta_n + o(n^{-\epsilon}) \text{ a.s. for some } \epsilon > 0. \quad (3.4)$$

Hence

$$n^{\rho/2}(\psi_n - \psi_\infty) \Big|_{\mathcal{F}_n} \xrightarrow{d} N\left(0, \frac{\sigma_1^2 m_1}{m_2^\rho} \psi_\infty(\omega)\right) \text{ a.s.} \quad (3.5)$$

and ψ_∞ has no point masses in $(0, \infty)$.

Proof. (3.3) follows from (3.2) and the law of iterated logarithm of the Brownian motion. (3.4) can be proved in the same way as proving Corollary 2.1. \square

Corollary 3.2 *Under the conditions in Theorem 3.2,*

$$n^{\rho/2} \left(\frac{N_{n,1}}{N_{n,2}^\rho} - \eta_\infty \right) \xrightarrow{d} N(0, 1) \cdot \sqrt{\eta_\infty (2\sigma_1^2 - 1)} \text{ stably} \quad (3.6)$$

and

$$\begin{cases} n^{\rho/2} \left(\frac{N_{n,1}}{N_{n,2}^\rho} - \eta_\infty \right) \xrightarrow{d} N(0, 1) \cdot \sqrt{\eta_\infty (2\sigma_1^2 - 1)} \text{ stably} & \text{if } \rho < 2/3 \\ n^{1-\rho} \left(\frac{N_{n,1}}{N_{n,2}^\rho} - \eta_\infty \right) \rightarrow -\rho \eta_\infty^2 \text{ a.s.} & \text{if } \rho > 2/3, \end{cases} \quad (3.7)$$

$$n^{1-\rho} \left(1 - \frac{N_{n,2}}{n} \right) \rightarrow \eta_\infty \text{ a.s.} \quad (3.8)$$

Proof. For (3.6), let $A_{n,k}$ be defined as in (2.10). Then

$$A_{n,2} = O(\sqrt{N_{n,2}^{-1} \log \log N_{n,2}}) = O(\sqrt{n^{-1} \log \log n}) = o(n^{-\rho/2-\epsilon}) \text{ a.s.},$$

$$A_{n,1} = O(\sqrt{N_{n,1}^{-1} \log \log N_{n,1}}) = O(\sqrt{n^{-\rho} \log \log n}) \text{ a.s.}$$

Note that $\eta_\infty = \psi_\infty m_2^\rho / m_1$. It follows that

$$\begin{aligned} \frac{N_{n,1}}{N_{n,2}^\rho} - \eta_\infty &= -\eta_\infty + \psi_n \frac{m_2^\rho (1 + A_{n,2})^\rho}{m_1 (1 + A_{n,1})} \\ &= -\eta_\infty + \psi_n \frac{m_2^\rho}{m_1} (1 - A_{n,1}) + o(n^{-\rho/2-\epsilon}) \\ &= (\psi_n - \psi_\infty) \frac{m_2^\rho}{m_1} - n^{-\rho/2} \psi_n \frac{m_2^\rho}{m_1} \sqrt{n^\rho / N_{n,1}} (\sqrt{N_{n,1}} A_{n,1}) + o(n^{-\rho/2-\epsilon}) \\ &= n^{-\rho/2} \left\{ \sqrt{\psi_n m_1 / m_2^\rho} \zeta_n - \psi_n \frac{m_2^\rho}{m_1} \sqrt{n^\rho / N_{n,1}} (\sqrt{N_{n,1}} A_{n,1}) + o(n^{-\epsilon}) \right\} \text{ a.s.} \end{aligned}$$

The proof of (3.6) is completed by noting that $\psi_n m_1 / m_2^\rho \rightarrow \eta_\infty$ a.s., $N_{n,1} / n^\rho \rightarrow \eta_\infty$ a.s., $\sqrt{N_{n,1}} A_{n,1} \xrightarrow{d} N(0, \sigma_1^2 - 1)$ mixing, and ζ_n is a standard normal random variable which is independent of \mathcal{F}_n .

For (3.7), it is sufficient to note that

$$\begin{aligned}\frac{N_{n,1}}{n^\rho} &= \frac{N_{n,1}}{N_{n,2}^\rho} \left(1 - \frac{N_{n,1}}{n}\right)^\rho = \frac{N_{n,1}}{N_{n,2}^\rho} \left(1 - \rho \frac{N_{n,1}}{n} + O\left(\frac{N_{n,1}}{n}\right)^2\right) \\ &= \frac{N_{n,1}}{N_{n,2}^\rho} - \rho \frac{N_{n,1}}{N_{n,2}^\rho} \frac{N_{n,1}}{n^\rho} n^{\rho-1} + O(n^{2(\rho-1)}) \\ &= \frac{N_{n,1}}{N_{n,2}^\rho} - \rho \eta_\infty^2 n^{\rho-1} + o(n^{\rho-1}) \quad a.s.\end{aligned}$$

(3.8) is obvious because

$$1 - \frac{N_{n,2}}{n} = \frac{N_{n,1}}{n} \sim \frac{\eta_\infty n^\rho}{n} \quad a.s.$$

The proof is now completed. \square .

Finally, we give the functional central limit theorem.

Corollary 3.3 *Define*

$$W_n(t) = n^{\rho/2} t^\rho (\psi_{[nt]} - \psi_\infty), t > 0.$$

Then

$$W_n(t) \xrightarrow{d} \sigma_1 \sqrt{\eta_\infty} B'(t^\rho) \quad \text{stably}, \quad (3.9)$$

in the Skorokhod Topological space $D(0, \infty)$, where $B'(t)$ is a standard Brownian motion which is independent of \mathcal{F}_∞ . In particular,

$$\lim_{n \rightarrow \infty} P \left(\frac{\max_{0 \leq l \leq n} l^\rho (\psi_l - \psi_n)}{\sigma_1 \sqrt{N_{n,1}}} \geq x \right) = e^{-2x^2}, \quad x > 0. \quad (3.10)$$

Proof. The proof of (3.9) is similar to that of (2.3) by noting that $\eta_\infty = \psi_\infty m_1 / m_2^\rho$. For (3.10), it is sufficient to see that

$$\max_{0 \leq l \leq n} \frac{l^\rho (\psi_\infty - \psi_n)}{n^{\rho/2}} = \sup_{0 < t \leq 1} (W_n(t) - t^\rho W_n(1))$$

and $N_{n,1} \sim \eta_\infty n^\rho$ a.s.. \square

4 Concluding Remark

We approximated $Z_n - Z_\infty$ and $\psi_n - \psi_\infty$ by a kind of Gaussian process $\int_t^\infty y^{-1} dB(y)$, which is a tail stochastic integral with respect to a Brownian motion, with time t stopping at a random variable nH or n/ψ_∞ , where $H^2 = (1 + \psi_\infty)(\sigma_1^2/\psi_\infty + \sigma_2^2)$. But this does not mean that $\int_{nH}^\infty y^{-1} dB(y)$ and $\int_{n/\psi_\infty}^\infty y^{-1} dB(y)$ are Gaussian random variables and their distributions are unknown because the mixing distribution of ψ_∞ is unknown. For deriving the asymptotic distributions, the approximations (2.6) and (3.4) seem more powerful than (2.1) and (3.1) because

the process for approximation is independent of other random variables considered. (2.1) and (3.1) are helpful for establishing the strong convergence such as the law of the iterated logarithm.

It is of interest to find the distribution of ψ_∞ . In the case that $Y_{0,k}$ and $U_{m,k}$, $k = 1, 2$, are all integers, in a recent manuscript of Zhang, et al (2010) it is proved that the distribution of ψ_∞ is absolutely continuous and is determined by $Y_{0,1}$, $Y_{0,2}$ and the distributions of $U_{1,1}$ and $U_{1,2}$, if $\mathbb{E}[U_{1,k} \log^+ U_{1,k}] < \infty$, $k = 1, 2$. In the general case, the distribution of ψ_∞ is still a open problem. In our Corollaries 2.3 and 3.1, by applying the Gaussian approximation and a clever idea of Aletti, May and Secchi (2009) we show that ψ_∞ has no point masses in $[0, \infty]$ under the assumption of finite $(2 + \epsilon)$ -th moments. The next step is to show that the distribution is absolutely continuous. Unfortunately, as discussed in Aletti, May and Secchi (2009), the almost-sure conditional central limit theorems (2.5) and (3.5) are not enough to prove the absolute continuity. Our method and that of Aletti, May and Secchi (2009) depends on the martingale approach, which is not a very powerful tool to derive the limit distribution which is not normal. To find the exact distribution of ψ_∞ needs new methods.

For a special case that $\mathbb{P}(U_{m,k} = 0) = p$ and $\mathbb{P}(U_{m,k} = \alpha) = 1 - p$ with $\alpha > 0$ and $0 < p \leq 1$, in Aletti, May and Secchi (2007) it is shown that the distribution of Z_∞ is a beta distribution, and hence the probability that $\psi_\infty = Z_\infty / (1 - Z_\infty)$ falls into any subset of $[0, \infty]$ will not be zero if the Lebesgue measure of this subset is positive. So, it is also of interest to prove in the general case that the probability of ψ_∞ falling into any nonempty subinterval of $(0, \infty)$ will not be zero.

Finally, this paper only consider the two-color urn model. In the manuscript of Zhang, et al (2010), the asymptotic properties for a multi-color reinforced urn model are studied. It is expected to approximate the urn components after being suitably normalized by a multi-dimensional Gaussian process. The Skorokhod embedding method used in this paper does not work for the multi-dimension case. Though strong approximations for multi-dimensional martingales can be found in literature, for example, Monrad and Philipp (1991), Eberlein (1986) and Zhang (2004), the martingales concerning to the reinforced urn model usually do not satisfied a condition that the asymptotic conditional variability is \mathcal{F}_k -measurable for some fixed k (c.f., (A.5)), which is needed in the approximation theorems for multi-dimensional martingales. A new approach is needed for approximating the multi-color reinforced urn models.

A Proof of the main results

Recall $|\mathbf{Y}_n| = Y_{n,1} + Y_{n,2}$, $\mathbb{P}(X_{n,k} = 1 | \mathcal{F}_{n-1}) = Y_{n-1,k} / |\mathbf{Y}_{n-1}|$. We first prove Theorems 2.1 and 3.1.

Proofs of Theorems 2.1 and 3.1. Without loss of generality, assume $1 < q \leq 2$. It is obvious that Theorem 2.1 follows from Theorem 3.1 with $Z_\infty =$

$\psi_\infty/(1 + \psi_\infty)$. For Theorem 3.1, let

$$Q_n = \frac{1}{m_1} \log Y_{n,1} - \frac{1}{m_2} \log Y_{n,2}.$$

Then $Q_n = \frac{1}{m_1} \log \psi_n$. So, it is sufficient to show that Q_n converges almost sure to a finite limit. Write

$$\begin{aligned} \Delta Q_n &= Q_n - Q_{n-1} \\ &= \frac{1}{m_1} X_{n,1} \log \left(1 + \frac{U_{n,1}}{Y_{n-1,1}} \right) - \frac{1}{m_2} X_{n,2} \log \left(1 + \frac{U_{n,2}}{Y_{n-1,2}} \right) \\ &= \left[X_{n,1} \frac{U_{n,1}/m_1}{Y_{n-1,1}} - X_{n,2} \frac{U_{n,2}/m_2}{Y_{n-1,2}} \right] \\ &\quad + \left[-\frac{1}{m_1} X_{n,1} f \left(\frac{U_{n,1}}{Y_{n-1,1}} \right) + \frac{1}{m_2} X_{n,2} f \left(\frac{U_{n,2}}{Y_{n-1,2}} \right) \right] \quad (\text{A.1}) \\ &:= \Delta Q_n^{(1)} + \Delta Q_n^{(2)}, \end{aligned}$$

where $f(x) = x - \log(1+x)$ satisfying $0 \leq f(x) \leq x^q$ for $x \geq 0$. We need to prove the almost sure convergence of the random series $\sum_{n=1}^{\infty} \Delta Q_n$. We first consider the terms in the second bracket above. Denote $\mathcal{A}_n = \sigma(\mathcal{F}_n, X_{n+1,1}, X_{n+1,2})$. Note that $Y_{n,k} \approx N_{n,k}$ a.s. by Lemma A.4 of Hu and Zhang (2004), and then $|\mathbf{Y}_n| \approx N_{n,1} + N_{n,2} = n$ a.s.. We have that

$$\begin{aligned} \sum_{n=1}^{\infty} \mathbb{E} \left[X_{n,k} f \left(\frac{U_{n,k}}{Y_{n-1,k}} \right) \middle| \mathcal{A}_{n-1} \right] &\leq \sum_{n=1}^{\infty} X_{n,k} \mathbb{E} \left[\left(\frac{U_{n,k}}{Y_{n-1,k}} \right)^q \middle| \mathcal{A}_{n-1} \right] \\ &\leq \sum_{n=1}^{\infty} X_{n,k} \frac{\mathbb{E} U_{n,k}^q}{Y_{n-1,k}^q} \leq C \sum_{n=1}^{\infty} \frac{X_{n,k}}{(1 + N_{n,k})^q} \\ &\leq C \sum_{n=1}^{\infty} \int_{N_{n-1,k}}^{N_{n,k}} \frac{1}{(1+x)^q} dx \leq C \int_0^{\infty} \frac{dx}{(1+x)^q} < \infty \text{ a.s.} \end{aligned}$$

Hence $\sum_{n=1}^{\infty} \Delta Q_n^{(2)}$ converges almost surely. For $\{\Delta Q_n^{(1)}\}$, it is easily seen that it is a sequence of martingale differences with respect to the σ -filtration $\{\mathcal{F}_n\}$, and

$$\mathbb{E} [|\Delta Q_n^{(1)}|^q | \mathcal{F}_{n-1}] \leq \frac{\mathbb{E} [|U_{1,1}/m_1|^q]}{|\mathbf{Y}_{n-1}| Y_{n-1,1}^{q-1}} + \frac{\mathbb{E} [|U_{1,2}/m_1|^q]}{|\mathbf{Y}_{n-1}| Y_{n-1,2}^{q-1}}.$$

So, it is sufficient to show that $\sum_{n=1}^{\infty} 1/(|\mathbf{Y}_{n-1}| Y_{n-1,k}^{q-1}) < \infty$ a.s., $k = 1, 2$. The proof will be completed if we have proven that $Y_{n,k} \geq n^\epsilon$ for some positive ϵ . Now, it is obvious that

$$\sum_{n=1}^{\infty} (\log n)^{-q} \mathbb{E} [|\Delta Q_n^{(1)}|^q | \mathcal{F}_{n-1}] \leq C \sum_{n=1}^{\infty} \frac{1}{|\mathbf{Y}_{n-1}| (\log n)^q} < \infty \text{ a.s.,}$$

which implies that

$$\frac{1}{\log n} \sum_{l=1}^n \Delta Q_n^{(1)} \rightarrow 0 \text{ a.s., and hence } \frac{Q_n}{\log n} \rightarrow 0 \text{ a.s.}$$

On the event $\{Y_{n,1} \geq Y_{n,2}\}$, we have that

$$\begin{aligned}\log Y_{n,2} &= -m_2 Q_n + \frac{m_2}{m_1} \log |\mathbf{Y}_n| + \frac{m_2}{m_1} \log \frac{Y_{n,1}}{|\mathbf{Y}_n|} \\ &\geq -m_2 Q_n + \frac{m_2}{m_1} \log |\mathbf{Y}_n| + \frac{m_2}{m_1} \log \frac{1}{2}.\end{aligned}$$

On the event $\{Y_{n,1} \leq Y_{n,2}\}$, we have $Y_{n,2} \geq |\mathbf{Y}_n|/2$. Note that $|\mathbf{Y}_n| \approx n$. It follows that

$$\liminf_{n \rightarrow \infty} \frac{\log Y_{n,2}}{\log n} \geq \frac{m_2}{m_1} \wedge 1 \quad a.s.$$

Similarly, we have that

$$\liminf_{n \rightarrow \infty} \frac{\log Y_{n,1}}{\log n} \geq \frac{m_1}{m_2} \wedge 1 \quad a.s.$$

The proof of Theorems 2.1 and 3.1 is now completed. \square .

Proof of Theorem 2.2. Let $f(x) = e^{m_1 x} / (1 + e^{m_1 x})$. Then $f(Q_n) = Z_n$ and $f'(Q_\infty) = m_1 Z_\infty (1 - Z_\infty)$. According to the Taylor expansion, it is sufficient to show that $B(t)$ and T_n can be constructed such that

$$Q_\infty - Q_n = \frac{H}{m_1} \int_{nH}^{\infty} \frac{dB(x)}{x} + o(\lambda_n) \quad a.s.. \quad (\text{A.2})$$

Recall $m_1 = m_2$. It is easily shown that $|\mathbf{Y}_n|/n \rightarrow m_1$ a.s.. According to Theorem 2.1, $Z_\infty \in (0, 1)$ a.s., which implies that $Y_{n,k} \approx n$ a.s., $k = 1, 2$. So, for $\Delta Q_n^{(2)}$ in (A.1) we have

$$\sum_{l=1}^{\infty} \lambda_l^{-1} \mathbb{E}[|\Delta Q_l^{(2)}| | \mathcal{F}_{l-1}] \leq \sum_{l=1}^{\infty} \frac{\lambda_l^{-1}}{|\mathbf{Y}_{n-1}|} \left(\frac{\sigma_1^2}{Y_{l-1,1}} + \frac{\sigma_2^2}{Y_{l-1,2}} \right) \leq C \sum_{l=1}^{\infty} \lambda_l^{-1} l^{-2} < \infty,$$

which implies that $\sum_{l=n+1}^{\infty} |\Delta Q_l^{(2)}| = o(\lambda_n)$ a.s..

For $\Delta Q_n^{(1)}$, we use the truncation method. Let $\tilde{U}_{n,k} = U_{n,k}/m_k$, $\bar{U}_{n,k} = \tilde{U}_{n,k} I\{\tilde{U}_{n,k} \leq n^{1/p}\}$, $\bar{\sigma}_{n,k}^2 = \mathbb{E}\bar{U}_{n,k}^2$, $\bar{m}_{n,k} = \mathbb{E}[\bar{U}_{n,k}]$, $k = 1, 2$, $\bar{m}_n = \bar{m}_{n,1} - \bar{m}_{n,2}$, and

$$\Delta M_n^{(1)} = m_1 n \left(X_{n,1} \frac{\bar{U}_{n,1}}{Y_{n-1,1}} - X_{n,2} \frac{\bar{U}_{n,2}}{Y_{n-1,2}} \right),$$

$$\Delta M_n = \Delta M_n^{(1)} - \mathbb{E}[\Delta M_n^{(1)} | \mathcal{F}_{n-1}] = \Delta M_n^{(1)} - \frac{m_1 n \bar{m}_n}{|\mathbf{Y}_{n-1}|}.$$

Then $\{\Delta M_n, \mathcal{F}_n\}$ is a sequence of martingale differences. Note that

$$\sum_{n=1}^{\infty} \mathbb{P}(U_{n,k}/m_k > n^{1/p}) \leq \mathbb{E}(U_{1,k}/m_k)^p < \infty, \quad k = 1, 2.$$

From the Borel-Cantelli lemma, it follows that

$$\mathbb{P}(\Delta Q_n^{(1)} \neq \frac{1}{m_1 n} \Delta M_n \text{ i.o.}) = 0.$$

Also,

$$\begin{aligned} & \sum_{l=n+1}^{\infty} \frac{1}{m_1 l} |\mathbb{E}[\Delta M_l^{(1)} | \mathcal{F}_{l-1}]| = \sum_{l=n+1}^{\infty} \frac{|\bar{m}_l|}{|\mathbf{Y}_{l-1}|} \\ & \leq C \sum_{l=n+1}^{\infty} \frac{1}{l} \sum_{k=1}^2 \mathbb{E}[|\tilde{U}_{1,k}| I\{\tilde{U}_{1,k} > l^{1/p}\}] \leq C n^{1/p-1} \sum_{k=1}^2 \mathbb{E}[|\tilde{U}_{1,k}|^p] = o(\lambda_n). \end{aligned}$$

Hence, we conclude that

$$\begin{aligned} \sum_{l=n+1}^{\infty} \Delta Q_l &= \sum_{l=n+1}^{\infty} \frac{1}{m_1 l} \Delta M_l + o(\lambda_n) \\ &= \frac{1}{m_1} \left(\sum_{l=n}^{\infty} \frac{1}{l(l+1)} M_l - \frac{M_n}{n} \right) + o(\lambda_n) \text{ a.s.} \end{aligned}$$

For the martingale $M_n = \sum_{l=1}^n \Delta M_l$, we have

$$\mathbb{E}[(\Delta M_n)^2 | \mathcal{F}_{n-1}] = \left(\frac{m_1 n}{|\mathbf{Y}_{n-1}|} \right)^2 \left(\frac{\mathbb{E}\bar{U}_{n,1}^2}{Z_{n-1}} + \frac{\mathbb{E}\bar{U}_{n,2}^2}{1 - Z_{n-1}} - \bar{m}_n^2 \right), \quad (\text{A.3})$$

$$\mathbb{E}[|\Delta M_n|^4 | \mathcal{F}_{n-1}] \leq \sum_{k=1}^2 \left(\frac{m_1 n}{Y_{n-1,k}} \right)^4 \mathbb{E}\bar{U}_{n,k}^4 \leq C(\omega) \sum_{k=1}^2 \mathbb{E}\bar{U}_{n,k}^4. \quad (\text{A.4})$$

By the Skorokhod embedding theorem (c.f., Theorem A.1 of Hall and Heyde (1980, page 269)), (possibly in an enlarged probability space) there is a standard motion $B(x)$ with a filtration $\{\mathcal{G}_n\}$ and a sequence of nonnegative stopping times τ_1, τ_2, \dots with the following properties

- (i) $M_n = B(T_n)$, where $T_n = \sum_{i=1}^n \tau_i$;
- (ii) $\mathcal{F}_n \subset \mathcal{G}_n$, τ_n is \mathcal{G}_n measurable, $\mathbb{E}[\tau_n | \mathcal{G}_{n-1}] = \mathbb{E}[(\Delta M_n)^2 | \mathcal{F}_{n-1}]$, $\mathbb{E}[\tau_n^r | \mathcal{G}_{n-1}] \leq C_r \mathbb{E}[(\Delta M_n)^{2r} | \mathcal{F}_{n-1}]$ for any $r \geq 1$;
- (iii) Conditional on \mathcal{G}_n , $B(T_n + x) - B(T_n)$, $x \geq 0$, is also a standard Brownian motion.

Now, we verify that the Brownian motion $B(x)$ and the stopping time T_n are desirable for Property (b) and (A.2). At first, we assume the following approximation for the conditional variance.

$$\mathbb{E}[(\Delta M_n)^2 | \mathcal{F}_{n-1}] = H(\omega) + o(n^{2/p-1}) \text{ a.s.} \quad (\text{A.5})$$

From (A.5) it follows that

$$\sum_{i=1}^n \mathbb{E}[\tau_n | \mathcal{G}_{i-1}] = nH(\omega) + o(n^{2/p}) \text{ a.s.}$$

On the other hand, by (ii) and (A.4) we have that

$$\sum_{n=1}^{\infty} \mathbb{E} \left[\left(\frac{\tau_n}{n^{2/p}} \right)^2 | \mathcal{G}_{n-1} \right] \leq C \sum_{n=1}^{\infty} \frac{\sum_{k=1}^2 \mathbb{E} \bar{U}_{n,k}^4}{n^{4/p}} \leq C \sum_{k=1}^2 \mathbb{E}[(U_{1,k}/m_k)^p] < \infty.$$

By the law of large numbers of martingale, it follows that

$$\sum_{i=1}^n (\tau_i - \mathbb{E}[\tau_i | \mathcal{G}_{i-1}]) = o(n^{2/p}) \text{ a.s.}$$

Hence

$$T_n = \sum_{i=1}^n \tau_i = nH + o(n^{2/p}) \text{ a.s.}$$

Property (b) is verified. Then it follows from the path properties of a Brownian motion (c.f., Theorem 1.2.1 of Csörgő and Révész (1981)) that

$$B(T_n) - B(nH) = o\left(\sqrt{n^{2/p} \left(\log \frac{n}{n^{2/p}} + \log \log n\right)}\right) = o(n\lambda_n) \text{ a.s.}$$

So

$$\begin{aligned} \sum_{l=n+1}^{\infty} \Delta Q_l &= \frac{1}{m_1} \left(\sum_{l=n}^{\infty} \frac{1}{l(l+1)} B(T_l) - \frac{B(T_n)}{n} \right) + o(\lambda_n) \\ &= \frac{1}{m_1} \left(\sum_{l=n}^{\infty} \frac{1}{l(l+1)} B(lH) - \frac{B(nH)}{n} \right) + \sum_{l=n}^{\infty} \frac{1}{l(l+1)} o(l\lambda_l) + o(\lambda_n) \\ &= \frac{1}{m_1} \left(\int_n^{\infty} \frac{B(xH)}{x^2} dx - \frac{B(nH)}{n} \right) + o(\lambda_n) \\ &= \frac{1}{m_1} \left(H \int_{nH}^{\infty} \frac{B(x)}{x^2} dx - \frac{B(nH)}{n} \right) + o(\lambda_n) \\ &= \frac{H}{m_1} \int_{nH}^{\infty} \frac{dB(x)}{x} + o(\lambda_n) \text{ a.s.} \end{aligned}$$

Finally, we verify (A.5). Note that $\mathbb{E}[\bar{U}_{n,k}^2] \rightarrow \sigma_k^2$, $|\mathbf{Y}_n|/n \rightarrow m_1$ a.s., $k = 1, 2$ and $Z_n \rightarrow Z_{\infty}$ a.s.. (A.5) is obvious for $p = 2$ by (A.3).

For $2 < p < 4$, we still have Corollary (2.1) due to the approximation for the case of $p = 2$. Hence

$$Z_n - Z_{\infty} = O(\sqrt{n^{-1} \log \log n}) = o(n^{2/p-1}) \text{ a.s.}$$

On the other hand,

$$\begin{aligned} \frac{Y_{n,1} + Y_{n,2}}{m_1 n} - 1 &= \frac{\sum_{k=1}^2 \sum_{i=1}^n X_{i,k} (U_{i,k} - \mathbb{E}[U_{i,k}])}{m_1 n} \\ &= O(\sqrt{n^{-1} \log \log n}) = o(n^{2/p-1}) \text{ a.s.} \end{aligned}$$

and

$$|\bar{m}_n| \leq \sum_{k=1}^2 \mathbb{E}[U_{1,k}/m_k I\{U_{1,k}/m_k > n^{1/p}\}] = o(n^{1/p-1}),$$

$$\sigma_k^2 - \mathbb{E}[\bar{U}_{n,k}^2] = \mathbb{E}[(U_{1,k}/m_k)^2 I\{U_{1,k}/m_k > n^{1/p}\}] = o(n^{2/p-1}), \quad k = 1, 2.$$

(A.5) follows by (A.3). The proof is now completed. \square

Proof of Theorem 3.2. As before, denote $|\mathbf{Y}_n| = Y_{n,1} + Y_{n,2}$, $Q_n = \frac{1}{m_1} \log Y_{n,1} - \frac{1}{m_2} \log Y_{n,2}$. According to the Taylor expansion, it is sufficient to show that $B(t)$ and T_n can be constructed such that

$$Q_\infty - Q_n = \frac{\sigma_1}{\sqrt{m_1 m_2} \psi_\infty} \int_{n^\rho/\psi_\infty}^\infty \frac{dB(y)}{y^2} + o(n^{-\rho(1+\delta)/2}) \text{ a.s.} \quad (\text{A.6})$$

Recall (A.1) and note that $Y_{n-1,2} \sim m_2 n$, $Y_{n-1,1} \sim \psi_\infty Y_{n,2}^\rho \sim \psi_\infty (m_2 n)^\rho$ a.s.. It can be show that for $0 < \delta_1 \leq 1/2$ and $\delta_1 < (1 - \rho)/\rho$,

$$\sum_{l=1}^\infty l^{\rho(1+\delta_1)/2} \mathbb{E}\left[X_{l,k} f\left(\frac{U_{l,k}}{Y_{l-1,k}}\right) \middle| \mathcal{F}_{l-1}\right] \leq \sum_{l=1}^\infty \frac{l^{\rho(1+\delta_1)/2}}{l \cdot l^\rho} < \infty,$$

which implies $\sum_{l=n+1}^\infty X_{l,k} f(U_{l,k}/Y_{l-1,k}) = o(n^{-\rho(1+\delta_1)/2})$ a.s., $k = 1, 2$. Also, for the martingale differences $X_{l,2} \frac{U_{l,2}/m_2}{Y_{l-1,2}} - \frac{1}{|\mathbf{Y}_{l-1}|}$, we have

$$\sum_{l=1}^\infty (l^{\rho(1+\delta_1)/2})^2 \mathbb{E}\left[\left(X_{l,2} \frac{U_{l,2}/m_2}{Y_{l-1,2}} - \frac{1}{|\mathbf{Y}_{l-1}|}\right)^2 \middle| \mathcal{F}_{l-1}\right] \leq \sum_{l=1}^\infty \frac{l^{\rho(1+\delta_1)}}{l^2} < \infty \text{ a.s.},$$

which implies $\sum_{l=n+1}^\infty (X_{l,2} \frac{U_{l,2}/m_2}{Y_{l-1,2}} - \frac{1}{|\mathbf{Y}_{l-1}|}) = o(n^{-\rho(1+\delta_1)/2})$ a.s.. Similarly, we can show that $\sum_{l=n+1}^\infty (X_{l,1} \frac{U_{l,1}/m_2}{Y_{l-1,1}} - \frac{1}{|\mathbf{Y}_{l-1}|}) = o(n^{-\rho/2} \log n)$ a.s. It follows that

$$Q_\infty - Q_n = \sum_{l=n+1}^\infty \left(X_{l,1} \frac{U_{l,1}/m_1}{Y_{l-1,1}} - \frac{1}{|\mathbf{Y}_{l-1}|} \right) + o(n^{-\rho(1+\delta_1)/2}) \text{ a.s.} \quad (\text{A.7})$$

and

$$Q_\infty - Q_n = o(n^{-\rho/2} \log n) \text{ a.s.} \quad (\text{A.8})$$

Define

$$\Delta M_n = \frac{\sqrt{\rho m_2} (m_2 n)^{\rho/2} n^{\rho/2}}{\sigma_1} \left(X_{n,1} \frac{U_{n,1}/m_1}{Y_{n-1,1}} - \frac{1}{|\mathbf{Y}_{n-1}|} \right). \quad (\text{A.9})$$

Then

$$\mathbb{E}[(\Delta M_n)^2 | \mathcal{F}_{n-1}] = \frac{\rho m_2 (m_2 n)^\rho n^\rho}{|\mathbf{Y}_{n-1}| Y_{n-1,1}} - \frac{\rho m_2 (m_2 n)^\rho n^\rho}{\sigma_1^2 |\mathbf{Y}_{n-1}|^2}.$$

Next, we first show that

$$\mathbb{E}[(\Delta M_n)^2 | \mathcal{F}_{n-1}] = \frac{\rho}{\psi_\infty} n^{\rho-1} (1 + o(n^{-\rho\delta_1})) \quad a.s. \quad (\text{A.10})$$

From (A.8) and the Taylor expansion, we conclude that

$$\frac{Y_{n,1}}{(Y_{n,2})^\rho} - \psi_\infty = o(n^{-\rho/2} \log n) \quad a.s.$$

On the other hand,

$$\begin{aligned} \frac{Y_{n,2}}{m_2 n} &= 1 - \frac{1}{m_1} \frac{Y_{n,1}}{n} + \sum_{k=1}^2 \frac{\sum_{l=1}^n X_{l,k} (U_{l,k}/m_k - \mathbb{E}[U_{l,k}/m_k])}{n} \\ &= 1 - \frac{1}{m_1} \frac{\psi_\infty (Y_{n,2})^\rho}{n} + o(n^{-\rho/2-1} \log n) + O(n^{-1/2} (\log \log n)^{1/2}) \\ &= 1 - O(n^{\rho-1}) + o(n^{-\rho\delta_1}) = 1 + o(n^{-\rho\delta_1}) \quad a.s. \end{aligned}$$

It follows that

$$\frac{|\mathbf{Y}_n|}{m_2 n} = \frac{Y_{n,2}}{m_2 n} + \frac{Y_{n,1}}{m_2 n} = 1 + o(n^{-\rho\delta_1}) \quad \text{and} \quad \frac{Y_{n,1}}{(m_2 n)^\rho} - \psi_\infty = o(n^{-\rho\delta_1}) \quad a.s.$$

(A.10) is verified. From (A.10), it follows that

$$\sum_{l=1}^n \mathbb{E}[(\Delta M_n)^2 | \mathcal{F}_{n-1}] = \frac{n^\rho}{\psi_\infty} (1 + o(n^{-\rho\delta_1})) \quad a.s.$$

On the other hand, for $0 < \delta_2 < 1/2 - 1/p$,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\mathbb{E}[|\Delta M_n|^p | \mathcal{F}_{n-1}]}{(n^{\rho(1-\delta_2)})^{p/2}} &\leq C \sum_{n=1}^{\infty} \frac{n^{\rho p}}{n^{\rho(1-\delta_2)\rho/2}} \frac{Y_{n-1,1}}{Y_{n-1,1}^\rho |\mathbf{Y}_{n-1}|} \\ &\leq C \sum_{n=1}^{\infty} \frac{1}{n^{\rho(1-\delta_2)\rho/2}} \frac{1}{n^{1-\rho}} < \infty. \end{aligned}$$

So, similarly as in the proof of Theorem 2.2, by the Skorokhod embedding theorem, the standard motion $B(x)$, the filtration $\{\mathcal{G}_n\}$ and the stopping times $\{T_n\}$ can be constructed such that $M_n = B(T_n)$ and

$$T_n = \sum_{l=1}^n \mathbb{E}[(\Delta M_l)^2 | \mathcal{F}_{l-1}] + o(n^{\rho(1-\delta_2)}) = \frac{1}{\psi_\infty} n^\rho + o(n^{\rho(1-\delta_1 \wedge \delta_2)}).$$

Denote $\delta_0 = \min\{(1 - \rho)/\rho, 1/2 - 1/p\}$. It is remained to verify (3.1). By the Properties (b') and the path properties of a Brownian motion, we have for any $0 < \delta < \delta_0$,

$$M_n - B(n^\rho/\psi_\infty) = o(n^{\rho(1-\delta)/2}) \text{ a.s.}$$

Hence

$$\begin{aligned} \sum_{l=n+1}^{\infty} \frac{\Delta M_l}{l^\rho} &= \sum_{l=n}^{\infty} \left(\frac{1}{l^\rho} - \frac{1}{(l+1)^\rho} \right) M_l - \frac{M_n}{n^\rho} \\ &= \sum_{l=n}^{\infty} \left(\frac{1}{l^\rho} - \frac{1}{(l+1)^\rho} \right) B(l^\rho/\psi_\infty) - \frac{B(n^\rho/\psi_\infty)}{n^\rho} \\ &\quad + \sum_{l=n}^{\infty} \frac{o(l^{\rho(1-\delta)/2})}{l^{1+\rho}} + \frac{o(n^{\rho(1-\delta)/2})}{n^\rho} \\ &= \int_n^\infty \frac{\rho B(x^\rho/\psi)}{x^{1+\rho}} dx - \frac{B(n^\rho/\psi_\infty)}{n^\rho} + o(n^{-\rho(1+\delta)/2}) \\ &= \frac{1}{\psi_\infty} \int_{n^\rho/\psi_\infty}^\infty \frac{B(x)}{x^2} dx - \frac{B(n^\rho/\psi_\infty)}{n^\rho} + o(n^{-\rho(1+\delta)/2}) \\ &= \frac{1}{\psi_\infty} \int_{n^\rho/\psi_\infty}^\infty \frac{dB(x)}{x} dx + o(n^{-\rho(1+\delta)/2}) \text{ a.s.} \end{aligned}$$

(A.6) is now proved by noting that (A.7), (A.8) and $\rho m_2 = m_1$. And hence (3.1) is verified. \square

REMARK A.1 *Using the truncation method as in the proof of Theorem 2.2, we can prove that (3.1) remains true under the assumption of only finite second moments if $n^{-\rho(1+\delta)/2}$ is replaced by $n^{-\rho/2}(\log \log n)^{1/2}$. This implies that the law of iterated logarithm (3.3) remains true when $EU_{1,k}^2 < \infty$, $k = 1, 2$.*

References

- [1] ALETTI, G., MAY, C. and SECCHI, P. (2007). On the distribution of the limit proportion for a two-color, randomly reinforced urn with equal reinforcement distributions. *Adv. Appl. Probab.*, **39**: 690-707.
- [2] ALETTI, G., MAY, C. and SECCHI, P. (2009). A central limit theorem, and related results, for a two-color randomly reinforced urn. *Adv. Appl. Probab.*, **41**: 829-844.
- [3] BAI, Z. D. and HU, F. (2005). Strong consistency and asymptotic normality for urn models. *Ann. Appl. Probab.*, **12**: 914-940.
- [4] BAI, Z. D., HU, F. and ROSENBERGER, W. F. (2002). Asymptotic properties of adaptive designs for clinical trials with delayed response. *Ann. Statist.*, **30**: 122-139.
- [5] BAI, Z. D., HU, F. and ZHANG, L. X. (2002). The Gaussian approximation theorems for urn models and their applications. *Ann. Appl. Probab.*, **12**: 1149-1173.

- [6] BEGGS, A. W. (2005). On the convergence of reinforcement learning. *J. Econom. Theory*, **122**(1): 1-36.
- [7] CHAUVIN, B., POUYANNE, N. and SAHNOUN, R. (2009). Limit distributions for large Pólya urns. <http://arxiv.org/abs/0907.1477>
- [8] CRIMALDI, I. (2009). An almost sure conditional convergence result and an application to a generalized Pólya Urn. *International Mathematical Forum*, **4** (23): 1139–1156
- [9] DURHAM, S. D., FLOURNOY, N. and LI, W. (1998). A sequential design for maximizing the probability of a favourable response. *Canad. J. Statist.*, **26** (3): 479-495.
- [10] DURHAM, S. D. and YU, K. F. (1990). Randomized play-the leader rules for sequential sampling from two populations. *Probability in Engineering and Information Science*, **26** (4): 355-367.
- [11] EBERLEIN, E. (1986). On strong invariance principles under dependence. *Ann. Probab.*, **14**: 260C270.
- [12] EGGENBERGER, F. and POLYA, G. (1923). Über die Statistik verketteter Vorgänge. *Zeitschrift Angew. Math. Mech.*, **3**: 279C289.
- [13] EREV, I. and ROTH, A. (1998). Predicting how people play games: reinforcement learning in experimental games with unique, mixed strategy equilibria. *Amer. Econ. Rev.*, **88**: 848-881.
- [14] HALL, P. and HEYDE, C. C. (1980). *Martingale Limit Theory and its Applications*. Academic Press, London.
- [15] HOPKINS, E. and POSCH, M. (2005). Attainability of boundary points under reinforcement learning. *Games Econom. Behavior*, **53**: 110-125.
- [16] HANSON, D. L. and RUSSO, RALPH P. (1983). Some results on increments of the Wiener process with applications to lag sums of i.i.d.r.v. *Ann. Probab.*, **11**: 609-623.
- [17] HU, F. and ROSENBERGER, W. F. (2006). *The Theory of Response-Adaptive Randomization in Clinical Trials*, John Wiley and Sons, Inc., New York.
- [18] HU, F. & ZHANG, L.-X. (2004). Asymptotic properties of doubly adaptive biased coin designs for multi-treatment clinical trials. *Ann. Statist.*, **32**: 268-301.
- [19] JANSON, S. (2004). Functional limit theorems for multitype branching processes and generalized Pólya urns. *Stochastic Process. Appl.*, **110**: 177-245.
- [20] JANSON, S. (2006). Limit theorems for triangular urn schemes. *Probab. Theory Relat. Fields*, **134**:417C452.
- [21] LI, W., DURHAM, S. D. and FLOURNOY, N. (1996). Randomized polya urn designs. *Proceedings of the Biometric Section of the Statistical Association*: 166-170.
- [22] MAY, C. and FLOURNOY, N. (2009). Asymptotics in response-adaptive designs generated by a two-color, randomly reinforced urn. *Ann. Statist.*, **37**(2): 1058-1078.
- [23] MARTIN, C. F. and HO, Y. C. (2002). Value of information in the Polya urn process. *Information Sciences*, **147**: 65-90.
- [24] MELFI, V. F. and PAGE, C. (2000). Estimation after adaptive allocation. *J. Statist. Plann. Inf.*, **87**: 353–363.
- [25] MONRAD, D. and PHILIPP, W. (1991). Nearby variables with nearby conditional laws and a strong approximation theorem for Hilbert space valued martingales. *Probab. Theory Relat. Fields*, **88**: 381C-404.

- [26] MULIERE, P., PAGANONI, A. M. and SECCHI, P. (2006a). A randomly reinforced urns. *J. Statist. Plann. Inference*, **136**(6):1853-1874.
- [27] MULIERE, P., PAGANONI, A. M. and SECCHI, P. (2006b). Randomly reinforced urns for clinical trials with continuous responses. In *SISProceedings of the XLIII Scientific Meeting*, 403C 414. Cleup, Padova.
- [28] PAGANONI, A. and SECCHI, P. (2007). A numerical study for comparing two response-adaptive designs for continuous treatment effects. *Statistical Methods and Applications*, **16**: 321C346.
- [29] PÓLYA, G. (1931). Sur quelques points de la théorie des probabilités. *Ann. Inst. Poincaré*, **1**: 117C161.
- [30] ZHANG, L. X. (2004). Strong approximations of martingale vectors and its applications in Markov-Chain adaptive designs. *ACTA MATH. APPL. SINICA, ENGLISH SERIES*, **20**(2): 337-352.
- [31] ZHANG, L. X. and HU, F. (2009). The Gaussian approximation for multi-color generalized Friedman's urn model. *Science in China, Ser. A*, **52** (6): 1305-1326.
- [32] ZHANG, L. X., HU, F. and CHEUNG, S. H. (2006). Asymptotic theorems of sequential estimation-adjusted urn models for clinical trials. *Ann. Appl. Probab.*, **16**(1): 340-369
- [33] ZHANG, L. X., HU, F., CHEUNG, S. H. and CHAN, W. S. (2010). Asymptotic properties of multi-color randomly reinforced Pólya urns. *Manuscript*.