

# Light Stop from $b$ - $\tau$ Yukawa Unification

Ilia Gogoladze,<sup>\*</sup> Shabbar Raza,<sup>†</sup> and Qaisar Shafi  
*Bartol Research Institute, Department of Physics and Astronomy  
 University of Delaware, Newark, Delaware 19716*

We show that  $b$ - $\tau$  Yukawa unification can be successfully implemented in the constrained minimal supersymmetric model and it yields the stop co-annihilation scenario. The lightest supersymmetric particle is a bino-like dark matter neutralino, which is accompanied by a 10-20% heavier stop of mass  $\sim 100 - 330$  GeV. We highlight some benchmark points which show a gluino with mass  $\sim 0.6 - 1.7$  TeV, while the first two family squarks and all sleptons have masses in the multi-TeV range.

PACS numbers: 12.60.Jv, 12.10.Dm, 14.80.Ly

The apparent unification at  $M_G \approx 2 \times 10^{16}$  GeV of the three Standard Model (SM) gauge couplings, assuming TeV scale supersymmetry (SUSY), strongly suggests the existence of an underlying grand unified theory with a single coupling constant. The minimal supersymmetric SU(5) model, in addition to unifying the gauge couplings, also predicts unification at  $M_G$  of the third family bottom ( $b$ ) quark and tau lepton ( $\tau$ ) Yukawa couplings [1]. This  $b$ - $\tau$  Yukawa unification (YU) is to be contrasted with the minimal supersymmetric SO(10) and  $SU(4)_c \times SU(2)_L \times SU(2)_R$  models which predict  $t$ - $b$ - $\tau$  YU [2], where  $t$  denotes the top quark. The low energy implications of  $b$ - $\tau$  [3] and  $t$ - $b$ - $\tau$  [4] YU have been discussed in the recent literature. For instance,  $t$ - $b$ - $\tau$  YU is not realized in the mSUGRA/constrained minimal supersymmetric standard model (CMSSM) [5] because of the difficulty of implementing radiative electroweak symmetry breaking (REWSB) [6].

In this paper we explore the low energy consequences of implementing  $b$ - $\tau$  YU in the CMSSM framework. REWSB in this case is not an issue anymore. We refer to this combination of  $b$ - $\tau$  YU and CMSSM as YCMSSM, the ‘Yukawa’ constrained version of CMSSM. Among other things, we require that YCMSSM delivers a viable cold dark matter (DM) candidate (lightest stable neutralino) whose relic energy density is compatible with the WMAP measured value [7]. One of our main observations is that the allowed fundamental parameter space of CMSSM is strikingly reduced in the YCMSSM setup. We find that  $M_{1/2} \ll m_0$ , where  $M_{1/2}$  and  $m_0$  denote universal gaugino and scalar soft SUSY breaking masses respectively. Furthermore,  $b$ - $\tau$  YU at the level of 10% or better yields the constraint  $5 \text{ TeV} \lesssim m_0 \lesssim 20 \text{ TeV}$ . The supersymmetric threshold corrections including finite loop corrections to the  $b$  quark mass play an essential role here [8].

The lightest supersymmetric particle (LSP) neutralino is essentially a bino, the spin 1/2 supersymmetric partner of the  $U(1)_Y$  gauge boson, which is closely followed in mass by a slightly heavier (next to lightest sparticle for short NLSP) stop, a scalar partner of the top quark. The desired LSP relic abundance is achieved via neutralino stop co-annihilation, which in our case requires that the NLSP stop is about 10-20% heavier than the neutralino. The parameter  $\tan \beta$ , the ratio of the up and down Higgs VEVs, turns out to lie in a narrow range  $35 \lesssim \tan \beta \lesssim 40$ . The universal trilinear scalar

coupling ( $A_0$ ) is found to satisfy  $|A_0/m_0| \sim 2.3$ .

We highlight some LHC testable benchmark points with comparable LSP neutralino and light stop masses of around 100 – 330 GeV, while the corresponding chargino and second neutralino masses are 200 – 600 GeV and gluino mass  $\sim 0.6 - 1.7$  TeV. Together with the lightest SM-like Higgs with mass 114 – 124 GeV, these are the only ‘light’ (LHC accessible) particles predicted in this NLSP stop scenario with  $b$ - $\tau$  Yukawa unification and neutralino DM. The squarks of the first two families, the heavy stop, the two sbottom particles, and the charged sleptons, all have large (multi-TeV) masses.

The fundamental parameters of CMSSM are:

$$m_0, M_{1/2}, \tan \beta, A_0, \text{sgn}(\mu) \quad (1)$$

where  $\text{sgn}(\mu)$  is the sign of supersymmetric bilinear Higgs parameter. All mass parameters are specified at  $M_G$ .

We use the ISAJET 7.80 package [9] to perform random scans over the CMSSM parameter space. ISAJET employs two-loop renormalization group equations (RGEs) and defines  $M_G$  to be the scale at which  $g_1 = g_2$ . This is more than adequate as a few percent deviation from the exact unification condition  $g_3 = g_1 = g_2$  can be assigned to unknown GUT scale threshold corrections [10]. The random scans cover the following parameter range:

$$\begin{aligned} 0 \leq m_0 \leq 25 \text{ TeV}, & \quad 0 \leq M_{1/2} \leq 2 \text{ TeV}, \\ 1.1 \leq \tan \beta \leq 60, & \quad -3 \leq A_0/m_0 \leq 3, \end{aligned} \quad (2)$$

with  $\mu > 0$  and  $m_t = 173.3$  GeV [11]. The results are not too sensitive to one or two sigma variation in the value of  $m_t$  [12].

In scanning the parameter space, we employ the Metropolis-Hastings algorithm as described in [13]. All of the collected data points satisfy the requirement of REWSB, with the neutralino in each case being the LSP. We direct the Metropolis-Hastings algorithm to search for solutions with 10% or better  $b$ - $\tau$  Yukawa unification (YU). After collecting the data, we impose the experimental mass bounds on all particles [14], and use the IsaTools package [15] to implement the following phenomenological constraints:

As far as muon anomalous magnetic moment is concerned, we




FIG. 1: Evolution of bottom (green) and  $\tau$  (red) Yukawa couplings without (a) and with (b) finite SUSY threshold corrections.

$$\begin{aligned}
 m_h \text{ (lightest Higgs mass)} &\geq 114.4 \text{ GeV} & [16] \\
 BR(B_s \rightarrow \mu^+ \mu^-) &< 5.8 \times 10^{-8} & [17] \\
 2.85 \times 10^{-4} \leq BR(b \rightarrow s\gamma) &\leq 4.24 \times 10^{-4} \text{ (} 2\sigma \text{)} & [18] \\
 0.15 \leq \frac{BR(B_u \rightarrow \tau \nu_\tau)_{\text{MSSM}}}{BR(B_u \rightarrow \tau \nu_\tau)_{\text{SM}}} &\leq 2.41 \text{ (} 3\sigma \text{)} & [19] \\
 \Omega_{\text{CDM}} h^2 &= 0.111^{+0.028}_{-0.037} \text{ (} 5\sigma \text{)} & [7]
 \end{aligned}$$

only require that the model does no worse than the standard model (SM). In Fig. 1(a) we show the evolution of  $b$  and  $\tau$  Yukawa couplings without the SUSY threshold corrections for a representative  $b$ - $\tau$  YU solution. It is evident that without suitable the SUSY threshold corrections,  $b$ - $\tau$  YU occurs at around  $10^8$  GeV. In Fig. 1(b) we show the need for SUSY threshold corrections in order to achieve  $b$ - $\tau$  YU.

The SUSY correction  $\delta m_\tau$  to the  $\tau$  lepton mass is given by  $\delta m_\tau = v \cos \beta \delta y_\tau$ . The finite and logarithmic corrections [20] to the  $\tau$  lepton mass are typically small, so that the value of  $y_\tau$  at  $M_G$  is more or less fixed. From Fig. 1(a), to implement  $y_\tau = y_b$  at  $M_G$  therefore requires suitable threshold correction ( $\delta y_b$ ) to the bottom quark [8]. The dominant contribution to  $\delta y_b$  comes from the gluino and chargino loops [8, 20]. With our sign conventions (the sign of  $\delta y_b$  is fixed as  $y_b$  evolves from  $M_G$  to  $M_Z$ ), a useful approximate formula for the finite one loop correction to  $\delta y_b$  is given by

$$\delta y_b^{\text{finite}} \approx \frac{\mu}{4\pi^2} \left( \frac{g_3^2}{3} \frac{m_{\tilde{g}}}{m_1^2} + \frac{y_t^2}{8} \frac{A_t}{m_2^2} \right) \tan \beta. \quad (3)$$

Here  $g_3$  is the strong gauge coupling,  $m_{\tilde{g}}$  is the gluino mass,  $A_t$  is the stop trilinear coupling, and  $m_1 \approx (m_{\tilde{b}_1} + m_{\tilde{b}_2})/2$ ,  $m_2 \approx (m_{\tilde{t}_2} + \mu)/2$ .  $\tilde{b}_1, \tilde{b}_2$  denote the two bottom squarks,  $\tilde{t}_2$  is the heavier stop, and we assume that  $m_{\tilde{g}} \ll m_{\tilde{b}_1, \tilde{b}_2}$  and  $m_{\tilde{t}_1} \ll \mu, m_{\tilde{t}_2}$ .

In order to achieve  $b$ - $\tau$  YU,  $y_b$  must receive a negative contribution ( $-0.2 \lesssim \delta y_b/y_b \lesssim -0.07$ ) from threshold corrections. This is a relatively narrow interval compared to the full range  $-0.2 \lesssim \delta y_b/y_b \lesssim 0.25$  in the data that we have collected. The logarithmic corrections to  $y_b$  are in fact, positive, which leaves the finite corrections to provide for the correct  $\delta y_b$ . Since  $\mu > 0$ , the gluino contribution is positive, and so the contribution from the chargino loop not only has to cancel out the contributions from the gluino loop and the logarithmic correction, it also must provide the correct overall (negative)

contribution to  $\delta y_b$ . This is achieved only for suitable large  $m_0$  values and large negative  $A_t$ , for which the gluino contribution scales as  $M_{1/2}/m_0^2$  while the chargino contribution scales as  $A_t/m_0^2$ . Note that large values of  $m_0$  imply heavy slepton masses, and so  $b$ - $\tau$  YU does not provide any significant SUSY contributions to the muon anomalous magnetic moment.

In order to quantify  $b$ - $\tau$  YU, we define the quantity  $R_{b\tau}$  as

$$R_{b\tau} = \frac{\max(y_b, y_\tau)}{\min(y_b, y_\tau)}. \quad (4)$$

In Fig. 2 we present our results in the  $R_{b\tau}$  -  $m_0$ ,  $R_{b\tau}$  -  $\tan \beta$ ,  $M_{1/2}$  -  $m_0$  and  $R_{b\tau}$  -  $\mu$  planes. In panels (a, b, d) the gray points are consistent with REWSB and  $\tilde{\chi}_1^0$  LSP. The orange points satisfy the particle mass bounds, constraints from  $BR(B_s \rightarrow \mu^+ \mu^-)$ ,  $BR(B_u \rightarrow \tau \nu_\tau)$  and  $BR(b \rightarrow s\gamma)$ . The blue points form a subset of orange points that satisfies WMAP bounds on  $\tilde{\chi}_1^0$  DM abundance. The dashed (red) line represents 10% or better  $b$ - $\tau$  YU. In the  $M_{1/2}$  -  $m_0$  plane the orange color has the same meaning as described earlier, red color represents solutions with 10% or better  $b$ - $\tau$  YU, and the blue points satisfy in addition the WMAP bounds. In the  $R_{b\tau}$  -  $m_0$  plane of Fig. 2, we show that in order to have 10% or better  $b$ - $\tau$  YU consistent with the collider bounds, we require  $m_0 \gtrsim 5$  TeV. But if  $b$ - $\tau$  YU is to be compatible with neutralino DM relic density, represented by the blue points in the figure, we can see that  $m_0 \gtrsim 8$  TeV. To better appreciate this




FIG. 2: Plots in  $R_{b\tau}$  -  $m_0$ ,  $R_{b\tau}$  -  $\tan \beta$ ,  $M_{1/2}$  -  $m_0$  and  $R_{b\tau}$  -  $\mu$  planes. In panels (a, b, d) the gray points are consistent with REWSB and  $\tilde{\chi}_1^0$  LSP. The orange points satisfy in addition the particle mass bounds, constraints from  $BR(B_s \rightarrow \mu^+ \mu^-)$ ,  $BR(B_u \rightarrow \tau \nu_\tau)$  and  $BR(b \rightarrow s\gamma)$ . The blue points form a subset of orange points that satisfies WMAP bounds on  $\tilde{\chi}_1^0$  DM abundance. The dashed (red) line represents 10% or better  $b$ - $\tau$  YU. In  $M_{1/2}$  -  $m_0$  plane the orange color has the same meaning as described above, red color represents the solution with 10% or better  $b$ - $\tau$  YU, and blue points satisfy the WMAP bounds.

result we consider the  $M_{1/2} - m_0$  plane. As we will see later, the neutralino-stop co-annihilation channel is the only solution we have found for neutralino DM compatible with  $b$ - $\tau$  YU. The lighter stop mass can be as light as 100 GeV [14], which means that the neutralino mass has to be of comparable order, in order to implement the stop co-annihilation solution. An ISAJET two loops analysis yields  $M_{1/2} \gtrsim 150$  GeV. We can see from Fig. 2(c) that  $M_{1/2} \gtrsim 150$  GeV is compatible with 10% or better YU (points in red) for  $m_0 \gtrsim 8$  TeV.

In the  $R_{b\tau} - \tan \beta$  plane (Fig. 2(b)) we have two regions with 10% or better  $b$ - $\tau$  YU. In one region  $\tan \beta \approx 1.1$ , but this is ruled out by the light CP-even higgs mass bound. The second region with successful  $b$ - $\tau$  YU occurs for  $34 \lesssim \tan \beta \lesssim 45$ . Beyond this range we do not have REWSB if we require 10% or better  $b$ - $\tau$  YU. Imposing, in addition, the neutralino DM requirement (represented by blue points), the allowed region shrinks to  $35 \lesssim \tan \beta \lesssim 40$ .

The plot in  $R_{b\tau} - \mu$  plane shows that for 10% or better YU,  $\mu \gtrsim 3$  TeV (orange points). The need for large values of  $\mu$  can be understood by analyzing the threshold effects associated with the  $b$  quark. As previously mentioned, the second term in Eq. (3) has to be larger than the first one for successful  $b$ - $\tau$  YU. With  $\tan \beta$  squeezed in a relatively narrow interval, and the magnitude of  $A_t$  bounded to avoid color and charge breaking, needs up with a large  $\mu$  term to obtain the required  $\delta y_b$ . On the other hand, a bino-Higgsino mixed DM scenario requires  $m_{\tilde{\chi}_1^0} \approx |\mu|$ . From Fig. 2(c), the neutralino cannot be much heavier than 600 GeV or so for successful  $b$ - $\tau$  YU. But with  $\mu > 4$  TeV that a bino-Higgsino mixed DM scenario is not realized here.

Next let us consider the A-funnel scenario of CMSSM where one needs  $m_A \approx 2m_{\tilde{\chi}_1^0}$ . For large  $\tan \beta$ ,  $m_A^2 = 2b/\sin(2\beta) \approx b \tan \beta$ , where  $b$  is the soft SUSY breaking Higgs bilinear term. We have  $b \tan \beta \approx 4$  TeV for good  $b$ - $\tau$  unification, which implies that  $m_A \gg 2m_{\tilde{\chi}_1^0}$  and thus, the A-funnel scenario is not consistent with 10% or better  $b$ - $\tau$  unification.

The neutralino-stop co-annihilation channel is compatible with 10% or better  $b$ - $\tau$  unification, as shown by the blue points in Fig. 2. Let see how a light stop mass ( $m_{\tilde{t}_1}$ ) of  $O(100)$  GeV,




FIG. 3: Plots in  $m_{\tilde{t}_1}/m_{\tilde{t}_2} - A_0$  and  $m_{\tilde{t}_1}/m_{\tilde{t}_2} - A_t$  planes. The orange points satisfy the particle mass bounds, constraints from  $BR(B_s \rightarrow \mu^+ \mu^-)$ ,  $BR(B_u \rightarrow \tau \nu_\tau)$  and  $BR(b \rightarrow s \gamma)$ . Red color represents the solution with 10% or better  $b$ - $\tau$  YU. Blue color represents the neutralino-stop co-annihilation solutions.

	Point 1	Point 2	Point 3
$m_0$	15220	10040	17920
$M_{1/2}$	177	152	521
$\tan \beta$	37	39	37
$A_0/m_0$	-2.36	-2.32	-2.33
$sgn(\mu)$	+1	+1	+1
$m_h$	115	120	115
$m_H$	6036	4566	9752
$m_A$	5997	4537	9688
$m_{H^\pm}$	6037	4568	9753
$m_{\tilde{\chi}_{1,2}^0}$	124,272	97,209	290,592
$m_{\tilde{\chi}_{3,4}^0}$	10379,10379	6836,6836	12347,12347
$m_{\tilde{\chi}_{1,2}^\pm}$	275,10406	211,6840	598,1239
$m_{\tilde{g}}$	796	640	1680
$m_{\tilde{u}_{L,R}}$	15170,15214	10000,10030	17892,17942
$m_{\tilde{t}_{1,2}}$	153,5930	114,4076	328,7894
$m_{\tilde{d}_{L,R}}$	15170,15222	10000,10036	17892,17951
$m_{\tilde{b}_{1,2}}$	6060,8357	4152,5752	8097,11159
$m_{\tilde{\nu}_1}$	15223	10041	17929
$m_{\tilde{\nu}_3}$	12744	8453	15082
$m_{\tilde{e}_{L,R}}$	15211,15208	10032,10032	17911,17909
$m_{\tilde{\tau}_{1,2}}$	9843,12771	6619,8474	11801,15130
$\sigma_{SI}(\text{pb})$	$3.28 \times 10^{-12}$	$5.85 \times 10^{-12}$	$7.93 \times 10^{-13}$
$\sigma_{SD}(\text{pb})$	$3.90 \times 10^{-12}$	$2.39 \times 10^{-11}$	$1.77 \times 10^{-12}$
$\Omega_{CDM} h^2$	0.11	0.09	0.1
$R_{b\tau}$	1.00	1.02	1.09

TABLE I: Point 1 has perfect  $b$ - $\tau$  unification with  $R_{b\tau} = 1$ , point 2 represents a solution with relatively light stop mass ( $\sim 114$  GeV), and point 3 represents a solution with a heavier stop mass ( $\sim 330$  GeV). The remaining squarks and sleptons all have masses in the multi-TeV range.

is realized for  $m_0 \gtrsim 8$  TeV and  $M_{1/2} \lesssim 600$  GeV. In this region of the parameter space the diagonal entries of the stop mass matrix to pick up dominant contribution from  $y_t$  and  $A_t$  couplings, thus making the  $m_{\tilde{t}_R}$  entry lighter than its value at  $M_G$ . In Fig. 3(a) we see that the ratio  $m_{\tilde{t}_R}/m_0$  can be as small as 0.05 for large values of  $A_0$ , which means that at the SUSY scale  $m_{\tilde{t}_R}$  can be as light as 1 TeV or so, despite the large  $\sim (8 - 20)$  TeV  $m_0$  values. On the other hand, from Fig. 3(b) we see that at SUSY scale, the value of  $|A_t|$  can be  $O(7 - 20)$  TeV. The off-diagonal entries  $-m_t(A_t + \mu \cot \beta)$  for the stop quark mass matrix can therefore be of comparable magnitude to the diagonal entries. Because of this, as seen in Fig. 3(b), one of the eigenvalues ( $m_{\tilde{t}_1}$ ) of the stop quark mass matrix can be 10-20 times smaller than  $m_{\tilde{t}_R}$  at the SUSY scale. Thus, with  $m_{\tilde{t}_1} \sim O(100)$  GeV, we can have neutralino-stop co-annihilation compatible with  $b$ - $\tau$  YU. A similar discussion for the stau mass matrix shows that the stau co-annihilation scenario is not realized in YCMSSM.

In Table I we present three characteristic benchmark points which satisfy all, especially the dark matter, constraints. Point 1 displays essentially perfect  $b$ - $\tau$  YU solution with  $R_{b\tau} = 1$ , point 2 represents a solution with a relatively light stop mass ( $\sim 114$  GeV), and finally point 3 represents a solution with a heavier stop mass ( $\sim 330$  GeV). We note that the remaining

squarks and sleptons all have masses in the multi-TeV range.

Since the LSP is essentially a pure bino, both its spin-independent and spin-dependent cross sections on nucleons are rather small [21],  $\sim 10^{-47} - 10^{-48} \text{cm}^2$ . Consequently it won't be easy to detect the LSP in direct and indirect experiments.

In summary, we have investigated  $b$ - $\tau$  Yukawa unification in the mSUGRA/CMSSM framework and find that it is consistent with the NLSP stop scenario and yields the desired LSP neutralino relic abundance. This YCMSSM predicts that there are just two 'light' (LHC accessible) colored sparticles, namely the NLSP stop with mass  $\sim 100 - 330 \text{ GeV}$ , and the gluino which is  $\sim 600 - 1700 \text{ GeV}$ . The chargino and a second neutralino are about a factor 2-3 lighter than the gluino. The remaining squarks as well as all sleptons have masses in the multi-TeV range. Regarding the fundamental CMSSM parameters, we find that  $5 \text{ TeV} \lesssim m_0 \lesssim 20 \text{ TeV}$ ,  $m_0/M_{1/2} \approx 30 - 50$ ,  $\tan \beta \approx 35 - 40$ ,  $|\mu| \sim 3 - 15 \text{ TeV}$  and  $|A_0/m_0| \sim 2.2 - 2.4$ .

We thank Rizwan Khalid, Azar Mustafayev and Mansoor Ur Rehman for valuable discussions. This work is supported in part by the DOE Grant No. DE-FG02-91ER40626 (I.G., S.R. and Q.S.) and GNSF Grant No. 07\_462\_4-270 (I.G.).

---

\* On leave of absence from: Andronikashvili Institute of Physics, GAS, Tbilisi, Georgia.

† On study leave from: Department of Physics, FUUAST, Islamabad, Pakistan; E-mail: [shabbar@udel.edu](mailto:shabbar@udel.edu)

- [1] M. S. Chanowitz, J. Ellis and M. K. Gaillard, Nucl. Phys. B **128** (1977) 506; A. J. Buras, J. Ellis, M. K. Gaillard and D. V. Nanopoulos, Nucl. Phys. B **135** (1978) 66; C. Panagiotakopoulos and Q. Shafi, Phys. Rev. Lett. **52**, 2336 (1984).
- [2] B. Ananthanarayan, G. Lazarides and Q. Shafi, Phys. Rev. D **44**, 1613 (1991) and Phys. Lett. B **300**, 24 (1993); Q. Shafi and B. Ananthanarayan, Trieste HEP Cosmol. 1991:233-244.
- [3] U. Chattopadhyay and P. Nath, Phys. Rev. D **65**, 075009 (2002); S. Komine and M. Yamaguchi, Phys. Rev. D **65**, 075013 (2002); S. Profumo, Phys. Rev. D **68**, 015006 (2003); C. Pallis, Nucl. Phys. B **678**, 398 (2004); C. Balazs and R. Dermisek, JHEP **0306**, 024 (2003); W. Altmannshofer, D. Guadagnoli, S. Raby and D. M. Straub, Phys. Lett. B **668**, 385 (2008); I. Gogoladze, R. Khalid, N. Okada and Q. Shafi, Phys. Rev. D **79**, 095022 (2009); S. Antusch and M. Spinrath, Phys. Rev. D **79**, 095004 (2009).
- [4] K. Tobe and J. D. Wells, Nucl. Phys. B **663**, 123 (2003); M. Gomez, T. Ibrahim, P. Nath and S. Skadhauge, Phys. Rev. D **72**, 095008 (2005); D. Guadagnoli, S. Raby and D. M. Straub, JHEP **0910**, 059 (2009); H. Baer, S. Kraml, S. Sekmen and H. Summy, JHEP **0803**, 056 (2008); I. Gogoladze, R. Khalid and Q. Shafi, Phys. Rev. D **79**, 115004 (2009) [[arXiv:0903.5204](https://arxiv.org/abs/0903.5204) [hep-ph]]. K. Choi, D. Guadagnoli, S. H. Im and C. B. Park, JHEP **1010**, 025 (2010).
- [5] A. Chamseddine, R. Arnowitt and P. Nath, Phys. Rev. Lett. **49** (1982) 970; R. Barbieri, S. Ferrara and C. Savoy, Phys. Lett. **B119** (1982) 343; N. Ohta, Prog. Theor. Phys. **70** (1983) 542; L. J. Hall, J. D. Lykken and S. Weinberg, Phys. Rev. **D27** (1983) 2359; for a review see S. Weinberg, *The Quantum Theory of Fields: Volume 3, Supersymmetry*, Cambridge University Press (2000) 442p.
- [6] For a recent discussion and additional references see H. Baer, S. Kraml and S. Sekmen, JHEP **0909**, 005 (2009).
- [7] E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **180**, 330 (2009).
- [8] L. J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. D **50**, 7048 (1994).
- [9] H. Baer, F. E. Paige, S. D. Protopopescu and X. Tata, [arXiv:hep-ph/0001086](https://arxiv.org/abs/hep-ph/0001086).
- [10] J. Hisano, H. Murayama, and T. Yanagida, Nucl. Phys. **B402** (1993) 46; Y. Yamada, Z. Phys. **C60** (1993) 83; J. L. Chkareuli and I. G. Gogoladze, Phys. Rev. D **58**, 055011 (1998).
- [11] [Tevatron Electroweak Working Group and CDF Collaboration and D0 Collab], [arXiv:0903.2503](https://arxiv.org/abs/0903.2503) [hep-ex].
- [12] I. Gogoladze, R. Khalid, S. Raza and Q. Shafi, [arXiv:1102.0013](https://arxiv.org/abs/1102.0013) [hep-ph].
- [13] G. Belanger, F. Boudjema, A. Pukhov and R. K. Singh, JHEP **0911**, 026 (2009); H. Baer, S. Kraml, S. Sekmen and H. Summy, JHEP **0803**, 056 (2008).
- [14] K. Nakamura *et al.* [Particle Data Group Collaboration], J. Phys. G **G37**, 075021 (2010).
- [15] H. Baer, C. Balazs, and A. Belyaev, JHEP **03** (2002) 042; H. Baer, C. Balazs, J. Ferrandis, and X. Tata Phys. Rev. **D64** (2001) 035004.
- [16] S. Schael *et al.* Eur. Phys. J. C **47**, 547 (2006).
- [17] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **100**, 101802 (2008).
- [18] E. Barberio *et al.* [Heavy Flavor Averaging Group (HFAG) Collaboration], [arXiv:0704.3575](https://arxiv.org/abs/0704.3575) [hep-ex].
- [19] E. Barberio *et al.* [Heavy Flavor Averaging Group], [arXiv:0808.1297](https://arxiv.org/abs/0808.1297) [hep-ex].
- [20] D. M. Pierce, J. A. Bagger, K. T. Matchev, and R.-j. Zhang, Nucl. Phys. **B491** (1997) 3.
- [21] I. Gogoladze, R. Khalid, Y. Mimura and Q. Shafi, [arXiv:1012.1613](https://arxiv.org/abs/1012.1613) [hep-ph].