

A Mellin Space Program for W^\pm and Z^0 Production at NNLO

Petra Kovačiková *

Deutsches Elektronen-Synchrotron, Platanenallee 6, D-15738 Zeuthen, Germany

Key words QCD, perturbative expansion, factorisation, PDFs, Mellin transform

We present a program for the evaluation of full unpolarized cross sections for the W^\pm and Z^0 production in the narrow width approximation at NNLO in perturbative QCD using Mellin space techniques.

Copyright line will be provided by the publisher

1 Introduction

The Drell-Yan process, originally described in the context of the parton model [1], concerns the production of a lepton pair of large invariant mass in hadron-hadron collisions. With the increase of the centre of mass energy at particle accelerators, the Drell-Yan process led to the discovery of W^\pm and Z^0 bosons at UA1 and UA2 experiments [2, 3]. Since then the properties of massive vector bosons have been studied in great detail. At present the production of W^\pm and Z^0 provides an important benchmark for the LHC and a test of the Standard Model (SM) in a new range of centre of mass energies [4].

As guaranteed by the factorisation theorem [5], one can separate the physics of soft energy scales from the physics at hard energy scales where perturbation theory applies. The higher order QCD corrections to the Drell-Yan process have been calculated up to next-to-next-to-leading order (NNLO), see [6–8] and references therein. The full cross section is obtained as a convolution with the parton distribution functions (PDFs) that encode the non-perturbative information.

In this paper, we present a program for evaluation of the full inclusive cross section for W^\pm and Z^0 production in a fast and accurate way using a Mellin space approach. After a brief description of the basic ingredients of the calculation we give formulae for the Mellin transforms. We then present a comparison with the code ZWPROD [7, 8] and discuss possible applications and extensions within this framework.

2 Formalism

We consider the inclusive production of a single vector boson $V = W^+, W^-$ or Z^0 in hadron-hadron collision with a centre of mass energy s which subsequently decays into a lepton pair of an invariant mass Q^2 . The decay of the vector boson is treated within the narrow width approximation which replaces the propagator by a delta function such that $Q^2 = M_V^2$. We consider massless quarks. The cross section for this process can be expressed as

$$\sigma^{h_1 h_2 \rightarrow V \rightarrow l_1 l_2}(s) = x \sigma^{V \rightarrow l_1 l_2} W^V(x, Q^2), \quad x = Q^2/s, \quad (1)$$

where $\sigma^{V \rightarrow l_1 l_2}$ represents the kinematically independent part of the Born level subprocess $q\bar{q} \rightarrow V \rightarrow l_1 l_2$ (the point-like cross section) multiplied by the appropriate branching ratio. The exact form of the point like cross section can be found in Ref. [7], formulae (A.10) and (A.11).¹

* Corresponding author E-mail: petra.kovacikova@desy.de, Phone: +00 493 376 277 193

¹ There is an extra factor of 2 in the denominator of the formula (A.11), corrected by [9]

The structure function $W^V(x, Q^2)$ is written as a convolution of two parton distribution functions $f_a(\mu_f^2)$ and $f_b(\mu_f^2)$ and a hard scattering cross section represented by coefficient functions Δ_{ab} ,

$$W^V(x, Q^2) = \sum_{a,b=q,\bar{q},g} C_{a,b}^V \left[f_a(\mu_f^2) \otimes f_b(\mu_f^2) \otimes \Delta_{ab}(Q^2, \mu_f^2, \mu_r^2) \right](x). \quad (2)$$

The perturbative coefficients are known up to NNLO [7, 8],

$$\Delta_{ab}^{\text{NkLO}}(x, Q^2, \mu_f^2, \mu_r^2) = \sum_{n=0}^k \frac{\alpha_s^n(\mu_r^2)}{4\pi} \Delta_{ab}^{(k)}(x, Q^2, \mu_f^2, \mu_r^2). \quad (3)$$

The factor $C_{a,b}^V$ in Eq.(2) contains information about couplings of vector bosons to partons a and b . For the detailed form of the Eq.(2) we refer the reader to the paper of Hamberg, Matsuura and van Neerven [7] whose notation we follow closely². The convolution sign represents an integral

$$\begin{aligned} & (f_1 \otimes f_2 \otimes \dots \otimes f_k)(x) \\ &= \int_0^1 dx_1 \int_0^1 dx_2 \dots \int_0^1 dx_k \delta(x - x_1 x_2 \dots x_k) f_1(x_1) f_2(x_2) \dots f_k(x_k). \end{aligned} \quad (4)$$

In principle one can perform the integrals in Eq.(2) directly however, the problem is much better addressed after transforming to Mellin space,

$$f(N) = \int_0^1 dx x^{N-1} f(x), \quad (5)$$

This transformation turns the integrals in Eq.(2) into ordinary products such that the structure function reads

$$W^V(N, Q^2) = \sum_{a,b=q,\bar{q},g} C_{a,b}^V f_a(N, Q^2) f_b(N, Q^2) \Delta_{ab}(N, Q^2) \quad \mu_f = \mu_r = Q^2, \quad (6)$$

and therefore it is possible to evaluate it in a fast and efficient way. The formula for the inverse Mellin transform defines how to recover the original momentum space result,

$$W^V(x, Q^2) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN x^{-N} W^V(N, Q^2), \quad (7)$$

where c represents a point on the real axis such that all poles N_i in the function $W(N, Q^2)$ lie to the left from c . Further on, we will refer to functions in Mellin space as N space functions and functions from momentum space as x space functions.

3 Implementation

The main ingredients of the calculation are the coefficient functions up to NNLO and the parton distribution functions in Mellin space in terms of a complex variable N . The condition $N \in \mathbb{C}$ is required for the numerical evaluation of the inversion formula (7). For this we adopted the technique implemented in QCD-PEGASUS [10]. The complex integral (7) is rewritten in terms of an integral over a real variable z

$$W(x, Q^2) = \frac{1}{\pi} \int_0^\infty dz \text{Im}[e^{i\varphi} x^{-c-z e^{i\varphi}} W(N, Q^2)] \quad N = c + z \exp^{i\phi} \in \mathbb{C} \quad (8)$$

and evaluated using Gaussian quadratures. The parameter $\phi > \pi/2$ represents the angle with respect to the positive real axis. Since the rightmost pole of the structure function is $N_{\text{max}} = 1$, we chose $c = 1.5$. These

² Several typos appearing in Ref. [7] have been pointed out in [9]

values as well as the maximum value of the integration variable z are flexible and can be modified by the user in the main program if desired. For a more detailed description of the shape of the integration contour we refer to the QCD-PEGASUS manual [10].

The coefficient functions in N space were published previously in Ref. [11], including corrections to the previous literature. The corresponding FORTRAN code is `DY.f` used together with `ANCONT` [12]. We performed the Mellin transforms starting from the x space expressions [13] using the `harmol` package [14]. The results can be expressed mostly in terms of complex-valued simple harmonic sums [15, 16] and several more complicated ones which we approximated by using the minimax method³ worked out in detail in [12] previously.⁴ The absolute accuracy of our approximation is better than 10^{-9} over the whole kinematic range.

At the moment there are two options for the input parton distribution functions in N space. A toy input corresponds to the one used for the 2001/2 benchmark tables [18] and is used for comparisons with ZWPROD [7, 8] assuming no evolution of PDFs. The general form reads

$$x f_{i,\text{toy}}(x, \mu_0^2) = n x^a (1-x)^b, \quad i = q, \bar{q}, g, \quad n, a, b \in \mathbb{R}, \quad (9)$$

which is in Mellin space represented by an Euler beta function

$$f_{i,\text{toy}}(N, \mu_0^2) = n \beta(a + N, b + 1). \quad (10)$$

The second option for the PDF input is using the FORTRAN code QCD-PEGASUS [10] which can be linked to our program.




Fig. 1 Cross section for W^- production up to NNLO in the narrow width approximation using the toy parton distribution functions and a fixed value of the strong coupling constant. Upper part: The full cross section. Lower part: Relative accuracy with respect to the ZWPROD.

4 Results And Outlook

There are several programs on the market using the standard momentum space evaluation [19–21] which can provide a cross-check for our N space calculation. We performed comparisons of the full cross sections with a program ZWPROD written by the authors of the original calculation of the NNLO Drell-Yan

³ We used the MINIMAX routine implemented in Maple

⁴ Exact expressions were given in [17].

coefficient functions [7, 8]. The Fig. (1) shows a comparison for the W^- cross section using toy input for PDFs corresponding to the Eq.(10) with no evolution and a fixed value of the coupling constant $\alpha_s = 0.1$. The relative accuracy is better than 6×10^{-6} in the relevant kinematical range $x \in (10^{-4}, 0.8)$. As an intermediate check, we compared the Mellin inversion of N space coefficient functions against the x space expressions using a program of Gehrmann and Remiddi [22] for the numerical evaluation of harmonic polylogarithms. The framework presented here is suitable for a further implementation of those cross sections where N space coefficient functions are also available, like Higgs production and deep inelastic scattering (DIS) [11, 23–26]. The setup is well suited for merging the program with threshold resummation calculations which are typically performed in Mellin space (see e.g. [27]). For the extraction of PDFs from W^\pm and Z production it would be desirable to have an access to the rapidity distributions in which case one will need to apply double Mellin transforms of two variables N_1 and N_2 however, this is a subject to further study. On the side of PDFs we aim for a direct interface to the LHAPDF grids [28] which will allow the user to freely choose any particular PDF set provided within this framework. Recent results [29] on N space input parametrizations also allow for more flexible input PDF parametrisations in QCD-PEGASUS. Further improvements with respect to the speed of the code are foreseen and together with an upgrade on the input PDFs this code can become a tool for PDF fits, where fast and accurate evaluations of cross sections are needed. The current version of the c++ code can be downloaded from <http://www-zeuthen.desy.de/~kpetra/sbp>.

Acknowledgements I would like to thank my advisor Sven-Olaf Moch for continuous support and advice throughout the course of this work and for careful reading of the manuscript. I am grateful to Prof. Johannes Blümlein for useful discussions and comments. This work was supported by the Marie-Curie Research Training Networks MRTN-CT-2006-035505 HEPTOOLS.

References

- [1] S. D. Drell and T. M. Yan, Phys. Rev. Lett. **25**, 316–320 (1970).
- [2] C. Albajar et al., Phys.Lett. **B198**, 271 (1987).
- [3] J. Alitti et al., Phys.Lett. **B276**, 365–374 (1992).
- [4] S. Alekhin, J. Blümlein, P. Jimenez-Delgado, S. Moch, and E. Reya, Phys.Lett. **B697**, 127–135 (2011).
- [5] J. C. Collins, D. E. Soper, and G. F. Sterman, Nucl.Phys. **B261**, 104 (1985).
- [6] G. Altarelli, R. Ellis, and G. Martinelli, Nucl.Phys. **B157**, 461 (1979).
- [7] R. Hamberg, W. van Neerven, and T. Matsuura, Nucl.Phys. **B359**, 343–405 (1991).
- [8] R. Hamberg, W. L. van Neerven, and T. Matsuura, Nuclear Physics B **644**(1-2), 403 – 404 (2002).
- [9] C. Anastasiou, L. J. Dixon, K. Melnikov, and F. Petriello, Phys.Rev. **D69**, 094008 (2004).
- [10] A. Vogt, Comput.Phys.Comm. **170**, 65–92 (2005).
- [11] J. Blümlein and V. Ravindran, Nucl.Phys. **B716**, 128–172 (2005).
- [12] J. Blümlein, Comput.Phys.Comm. **133**, 76–104 (2000).
- [13] V. Ravindran, unpublished.
- [14] E. Remiddi and J. Vermaseren, Int.J.Mod.Phys. **A15**, 725–754 (2000).
- [15] J. Vermaseren, Int.J.Mod.Phys. **A14**, 2037–2076 (1999).
- [16] J. Blümlein and S. Kurth, Phys.Rev. **D60**, 014018 (1999).
- [17] J. Blümlein, Comput.Phys.Comm. **180**, 2218–2249 (2009).
- [18] W. Giele, E. Glover, I. Hinchliffe, J. Huston, E. Laenen et al., arXiv:hep-ph/0204316 pp. 275–426 (2002).
- [19] S. Catani and M. Grazzini, Phys.Rev.Lett. **98**, 222002 (2007).
- [20] S. Catani, L. Cieri, G. Ferrera, D. de Florian, and M. Grazzini, Phys.Rev.Lett. **103**, 082001 (2009).
- [21] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, arXiv:1011.3540[hep-ph] (2010).
- [22] T. Gehrmann and E. Remiddi, Comput.Phys.Comm. **141**, 296–312 (2001).
- [23] W. van Neerven and A. Vogt, Nucl.Phys. **B588**, 345–373 (2000).
- [24] A. Vogt, S. Moch, and J. Vermaseren, Nucl.Phys.Proc.Suppl. **160**, 44–50 (2006).
- [25] S. Moch, J. Vermaseren, and A. Vogt, Phys.Lett. **B606**, 123–129 (2005).
- [26] J. Vermaseren, A. Vogt, and S. Moch, Nucl.Phys. **B724**, 3–182 (2005).
- [27] S. Moch, J. Vermaseren, and A. Vogt, Nucl.Phys. **B726**, 317–335 (2005).
- [28] M. R. Whalley, D. Bourilkov, and R. C. Group, arXiv:hep-ph/0508110 (2005), <http://hepforge.cedar.ac.uk/lhapdf/>.
- [29] J. Blümlein, A. Hasselhuhn, P. Kovacikova, and S. Moch, arXiv:1104.3449v1.