

Three-dimensional parton distribution functions g_{1T} and h_{1L}^\perp in the polarized proton-antiproton Drell-Yan process

Jiacai Zhu¹ and Bo-Qiang Ma^{1,2,a}

¹ School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China

² Center for High Energy Physics, Peking University, Beijing 100871, China

Received: date / Revised version: date

Abstract. We present predictions of the unweighted and weighted double spin asymmetries related to the transversal helicity distribution g_{1T} and the longitudinal transversity distribution h_{1L}^\perp , two of eight leading-twist transverse momentum dependent parton distributions (TMDs) or three-dimensional parton distribution functions (3dPDFs), in the polarized proton-antiproton Drell-Yan process at typical kinematics on the Facility for Antiproton and Ion Research (FAIR). We conclude that FAIR is ideal to access the new 3dPDFs towards a detailed picture of the nucleon structure.

1 Introduction

The nucleon spin structure is an active direction under both theoretical and experimental investigations. A number of new physical quantities of the nucleon, which can provide a detailed picture of the nucleon with more detailed information in three dimensional momentum space [1, 2], have been introduced. At leading twist, the quark-quark correlation matrix [3, 4] can be decomposed into eight transverse momentum dependent parton distributions (TMDs), or we call them three-dimensional parton distribution functions (3dPDFs). Besides the three usual parton distribution functions, *i.e.*, the unpolarized one with light-cone longitudinal momentum and transverse momentum distribution f_1 , the helicity one with longitudinal helicity distribution g_{1L} , and the transversity one with transversal spin distributions h_1 , there are five new ones. Among them the Sivers distribution f_{1T}^\perp and its chiral-odd partner, the Boer-Mulders distribution h_1^\perp , are well known for their T-odd property, *i.e.*, they change sign under naive time reversal. The other three of the new 3dPDFs, *i.e.*, the pretzelosity distribution h_{1T}^\perp , the transversal helicity distribution g_{1T} , and the longitudinal transversity distribution h_{1L}^\perp , are T-even. They can be measured through the semi-inclusive deep inelastic scattering [5, 6]. However, the single spin asymmetry related to the pretzelosity distribution h_{1T}^\perp in the semi-inclusive deep inelastic scattering is rather small. For the chiral-odd distributions h_{1T}^\perp and h_{1L}^\perp , they can be probed through the single spin asymmetries when combined with another chiral-odd distribution h_1^\perp in the pion-nucleon Drell-Yan process [7, 8]. In Ref. [9], it has been shown that the polarized proton-antiproton Drell-Yan process is ideal to probe the pretzelosity distribution

h_{1T}^\perp and the magnitudes of the related spin asymmetries are significantly large. Thus it is natural to discuss the spin asymmetries related to the rest two 3dPDFs, g_{1T} and h_{1L}^\perp , in the polarized proton-antiproton Drell-Yan process.

Since the proposal on measuring the transversity distributions via the polarized proton-antiproton Drell-Yan process by the polarized antiproton experiment (PAX) collaboration [10, 11], there have been some other experiments [12, 13, 14, 15, 16, 17, 18] on the measurements of the transversity distributions. Recently, there has been new technical progress [19] towards the goal for a proton-antiproton collider with both beams polarized [20], and such plan has the potential to be realized at FAIR (Facility for Antiproton and Ion Research) in GSI Helmholtzzentrum für Schwerionenforschung. The expected antiproton beam polarizations might be $0.15 \sim 0.20$ (spin filtering with transverse target orientation) or $0.35 \sim 0.40$ (longitudinal) [21, 22]. Thus the three new 3dPDFs, h_{1T}^\perp , g_{1T} , and h_{1L}^\perp , with important information on the quark spin and orbital correlation of the nucleon, could be measured on FAIR. Our calculation below shows that some of the double spin asymmetries related to g_{1T} and h_{1L}^\perp in the polarized proton-antiproton Drell-Yan process are large, and thus FAIR is an ideal facility to access the new 3dPDFs for revealing more information towards a detailed picture of the nucleon.

2 T-even 3dPDFs in the light-cone quark-diquark model

In the light-cone quark-diquark model [23], the Melosh-Wigner rotation plays an important role to understand the proton spin puzzle [24, 25] due to the relativistic effect of quark transversal motions. The T-even 3dPDFs have

^a Corresponding author. Email address: mabq@pku.edu.cn

been calculated [5, 6, 23, 26, 27, 28]:

$$\begin{aligned} f_1^{(uv)}(x, k_T^2) &= \frac{1}{16\pi^3} \left(\frac{1}{3} \sin^2 \theta_0 \varphi_V^2 + \cos^2 \theta_0 \varphi_S^2 \right), \\ f_1^{(dv)}(x, k_T^2) &= \frac{1}{8\pi^3} \frac{1}{3} \sin^2 \theta_0 \varphi_V^2, \end{aligned} \quad (1)$$

and

$$\begin{aligned} j^{(uv)}(x, k_T^2) &= -\frac{1}{16\pi^3} \times \left(\frac{1}{9} \sin^2 \theta_0 \varphi_V^2 W_V^j - \cos^2 \theta_0 \varphi_S^2 W_S^j \right), \\ j^{(dv)}(x, k_T^2) &= -\frac{1}{8\pi^3} \times \frac{1}{9} \sin^2 \theta_0 \varphi_V^2 W_V^j, \end{aligned} \quad (2)$$

with the notation

$$j = g_{1L}, g_{1T}, h_1, h_{1T}^\perp, h_{1L}^\perp,$$

and the superscripts “ uv ” and “ dv ” stand for the valence up and down quark distributions respectively. φ_D ($D = V, S$) is the wave function in the momentum space for the quark-diquark, and for which we can use the Brodsky-Huang-Lepage (BHL) prescription [29, 30]:

$$\varphi_D(x, k_T^2) = A_D \exp \left\{ -\frac{1}{8\alpha_D^2} \left[\frac{m_q^2 + k_T^2}{x} + \frac{m_D^2 + k_T^2}{1-x} \right] \right\}. \quad (3)$$

The parameters $\alpha_D = 0.33$ GeV (which is the same for $D = V, S$), the quark mass $m_q = 0.33$ GeV, the diquark mass $m_S = 0.60$ GeV, $m_V = 0.80$ GeV, and $\theta_0 = \pi/4$ are adopted for numerical calculation. θ_0 is the mixing angle that breaks the SU(6) symmetry when $\theta_0 \neq \pi/4$. The Melosh-Wigner rotation factors W_D ($D = V, S$) are

$$W_D^{g_{1L}}(x, k_T^2) = \frac{(x\mathcal{M}_D + m_q)^2 - k_T^2}{(x\mathcal{M}_D + m_q)^2 + k_T^2}, \quad (4)$$

$$W_D^{g_{1T}}(x, k_T^2) = \frac{2M_N(x\mathcal{M}_D + m_q)}{(x\mathcal{M}_D + m_q)^2 + k_T^2}, \quad (5)$$

$$W_D^{h_1}(x, k_T^2) = \frac{(x\mathcal{M}_D + m_q)^2}{(x\mathcal{M}_D + m_q)^2 + k_T^2}, \quad (6)$$

$$W_D^{h_{1T}^\perp}(x, k_T^2) = -\frac{2M_N^2}{(x\mathcal{M}_D + m_q)^2 + k_T^2}, \quad (7)$$

$$W_D^{h_{1L}^\perp}(x, k_T^2) = -\frac{2M_N(x\mathcal{M}_D + m_q)}{(x\mathcal{M}_D + m_q)^2 + k_T^2}, \quad (8)$$

where

$$\mathcal{M}_D = \sqrt{\frac{m_q^2 + k_T^2}{x} + \frac{m_D^2 + k_T^2}{1-x}}. \quad (9)$$

Using Eqs. (1) and (2), the polarized distributions can be given by the unpolarized distributions as

$$\begin{aligned} j^{(uv)}(x, k_T^2) &= [f_1^{(uv)}(x, k_T^2) - \frac{1}{2} f_1^{(dv)}(x, k_T^2)] W_S^j(x, k_T^2) \\ &\quad - \frac{1}{6} f_1^{(dv)}(x, k_T^2) W_V^j(x, k_T^2), \\ j^{(dv)}(x, k_T^2) &= -\frac{1}{3} f_1^{(dv)}(x, k_T^2) W_V^j(x, k_T^2). \end{aligned} \quad (10)$$

3 The g_{1T} and h_{1L}^\perp related asymmetries in the polarized proton-antiproton Drell-Yan process

In the polarized proton-antiproton Drell-Yan process, the cross section is [31]

$$\begin{aligned} \frac{d\sigma}{dx_a dx_b d\mathbf{q}_T d\Omega} &= \frac{\alpha_{em}^2}{4Q^2} \left\{ (1 + \cos^2 \theta) F_{UU}^1 + S_{aL} S_{bL} \sin^2 \theta \cos 2\phi F_{LL}^{\cos 2\phi} \right. \\ &\quad + |S_{aT}| S_{bL} (1 + \cos^2 \theta) \cos \phi_a F_{TL}^{\cos \phi_a} \\ &\quad + S_{aL} |S_{bT}| \sin^2 \theta [\cos(2\phi + \phi_b) F_{LT}^{\cos(2\phi + \phi_b)} \\ &\quad + \cos(2\phi - \phi_b) F_{LT}^{\cos(2\phi - \phi_b)}] \\ &\quad + |S_{aT}| |S_{bT}| (1 + \cos^2 \theta) [(\cos(\phi_a + \phi_b) F_{TT}^{\cos(\phi_a + \phi_b)} \\ &\quad + \cos(\phi_a - \phi_b) F_{TT}^{\cos(\phi_a - \phi_b)})] + \dots \left. \right\}. \end{aligned} \quad (11)$$

The subscripts a and b stand for the incoming hadrons in the Drell-Yan process, and ϕ_a and ϕ_b are the angles of \mathbf{S}_{aT} and \mathbf{S}_{bT} respectively. Other terms will not contribute in our analysis below and we should note that although some of these terms involve g_{1T} or h_{1L} , they can be obtained from the terms above by switching the subscript labels a and b . The structure functions are

$$F_{UU}^1 = C[f_1 \bar{f}_1], \quad (12)$$

$$F_{LL}^{\cos 2\phi} = C \left[\frac{2(\mathbf{h} \cdot \mathbf{k}_{aT})(\mathbf{h} \cdot \mathbf{k}_{bT}) - \mathbf{k}_{aT} \cdot \mathbf{k}_{bT}}{M_N^2} h_{1L}^\perp \bar{h}_{1L}^\perp \right], \quad (13)$$

$$F_{TL}^{\cos \phi_a} = C \left[-\frac{\mathbf{h} \cdot \mathbf{k}_{aT}}{M_N} g_{1T} \bar{g}_{1L} \right], \quad (14)$$

$$F_{LT}^{\cos(2\phi - \phi_b)} = C \left[\frac{\mathbf{h} \cdot \mathbf{k}_{aT}}{M_N} h_{1L}^\perp \bar{h}_1 \right], \quad (15)$$

$$\begin{aligned} F_{LT}^{\cos(2\phi + \phi_b)} &= C \left[\frac{2(\mathbf{h} \cdot \mathbf{k}_{bT})[2(\mathbf{h} \cdot \mathbf{k}_{aT})(\mathbf{h} \cdot \mathbf{k}_{bT}) - \mathbf{k}_{aT} \cdot \mathbf{k}_{bT}] - k_{bT}^2(\mathbf{h} \cdot \mathbf{k}_{aT})}{2M_N^3} \right. \\ &\quad \left. h_{1L}^\perp \bar{h}_{1T}^\perp \right], \end{aligned} \quad (16)$$

$$F_{TT}^{\cos(\phi_a + \phi_b)} = C \left[\frac{2(\mathbf{h} \cdot \mathbf{k}_{aT})(\mathbf{h} \cdot \mathbf{k}_{bT}) - \mathbf{k}_{aT} \cdot \mathbf{k}_{bT}}{2M_N^2} (f_{1T}^\perp \bar{f}_{1T}^\perp - g_{1T} \bar{g}_{1T}) \right], \quad (17)$$

$$F_{TT}^{\cos(\phi_a - \phi_b)} = C \left[-\frac{\mathbf{k}_{aT} \cdot \mathbf{k}_{bT}}{2M_N^2} (f_{1T}^\perp \bar{f}_{1T}^\perp + g_{1T} \bar{g}_{1T}) \right], \quad (18)$$

where we use the shorthand notation

$$\begin{aligned} &\mathcal{C}[w(\mathbf{k}_{aT}, \mathbf{k}_{bT}) f_a \bar{f}_b] \\ &= \frac{1}{N_c} \sum_q e_q^2 \int d\mathbf{k}_{aT} d\mathbf{k}_{bT} \delta^{(2)}(\mathbf{q}_T - \mathbf{k}_{aT} - \mathbf{k}_{bT}) \\ &\quad \times w(\mathbf{k}_{aT}, \mathbf{k}_{bT}) f_a^q(x_a, k_{aT}^2) f_b^q(x_b, k_{bT}^2), \end{aligned} \quad (19)$$




Fig. 1. The double spin asymmetries related to h_{1L}^\perp shown in Eqs. (30) and (34) as functions of x_F for $s = 45 \text{ GeV}^2$ and $Q^2 = 12 \text{ GeV}^2$. The dashed curves correspond to approach 1, while the solid curves correspond to approach 2.

where both quarks and antiquarks of all flavors are taken into account during the summation over q . The unit vector is defined as $\mathbf{h} \equiv \mathbf{q}_T/q_T$.

Considering the charge conjugation invariance

$$f_p^q(x, k_T^2) = f_{\bar{p}}^{\bar{q}}(x, k_T^2), \quad (20)$$

$$f_{\bar{p}}^{\bar{q}}(x, k_T^2) = f_p^q(x, k_T^2), \quad (21)$$

with p for proton and \bar{p} for antiproton, and using the method introduced in Refs. [32,33], we get

$$\int d\mathbf{q}_T F_{UU}^1 = \frac{1}{N_c} \sum_q e_q^2 f_1^q(x_a) f_1^q(x_b), \quad (22)$$

$$\int d\mathbf{q}_T \frac{q_T^2}{8M_N^2} F_{LL}^{\cos 2\phi} = \frac{1}{N_c} \sum_q e_q^2 h_{1L}^{\perp(1)q}(x_a) h_{1L}^{\perp(1)q}(x_b), \quad (23)$$

$$\int d\mathbf{q}_T \frac{q_T}{2M_N} F_{TL}^{\cos \phi_a} = -\frac{1}{N_c} \sum_q e_q^2 g_{1T}^{(1)q}(x_a) g_{1L}^q(x_b), \quad (24)$$

$$\int d\mathbf{q}_T \frac{q_T}{2M_N} F_{LT}^{\cos(2\phi-\phi_b)} = \frac{1}{N_c} \sum_q e_q^2 h_{1L}^{\perp(1)q}(x_a) h_{1L}^q(x_b), \quad (25)$$

$$\int d\mathbf{q}_T \frac{q_T^3}{12M_N^3} F_{LT}^{\cos(2\phi+\phi_b)} = \frac{1}{N_c} \sum_q e_q^2 h_{1L}^{\perp(1)q}(x_a) h_{1T}^{\perp(2)q}(x_b), \quad (26)$$

$$\begin{aligned} & \int d\mathbf{q}_T \frac{q_T^2}{4M_N^2} F_{TT}^{\cos(\phi_a+\phi_b)} \\ &= \frac{1}{N_c} \sum_q e_q^2 [f_{1T}^{\perp(1)q}(x_a) f_{1T}^{\perp(1)q}(x_b) - g_{1T}^{(1)q}(x_a) g_{1T}^{(1)q}(x_b)], \end{aligned} \quad (27)$$

$$\begin{aligned} & \int d\mathbf{q}_T \frac{q_T^2}{2M_N^2} F_{TT}^{\cos(\phi_a-\phi_b)} \\ &= -\frac{1}{N_c} \sum_q e_q^2 [f_{1T}^{\perp(1)q}(x_a) f_{1T}^{\perp(1)q}(x_b) + g_{1T}^{(1)q}(x_a) g_{1T}^{(1)q}(x_b)], \end{aligned} \quad (28)$$

with

$$j^{(n)}(x) \equiv \int d\mathbf{k}_T \left(\frac{k_T^2}{2M_N^2} \right)^n j(x, k_T^2), \quad (29)$$

for 3dPDF j . The unweighted double spin asymmetries are (Q^2 is fixed)

$$\begin{aligned} A_{LL}^{\cos 2\phi}(x_F) &= \frac{\int d\mathbf{q}_T F_{LL}^{\cos 2\phi}(x_a, x_b, \mathbf{q}_T)}{\int d\mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} \\ &= \frac{N_c \int d\mathbf{q}_T F_{LL}^{\cos 2\phi}(x_a, x_b, \mathbf{q}_T)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}, \end{aligned} \quad (30)$$

$$\begin{aligned} A_{TL}^{\cos \phi_a}(x_F) &= \frac{\int d\mathbf{q}_T F_{TL}^{\cos \phi_a}(x_a, x_b, \mathbf{q}_T)}{\int d\mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} \\ &= \frac{N_c \int d\mathbf{q}_T F_{TL}^{\cos \phi_a}(x_a, x_b, \mathbf{q}_T)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}, \end{aligned} \quad (31)$$

$$\begin{aligned} A_{LT}^{\cos(2\phi-\phi_b)}(x_F) &= \frac{\int d\mathbf{q}_T F_{LT}^{\cos(2\phi-\phi_b)}(x_a, x_b, \mathbf{q}_T)}{\int d\mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} \\ &= \frac{N_c \int d\mathbf{q}_T F_{LT}^{\cos(2\phi-\phi_b)}(x_a, x_b, \mathbf{q}_T)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}, \end{aligned} \quad (32)$$

$$\begin{aligned} A_{LT}^{\cos(2\phi+\phi_b)}(x_F) &= \frac{\int d\mathbf{q}_T F_{LT}^{\cos(2\phi+\phi_b)}(x_a, x_b, \mathbf{q}_T)}{\int d\mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} \\ &= \frac{N_c \int d\mathbf{q}_T F_{LT}^{\cos(2\phi+\phi_b)}(x_a, x_b, \mathbf{q}_T)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}. \end{aligned} \quad (33)$$

(27) The weighted double spin asymmetries are (Q^2 is fixed)

$$\begin{aligned} A_{LL}^{\frac{W^2}{8} \cos 2\phi}(x_F) &= \frac{\int d\mathbf{q}_T \frac{W^2}{8} F_{LL}^{\cos 2\phi}(x_a, x_b, \mathbf{q}_T)}{\int d\mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} \\ &= \frac{\sum_q e_q^2 h_{1L}^{\perp(1)q}(x_a) h_{1L}^{\perp(1)q}(x_b)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}, \end{aligned} \quad (34)$$




Fig. 2. The double spin asymmetries related to g_{1T} and g_{1L} shown in Eqs. (31) and (35) as functions of x_F for $s = 45 \text{ GeV}^2$ and $Q^2 = 12 \text{ GeV}^2$. The dashed curves correspond to approach 1, while the solid curves correspond to approach 2.

$$A_{TL}^{\frac{W}{2} \cos \phi_a}(x_F) = \frac{\int d\mathbf{q}_T \frac{W}{2} F_{TL}^{\cos \phi_a}(x_a, x_b, \mathbf{q}_T)}{\int d\mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} = -\frac{\sum_q e_q^2 g_{1T}^{(1)q}(x_a) g_{1L}^q(x_b)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}, \quad (35)$$

$$A_{LT}^{\frac{W}{2} \cos(2\phi - \phi_b)}(x_F) = \frac{\int \mathbf{q}_T \frac{W}{2} F_{LT}^{\cos(2\phi - \phi_b)}(x_a, x_b, \mathbf{q}_T)}{\int \mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} = \frac{\sum_q e_q^2 h_{1L}^{\perp(1)q}(x_a) h_1^q(x_b)}{\sum_q e_q^2 [f_1^q(x_a) f_1^q(x_b)]}, \quad (36)$$

$$A_{LT}^{\frac{W^3}{12} \cos(2\phi + \phi_b)}(x_F) = \frac{\int \mathbf{q}_T \frac{W^3}{12} F_{LT}^{\cos(2\phi + \phi_b)}(x_a, x_b, \mathbf{q}_T)}{\int \mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} = \frac{\sum_q e_q^2 h_{1L}^{\perp(1)q}(x_a) h_{1T}^{\perp(2)q}(x_b)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}, \quad (37)$$

$$A_{TT}^g(x_F) = -\frac{\int \mathbf{q}_T \frac{W^2}{8} F_{TT}^{\cos(\phi_a + \phi_b)}(x_a, x_b, \mathbf{q}_T)}{\int \mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} - \frac{\int \mathbf{q}_T \frac{W^2}{4} F_{TT}^{\cos(\phi_a - \phi_b)}(x_a, x_b, \mathbf{q}_T)}{\int \mathbf{q}_T F_{UU}^1(x_a, x_b, \mathbf{q}_T)} = \frac{\sum_q e_q^2 g_{1T}^{(1)q}(x_a) g_{1T}^{(1)q}(x_b)}{\sum_q e_q^2 f_1^q(x_a) f_1^q(x_b)}, \quad (38)$$

where $W = q_T/M_N$. In Eq. (38), the weight function is complicated and we use “ g ” to denote it in Fig. 5. x_a and x_b are given by

$$x_F = x_a - x_b, \quad x_a x_b = \frac{Q^2}{s}. \quad (39)$$

In Ref. [9], the h_{1T}^\perp related double spin asymmetries in the polarized proton-antiproton Drell-Yan process have been calculated at typical kinematics on FAIR [10]. In

this paper, we focus on predictions of the unweighted and weighted double spin asymmetries related to g_{1T} and h_{1L}^\perp in the same process. We present numerical calculations in two different approaches as described in Ref. [9]:

- *Approach 1.* We use Eqs. (1) and (2) directly to calculate and only sum over the valence quark distributions.
- *Approach 2.* For the unpolarized quark and antiquark distributions, we use the CTEQ6L parametrization [34], and adopt a Gaussian form factor for the transverse momentum dependence which has been adopted in many phenomenological analysis [35]:

$$f_1(x, k_T^2) = f_1(x) \frac{\exp(-k_T^2/k_{un}^2)}{\pi k_{un}^2} \quad (40)$$

with $k_{un}^2 = 0.25 \text{ GeV}^2$. For the polarized distributions, we keep the relations in Eqs. (10) which we get in the light-cone quark-diquark model so that the Melosh-Wigner rotation factors remain as the relativistic effect of quark transversal motions.

The kinematics on FAIR are chosen as $s = 45 \text{ GeV}^2$ and $Q^2 = 12 \text{ GeV}^2$ [10]. The magnitudes of the helicity and transversity distributions in approach 2 are comparable with the global analysis results [36, 37, 38, 39, 40] for helicity and [35] for transversity at the middle x region. Besides, the quark-diquark model gives a good description of the nucleon form-factors [41, 42]. The quark-diquark model realized in approach 2 also provides reasonable descriptions of many experiments related to helicity distributions [43], transversity distributions [44, 18], together with some new 3dPDFs [45]. The effect of the CTEQ6L parametrization, which has been well verified and constrained by many experiments concerning the unpolarized quark distributions, has been taken into account in approach 2. Thus approach 2 might give more reasonable predictions for future experiments.

We plot the unweighted and weighted asymmetries related to g_{1T} and h_{1L}^\perp as show in Eqs. (30), (31), (32), (33), (34), (35), (36), (37), and (38). The results are shown in Figs. 1, 2, 3, 4, and 5. In the light-cone quark-diquark model, the ratios of $h_{1L}^{\perp(1/2)}(x)/f_1(x)$ as shown in Ref. [6]




Fig. 3. The double spin asymmetries related to h_{1L}^\perp and h_1 shown in Eqs. (32) and (36) as functions of x_F for $s = 45 \text{ GeV}^2$ and $Q^2 = 12 \text{ GeV}^2$. The dashed curves correspond to approach 1, while the solid curves correspond to approach 2.




Fig. 4. The double spin asymmetries related to h_{1L}^\perp and h_{1T}^\perp shown in Eqs. (33) and (37) as functions of x_F for $s = 45 \text{ GeV}^2$ and $Q^2 = 12 \text{ GeV}^2$. The dashed curves correspond to approach 1, while the solid curves correspond to approach 2.




Fig. 5. The double spin asymmetry related to g_{1T} shown in Eq. (38) as a function of x_F for $s = 45 \text{ GeV}^2$ and $Q^2 = 12 \text{ GeV}^2$. The dashed curve corresponds to approach 1, while the solid curve corresponds to approach 2.

are larger compared with those of $h_{1T}^{\perp(1)}(x)/f_1(x)$ as shown in Ref. [5], so the unweighted double spin asymmetry related to h_{1L}^\perp and h_1 in Fig. 3 is correspondingly larger compared with the unweighted double spin asymmetry re-

lated to h_{1T}^\perp and h_1 that we obtained in Ref. [9]. The unweighted asymmetries are larger compared with the corresponding weighted ones. In Figs. 1, 2 and 3, the unweighted asymmetries are significantly large. Besides, the unique feature of the unweighted double spin asymmetry $A_{LL}^{\cos 2\phi}$ in Eq. (30) as shown in Fig. 1 is that only the new 3dPDF h_{1L}^\perp is involved. Thus it is ideal to measure the new 3dPDFs g_{1T} and h_{1L}^\perp through the double spin asymmetries in the polarized proton-antiproton Drell-Yan process. Moreover, the unweighted double spin asymmetry related to h_{1L}^\perp and h_{1T}^\perp in Fig. 4 is about several percent. This provides us a way to measure the pretzelosity distributions h_{1T}^\perp in the proton-antiproton Drell-Yan process with one nucleon longitudinal polarized and another one transversal polarized, which is different from the unweighted double spin asymmetry related to h_{1L}^\perp and h_1 in the proton-antiproton Drell-Yan process with both nucleons transversal polarized in Ref. [9].

4 Summary

g_{1T} and h_{1L}^\perp , *i.e.*, the transversal helicity and the longitudinal transversity, are two of the eight leading-twist

3-dimensional parton distributions (3dPDFs). We present predictions of the unweighted and weighted double spin asymmetries related to them in the polarized proton-antiproton Drell-Yan process at typical kinematics on FAIR respectively. We conclude that the Facility for Antiproton and Ion Research (FAIR) is ideal to access the new 3dPDFs towards a detailed picture of the nucleon structure.

This work is supported by National Natural Science Foundation of China (Grants No. 11021092, No. 10975003, No. 11035003, and No. 11120101004).

References

1. V. Barone, A. Drago, P. G. Ratcliffe, Phys. Rept. **359**, 1 (2002)
2. V. Barone, F. Bradamante, A. Martin, Prog. Part. Nucl. Phys. **65**, 267 (2010)
3. P. J. Mulders, R. D. Tangerman, Nucl. Phys. B **461**, 197 (1996)
4. D. Boer, P. J. Mulders, Phys. Rev. D **57**, 5780 (1998)
5. J. She, J. Zhu, B.-Q. Ma, Phys. Rev. D **79**, 054008 (2009)
6. J. Zhu, B.-Q. Ma, Phys. Lett. B **696**, 246 (2011)
7. Z. Lu, B.-Q. Ma, J. She, Phys. Lett. B **696**, 513 (2011)
8. Z. Lu, B.-Q. Ma, J. She, Phys. Rev. D **84**, 034010 (2011)
9. J. Zhu, B.-Q. Ma, Phys. Rev. D **82**, 114022 (2010)
10. V. Barone et al. (PAX Collaboration), arXiv:hep-ex/0505054 (2005)
11. A. V. Efremov, K. Goeke, P. Schweitzer, Eur. Phys. J. C **35**, 207 (2004)
12. A. Airapetian et al. (HERMES Collaboration), Phys. Rev. Lett. **94**, 012002 (2005)
13. A. Airapetian et al. (HERMES Collaboration), Phys. Lett. B **693**, 11 (2010)
14. V. Y. Alexakhin et al. (COMPASS Collaboration), Phys. Rev. Lett. **94**, 202002 (2005)
15. M. G. Alekseev et al. (COMPASS Collaboration), Phys. Lett. B **692**, 240 (2010)
16. X. Jiang, J. P. Chen, E. Cisbani, H. Gao, J. C. Peng, spokespersons, JLab Report No. E06-010/E06-011
17. H. Gao et al., Eur. Phys. J. Plus **126**, 1 (2011)
18. X. Qian et al. (Jefferson Lab Hall A Collaboration), Phys. Rev. Lett. **107**, 072003 (2011)
19. P. Lenisa, F. Rathmann, CERN Cour. **50N6**, 21 (2010)
20. F. Rathmann et al., Phys. Rev. Lett. **94**, 014801 (2005)
21. C. Barschel et al. (PAX Collaboration), arXiv:0904.2325 [nucl-ex] (2009)
22. V. F. Dmitriev, A. I. Milstein, S. G. Salnikov, Phys. Lett. B **690**, 427 (2010)
23. B.-Q. Ma, Phys. Lett. B **375**, 320 (1996)
24. B.-Q. Ma, J. Phys. G **17**, L53 (1991), [arXiv:0711.2335 [hep-ph]]
25. B.-Q. Ma, Z. Phys. C **58**, 479 (1993)
26. I. Schmidt, J. Soffer, Phys. Lett. B **407**, 331 (1997)
27. B.-Q. Ma, I. Schmidt, J. Soffer, Phys. Lett. B **441**, 461 (1998)
28. B.-Q. Ma, I. Schmidt, J. Soffer, J.-J. Yang, Phys. Rev. D **62**, 114009 (2000)
29. S. J. Brodsky, T. Huang, and G. P. Lepage, in *Quarks and Nuclear Forces*, edited by D. Fries and B. Zeitnitz (Springer, Tracts in Modern Physics, Vol. 100) (Springer, New York, 1982)
30. T. Huang, B.-Q. Ma, Q.-X. Shen, Phys. Rev. D **49**, 1490 (1994)
31. S. Arnold, A. Metz, M. Schlegel, Phys. Rev. D **79**, 034005 (2009)
32. A. M. Kotzinian, P. J. Mulders, Phys. Lett. B **406**, 373 (1997)
33. Z. Lu, B.-Q. Ma, I. Schmidt, Phys. Rev. D **75**, 094012 (2007)
34. J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky, W. K. Tung, J. High Energy Phys. **0207**, 012 (2002)
35. M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, A. Prokudin, S. Melis, Nucl. Phys. Proc. Suppl. **191**, 98 (2009)
36. M. Hirai, S. Kumano, N. Saito, Phys. Rev. D **74**, 014015 (2006)
37. D. de Florian, R. Sassot, M. Stratmann, W. Vogelsang, Phys. Rev. Lett. **101**, 072001 (2008)
38. D. de Florian, R. Sassot, M. Stratmann, W. Vogelsang, Phys. Rev. D **80**, 034030 (2009)
39. E. Leader, A. V. Sidorov, D. B. Stamenov, Phys. Rev. D **82**, 114018 (2010)
40. J. Blumlein, H. Bottcher, Nucl. Phys. B **841**, 205 (2010)
41. B.-Q. Ma, D. Qing, I. Schmidt, Phys. Rev. C **65**, 035205 (2002)
42. B.-Q. Ma, D. Qing, I. Schmidt, Phys. Rev. C **66**, 048201 (2002)
43. X. Chen, Y. Mao, B.-Q. Ma, Nucl. Phys. A **759**, 188 (2005)
44. Y. Huang, J. She, B.-Q. Ma, Phys. Rev. D **76**, 034004 (2007)
45. J. Huang et al. (Jefferson Lab Hall A Collaboration), arXiv:1108.0489 [nucl-ex] (2011)