

Looking for a charge asymmetry in cosmic rays

I Masina^{1,2}

¹ Dip. di Fisica dell'Università di Ferrara and INFN Sez. di Ferrara, Via Saragat, Ferrara, Italy

² CP3-Origins, University of Southern Denmark, Campusvej, Odense, Denmark

E-mail: masina@fe.infn.it

Abstract. We combine the data from PAMELA and FERMI-LAT cosmic ray experiments by introducing a simple sum rule. This allows to investigate whether the lepton excess observed by these experiments is charge symmetric or not. We also show how the data can be used to predict the positron fraction at energies yet to be explored by the AMS-02 experiment.

1. Introduction

The data collected by PAMELA [1] indicate that there is a positron excess in the cosmic ray (CR) energy spectrum above 10 GeV. The rising behavior observed by PAMELA does not fit previous estimates of the CR formation and propagation implying the possible existence of a direct excess of CR positrons of unknown origins. Interestingly PAMELA's data show no excess in the anti-protons.

While ATIC [2] and PPB-BETS [3] reported unexpected structure in the all-electron spectrum in the range 100 GeV- 1 TeV, the picture has changed with the higher-statistics measurements by FERMI-LAT [4] and HESS [5], leading to a possible slight additional unknown component in the CR e^\pm flux over and above the specific Moskalenko and Strong model prediction [6, 7] which assumes a single-power-law injection spectrum.

These interesting features have drawn much attention, and many explanations have been proposed: For example, these excesses could be due to an inadequate account of the CR background in previous modeling; The presence of new astrophysical sources; They could also originate from annihilations and/or decays of dark matter. We refer to [8] for a recent review.

Whatever the origin of these excesses might be, following Ref. [9], we show how to derive constraints about their charge asymmetry. Clearly, this will eventually shed some light on the physical nature of their source.

2. A useful sum rule

We start by writing the observed flux of electrons and positrons as the sum of two contributions: A background component, ϕ_\pm^B , due to known astrophysical sources (which actually are better known for the electrons rather than the positrons), and an unknown component, ϕ_\pm^U , in formulae:

$$\phi_\pm = \phi_\pm^U + \phi_\pm^B . \quad (1)$$

The component ϕ_\pm^U is the one needed to explain the features in the spectra observed by PAMELA and FERMI-LAT.

These experiments measure respectively the positron fraction and the total electron and positron fluxes as a function of the energy E of the detected e^\pm , i.e.:

$$P(E) = \frac{\phi_+(E)}{\phi_+(E) + \phi_-(E)}, \quad F(E) = \phi_+(E) + \phi_-(E). \quad (2)$$

The left-hand side of the equations above refer to the experimental measures. The contribution from the unknown source is then:

$$\phi_+^U(E) = P(E) F(E) - \phi_+^B(E), \quad (3)$$

$$\phi_-^U(E) = F(E) (1 - P(E)) - \phi_-^B(E). \quad (4)$$

In terms of their difference and sum:

$$\begin{aligned} \phi_+^U(E) - \phi_-^U(E) &= F(E) (2P(E) - 1) + (\phi_-^B(E) - \phi_+^B(E)), \\ \phi_+^U(E) + \phi_-^U(E) &= F(E) - (\phi_-^B(E) + \phi_+^B(E)). \end{aligned} \quad (5)$$

The latter equation implies $F(E) \geq \phi_-^B(E) + \phi_+^B(E)$. We model the background spectrum using $\phi_\pm^B(E) = N_B B^\pm(E)$, where N_B is a normalization coefficient such that $F(E)/(B^-(E) + B^+(E)) \geq N_B$ and $B^\pm(E)$ are provided using specific astrophysical models. In this paper we adopt the popular Moskalenko and Strong model [6, 7], for which N_B is less than 0.75 and $B^\pm(E)$ are given by:

$$B^+ = \frac{4.5E^{0.7}}{1 + 650E^{2.3} + 1500E^{4.2}}, \quad (6)$$

$$B^- = B_1^- + B_2^-, \quad (7)$$

$$B_1^- = \frac{0.16E^{-1.1}}{1 + 11E^{0.9} + 3.2E^{2.15}}, \quad (8)$$

$$B_2^- = \frac{0.70E^{0.7}}{1 + 110E^{1.5} + 600E^{2.9} + 580E^{4.2}}, \quad (9)$$

where E is measured in GeV and the B s in $\text{GeV}^{-1}\text{cm}^{-2}\text{sec}^{-1}\text{sr}^{-1}$ units. We checked that our results remain unchanged when replacing the parameterization above with the one adopted by the Fermi Collaboration (model zero) [10, 11].

It is convenient to introduce the following parameter:

$$r_U(E) \equiv \frac{\phi_-^U(E)}{\phi_+^U(E)} = \frac{F(E) (1 - P(E)) - \phi_-^B(E)}{P(E) F(E) - \phi_+^B(E)}, \quad (10)$$

which quantifies the level of charge asymmetry in the unknown component of electrons and positrons. The equation above can be rewritten as

$$R(E) \equiv \frac{F(E)}{B^-(E)} \frac{1 - (1 + r_U(E))P(E)}{1 - r_U(E) \frac{\phi_+^B(E)}{\phi_-^B(E)}} = N_B. \quad (11)$$

Although the sum rule $R(E)$ seems to depend on the energy it should, in fact, be a constant as is clear from the right hand side of the previous equation. *This leads to a nontrivial constraint linking together in an explicit form the experimental results, the model of the backgrounds and the dependence on the energy of the unknown components.*

We now turn to the actual data and show in which way the sum rule provides essential information on the unknown components of the CRs.

Since we use simultaneously the results of FERMI-LAT and PAMELA we are obliged to consider only the common energy range. Note that the CRs energy range below $10 - 20$ GeVs, where the spectrum is affected by the Sun, is outside the common range. Within the relevant but limited range of energies we will consider here it is therefore sensible to assume r_U to be nearly constant.

We find useful to plot the function $R(E)$ for different values of r_U in order to test the sum rule. This would imply that this function is independent of the energy. The associated constant value would then be identified with the background CRs normalization factor N_B . We report the results in Fig. 1. The straight (red) line is the $N_B = 0.75$ value which is the largest one can assume for the background not to be larger than the FERMI-LAT results. The shaded region is obtained by using the PAMELA and FERMI-LAT data released in 2010, assuming a 1σ error band for both. We observe that there is a clear tendency for the combined data to predict a lower value of the constant N_B , say about 0.66, 0.64, 0.62, 0.55 for increasing value of the ratio $r_U = 0, 1, 2, 4$. This is clear when looking, from top to bottom, at the different panels of Fig. 1. It is interesting to note that we find a plateau, in the relevant energy range, up to r_U near the value of 4 when $R(E)$ starts showing some deviation.




Figure 1. Ratio $R(E)$ as a function of the energy E of electrons and positrons and for values of $r_U = 0, 1, 2, 4$. The shaded region accounts for the 1σ error in PAMELA and FERMI-LAT data (2010 release). Secondaries are estimated according to the expressions in [6, 7]. The value of $R(E)$ thus corresponds to the normalization factor for the background, called N_B in the text.

Given the large uncertainties (for PAMELA, the 2010 1σ error bands are larger than in 2008) we cannot yet provide a more solid conclusion, however we can use the derived normalization N_B

for each different ratios of the unknown components to *predict* the positron fraction at energies higher than the ones provided so far by PAMELA.

3. PAMELA and the charge asymmetry of the unknown component

In order to be able to make such a prediction we first rewrite eq. (11) as follows:

$$P(E) = \frac{1}{1+r_U} \left(1 - \frac{\phi_-^B(E)}{F(E)} \left(1 - r_U \frac{\phi_+^B(E)}{\phi_-^B(E)} \right) \right), \quad (12)$$

where we use for each r_U the estimated associated N_B (referring to Fig. 1, we consider the range of N_B values allowed for $E \approx 25$ GeV). The different predictions for the positron fraction, assuming that r_U remains constant over the entire energy range, up to 1000 GeV are shown in Fig. 2. The resulting picture disfavors both very small and very large values of r_U which, in




Figure 2. Model independent prediction for the positron fraction $P(E)$ as a function of the energy E of electrons and positrons and for values of $r_U = 0, 1/2, 1, 2, 4$, from top to bottom. Secondaries are estimated according to the expressions in [6, 7] reported in the main text. The shaded regions account for the range of the allowed N_B values (see Fig. 1), obtained in turn by considering the 2010 PAMELA and FERMI-LAT data at 1σ . We also display the 2010 (red) and 2008 (black) PAMELA data, with 1σ error bars.

turn, means that one expects the electron fraction to be neither be small nor too large with respect to the positron fraction.

As the experimental data on the positron fraction should hopefully soon become more accurate [12], the method outlined above will allow to extract crucial information about the charge asymmetry of the unknown component of the electrons and positrons in CRs.

4. Conclusions

The model independent analysis of the combined PAMELA and FERMI-LAT data proposed in ref. [9] shows that current data still allow for approximately equal contributions of the electrons

and positrons from the unknown components of the associated CRs, but disfavor electron to positron fractions much smaller than 1/2 or larger than 4. Hence, the typical oversimplifying model assumption of charge symmetry ($r_U = 1$) used so far, actually constitutes just a small portion of the allowed models still left unconstrained by the present data.

Interestingly enough, we emphasize that the planned improvements in the experimental sensitivity to the positron fraction, as well as further data at energies higher than the one explored so far by PAMELA, could reveal whether the unknown component of the CR electrons and positrons is charge asymmetric or not. If it turned out to be charge asymmetric, an explanation in terms of new physics beyond the Standard Model would clearly become more compelling with respect to an astrophysical one.

Acknowledgments

We thank F.Palmonari and R.Sparvoli for useful comments and suggestions.

References

- [1] Adriani O *et al.* 2009 *Nature* **458** 607 (*Preprint* arXiv:0810.4995 [astro-ph]). Adriani O *et al.* 2010 *Astropart. Phys.* **34** 1-11 (*Preprint* arXiv:1001.3522 [astro-ph.HE]). Adriani O *et al.* 2010 *Phys. Rev. Lett.* **105** 121101 (*Preprint* arXiv:1007.0821 [astro-ph.HE]).
- [2] Chang J *et al.* 2008 *Nature* **456** 362.
- [3] Yoshida K *et al.* 2008 *Adv. Space Res.* **42** 1670. Torii S *et al.* *Preprint* arXiv:0809.0760 [astro-ph].
- [4] Abdo A *et al.* 2009 *Phys. Rev. Lett.* **102** 181101 (*Preprint* arXiv:0905.0025 [astro-ph.HE]).
- [5] Aharonian F *et al.* 2008 *Phys. Rev. Lett.* **101** 261104 (*Preprint* arXiv:0811.3894 [astro-ph]). Aharonian F *et al.* 2009 *Astron. Astrophys.* **508** 561 (*Preprint* arXiv:0905.0105 [astro-ph.HE]).
- [6] Strong A W and Moskalenko I V 1998 *Astrophys. J.* **509** 212 (*Preprint* arXiv:astro-ph/9807150).
- [7] Baltz E A and Edsjo E 1998 *Phys. Rev. D* **59** 023511 (*Preprint* arXiv:astro-ph/9808243).
- [8] Fan Y Z Zhang B and Chang J 2010 *Int. J. Mod. Phys. D* **19** 2011-2058 (*Preprint* arXiv:1008.4646 [astro-ph.HE]).
- [9] Frandsen M T Masina I and Sannino F *Preprint* arXiv:1011.0013 [hep-ph].
- [10] Grasso D *et al.* 2009 *Astropart. Phys.* **32** 140 (*Preprint* arXiv:0905.0636 [astro-ph.HE]).
- [11] Ibarra A Tran D and Weniger C 2010 *J. High Energy Phys.* JCAP01(2010)009 (*Preprint* arXiv:0906.1571 [hep-ph]).
- [12] The AMS-02 webpage: <http://ams.cern.ch/>