

JET QUENCHING FROM RHIC TO LHC

B.G. ZAKHAROV

*L.D. Landau Institute for Theoretical Physics, GSP-1, 117940,
Kosygina Str. 2, 117334 Moscow, Russia*

We perform a joint analysis of the data from PHENIX at RHIC and ALICE at LHC on the nuclear modification factor R_{AA} . The computations are performed within the light-cone path integral approach to induced gluon emission. Our results show that slow variation of R_{AA} from RHIC to LHC energies indicates that the QCD coupling constant is suppressed in the quark-gluon plasma produced at LHC.

1. One of the most striking results of experiments at RHIC is strong suppression of high- p_T hadrons in AA -collisions¹ (called “jet quenching”). Recently, a similar effect has been observed in the ALICE experiment at LHC² for $Pb + Pb$ collisions at $\sqrt{s} = 2.76$ TeV. The most natural reason for this phenomenon is parton energy loss (radiative and collisional) in the hot quark-gluon plasma (QGP) produced in the initial stage of AA -collisions. It is of great interest to perform a joint analysis of the RHIC and LHC data. It is interesting since variation of the nuclear modification factor R_{AA} from RHIC to LHC energies should not be very sensitive to the systematic theoretical uncertainties that are rather large. These uncertainties come mostly from multiple induced gluon emission. The available theoretical approaches to radiative induced gluon emission^{3,4,5,6,7,8} are restricted to one gluon emission, and the multiple gluon emission is usually evaluated in the approximation of independent gluon radiation⁹.

In this talk, I will present results of an analysis of the data on R_{AA} for $Au + Au$ collisions at $\sqrt{s} = 200$ GeV from PHENIX¹⁰ and for $Pb + Pb$ collisions at $\sqrt{s} = 2.76$ TeV from ALICE². The analysis is based on the light-cone path integral (LCPI) approach⁴. We evaluate the nuclear modification factor using the method developed in¹¹. A major purpose of this analysis is to decide whether the variation of R_{AA} from RHIC to LHC indicates that the QCD coupling constant becomes smaller in the plasma produced at LHC, which is hotter than that at RHIC.

2. The nuclear modification factor R_{AA} for a given impact parameter b can be written as

$$R_{AA}(b) = \frac{dN(A + A \rightarrow h + X)/d\mathbf{p}_T dy}{T_{AA}(b)d\sigma(N + N \rightarrow h + X)/d\mathbf{p}_T dy}. \quad (1)$$

Here \mathbf{p}_T is the hadron transverse momentum, y is rapidity (we consider the central region $y = 0$), $T_{AA}(b) = \int d\boldsymbol{\rho} T_A(\boldsymbol{\rho}) T_A(\boldsymbol{\rho} - \mathbf{b})$, T_A is the nucleus profile function. The differential yield for high- p_T hadron production in AA -collision can be written in the form

$$\frac{dN(A + A \rightarrow h + X)}{d\mathbf{p}_T dy} = \int d\boldsymbol{\rho} T_A(\boldsymbol{\rho}) T_A(\boldsymbol{\rho} - \mathbf{b}) \frac{d\sigma_m(N + N \rightarrow h + X)}{d\mathbf{p}_T dy}, \quad (2)$$

where $d\sigma_m(N + N \rightarrow h + X)/d\mathbf{p}_T dy$ is the medium-modified cross section for the $N+N \rightarrow h+X$ process. Similarly to the ordinary pQCD formula, we write it as

$$\frac{d\sigma_m(N + N \rightarrow h + X)}{d\mathbf{p}_T dy} = \sum_i \int_0^1 \frac{dz}{z^2} D_{h/i}^m(z, Q) \frac{d\sigma(N + N \rightarrow i + X)}{d\mathbf{p}_T^i dy}. \quad (3)$$

Here $\mathbf{p}_T^i = \mathbf{p}_T/z$ is the parton transverse momentum, $d\sigma(N + N \rightarrow i + X)/d\mathbf{p}_T^i dy$ is the hard cross section, $D_{h/i}^m$ is the medium-modified fragmentation function (FF) for transition of a parton i into the observed hadron h . For the parton virtuality scale Q we take the parton transverse momentum p_T^i . We assume that hadronization occurs outside of the QGP. For jets with $E \lesssim 100$ GeV the hadronization scale, μ_h , is relatively small. Indeed, one can easily show that the L dependence of the parton virtuality reads $Q^2(L) \sim \max(Q/L, Q_0^2)$, where $Q_0 \sim 1 - 2$ GeV is some minimal nonperturbative scale. For RHIC and LHC, when $\tau_{QGP} \sim R_A$ (τ_{QGP} is the typical lifetime/size of the QGP, R_A is the nucleus radius), it gives $\mu_h \sim Q_0$ (for $E \lesssim 100$ GeV). Then we can write

$$D_{h/i}^m(z, Q) \approx \int_z^1 \frac{dz'}{z'} D_{h/j}(z/z', Q_0) D_{j/i}^m(z', Q_0, Q), \quad (4)$$

where $D_{h/j}(z, Q_0)$ is the vacuum FF, and $D_{j/i}^m(z', Q_0, Q)$ is the medium-modified FF for transition of the initial parton i with virtuality Q to a parton j with virtuality Q_0 . For partons with $E \lesssim 100$ GeV the typical length scale dominating the energy loss in the DGLAP stage is relatively small $\sim 0.3 - 1$ fm¹¹. This length is of the order of the formation time of the QGP $\tau_0 \sim 0.5$ fm. Since the induced radiation stage occurs at larger length scale $l \sim \tau_0 \div \tau_{QGP}$, to the first approximation one can ignore the overlap of the DGLAP and induced radiation stages at all¹¹. Then we can write

$$D_{j/i}^m(z, Q_0, Q) = \int_z^1 \frac{dz'}{z'} D_{j/l}^{ind}(z/z', E_l) D_{l/i}^{DGLAP}(z', Q_0, Q), \quad (5)$$

where $E_l = Qz'$, $D_{j/l}^{ind}$ is the induced radiation FF (it depends on the parton energy E , but not virtuality), and $D_{l/i}^{DGLAP}$ is the vacuum DGLAP FF.

We have computed the DGLAP FFs with the help of the PYTHIA event generator¹². One gluon induced emission has been computed within the LCPI formalism⁴ using the method elaborated in¹³. As in^{13,11} we take $m_q = 300$ and $m_g = 400$ MeV for the quark and gluon quasiparticle masses. Our method of calculation of the in-medium FF via the one gluon probability distribution is described in detail in¹¹, and need not to be repeated here. We just enumerate its basic aspects. The multiple gluon emission is accounted for employing Landau's method as in⁹. For quarks the leakage of the probability to the unphysical region of $\Delta E > E$ is accounted for by renormalizing the FF. We also take into account the $q \rightarrow g$ FF. Its normalization is fixed from the momentum conservation for $q \rightarrow q$ and $q \rightarrow g$ transitions. The normalization of the $g \rightarrow g$ FF is also fixed from the momentum sum rule. The collisional energy loss, which is small¹⁴, is taken into account by renormalizing the temperature of the QGP for the radiative FFs using the condition: $\Delta E_{rad}(T_0') = \Delta E_{rad}(T_0) + \Delta E_{col}(T_0)$, where $\Delta E_{rad/col}$ is the radiative/collisional energy loss, T_0 is the real initial temperature of the QGP, and T_0' is the renormalized temperature.

We calculate the hard cross sections using the LO pQCD formula. To simulate the higher order K -factor we take for the virtuality scale in α_s the value cQ with $c = 0.265$ as in the PYTHIA event generator¹². We account for the nuclear modification of the parton densities (which leads to some small deviation of R_{AA} from unity even without parton energy loss) with the help of the EKS98 correction¹⁵. For the vacuum FFs we use the KKP parametrization¹⁶.

As in¹¹, we evaluate the induced gluon emission and the collisional energy loss for the running α_s frozen at some value α_s^{fr} at low momenta. For vacuum a reasonable choice is

$\alpha_s^{fr} \approx 0.7$. This value was previously obtained by fitting the low- x proton structure function F_2 within the dipole BFKL equation¹⁷. To study the role of the in-medium suppression of α_s we perform the computations for several smaller values of α_s^{fr} .

3. We describe the QGP in the Bjorken model¹⁸ which gives $T_0^3 \tau_0 = T^3 \tau$. We take $\tau_0 = 0.5$ fm. To simplify numerical computations for each impact parameter b we neglect variation of the initial temperature T_0 in the transverse directions. We evaluate its value using the entropy/multiplicity ratio $dS/dy/dN_{ch}/d\eta \approx 7.67$ obtained in¹⁹. For the central $Au + Au$ collisions at $\sqrt{s} = 200$ GeV $T_0 \approx 300$ MeV and for $Pb + Pb$ collisions at $\sqrt{s} = 2.76$ TeV $T_0 \approx 400$ MeV. For the nuclear density we use the Woods-Saxon nucleus density with parameters as in². The fast parton path length in the QGP, L , in the medium has been calculated according to the position of the hard reaction in the impact parameter plane. To take into account the fact that at times about 1–2 units of R_A the transverse expansion should lead to fast cooling of the hot QCD matter¹⁸ we also impose the condition $L < L_{max}$. We performed the computations for $L_{max} = 8$ and 10 fm. The difference between these two versions is small.

4. In Fig. 1 the theoretical R_{AA} obtained for $\alpha_s^{fr} = 0.7, 0.6,$ and 0.5 for the chemically equilibrium and purely gluonic plasmas is compared to the PHENIX data¹⁰ on π^0 production in the 0-5% central $Au + Au$ collisions at $\sqrt{s} = 200$ GeV. The results are presented for radiative energy




Figure 1: The factor R_{AA} for π^0 production in the 0-5% central $Au + Au$ collisions at $\sqrt{s} = 200$ GeV for $\alpha_s^{fr} = 0.7, 0.6,$ and 0.5 . The upper panels are for the chemically equilibrium plasma, and the lower ones for purely gluonic plasma. Black line: the total radiative part (quarks plus gluons); red line: the radiative quark part; green line: the radiative gluon part; blue line: the radiative (quarks and gluons) plus collisional, and plus energy gain due to gluon absorption. The theoretical curves obtained for $L_{max} = 8$ fm. The experimental points are the PHENIX data¹⁰.

loss and with inclusion of collisional energy loss and radiative energy gain. The effect of the radiative energy gain on R_{AA} is practically negligible and can be safely neglected. The growth of R_{AA} for gluons in Fig. 1 is due to the $q \rightarrow g$ transition which is usually neglected. However, it does not affect strongly the total R_{AA} since for $\sqrt{s} = 200$ GeV the gluon contribution to the hard cross section is small at $p_T \gtrsim 15$ GeV. In Fig. 2 we compare our results for $\alpha_s^{fr} = 0.7, 0.5,$ and 0.4 with the ALICE data² for charged hadrons in $Pb + Pb$ collisions at $\sqrt{s} = 2.76$ TeV.

As can be seen from Figs. 1, 2, the collisional energy loss suppresses R_{AA} only by about 15-25%. For the equilibrium plasma the data for $\sqrt{s} = 200$ GeV can be described with $\alpha_s^{fr} \approx 0.6 \div 0.7$. The data for $\sqrt{s} = 2.76$ TeV agree better with $\alpha_s^{fr} \approx 0.4 \div 0.5$. It provides evidence for the thermal suppression of α_s at LHC due to higher temperature of the QGP.

5. In summary, we have analyzed the data on R_{AA} obtained in the PHENIX experiment on




Figure 2: The same as in Fig. 1 for the charged hadrons in $Pb + Pb$ collisions at $\sqrt{s} = 2.76$ TeV for $\alpha_s^{fr} = 0.7$, 0.5 and 0.4. The experimental points are the ALICE data², as in² the boxes contain the systematic errors.

$Au + Au$ collisions at $\sqrt{s} = 200$ GeV¹⁰ at RHIC and in the ALICE experiment on $Pb + Pb$ collisions at $\sqrt{s} = 2.76$ TeV² at LHC. Our results show that slow variation of R_{AA} from RHIC to LHC supports that the QCD coupling constant becomes smaller in the hotter QGP at LHC.

Acknowledgments

I am grateful to the organizers for such an enjoyable and stimulating meeting and for financial support of my participation.

References

1. P.M. Jacobs, M. van Leeuwen, Nucl. Phys. **A774**, 237 (2006) and references therein.
2. K. Aamodt *et al.* [ALICE Collaboration], Phys. Lett. **B696**, 30 (2011).
3. R. Baier, Y.L. Dokshitzer, A.H. Mueller, S. Peigné, and D. Schiff, Nucl. Phys. **B483**, 291 (1997); *ibid.* **B484**, 265 (1997); R. Baier, Y.L. Dokshitzer, A.H. Mueller, and D. Schiff, Nucl. Phys. **B531**, 403 (1998).
4. B.G. Zakharov, JETP Lett. **63**, 952 (1996); *ibid* **65**, 615 (1997); **70**, 176 (1999); Phys. Atom. Nucl. **61**, 838 (1998).
5. R. Baier, D. Schiff, and B.G. Zakharov, Ann. Rev. Nucl. Part. **50**, 37 (2000).
6. U.A. Wiedemann, Nucl. Phys. **A690**, 731 (2001).
7. M. Gyulassy, P. Lévai, and I. Vitev, Nucl. Phys. **B594**, 371 (2001).
8. P. Arnold, G.D. Moore, and L.G. Yaffe, JHEP **0206**, 030 (2002).
9. R. Baier, Yu.L. Dokshitzer, A.H. Mueller, and D. Schiff, JHEP **0109**, 033 (2001).
10. A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **101**, 232301 (2008).
11. B.G. Zakharov, JETP Lett. **88**, 781 (2008) [arXiv:0811.0445 [hep-ph]].
12. T. Sjostrand, L. Lonnblad, S. Mrenna, and P. Skands, arXiv:hep-ph/0308153.
13. B.G. Zakharov, JETP Lett. **80**, 617 (2004) [arXiv:hep-ph/0410321].
14. B.G. Zakharov, JETP Lett. **86**, 444 (2007) [arXiv:0708.0816 [hep-ph]].
15. K.J. Eskola, V.J. Kolhinen, and C.A. Salgado, Eur. Phys. J. **C9**, 61 (1999).
16. B. A. Kniehl, G. Kramer, and B. Potter, Nucl. Phys. **B582**, 514 (2000).
17. N.N. Nikolaev and B.G. Zakharov, Phys. Lett. **B327**, 149 (1994).
18. J.D. Bjorken, Phys. Rev. **D27**, 140 (1983).
19. B. Müller and K. Rajagopal, Eur. Phys. J. **C43**, 15 (2005).