

Electromagnetic properties of baryons

T. Ledwig,¹ J. Martin-Camalich,² V. Pascalutsa,¹ and M. Vanderhaeghen¹

¹*Institut für Kernphysik, Universität Mainz, D-55099 Mainz, Germany*

²*Department of Physics and Astronomy, University of Sussex, BN1 9Qh, Brighton, UK.*

Departamento de Física Teórica and IFIC, Universidad de Valencia-CSIC, Spain

We discuss the chiral behavior of nucleon and $\Delta(1232)$ electromagnetic properties within the framework of a $SU(2)$ covariant baryon chiral perturbation theory. Our one-loop calculation is complete to the order p^3 and p^4/Δ with Δ as the $\Delta(1232)$ -nucleon energy gap. We show that the magnetic moment of a resonance can be defined through the linear energy shift only when an additional relation between the involved masses and the applied magnetic field strength is fulfilled. Singularities and cusps in the pion mass dependence of the $\Delta(1232)$ electromagnetic moments reflect a non-fulfillment. We show results for the pion mass dependence of the nucleon iso-vector electromagnetic quantities and present results for finite volume effects on the iso-vector anomalous magnetic moment.

Keywords: electromagnetic moments, resonances, chiral perturbation theory, chiral behavior

I. ANOMALOUS MAGNETIC MOMENT OF A RESONANCE

We consider the energy shift $\Delta E(\vec{B})$ of an unstable spin-1/2 particle Ψ of mass M_* in an external magnetic field \vec{B} . The decay products are a spin-1/2 particle ψ and a scalar particle ϕ of masses M and m , respectively. We take a Yukawa interaction of strength g given by: $\mathcal{L}_{\text{int}} = g \left(\bar{\Psi} \psi \phi + \bar{\psi} \Psi \phi^* \right)$. For stable particles with spin $s_z = +1/2$ and $\vec{B} = B e_z$, we can normally write the energy shift induced by the anomalous magnetic moment (a.m.m.) κ_* as $\Delta \tilde{E} = \frac{\Delta E}{M_*} + \frac{1}{2} \tilde{B} = -\frac{\kappa_*}{2} \tilde{B}$ with $\tilde{B} = e B_z / M_*^2$. This is not generally possible for unstable particles [1]. To see this, we calculate $\Delta \tilde{E}$ by the background field method of [2]:

$$\Delta \tilde{E} = \frac{g^2}{(4\pi)^2} \int_0^1 dx (r+x) \ln \left[1 + \frac{x(1-x)\tilde{B}}{x\mu^2 - x(1-x) + (1-x)r^2} \right], \quad (1)$$

with $r = M/M_*$, $\mu = m/M_*$. After integrating the Feynman parameter, we obtain contributions which are non-analytic in \tilde{B} . These terms can only be expanded, i.e. a magnetic moment defined by $\Delta E = -\vec{\mu} \cdot \vec{B}$, when the following condition is met:

$$|eB|/2M_* \ll |M_* - M - m|. \quad (2)$$

A mass of $m = M_* - M$ violates the above condition which is reflected by singularities and cusps in electromagnetic (em) moments. Examples are the em moments of the W -boson [3]. The loop contributions with top and bottom quarks are singular when the masses are tuned to $m_t = m_W - m_b$. Another example are the em moments of the $\Delta(1232)$. We obtain singularities and cusps in the pion mass dependence of the $\Delta(1232)$ magnetic dipole, electric quadrupole and magnetic octupole moments at $m_\pi = M_\Delta - M_N$ [4].

II. NUCLEON ELECTROMAGNETIC PROPERTIES

The nucleon Dirac $F_1(Q^2)$ and Pauli $F_2(Q^2)$ form factors are defined by the following matrix element:

$$\langle N(p') | \bar{\Psi}(0) \gamma^\mu \Psi(0) | N(p) \rangle = \bar{u}(p') \left[\gamma^\mu F_1(Q^2) + \frac{i\sigma^{\mu\nu} q_\nu}{2m_N} F_2(Q^2) \right] u(p), \quad (3)$$

with $q = p' - p$, $Q^2 = -q^2$ as the momentum transfer and $u(p)$ as the nucleon spinor with mass m_N . At $Q^2 = 0$ these form factors are the nucleon charge e_N , anomalous magnetic moment κ_N and magnetic moment μ_N : $F_1(0) = e_N$, $F_2(0) = \kappa_N$ and $\mu_N = (e_N + \kappa_N)e/2m_N$.

We concentrate on the chiral behavior of the nucleon iso-vector anomalous magnetic moment κ_V , Dirac $\langle r_1^2 \rangle_V$ and Pauli $\langle r_2^2 \rangle_V$ radii that are defined by:

$$\kappa_V = F_2^p(0) - F_2^n(0) \quad , \quad \langle r_i^2 \rangle_V = -\frac{6}{F_i^V(0)} \frac{dF_i^V(Q^2)}{dQ^2}. \quad (4)$$

We use the chiral perturbation theory given by the $B\chi$ PT Lagrangian of [6] with the $\Delta(1232)$ -isobar included by the δ -power counting scheme of [7]. The explicit Lagrangian consisting of pion, nucleon, $\Delta(1232)$ -isobar and photon fields can be found in [5, 8]. The power counting breaking terms as found in [6] are treated by the renormalization prescription of [9]. Our $B\chi$ PT investigation is complete to the order p^3 with inclusion of $\Delta(1232)$ effects up to p^4/Δ where Δ is the $\Delta(1232)$ -nucleon energy gap.

In fig. 1 we depict all one-loop contributions originating at the considered order. For the diagram (N4) we also investigate contributions from a non-minimally coupled photon to the $\Delta\Delta\gamma$ vertex as discussed in [10].




FIG. 1: Diagrams contributing to the nucleon electromagnetic properties. Solid-single lines represent nucleons, solid-double lines the $\Delta(1232)$ -isobar and dashed lines the pions. The photon coupling is denoted by the blue squares while the $NN\pi$ and $\Delta N\pi$ vertices by black dots. For the Dirac radius we also include the tadpole graphs that are not depicted here.




FIG. 2: Nucleon iso-vector anomalous magnetic moment κ_V , Dirac $\langle r_1^2 \rangle_V$ and Pauli $\langle r_2^2 \rangle_V$ radii as function of the pion mass squared. The lattice data are taken from: brown up-triangles [11], red rectangles [12], blue down-triangles [13], green squares [14]. The green dotted line corresponds to contributions coming from the graphs (N1) and (N2). The red solid line corresponds to the contributions of all graphs with a non-minimal Δ -photon coupling in (N4) while the blue dotted line is with a minimal Δ -photon coupling. We constrain our results for κ_V and $\langle r_1^2 \rangle_V$ to the phenomenological values $\kappa_V = 3.7$ and $\langle r_1^2 \rangle_V = (0.765)^2 \text{ fm}^2$ at $m_\pi = 139 \text{ MeV}$ which are denoted by blue circles.

In fig. 2 we show our results [4] for the pion mass dependence of the nucleon iso-vector a.m.m., Dirac $\langle r_1^2 \rangle_V$ and Pauli $\langle r_2^2 \rangle_V$ radii. To the order p^3 there appear counter terms for the a.m.m. and $\langle r_1^2 \rangle_V$ while one for the $\langle r_2^2 \rangle_V$ appears at higher orders.

III. FINITE VOLUME EFFECTS FOR κ_V

We now discuss preliminary results for finite volume contributions to the nucleon iso-vector a.m.m. coming from the graphs (N1) and (N2) of fig. 1. For this, we calculate the self-energy of the nucleon in a constant magnetic field $\vec{B} = B e_z$. We take here the pseudo-scalar $NN\pi$ vertex. We obtain for the self-energies of the proton and neutron the following results:

$$\left. \begin{aligned} \Sigma_p &= 2\Sigma_1 + \Sigma_2 \\ \Sigma_n &= 2\Sigma_3 \end{aligned} \right\} \text{ with } \Sigma_i = \frac{g^2 m_N}{i(4\pi)^4} \int_0^1 dz \int d^4 l \frac{1-z}{[l^2 + m_\pi^2 z - (1-z)^2 m_N^2 - \beta_i]^2} \quad (5)$$

and $\beta_1 = z(1-z)B$, $\beta_2 = -(1-z)^2 B$, $\beta_3 = -(1-z)B$. In a finite volume the loop momentum \vec{l} is discretized by $\int \frac{d^3 \vec{l}}{(2\pi)^3} \rightarrow \frac{1}{L^3} \sum_{\vec{n}}$ with $\vec{l} = \frac{2\pi}{L} \vec{n}$ and $\vec{n} \in Z^3$. We extract the a.m.m. through the linear coefficient in an expansion

in B . The a.m.m. infinite volume contribution obtained from (5) coincides with the one discussed in the previous section.

In fig. 3 we depict our results for the nucleon iso-vector a.m.m. for a volume of size $L = 3$ fm with $n_i \leq 1$ and $n_i \leq 2$. The main effect comes from the $n_i \leq 1$ contribution which adds negatively to the infinite volume result.




FIG. 3: Nucleon iso-vector anomalous magnetic moment. The lattice data is the same as in fig. 2. The dotted green line is our infinite volume result obtained from (5) which coincide with the dotted green line of fig. 1. The other two lines correspond to the finite volume contributions for a box size of $L = 3$ fm and $n_i \leq 1$ for the red line and $n_i \leq 2$ for the dashed black one.

IV. SUMMARY

We first discussed the anomalous magnetic moment of a resonance. In contrast to stable particles, the self-energy of unstable particles depends non-analytically on the applied external magnetic field. A resonance magnetic moment is defined by the linear energy shift $\Delta E = -\vec{\mu} \cdot \vec{B}$ only when condition (2) is met. We calculated the pion mass dependence of the $\Delta(1232)$ electromagnetic moments which reveal singularities and cusps for the pion mass at which (2) is not fulfilled. We studied the chiral behavior of the nucleon iso-vector a.m.m., Dirac- and Pauli-radii in a manifestly covariant $SU(2)$ baryon chiral perturbation theory with inclusion of $\Delta(1232)$ -isobar degrees of freedom. Our results are complete to the order p^3 and p^4/Δ . We also showed preliminary results for finite volume effects on the iso-vector a.m.m. .

Acknowledgments

The work of T.L. was partially supported by the Research Centre Elementarkraefte und Mathematische Grundlagen at the Johannes Gutenberg University Mainz. JMC acknowledges the MEC contract FIS2006-03438, the EU Integrated Infrastructure Initiative Hadron Physics Project contract RII3-CT-2004-506078 and the Science and Technology Facilities Council [grant number ST/H004661/1] for support.

-
- [1] T. Ledwig, V. Pascalutsa, M. Vanderhaeghen, Phys. Rev. D **82**, 091301 (2010).
 - [2] C. M. Sommerfield, Annals of Physics **5**, 26 (1958).
 - [3] G. Couture, J. N. Ng, Z. Phys. C **35**, 65 (1987).
 - [4] T. Ledwig, J. Martin-Camalich, V. Pascalutsa, M. Vanderhaeghen, in preparation.
 - [5] V. Pascalutsa and M. Vanderhaeghen, Phys. Rev. Lett. **94**, 102003 (2005).
 - [6] J. Gasser, M. E. Sainio, A. Svarc, Nucl. Phys. B **307**, 779 (1988).
 - [7] V. Pascalutsa and D. R. Phillips, Phys. Rev. C **67**, 055202 (2003).
 - [8] V. Pascalutsa, M. Vanderhaeghen, Phys. Rev. D **73**, 034003 (2006); Phys. Rev. D **77**, 014027 (2008); V. Pascalutsa, M. Vanderhaeghen, S. N. Yang, Phys. Rept. **437**, 125 (2007).
 - [9] J. Gegelia, G. Japaridze, Phys. Rev. D **60**, 114038 (1999); J. Gegelia, G. Japaridze, X.Q. Wang, J. Phys. G **29**, 2303 (2003); T. Fuchs, J. Gegelia, G. Japaridze, S. Scherer, Phys. Rev. D **68**, 056005 (2003).
 - [10] C. Alexandrou *et al.*, Nucl. Phys. A **825**, 115 (2009); Phys. Rev. D **79**, 014507 (2009).
 - [11] T. Yamazaki *et al.* (RBC and UKQCD Collab.), Phys. Rev. D **79**, 114505 (2009).

- [12] H.-W. Lin, K. Orginos, Phys. Rev. D **79**, 074507 (2009).
- [13] S. N. Syritsyn *et al.* (LHPC Collab.), Phys. Rev. D **81**, 034507 (2010).
- [14] J. D. Bratt *et al.* (LHPC Collab.), Phys. Rev. D **82**, 094502 (2010).