Higgs Physics at the Large Hadron Collider¹

Rohini M. Godbole

Center for High Energy Physics, Indian Institute of Science, Bangalore 560 012, India. &

Institute for Theoretical Physics and Spinoza Institute, Utrecht University, 3508 TD Utrecht, The Netherlands.

Abstract

In this talk I will begin by summarising the importance of the Higgs physics studies at the LHC. I will then give a short description of the pre-LHC constraints on the Higgs mass and the theoretical predictions for the LHC along with a discussion of the current experimental results, ending with prospects in the near future at the LHC. In addition to the material covered in the presented talk, I have included in the writeup, a critical appraisal of the theoretical uncertainties in the Higgs cross-sections at the Tevatron as well as a discussion of the recent experimental results from the LHC which have become available since the time of the workshop.

1 Introduction

It goes without saying that establishing the exact nature of the mechanism of electroweak symmetry breaking is perhaps 'THE' most important issue in the subject of particle physics at present and arguably the 'raison d'être' for the Large Hadron Collider (LHC). The excellent agreement of the LEP data on $\sigma(e^+e^- \rightarrow W^+W^-)$ with predictions of the Standard Model (SM) shown in Fig. 1 gives us a direct confirmation of the triple gauge boson (ZWW) coupling as predicted by the $SU(2) \times U(1)$ symmetry. At the same time the observed nonzero mass of the W-boson confirms that the same EW symmetry is broken as well. The Higgs mechanism [1,2] is one way of achieving the desired breakdown of the EW symmetry. This predicts the existence of a $J^{PC} = 0^{++}$ state, as the remnant of the $SU(2)_L$ doublet, with precise predictions for the coupling of this state to all the SM particles, but is able to give only very weak theoretical constraints on its mass. Since this is the only particle of the SM [2] still lacking confirmation by direct experimental observation, it is clear that discovery of the Higgs boson and a study of its properties are at the heart of the LHC program which has begun operations at 7 TeV since February 2010.

¹Written version of the talk presented at the 11th Workshop on High Energy Physics Phenomenology (WHEPP), Allahabad, India.



Figure 1: Comparison of the LEP data (taken from LEPEWWG) with the SM prediction shown in (b), the contributory processes being depicted in (a).

A few remarks are in order here. Theoretical ideas of electro weak symmetry breaking (EWSB) span a large range, beginning from the weakly coupled Higgs to those of strong interaction dynamics which can involve a composite (or worse no) Higgs boson. All of these, including the SM, of course have had to pass the acid test of the EW precision measurements, at the Z pole at the LEP collider. The latter class of models, involving dynamical symmetry breaking triggered by strong dynamics, have got a new lease of life due to theoretical developments in the context of models with extra dimensions. Needless also to say that the contents of the Higgs sector and the properties of the said particles, both can differ from the SM in the many different proposals of going beyond the standard model (BSM). Further, CP violation in the Higgs sector can be the possible BSM physics that seems to be necessary for a *quantitative* explanation of the Baryon Asymmetry in the Universe. In addition to this, the dark matter in the Universe, seems to also not consist of any of the known particles in the SM. Interestingly, almost all the extensions of the SM, always have a particle which has all the right properties to be a dark matter (DM) candidate. Since most of the extensions of the SM are introduced to deal with some of the, not yet completely understood and/or unsatisfactory, features of the EWSB, almost always this candidate DM particle has interesting connections to Higgs physics as well.

We expect the LHC to unravel the secrets of the physics of the EWSB as well as to provide pointers to the BSM physics which, we hope, in turn will provide key to the explanation of the issues of cosmological importance, viz. the Baryon Asymmetry in the Universe (BAU) and the DM in the Universe. The discussion preceding these few lines, should then convince us that 'Higgs Physics at the LHC', will indeed touch upon almost all the aspects of active investigation in theoretical and experimental particle physics.

While discussing 'Higgs Physics at the LHC' the different issues that need be addressed are:

- 1. Discovering the spin 0 state(s) and measuring their mass as well as couplings to fermions and gauge bosons.
- 2. Can these measurements uniquely decide the gauge group representation to which these

scalar(s) belong? Can they give information about whether the SM is a strongly coupled theory with (perhaps) a composite Higgs boson or a weakly coupled theory with an elementary Higgs boson?

- 3. Is there a CP violation in the Higgs sector?
- 4. Can the LHC give any information on the triple Higgs coupling, which is present as a result of the spontaneous symmetry breaking?

LHC is capable of answering these questions to different degree of completeness, some early and some in the far future.

Clearly, a short survey such as this can not do justice to the enormous amount of work done on the subject [3,4]. The discussion here will hence only focus on a few issues. I shall summarise first the current constraints on the mass and then go on to discuss the status of theoretical predictions for the LHC. I will then present the current projections for discovery and exclusions made by the two LHC experiments. I will then discuss briefly two new developments in the subject: 1) the jet substructure technique which enables use of the $b\bar{b}$ final state arising from the Higgs decay. Due to the large QCD backgrounds this final state could not always be utilised in the analyses hitherto, 2) the possibility of obtaining spin and parity of the observed scalar state even in the early data.

2 SM Higgs: profile and current constraints

As is well known, theoretical considerations are capable of only giving bounds on the Higgs mass. These bounds arise from considerations of triviality and vacuum stability of the Higgs sector and are shown in Fig. 2. To be specific, these limits are obtained from requiring that the Landau pole



Figure 2: Theoretical upper and lower bounds on the Higgs mass in the SM from the assumption of the validity of the SM up to a cut-off scale Λ and vacuum stability [5].

in the evolution of the Higgs self coupling lie above a cut off scale Λ , as well as by demanding that the scalar potential be bounded from below. The limit on the Higgs mass of about 800 GeV for $\Lambda \sim 1$ TeV seen in Fig. 2, is also consistent with the bound that one obtains on very general grounds by demanding that the tree level scattering amplitudes involving the gauge bosons W and Z satisfy unitarity [6]. In fact it is interesting to know that this unitarity demand had been used to derive successfully, the gauge and Higgs boson content of the SM and the coupling structure of these bosons among themselves and the fermions in the SM [7].

The radiative corrections to the W/Z boson masses coming from the Higgs boson are proportional to $\log(M_H/M_W)$. This logarithmic dependence is related to the 'custodial' symmetry which the SM possesses. As a result the precision measurements of the gauge boson masses already can put *indirect* constraints on the allowed value of M_H . Further the 'direct' searches at LEP [8] and Tevatron [9] also exclude the existence of a Higgs boson, in certain mass regions. Fig. 3 shows a compilation of these indirect constraints obtained by the LEPEWWG and Gfitter groups, along with the limits from the direct searches. The two panels show $\Delta \chi^2$ as a function of M_H for a SM fit to the various EW precision observables. Two comments are in order here. Firstly the quality of the SM fit indicates that every candidate for the BSM physics must have at least an approximate custodial symmetry. Secondly, the exclusion of an SM Higgs in the mass range around 160 GeV, coming from direct searches at the Tevatron, has a nontrivial dependence on the nonperturbative knowledge of the proton [10, 11]. These different issues can affect the estimate of uncertainties in the theoretical predictions for the cross-sections for the signal as well as the background. These issues can thus affect the significance of this limit. I will comment upon it later.



Figure 3: The two panels show a summary of the current, direct and indirect, experimental constraints on the Higgs mass from the collider experiments, taken from the web pages of the LEPEWWG and the Gfitter group. Both the panels show $\Delta \chi^2$ as a function of M_H for a SM fit to a variety of precisely measured electroweak observables.

These results shown in Fig. 3 tell us that in the SM, current data prefer a light Higgs and on inclusion of the direct limits from the collider searches, one gets $M_H < 185$ GeV, at 95% confindence level (CL). The closeness of this bound with that seen in the theoretical analysis presented in Fig. 2 in fact raises the hairy prospect that we might find only such a light Higgs with $M_H \leq 180$ –185 GeV and nothing else at the LHC. It should be mentioned here however, that some of the details of the analyses of the EW observables, are quite sensitively dependent on the way in which the theoretical and experimental errors are accounted for therein. This knowledge thus sets now the stage for the LHC Higgs searches.

A remark is in order before we proceed with a discussion of Higgs searches at the LHC in the mass range allowed by the current direct and indirect constraints. These indirect and direct bounds of Figs. 2 and 3 indicate that just the mass of the observed scalar state, should be able to give information about the energy scale at which new physics may appear. For example, a Higgs boson with $m_H \sim 130-140$ GeV, while not providing any 'proof' for TeV scale physics, will indicate strong possibility of TeV scale physics which can keep the Higgs 'naturally' light. Supersymmetry is an example of one BSM physics which can achieve this. Observation of a scalar state with mass in the region of $\sim 180 \text{ GeV}$ will indicate possible compatibility with the absence of any new physics up to such high scales as may not be accessible in direct searches at the LHC. On the other hand a heavier Higgs (say $\sim 300 \text{ GeV}$) would necessarily mean new physics around TeV scale. Specific examples of such new physics can be, e.g., a sequential fourth generation of fermions and/or presence of strong interaction dynamics around TeV scale. An interesting question to ask in the context of Higgs and BSM searches at the LHC is about the time line of possible discovery of a Higgs of a particular mass and that of the corresponding new physics which has been postulated to make the existence of a Higgs boson with that mass consistent with the different theoretically motivated requirements such as naturalness and/or better consistency with the EW precision measurements used in arriving at Fig. 3.

We begin by summarising some of the Higgs properties, relevant for its search at the LHC, such as its width and branching ratio into different channels. Fig. 4 shows the branching ratios for the SM Higgs over the entire mass range that is consistent with the theoretical constants mentioned earlier. Thus we see that for the light Higgs, such as the one indicated by above constraints, the



Figure 4: The decay branching ratios and the total decay width of the SM Higgs boson as a function of its mass, taken from [3].

width of the Higgs boson is expected to be ≤ 1 GeV. For a Higgs with mass ≤ 130 GeV branching ratio into the $b\bar{b}$ channel is expected to be large, with that in the $\gamma\gamma$ channel ~ 10^{-3} . For larger values of the Higgs mass, the VV decay modes are dominant, with WW and ZZ sharing it in the ratio 2:1. For the Higgs in the mass range ≥ 135 GeV the four fermion decay mode is the most important one. Combined QCD and EW corrections can change this by upto a few percent. Due to the large QCD backgrounds, the $\gamma\gamma$ mode is considered optimal for the Higgs in the mass range 115–135 GeV. However, there has been a major change in the attitude since it has been pointed out that the use of $b\bar{b}$ final states can be made possible using jet substructure methods [12]. I will give a short description of these methods in the later discussions.

3 Production of the Higgs at the LHC

Since LHC is a hadronic collider, one of the most relevant activity is accurate predictions of the expected cross-sections as well as differential distributions in important kinematical variables such as, eg., p_T^H for various Higgs production processes. QCD factorisation theorem at short distances tells us that this cross-section can be calculated in the following formalism:

$$\sigma(pp \to X + ..) = \sum_{a,b} \int_0^1 dx_1 dx_2 f_a(x_1, \mu_F^2) f_b(x_2, \mu_F^2) \\ \times \sigma(a + b \to X) \left(x_1, x_2, \mu_R^2, \alpha_s(\mu_R^2), \alpha(\mu_R^2), \frac{Q^2}{\mu_R^2}, \frac{Q^2}{\mu_F^2} \right)$$
(1)

An accurate calculation requires precise inputs on two non perturbative quantities α_s and the parton density functions, PDF's, along with an accurate evaluation of the subprocess cross-sections. An enormous amount of work has been done on the subject. An evaluation of the theoretical uncertainties in the predictions for the Higgs cross-section at the LHC was presented in [13]. In fact a joint collective effort [14], involving experimentalists and theorists, has been made recently to make the most accurate predictions for total observable cross-sections, taking into account most of the current theoretical uncertainties, both in the calculation of production cross-sections and the branching ratios. The next step is to do the same for exclusive distributions. Since all the experimental searches necessarily require cuts on the phase space variables, predictions for exclusive distributions for both the signal and dominant background are extremely crucial. Below, I summarise the main features and refer the reader to [13, 14] and references therein for further details.

The most important mode of production at the LHC is the gg fusion, dominated by the top loop. The next to leading order (NLO) corrections have been computed both in the Effective Field Theory (EFT) approach in the limit of infinite top mass and for finite heavy quark mass. Further, the next to next leading order (NNLO) corrections have been computed doing the three loop calculation. On top of it, the resummation of soft and collinear corrections has been performed at the next to next to leading logs (NNLL). The non factorisable EW and QCD corrections to the process have also been computed and shown to be \sim 5%. The K-factor, defined as the ratio to the leading order (LO) crosssection, for the dominant gg fusion process for low Higgs masses, at the LHC, is 1.7 at NLO and grows to about 2 at NNLO, thus showing a good convergence of the perturbation series. The NNLO result has small dependence on the renormalisation and factorisation scale variations, the hallmark of stability of a perturbative QCD calculation. The cleanest prediction is for the WH/ZH production, where both the QCD and EW corrections have been computed and the resulting cross-section has a K-factor $\sim 1.2 - 1.3$ at NNLO. The WW/ZZ fusion mechanism has the second largest cross-section at the LHC and would be very important for coupling/quantum number measurements once the Higgs boson has been discovered. In this case, the extraction of the signal for precision measurements requires extensive cuts on the phase space and hence calculation of higher order corrections to exclusive distributions is very important. Both the QCD and EW corrections have been computed and the K-factors are found to be modest. Equally important for the measurements of the couplings is the the ttH production. Use of jet substructure method [15] may vet revive the utility of this channel. The NLO corrections to this $2 \rightarrow 3$ processes are now available and the scale variation for the NLO result for $\sigma(pp \rightarrow t\bar{t}H)$ at the LHC is found to be rather modest ~ 10–20%. The two panels of Fig. 5 summarise the state of art predictions for the Higgs production cross section in gg fusion for $\sqrt{s} = 14$ TeV from [13] and $\sqrt{s} = 7$ TeV for all the relevant production processes from [14].



Figure 5: Cross-sections for the gg fusion process with all errors [13] for $\sqrt{s} = 14$ TeV in the left panel and cross-sections for all the relevant processes for $\sqrt{s} = 7$ TeV [14] in the right panel.

It is worth mentioning here that the situation about the theoretical uncertainties in the production cross-section of the gg fusion process at the Tevatron [10,16] is quite different. In fact this process is observable at the Tevatron only because of the rather large NLO/NNLO corrections it receives corresponding to a K-factor, defined as the ratio of the result of the higher order calculation to that of the LO, of 2(3) at NLO (NNLO). This thus means that the range of variation of the common factorisation and renormalisation scales in this case has to be somewhat larger than for the LHC case, possibly leading to a larger scale variation uncertainty in the cross-section. Further, the different parametrisations for the PDF's which correspond to different assumptions on these nonperturbative inputs, can differ in the central value of the predicted cross-section [10, 11] by up to 40% for Higss masses where the sensitivity is maximal. The left panel in the Fig. 6 taken from [10] shows the Higgs production cross section $\sigma_{gg \to H}^{NNLO}$ at the Tevatron with the uncertainty band when all theoretical uncertainties are added as suggested in Ref. [16]. This uncertainty is compared the uncertainties quoted by the CDF and D0 experiments [9] as well as the uncertainty that results when the LHC procedure as suggested [14] is adopted. In the insert, the relative size of the uncertainties compared to the central value are shown. Thus we see that if one were to evaluate the theoretical uncertainties for the Tevatron by the method prescribed in [14] one would get about 35% uncertainty in the cross-section as opposed to the 20% and 10% assumed in the CDF and D0 analyses respectively [9]. These differences as well as the strong dependence of the normalisation of the Higgs cross-section on the PDF, suggest that one should critically evaluate the dependence of the exclusion bounds from the Tevatron shown in Fig. 3, on the same. In fact analysis of Ref. [10] shows that if the true normalisation is indeed smaller by $\sim 40\%$ than that for the used MSTW parameterisation, then one might need up to a factor 2 higher luminosity to achieve the same exclusion. This situation is illustrated in the right hand panel (b) of this figure where the needed luminosity to recover the current sensitivity, if the cross-section should be lowered by 20% and 40%, is indicated. This underlies the importance of having a complete assessment of the theoretical uncertainties as is presented in [10, 13, 14, 16].



Figure 6: The needed luminosity by the CDF experiment to recover the current sensitivity (with 5.9 fb⁻¹ data) when the $gg \to H \to \ell\ell\nu\nu$ signal is lowered by 20% and 40% and with a ±10% change in the dominant $p\bar{b} \to WW$ background, taken from [10].

4 LHC: projections and results

As said before at the LHC, gg fusion is the dominant production mechanism and the final state contributing to the discovery depends on the mass of the Higgs. Fig. 7 taken from the ATLAS TDR [4], shows the signal significance that one expected to achieve at the originally planned 14 TeV LHC, for an integrated luminosity of 100 fb⁻¹, neglecting all the K-factors. This corresponds to the assertion that used to be made that a single experiment can discover the Higgs over the entire mass range allowed by theoretical considerations at 5σ at the 14 TeV LHC.



Figure 7: The expected signal significance for different search channels at the LHC with 14 TeV, assuming no K-factors, for 100 fb^{-1} integrated luminosity. Taken from ATLAS TDR in [4].

Now the LHC has been running at the lower energy of 7 TeV, at a lower luminosity than planned but has already collected 35 pb^{-1} data per experiment thanks to the very good performance of the LHC machine. It will now continue to run at 7 TeV till end 2012.



Figure 8: Examples of the results for the SM Higgs and MSSM Higgs available from the 35 pb^{-1} luminosity at LHC taken from [18] and [19] respectively.

Plots in Fig. 8 show that even with the very small amount of data the LHC has started giving significant results. The left panel shows that the ATLAS collaboration is getting close to being sensitive to the SM Higgs in the mass range around 160 GeV and has put limits on the cross-section ,at 95% CL, of about 1.2 times the SM cross-section, for Higgs mass around 160 GeV. Clearly, one has to watch this space now closely for future news.

For the CMS results I have chosen the example of the SUSY Higgs about which I have not talked much in sections 2 and 3. Supersymmetry is one of the most popular and arguably the best motivated BSM physics candidate. In the MSSM [17] there exist 5 Higgs bosons, three neutrals and two charged, one of the three neutrals being a pseudoscalar. An important difference from the SM case is that the lightest Higgs mass is now constrained from above (~ 130–140 GeV) and the Higgs mass bound depends on some details of the specific SUSY model and the parameters thereof. The heavier neutral Higgs bosons decay mostly into b and τ 's and thus the phenomenology is quite distinct. The production cross-section for the inclusive production of the supersymmetric Higgs in the process $gg, b\bar{b} \rightarrow H$ with $H \rightarrow \tau \tau$, is considerably enhanced at large tan β citchiggs-review and is thus accessible even with low luminosity. The exclusion for the supersymmetric Higgs achieved by the CMS experiment is shown in the figure in the right hand panel. The ATLAS exclusion for the same seen in [18] is similar. These results have already led to theoretical analyses of their implications not just for SUSY searches in general but also for the search of a light, SM higgs at the LHC [22].

The LHC experiments seem to be performing amazingly well and the time gap between data taking and availability of results is indeed very short. It is therefore important to know what are their projections now for the Higgs searches. For more details I refer the readers to information available on the web pages in Refs. [20] and [21] respectively. The left panel of Fig. 9 shows the luminosity required for 5 (3) σ Higgs discovery and Higgs exclusion at 95% CL at centre of mass energy of 7 and 8 TeV respectively, whereas the right panel shows the CMS version of the plot of Fig. 7 but now for $\sqrt{s} = 7$ and 8 TeV, for few selected values of integrated luminosities. These figures show clearly that depending on the luminosity the LHC machine manages to deliver, we



Figure 9: ATLAS simulation for the required integrated Luminosity for exclusion at 95% CL and discovery at 3 and 5 σ level [20] (left panel) and the expected level of significance of observation at different integrated luminosities from CMS simulation [21], as a function of M_H . Results are shown for both $\sqrt{s} = 7$ and 8 TeV.

would have very significant information on the SM Higgs mass by the end of the 2012 run. This makes now for a very agonising wait indeed.

5 Determination of Higgs properties and couplings

As already stated, just discovering the Higgs at a particular mass and the simultaneous results from the associated searches for BSM physics, will begin to give indicative answers about whether the SM is a strongly coupled theory with a composite Higgs boson or a weakly coupled theory with and elementary Higgs boson. But for a good scrutiny of the gauge group representation to which the Higgs belongs and whether it is 'the' SM Higgs, measurements of its couplings to the other SM particles, determination of its spin and further determination of its CP property is quite essential. The standard wisdom [3] in this respect was that these are usually high luminosity measurements. For example, the studies of Ref. [23] had shown that with an integrated luminosity of about 600 fb⁻¹, at 14 TeV, it will be possible to measure various couplings at ~ 20–30% level for the SM Higgs. These results were confirmed with a more sophisticated analysis recently [24].

Another example is of investigations of Ref. [25] which indicated that at 14 TeV LHC, one would be able to establish some of the anomalous (CP violating) HZZ couplings at 3–5 σ level, with 100–300 fb⁻¹ integrated luminosity, if these couplings were of the same order of magnitude as the SM couplings. In Fig. 10 taken from Ref. [25], are shown regions in the |c|-a coupling plane that can be probed by just a measurement of the width of the Higgs boson. Here the HZZ vertex has been parametrized in the most general model independent way given by:

$$V_{HZZ}^{\mu\nu} = \frac{igm_Z}{\cos\theta_W} \left[a g_{\mu\nu} + b \frac{p_\mu p_\nu}{m_Z^2} + c \epsilon_{\mu\nu\alpha\beta} \frac{p^\alpha k^\beta}{m_Z^2} \right],\tag{2}$$

in obvious notation for the different quantities appearing therein, p, k representing the sum and difference of the four momenta of the two Z bosons.



Figure 10: The number of standard deviations from the SM which can be obtained in the process $gg \rightarrow H \rightarrow Z^*Z^* \rightarrow 4$ leptons, as a scan over the (a, |c|) plane. The Higgs mass has been chosen to be 150 GeV (*left*) and 200 GeV (*right*). The white region is where the deviation from the SM is less than 3σ ; in the light blue/light Grey region the deviation is between 3σ and 5σ ; while for the dark blue/dark Grey region the deviation is greater than 5σ for an integrated luminosity of 300 fb⁻¹, taken from [25].



Figure 11: Distribution in M_{Z^*} for $H \to ZZ^*$ taken from [26].

It was also demonstrated in [26] that for a Higgs heavy enough to have a reasonable branching ratio in the ZZ^* channel, the shape of the distribution in the invariant mass of the $\ell^+\ell^-$ pair coming from the Z^* decay, can give clear information about the spin of the Higgs boson. The plot in Fig. 11 taken from Ref. [26], shows the measurement possible for an integrated luminosity of 100 fb⁻¹, at $\sqrt{s} = 14$ TeV, the histogram showing the expected statistical error. It had also been shown that the distribution in the azimuthal angle between the planes of the two pairs of the decay leptons can also carry information about the spin and the parity of the decaying resonance [27]. Recently there were investigations [28,29] which showed that more complicated, multivariate analyses might be able to do the job of establishing the J^{PC} to be 0^{++} with ~ 3σ significance for ≤ 10 fb⁻¹ luminosity. Apart from the high luminosity, for the coupling measurements use of the bb final state is also essential and so is the possibility to make a good measurement of the $t\bar{t}H$, $H \rightarrow b\bar{b}$ process. The $t\bar{t}jj$ background seems to make the use of this channel very difficult [4]. Hence any methods to improve the visibility of this channel are welcome. As mentioned before, methods using substructure of jets have given new hope in both these issues [12, 15]. I would therefore describe briefly this method.

The idea here is based on the fact that for high p_T Higgs bosons, the $b\bar{b}$ decay products would emerge close to each other and hence will look in the detector to be a single, fat jet with large invariant mass. The jets produced by QCD emission will not have this feature. Thus if one can develop an algorithm to see if a fat, heavy jet is made of two fast objects emitted close to each other, one can then reduce the QCD backgrounds to a low level. For production processes like WH, ZHwhere one has to select large p_T Higgs bosons to get rid of the irreducible SM background anyway, this technique seems to work quite nicely. The left panel in Fig. 12 shows a cartoon which illustrates this kinematical fact and in the right panel is shown a plot from [12], which indicates the clean way in which the signal can be separated from the background for the WH case and S/\sqrt{B} as high as 4.5 can be reached in this channel, with $b\bar{b}$ final state, for 30 fb⁻¹ of integrated luminosity for a 120 GeV Higgs boson. These kinds of studies would be the future of Higgs physics at the LHC once it has been discovered through any one channel.



Figure 12: Left panel shows a cartoon of a 'fat' jet from the $b\bar{b}$ decay of a large P_T Higgs and the right panel shows how the substructure analysis can help increase the S/\sqrt{B} in the WH channel, taken from [12].

6 Conclusion

Thus we are now at a very exciting stage where in the next two years we should expect either a 3σ signal or a 95% exclusion over a very large range of the Higgs masses at the LHC. As said already the mass of the Higgs boson alone can give completely non trivial indications of the presence or absence of BSM physics. Should the Higgs mass give an indication of the BSM physics, in most cases the corresponding BSM physics should also reveal itself in the direct searches which would be taking place alongside, for the value of the integrated luminosity we expect to have. Hence, the next two years of the Higgs physics at the LHC should be very exciting indeed. Precision

measurements of the couplings, spin, parity, CP characteristic and determination of triple Higgs couplings all has to however, wait for higher luminosities, higher energies and perhaps even for a leptonic collider [30].

7 Acknowledgments

I wish to acknowledge the Department of Science and Technology of India, for financial support under the J.C. Bose Fellowship scheme under grant no. SR/S2/JCB-64/2007. I would like to thank Abdelhak Djouadi not only for an enjoyable collaboration but also for a careful reading of the manuscript and useful comments.

References

- P.W. Higgs, Phys. Lett. **12** (1964) 132; and Phys. Rev. **145** (1966) 1156; F. Englert and R. Brout, Phys. Rev. Lett. **13** (1964) 321; P. W. Higgs, Phys. Rev. Lett. **13** (1964) 508; G.S. Guralnik, C.R. Hagen and T.W. Kibble, Phys. Rev. Lett. **13** (1964) 585.
- [2] J. Goldstone, A. Salam and S. Weinberg, Phys. Rev. **127** (1962) 965; S. Weinberg, Phys. Rev. Lett. **19** (1967) 1264; S.L. Glashow, S. Weinberg, Phys. Rev. Lett. **20** (1968) 224; A. Salam, Proceedings Of The Nobel Symposium, Stockholm 1968, ed. N. Svartholm.
- [3] For a review, see for example, A. Djouadi, Phys. Rept. 457 (2008) 216 [hep-ph/0503172], Phys. Rept. 459 (2008) 1-241, [hep-ph/0503173] and references therein; A. Djouadi, R. M. Godbole, in "Physics at the Large Hadron Collider", Ed. A. Datta, B. Mukhopadhyaya and A. Raychaudhuri, Springer (India), New Delhi, pp. 47-74 [arXiv:0901.2030 [hep-ph]] and references therein.
- [4] ATLAS Collaboration, Technical Design Report, CERN-LHCC-14 and CERN-LHCC-15; CMS Collaboration, Technical Design Report, CMS-LHCC-2006-21.
- [5] T. Hambye and K. Riesselmann, Phys. Rev. D55, 7255 (1997).
- [6] B.W. Lee, C. Quigg and H.B. Thacker, Phys. Rev. D16 (1977) 1519.
- [7] C.H. Llewellyn Smith, Phys. Lett. B46 (1973) 233; J. S. Bell, Nucl. Phys. B60 (1973) 427
 ; J. Cornwall et al, Phys. Rev. Lett. 30 (1973) 1268, [Erratum-ibid. **31**, 572 (1973)]; Phys. Rev. D10 (1974) 1145, [Erratum-ibid. D 11 (1975) 972]; S. D. Joglekar, Annals Phys. **83**, 427 (1974).
- [8] LEP working group for Higgs searches, R. Barate et al., Phys. Lett. B 565 (2003) 61.
- [9] CDF and D0 collaborations, Phys. Rev. Lett. 104 (2010) 061802; updated in arXiv:1007.4587 [hep-ex].
- [10] J. Baglio, A. Djouadi, S. Ferrag, R. M. Godbole, [arXiv:1101.1832 [hep-ph]], to appear in Phys. Lett. B.

- [11] S. Alekhin, J. Blumlein and S. Moch, arXiv:1101.5261;
 S. Moch, talk at Kick-off meeting of the LHCPhenoNet, Feb. 2011, Valencia, (pp 8-9) http://indico.ific.uv.es/indico/contributionDisplay.py?contribId=5&sessionId=2&confId=339.
- [12] One of the pioneering work in this area is J. M. Butterworth, A. R. Davison, M. Rubin, G. P. Salam, Phys. Rev. Lett. 100 (2008) 242001. [arXiv:0802.2470 [hep-ph]].
- [13] J. Baglio, A. Djouadi, JHEP **1103**, 055 (2011), [arXiv:1012.0530 [hep-ph]].
- [14] S. Dittmaier et al, "Handbook of LHC Higgs cross sections", arXiv: 1101.0593 [hep-ph]
- [15] T. Plehn, G. P. Salam, M. Spannowsky, Phys. Rev. Lett. 104, 111801 (2010). [arXiv:0910.5472 [hep-ph]].
- [16] J. Baglio, A. Djouadi, JHEP 1010, 064 (2010). [arXiv:1003.4266 [hep-ph], arXiv:1009.1363 [hep-ph]].
- [17] For a pedagogic introduction see e.g., M. Drees, R.M. Godbole and P. Roy, "Theory and phenomenology of sparticles", World Scientific, 2005; H. Baer and X. Tata, "Weak scale Supersymmetry: From superfields to scattering events", Cambridge University Press, 2006.
- [18] ATLAS Collaboration, ATLAS-CONF-2011-005 for the SM Higgs search. For further information see also https://twiki.cern.ch/twiki/bin/view/AtlasPublic/HiggsPublicResults#Data_2011.
- [19] CMS collaboration, CMS PAS HIG-10-002, for more information see also, https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResults.
- [20] ATLAS Collaboration, ATL-PHYS-PUB-2011-001, for more inforamtion see also, https://twiki.cern.ch/twiki/bin/view/AtlasPublic/PublicHGPlotsAtlPhysPub2011001 and https://twiki.cern.ch/twiki/bin/view/AtlasPublic/PublicHGPlotsAtlPhysPub2010726
- [21] CMS-NOTE-2010/008, for more information see also, https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsHIGStandardModelProjections.
- [22] J. Baglio and A. Djouadi, arXiv:1103.6247 [hep-ph].
- [23] M. Duhrssen, S. Heinemeyer, H. Logan, D. Rainwater, G. Weiglein, D. Zeppenfeld, Phys. Rev. D70, 113009 (2004), [hep-ph/0406323].
- [24] R. Lafaye, T. Plehn, M. Rauch, D. Zerwas, M. Duhrssen, JHEP 0908, 009 (2009).
- [25] R. M. Godbole, D. J. Miller, 2, M. M. Muhlleitner, JHEP 0712, 031 (2007). [arXiv:0708.0458 [hep-ph]].
- [26] D. J. Miller, 2, S. Y. Choi, B. Eberle, M. M. Muhlleitner, P. M. Zerwas, Phys. Lett. B505, 149-154 (2001). [hep-ph/0102023].
- [27] S. Y. Choi, D. J. Miller, 2, M. M. Muhlleitner, P. M. Zerwas, Phys. Lett. B553, 61-71 (2003). [hep-ph/0210077].

- [28] A. De Rujula, J. Lykken, M. Pierini, C. Rogan, M. Spiropulu, Phys. Rev. D82, 013003 (2010). [arXiv:1001.5300 [hep-ph]].
- [29] Y. Gao, A. V. Gritsan, Z. Guo, K. Melnikov, M. Schulze, N. V. Tran, Phys. Rev. D81, 075022 (2010). [arXiv:1001.3396 [hep-ph]].
- [30] G. Aarons et al. [ILC Collaboration], [arXiv:0709.1893 [hep-ph]].