

Electroweak Precision Observables in a Fourth Generation Model with General Flavour Structure

P. GONZÁLEZ^{a*} J. ROHRWILD^{a†} M. WIEBUSCH^{b‡}

^a*Institute for Theoretical Particle Physics and Cosmology,
RWTH Aachen, D-52056 Aachen, Germany*

^b*Institute for Theoretical Particle Physics,
Karlsruhe Institute of Technology (KIT), D-76128 Karlsruhe, Germany*

Abstract

We calculate the contributions to electroweak precision observables (EWPOs) due to a fourth generation of fermions with the most general (quark-) flavour structure (but assuming Dirac neutrinos and a trivial flavour structure in the lepton sector). The new-physics contributions to the EWPOs are calculated at one-loop order using automated tools (FeynArts/FormCalc). No further approximations are made in our calculation. We discuss the size of non-oblique contributions arising from Z -quark-anti-quark vertex corrections and the dependence of the EWPOs on all CKM mixing angles involving the fourth generation. We find that the electroweak precision observables are sensitive to two of the fourth-generation mixing angles and that the corresponding constraints on these angles are competitive with those obtained from flavour physics. For non-trivial 4×4 flavour structures, the non-oblique contributions lead to relative corrections of several permille and should be included in a global fit.

*email: gonzalez@physik.rwth-aachen.de

†email: rohrwild@physik.rwth-aachen.de

‡email: martin.wiebusch@kit.edu

1 Introduction

With the advent of LHC data the first direct tests for many models of new physics are within reach. Among the conceptionally simplest extensions of the Standard Model (SM3) are those which only add a minimal set of fermions to the SM particle content. This class encompasses both the additional vector-like quarks [1–4] and the fourth-generation scenario (SM4).

The SM4 was fairly popular in the 1980s until electroweak precision observables seemed to rule it out. In the last years models with an additional fourth generation experienced a renaissance as new analyses, e.g. [5–12], somewhat relaxed the electroweak tensions. This realisation also prompted numerous studies of the non-trivial flavour structure of the SM4 [13–17], as well as searches for specific signatures in new physics observables [18–24]. Currently, the progress of the LHC provides first direct constraints on new SM-like heavy quarks. CMS gives lower mass limits of 450 GeV for a t' decaying into bW [25] and 495 GeV for a b' decaying into tW [26]. Further constraints come from direct Higgs searches at the LHC as the presence of additional heavy quarks would dramatically increase the Higgs production cross section [27, 28]. However, this constraint can be avoided if the fourth generation neutrino is light enough to allow for an additional invisible decay channel of the Higgs boson [29–33].

Recently, some effort has been directed towards providing an actual *fit* of the parameters of the model. One of the first attempts in this direction primarily used the electroweak precision observables and restricted itself to only one CKM parameter [34]; still non-trivial correlations were found, for example, between the Higgs mass and the new mixing angle. More recent studies seek to constrain [35] or even determine [36] the full 4×4 CKM matrix. In this case the main challenge is the fact that, if one allows for a generic CKM structure, the flavour and electroweak sector are intertwined and have to be treated simultaneously.

Usually the effects of new physics in the electroweak sector are parametrised by the *oblique electroweak parameters* S , T and U , as introduced by Peskin and Takeuchi [37, 38]. These allow for fairly simple and straightforward estimates of new physics contribution to electroweak observables. However, the validity of this parametrisation relies on certain assumptions about the new physics model, which are, in principle, no longer satisfied in an SM4 with the most general flavour structure. Expressions for the leading non-oblique contributions were given in [15] for the special case where only the third and fourth generation quarks are allowed to mix.

In this letter we discuss the contributions to electroweak precision observables (EWPOs) due to a fourth generation with general 4×4 flavour mixing. For the sake of simplicity we assume Dirac neutrinos and a trivial flavour structure in the lepton sector. Our method for the computation of EWPOs in the SM4 uses a high level of automatisisation and can easily be used for other models. We break up the observables into ‘SM3 parts’ and ‘new physics contributions’. The former can be computed with

well-established programs like `ZFITTER` [39, 40]. The latter we calculate to one-loop order using `FeynArts/FormCalc` [41–43]. Unlike most other literature on EWPOs in the SM4, no further approximations are made in this paper. Our calculation thus includes all one-loop non-oblique contributions, i.e. those which are not captured by the S , T and U parameters. We discuss the importance of the non-oblique contributions and the impact of flavour mixing between the fourth and the first three generation in several SM4 scenarios.

In section 2 we briefly review the oblique parameters and their range of applicability. In section 3 we introduce our notations for the SM4 parameters and explain our method for calculating the corrections to the EWPOs. In section 4 we describe our treatment of the Fermi constant G_F , which is an observable and not a parameter in our analysis. Our numerical results are presented in section 5. We find that the EWPOs are equally sensitive to all three fourth-generation mixing angles and that the corresponding constraints on these angles are competitive with those obtained from flavour physics. For non-trivial 4×4 flavour structures, the non-oblique contributions lead to relative corrections of up to one permille for the hadronic Z width Γ_{had} and of several permille for the hadronic $Z \rightarrow b\bar{b}$ branching ratio R_b . A simultaneous fit of the SM4 masses, couplings and CKM matrix should therefore take into account all six SM4 CKM mixing angles and the non-oblique corrections to the EWPOs. We conclude in section 6.

2 Oblique Corrections and Electroweak Observables

The constraints imposed on new physics by EWPOs measured at LEP have already been discussed extensively in the literature. In 1992 Peskin and Takeuchi presented a model-independent way of parametrising the new physics contributions to the Z pole observables [37]. Their analysis was based on three assumptions:

1. The electroweak gauge group of the new-physics model is $SU(2)_L \times U(1)_Y$.
2. The new-physics couplings to light fermions (i.e. all SM3 fermions except the top-quark) are negligible.
3. The scale of new physics is much larger than the electroweak scale.

The first assumption forbids the existence of additional gauge bosons coupling directly to leptons. The second assumption guarantees that there are no additional vertex or box-diagrams contributing to the Drell-Yan process. Thus, the only way the new physics contribute to the Z pole observables is through the renormalisation of weak gauge boson wave functions, the electric charge or the Weinberg angle. The third assumption is needed to justify a step in the discussion in [37], where the gauge boson self-energies are expanded to first order around $q^2 = 0$ (q being the momentum flowing through the

	experiment	theory (SM3)
$\Gamma_{\text{had}} [\text{GeV}]$	1.7444 ± 0.002	1.7418 ± 0.0009
R_b	0.21629 ± 0.00066	0.21578 ± 0.00005
A_{FB}^b	0.0992 ± 0.0016	0.1034 ± 0.0007
\mathcal{A}_b	0.923 ± 0.020	0.9348 ± 0.0001
\mathcal{A}_e	0.15138 ± 0.00216	0.1475 ± 0.0010
$M_W [\text{GeV}]$	80.420 ± 0.031	80.384 ± 0.014

Table 1: Experimental results and Standard Model predictions for selected electroweak observables. All numbers were taken from [45].

self-energy graphs). In practice, it is usually sufficient to require that new particles coupling directly to weak gauge bosons are heavier than the Z boson.

In SM extensions that satisfy the criteria above, the new physics contributions to the Z pole observables can be expressed in terms of the *oblique electroweak parameters* S , T and U which were defined in [37] and represent different linear combinations of gauge boson self-energies and their derivatives. On the experimental side, the values of S , T and U can then be determined from data by performing a global fit of S , T , U and the SM3 parameters to the Z pole and possibly other low-energy observables. (See [44] for a recent analysis of this type.) On the theoretical side one can test to what extent a given model of new physics agrees with low-energy observables by computing S , T and U in this model and comparing the results with the best-fit values.

This method of testing an SM extension against constraints from low-energy experiments is very convenient since it only requires the computation of three quantities. It has been applied to a number of models including the SM4 [10]. One should, however, keep in mind that the validity of this method depends on the validity of the assumptions listed above. In the SM4 the second assumption is no longer valid if the fourth generation quarks are allowed to mix with the quarks of the first three generations. Furthermore, if one allows for a fourth-generation neutrino with a mass just above $m_Z/2$ the third assumption is no longer satisfied.¹ Hence, the validity of the “oblique method” must be checked explicitly if one attempts to constrain the new mixing angles of the SM4 CKM matrix.

3 The $Zq\bar{q}$ Vertex in the SM4

The properties of the Z boson and its couplings to fermions have been measured at LEP 1 with a very high accuracy. Table 1 shows the experimental values and accuracies for

¹If the first two assumptions are still valid, the case of new physics near the electroweak scale can be handled by introducing three additional oblique parameters. This was discussed in [38].

a selection of Z -pole observables as well as their theoretical predictions within the SM3. The observables are: the partial width for $Z \rightarrow \text{hadrons}$ (Γ_{had}), the hadronic branching fraction for $Z \rightarrow b\bar{b}$ (R_b), the forward-backward asymmetry for $Z \rightarrow b\bar{b}$ (A_{FB}^b) and the mass of the W (M_W). In the Z -pole approximation, the forward-backward asymmetry can be written as $\frac{3}{4}\mathcal{A}_e\mathcal{A}_b$, where the quantities \mathcal{A}_e and \mathcal{A}_b only depend on the Ze^+e^- and $Zb\bar{b}$ couplings, respectively. The relative precision of Γ_{had} is approximately 0.1% and R_b is known to an accuracy of 0.3%. The measured value of A_{FB}^b deviates from its SM3 prediction by more than two standard deviations. The discrepancy originates mainly from the factor \mathcal{A}_e . Oblique corrections due to a fourth generation of fermions affect all Z -pole observables, but only observables related to the Z -quark-anti-quark vertex are subject to non-oblique corrections; of the observables from table 1, only Γ_{had} , R_b and \mathcal{A}_b receive non-oblique contributions. Our discussion will therefore mainly focus on these quantities.²

Within the SM3 the couplings of Z bosons to quarks have been studied in great detail. Electroweak and QCD corrections to the gauge boson self-energies and the Z -quark-anti-quark vertex have been calculated at two-loop order [46–65] and the results have been implemented in public codes such as TOPAZO [66] or ZFITTER [39, 40]. Radiative corrections to the partial widths are of the order of 0.1% (QED) and 4% (QCD). To match the experimental accuracy of the Z -pole observables they must therefore be included in theoretical calculations. In this section we explain how predictions for the Z pole observables within the SM4 can be calculated at the required level of accuracy without the need to re-visit the SM3 calculations.

Before we begin, let us briefly explain our notations for the SM3 and SM4 parameters. For the SM3 CKM matrix we use the *standard parametrisation*. In this parametrisation the independent parameters are the three mixing angles θ_{12} , θ_{13} and θ_{23} and one complex phase δ_{13} . The explicit form of the SM3 CKM matrix in terms of the phase and mixing angles is given in appendix A.

In the SM4 the CKM matrix is a unitary 4×4 matrix. After absorbing unphysical complex phases into the definitions of the quark fields, its parametrisation requires only three additional mixing angles θ_{14} , θ_{24} and θ_{34} and two additional complex phases δ_{14} and δ_{24} . The explicit form of the SM4 CKM matrix is also given in appendix A. For the discussion below it is only important to know that for $\theta_{14} = \theta_{24} = \theta_{34} = \delta_{14} = \delta_{24} = 0$ the SM4 CKM matrix assumes a block-diagonal form with the SM3 CKM matrix in the first 3×3 block and a one in the last block.

To distinguish the phase δ_{13} and the mixing angles θ_{12} , θ_{13} and θ_{23} of the SM4 CKM matrix from their SM3 counterparts we will use superscripts ‘SM4’ and ‘SM3’, respectively. The same applies to other parameters like m_H or M_W , which exist in both models. We will also use the shorthands s_{ij} and c_{ij} for the sines and cosines of the mixing angles θ_{ij} . Finally, we denote the lepton, neutrino, up and down-type quark of

²The branching fraction R_c and asymmetry factor \mathcal{A}_c for the charm quark also receive non-oblique corrections, but these observables are less constraining due to their larger experimental error.

the fourth generation as ℓ_4 , ν_4 , t' and b' , respectively. Their masses m_{ℓ_4} , m_{ν_4} , $m_{t'}$ and $m_{b'}$ are independent parameters of the SM4.

Let us now proceed with the discussion of higher order corrections to the $Zq\bar{q}$ vertex. In the limit of vanishing external quark masses m_q , the on-shell $Zq\bar{q}$ vertex function only contains two Lorentz structures:

$$\Gamma_\mu^q = ie\gamma_\mu[F_V^q - F_A^q\gamma_5] \quad . \quad (1)$$

Here and in the following, $q = u, d, s, c, b$ denotes the quark flavour. The form factors F_V^q and F_A^q depend on the quark flavour, the external masses and the parameters of the model under consideration (SM3 or SM4). Following the discussion in [67], we express QCD and QED radiative corrections to F_V^q and F_A^q in terms of *radiator functions* \mathcal{R}_V^q and \mathcal{R}_A^q and write

$$F_V^q = g_V^q \sqrt{\mathcal{R}_V^q} \quad , \quad F_A^q = g_A^q \sqrt{\mathcal{R}_A^q} \quad . \quad (2)$$

In doing this, we neglect the non-factorisable contributions [68, 69], whose effect is below the permille level. The *effective couplings* g_V^q and g_A^q now only contain infrared finite contributions. At leading order $\mathcal{R}_V^q = \mathcal{R}_A^q = 1$ and g_V^q and g_A^q are the tree-level vector and axial couplings of the Z boson.

In this paper we are interested in the difference between predictions for Z pole observables within the SM3 and SM4. For this purpose we denote, for any quantity X , the *new physics correction* by

$$\delta X = X^{\text{SM4}} - X^{\text{SM3}} \quad , \quad (3)$$

where the superscripts ‘SM4’ and ‘SM3’ indicate that X is evaluated with a given set of SM4 or SM3 parameters, respectively. In principle, the two sets of parameters can be completely unrelated. It is, however, extremely convenient to use the same values of M_Z , M_W , m_t , α and α_s in both sets.³ In this case, $\delta\mathcal{R}_V^q = \delta\mathcal{R}_A^q = 0$ and the new physics corrections to any Z pole observable can be obtained by only computing the infrared finite quantities δg_V^q and δg_A^q . The form factors $F_V^{q,\text{SM4}}$ and $F_A^{q,\text{SM4}}$ (and thus for the Z -pole observables within the SM4) may then be calculated by scaling the corresponding SM3 form factors with the ratios $g_V^{q,\text{SM4}}/g_V^{q,\text{SM3}}$ and $g_A^{q,\text{SM4}}/g_A^{q,\text{SM3}}$, respectively. This way, factorisable QCD and QED corrections are included in F_V^{SM4} and F_A^{SM4} if they were included in the SM3 ‘reference values’ F_V^{SM3} and F_A^{SM3} . As we will see below, the ratios $\delta g_V^q/g_V^{q(0)}$ and $\delta g_A^q/g_A^{q(0)}$ (with $g_V^{q(0)}$ and $g_A^{q(0)}$ being the tree-level couplings) are typically below 1%. Thus, the approximation

$$F_V^{q,\text{SM4}} \approx F_V^{q,\text{SM3}} \left(1 + \frac{\delta g_V^q}{g_V^{q(0)}} \right) \quad , \quad F_A^{q,\text{SM4}} \approx F_A^{q,\text{SM3}} \left(1 + \frac{\delta g_A^q}{g_A^{q(0)}} \right) \quad (4)$$

³These are independent SM3 input parameters in the *on-shell* renormalisation scheme [70], which is the scheme we used in our calculations.

is generally valid with a relative precision of the order of 10^{-4} .

The difference between \mathcal{R}_V^q and \mathcal{R}_A^q is of the order of a few percent [67]. Thus, to estimate the size of the new physics contributions to the EWPOs we use the approximation

$$\mathcal{R}_V^q \approx \mathcal{R}_A^q \equiv \mathcal{R}^q \quad (5)$$

and obtain

$$\delta\Gamma_\mu^q = ie\mathcal{R}^q\gamma_\mu[\delta g_V^q - \delta g_A^q\gamma_5] \quad . \quad (6)$$

The hadronic Z partial widths and asymmetries are then given by

$$\Gamma(Z \rightarrow q\bar{q}) = \alpha M_Z \mathcal{R}^q [|g_V^q|^2 + |g_A^q|^2] \quad , \quad \mathcal{A}_q = \frac{2 \operatorname{Re} g_V^q \operatorname{Re} g_A^q}{(\operatorname{Re} g_V^q)^2 + (\operatorname{Re} g_A^q)^2} \quad . \quad (7)$$

The new physics corrections to these quantities are readily obtained by expanding the effective couplings to first order in δg_V^q and δg_A^q :

$$\begin{aligned} \frac{\delta\Gamma(Z \rightarrow q\bar{q})}{\Gamma^{\text{SM3}}(Z \rightarrow q\bar{q})} &= 2 \frac{\operatorname{Re}[g_V^{q(\text{SM3})*} \delta g_V^q] + \operatorname{Re}[g_A^{q(\text{SM3})*} \operatorname{Re} \delta g_A^q]}{|g_V^{q(\text{SM3})}|^2 + |g_A^{q(\text{SM3})}|^2} \\ &\approx 2 \frac{g_V^{q(0)} \operatorname{Re} \delta g_V^q + g_A^{q(0)} \operatorname{Re} \delta g_A^q}{(g_V^{q(0)})^2 + (g_A^{q(0)})^2} \quad , \end{aligned} \quad (8a)$$

$$\begin{aligned} \frac{\delta\mathcal{A}_q}{\mathcal{A}_q^{\text{SM3}}} &= \frac{\operatorname{Re} \delta g_V^q}{\operatorname{Re} g_V^{q(\text{SM3})}} + \frac{\operatorname{Re} \delta g_A^q}{\operatorname{Re} g_A^{q(\text{SM3})}} - 2 \frac{\operatorname{Re} g_V^{q(\text{SM3})} \operatorname{Re} \delta g_V^q + \operatorname{Re} g_A^{q(\text{SM3})} \operatorname{Re} \delta g_A^q}{(\operatorname{Re} g_V^{q(\text{SM3})})^2 + (\operatorname{Re} g_A^{q(\text{SM3})})^2} \\ &\approx \frac{\operatorname{Re} \delta g_V^q}{g_V^{q(0)}} + \frac{\operatorname{Re} \delta g_A^q}{g_A^{q(0)}} - 2 \frac{g_V^{q(0)} \operatorname{Re} \delta g_V^q + g_A^{q(0)} \operatorname{Re} \delta g_A^q}{(g_V^{q(0)})^2 + (g_A^{q(0)})^2} \quad . \end{aligned} \quad (8b)$$

Note that, as a result of approximating $\mathcal{R}_V^q \approx \mathcal{R}_A^q$, the radiator functions cancel in the ratios above.

If mixing between the fourth generation quarks and the quarks of the first three generations is neglected and the fourth-generation fermions are sufficiently heavy, the new physics corrections can be expressed in terms of the oblique electroweak parameters S , T and U [37, 38]. In this case, the relations between δg_V^q , δg_A^q and S , T and U are

$$\delta g_V^{q,(1)} = \frac{\alpha}{16c_W s_W^3} \left[2I_3^q S - 4[(c_W^2 - s_W^2)I_3^q + 2s_W^2 Q^q]T - \left(\frac{c_W^2 - s_W^2}{s_W^2} I_3^q + 2Q^q \right) U \right] \quad , \quad (9a)$$

$$\delta g_A^{q,(1)} = \frac{\alpha}{16c_W s_W^3} \left[2S - \frac{c_W^2 - s_W^2}{s_W^2} (4s_W^2 T + U) \right] I_3^q \quad , \quad (9b)$$

where Q^q and I_3^q are the electric charge and weak isospin of the quark q and s_W and c_W are the sine and cosine of the Weinberg angle, defined by $s_W^2 = 1 - M_W^2/M_Z^2$.

The superscript ‘(1)’ denotes the (electroweak) one-loop corrections to the effective couplings.⁴

If the fourth generation quarks are allowed to mix with the quarks of the first three generations one also needs to compute the vertex diagrams contributing to $\delta g_V^{q,(1)}$ and $\delta g_A^{q,(1)}$. We used the **FeynArts/FormCalc** package [41–43] to compute the required one-loop order diagrams. The renormalisation of the $Zq\bar{q}$ vertex was done in the on-shell scheme.⁵ Only diagrams involving W bosons, charged Goldstone bosons or Higgs bosons contribute to $\delta g_V^{q,(1)}$ and $\delta g_A^{q,(1)}$, as long as α , α_s , M_Z , M_W and m_t are chosen to be the same in the SM3 and SM4. The SM3 parameters and corresponding values for $\Gamma(Z \rightarrow q\bar{q})$ and \mathcal{A}_q were taken from [45]. Specifically, we use

$$\begin{aligned} 1/\alpha(m_Z) &= 128.892 \quad , \quad \alpha_s(m_Z) = 0.1185 \quad , \quad M_Z = 91.1875 \text{ GeV} \quad , \\ M_W^{\text{SM3}} &= 80.384 \text{ GeV} \quad , \quad m_t = 173.2 \text{ GeV} \quad , \quad m_H^{\text{SM3}} = 90 \text{ GeV} \quad , \\ \Gamma_{\text{had}}^{\text{SM3}} &= 1.7418 \text{ GeV} \quad , \quad R_b^{\text{SM3}} = 0.21578 \quad , \quad \mathcal{A}_b^{\text{SM3}} = 0.9348 \quad , \end{aligned} \quad (10)$$

where

$$\Gamma_{\text{had}} = \sum_{q=u,d,s,c,b} \Gamma(Z \rightarrow q\bar{q}) \quad , \quad R_q = \frac{\Gamma(Z \rightarrow q\bar{q})}{\Gamma_{\text{had}}} \quad . \quad (11)$$

The phase and mixing angles of the SM3 CKM matrix were also taken from [45]:

$$\theta_{12}^{\text{SM3}} = 0.2273 \quad , \quad \theta_{13}^{\text{SM3}} = 0.003466 \quad , \quad \theta_{23}^{\text{SM3}} = 0.04103 \quad , \quad \delta_{13}^{\text{SM3}} = 1.2020 \quad . \quad (12)$$

Note that the numerical values for $\Gamma_{\text{had}}^{\text{SM3}}$ and R_b^{SM3} are for a fixed “reference” Higgs mass $m_H^{\text{SM3}} = 90 \text{ GeV}$. In the SM4 the Higgs mass is treated as a free parameter.

4 A Note on G_F

As mentioned above, we use in this work the on-shell renormalisation scheme for the computation of new physics corrections. In this scheme, the quantities $\alpha(M_Z)$, M_Z and M_W are independent parameters. This parametrisation is very convenient for the computation of higher order corrections, but it has its disadvantages if one wants to compare it with experimental data. The Fermi constant G_F , which is determined from the muon lifetime, is a non-trivial function of $\alpha(M_Z)$, M_Z , M_W and the other model parameters. Since G_F is measured very accurately (namely, to a relative precision of 10^{-5}) it constrains the model to a non-trivial hyper-surface in its parameter space. In other words, one parameter of the model is fixed by the requirement that G_F assumes

⁴If $\alpha(M_Z)$, M_Z and M_W are the same in the SM3 and SM4, the new physics only enters at one-loop order. In the next section we deal with the case where a different value of M_W is chosen in the two models.

⁵See e.g. [71] for a detailed description.

its measured value. Typically, one adjusts the value of M_W to obtain the correct value of G_F .

The relation between G_F and M_W is conventionally written as [72]

$$G_F = \frac{\pi\alpha}{\sqrt{2}s_W^2 M_W^2} \frac{1}{1 - \Delta r} \quad , \quad (13)$$

where Δr encodes higher order corrections and is, in general, a function of all other parameters. New physics, like the existence of a fourth generation of fermions, changes the function Δr . Denoting, as before, the new physics correction to Δr as $\delta\Delta r$ and writing the solutions of (13) in the SM3 and SM4 as M_W^{SM3} and $M_W^{\text{SM4}} \equiv M_W^{\text{SM3}} + \delta M_W$, respectively, we find

$$\frac{\delta M_W}{M_W^{\text{SM3}}} = -\frac{s_W^2}{2(c_W^2 - s_W^2)} \delta\Delta r \quad . \quad (14)$$

However, since the parameters (10) already satisfy the G_F constraint we have $M_W^{\text{SM3}} = M_W$ with M_W from (10). If the SM4 is to agree with the measured value of the W mass, the ratio $\delta M_W/M_W^{\text{SM3}}$ cannot be much larger than one permille. Hence, the shift in M_W is unimportant for the purpose of computing $\delta\Delta r$ and the loop corrections to the effective couplings g_V^q and g_A^q . We can therefore safely use the value from (10) in these calculations. However, the change in M_W also affects the tree-level $Zq\bar{q}$ couplings $g_V^{q,(0)}$ and $g_A^{q,(0)}$, since $\sin^2\theta_W$ is defined as $1 - M_W^2/M_Z^2$ in the on-shell scheme. This effect is of the same order as the loop corrections to the effective couplings and must be included in the new-physics corrections δg_V^q and δg_A^q . This can be achieved by making the following substitution in (8):

$$\delta g_{V,A}^q = g_{V,A}^{q,(0)}|_{M_W=M_W^{\text{SM4}}} - g_{V,A}^{q,(0)}|_{M_W=M_W^{\text{SM3}}} + \delta g_{V,A}^{q,(1)}|_{M_W=M_W^{\text{SM3}}} \quad , \quad (15)$$

where $\delta g_{V,A}^{q,(1)}$ denotes the new-physics corrections of the one-loop contributions to $g_{V,A}^q$.

5 Numerical Results

In this section we study the dependence of the effective couplings and the Z pole observables on the masses and mixing angles of the fourth generation. To this end we set

$$\begin{aligned} \theta_{12}^{\text{SM4}} &= \theta_{12}^{\text{SM3}} \quad , \quad \theta_{13}^{\text{SM4}} = \theta_{13}^{\text{SM3}} \quad , \quad \theta_{23}^{\text{SM4}} = \theta_{23}^{\text{SM3}} \quad , \quad \delta_{13}^{\text{SM4}} = \delta_{13}^{\text{SM3}} \quad , \\ \delta_{14} &= \delta_{24} = 0 \quad , \\ m_{\ell_4} &= 101 \text{ GeV} \quad , \quad m_{\nu_4} = 50 \text{ GeV} \quad , \quad m_{t'} = m_{b'} = 500 \text{ GeV} \quad , \end{aligned} \quad (16)$$

with θ_{12}^{SM3} , θ_{13}^{SM3} , θ_{23}^{SM3} and δ_{13}^{SM3} from (12). In this scenario the values of $m_{t'}$ and $m_{b'}$ evade the current CMS limits on fourth-generation quarks [25, 26]. The constraints

from direct Higgs searches at the LHC are avoided by choosing a sufficiently small fourth-generation neutrino mass, so that the Higgs boson can decay invisibly into $\nu_4\bar{\nu}_4$ [29–32].

The fourth-generation phases δ_{14} and δ_{24} have no noticeable impact on the effective couplings g_V^q and g_A^q since they only depend on moduli of the V_{CKM4} matrix elements. We have checked explicitly that the variations in the EWPOs are far below their experimental uncertainties when the new CP phases are varied between 0 and 2π .

The absolute values of the entries in the first two rows of the 3×3 CKM matrix are strongly constrained by various flavour-observables. In the context of the SM4 these constraints were first discussed in [14] and it was found that scenarios are allowed where the mixing angles θ_{12} , θ_{13} and θ_{23} deviate substantially from their best-fit SM3 values. In [15] it was then pointed out that electroweak precision measurements rule out these large-mixing scenarios and the analysis [14] was updated in [17], where electroweak precision constraints were implemented via the oblique parameters S and T . Here we are mainly interested in the difference between the oblique and the full corrections to the EWPOs. To simplify the numerical discussion we therefore keep the SM3 mixing angles fixed according to (12) and only vary the new mixing angles. For these we will use the following independent limits from [17]:

$$|s_{14}| < 0.05 \quad , \quad |s_{24}| < 0.1 \quad , \quad |s_{34}| < 0.2 \quad , \quad (17)$$

with $s_{ij} \equiv \sin \theta_{ij}$. In a future combined fit of flavour and electroweak precision observables all mixing angles should be varied independently, and the non-oblique corrections will then also affect the best-fit values of the SM3 mixing angles.

Figure 1 and 2 show the relative new physics corrections to g_V^q and g_A^q for up-type and down-type quarks q , respectively, as functions of s_{34} and for different combinations of s_{14} and s_{24} . Varying s_{24} s_{34} in their allowed ranges leads to effects above the permille level. Hence, one should expect that EWPOs constrain θ_{24} and θ_{34} simultaneously. The contributions which depend on s_{14} and s_{24} are positive and the effects are additive: varying one mixing angle shifts the effective couplings by amounts which are mostly independent of the other mixing angles. This can be explained by the fact that the largest s_{14} or s_{24} -dependent contributions come from terms where one angle enters through a sine factor and the others through cosine factors. The highest sensitivity to s_{14} and s_{24} is observed in the vector couplings of up-type quarks (g_V^u and g_V^c). Varying s_{14} , s_{24} and s_{34} in the chosen ranges leads to corrections between -0.3 and 1.2% for up-type quarks and between -0.1 and 0.5% for down-type quarks.

The dashed lines in Figures 1 and 2 contain only the oblique corrections while the solid lines are the result of our “exact” calculation. As explained earlier there are two reasons for their discrepancy: the mixing between all four generations of quarks leads to vertex corrections that are not included in the oblique parameters and the small mass of the fourth-generation neutrino makes the expansions used in the derivation of S , T




Figure 1: The relative ‘new physics’ corrections to the effective couplings g_V^q and g_A^q for up-type quarks ($q = u, c$) as functions of s_{34} for the parameters (16) and different combinations of s_{14} and s_{24} . The dashed lines show the oblique corrections as computed from (9).




Figure 2: The relative ‘new physics’ corrections to the effective couplings g_V^q and g_A^q for down-type quarks ($q = d, s, b$) as functions of s_{34} for the parameters (16) and different combinations of s_{14} and s_{24} . The dashed lines show the oblique corrections as computed from (9).




Figure 3: The relative ‘new physics’ corrections to the observables Γ_{had} , R_b , M_W and \mathcal{A}_b as functions of s_{34} for the parameters (16) and different combinations of s_{14} and s_{24} . The dashed lines contain only the oblique corrections. The dotted lines indicate the range in which the observable would be in 1σ agreement with the experimental value. For \mathcal{A}_b the dotted lines are outside the displayed range and all values are in 1σ agreement with the measurement.




Figure 4: The dependence of Γ_{had} and R_b on m_{ν_4} for two different values of $m_{b'}$, $s_{14} = s_{24} = s_{34} = 0$ and all other parameters according to (16). The dashed lines only contain the oblique corrections while the solid lines are the results of the exact calculation.

and U invalid. We see that the non-oblique effects are at the level of a few permille and reduce the size of the effective couplings.

Figure 3 shows the relative new physics corrections to Γ_{had} , R_b , M_W and \mathcal{A}_b as functions of s_{34} for the same combinations of s_{14} and s_{24} . The corrections to Γ_{had} are between -0.1 and 0.5% . Varying s_{24} in its allowed range shifts the ratio $\delta\Gamma_{\text{had}}/\Gamma_{\text{had}}^{\text{SM3}}$ by approximately 0.2% . The constraint imposed by Γ_{had} on θ_{24} and especially θ_{34} is therefore competitive with those obtained from flavour physics. The oblique approximation over-estimates Γ_{had} by up to 0.4% . The hadronic branching fraction R_b is essentially insensitive to θ_{14} and θ_{24} . The dependence on θ_{34} is not captured at all by the oblique approximation and for $s_{34} = 0.2$ it differs from the exact result by 0.4% . The corrections to M_W lie between 0 and 0.5% . The corrections to \mathcal{A}_b are of the order of one permille and thus negligible compared to the experimental error on \mathcal{A}_b .

It is worth noting that agreement with the experimental value within approximately one standard deviation can be achieved simultaneously for the two most constraining observables, Γ_{had} and M_W , even for nonzero values of s_{24} and s_{34} . In the scenario discussed above this happens, for example, for $s_{24} = 0$, $s_{34} = 0.1$ or for $s_{24} = 0.1$ and $s_{34} = 0$. This indicates that non-trivial mixing scenarios, i.e. scenarios where all three fourth-generation mixing angles are nonzero, may well be in good agreement with electroweak precision measurements and that varying θ_{24} might even improve the electroweak fit. To quantify this statement it would be necessary to perform a global fit of all SM4 parameters (couplings, masses and CKM mixing angles) which takes into account electroweak precision measurements and flavour observables at the same time. In such a fit the non-oblique contributions to the EWPOs must be included since their effect is comparable to the experimental error.

Let us finally discuss the quality of the oblique approximation as a function of the

fourth-generation neutrino mass. Figure 4 shows the dependence of Γ_{had} and R_b on m_{ν_4} for two different values of $m_{b'}$, $s_{14} = s_{24} = s_{34} = 0$ and all other parameters according to (16). Again, the dashed lines only contain the oblique corrections while the solid lines are the results of the exact calculation. We see that R_b is insensitive to the neutrino mass and that (for vanishing mixing angles) the oblique corrections agree with the exact results. For Γ_{had} the oblique effects become relevant for neutrino masses below approximately 60 GeV. At the threshold $m_{\nu_4} = M_Z/2$ the full one-loop results diverge and become unreliable.

6 Conclusion

In this paper we studied the potential size of the corrections to the EWPOs that arise in a fourth-generation model with the most general (quark-)flavour structure (but assuming a trivial flavour-structure in the lepton-sector).

By computing ratios of form factors in the SM4 and SM3 to one-loop order and scaling results of precision calculations for the corresponding SM3 form factors with these ratios we can compute EWPOs within the SM4, including factorisable higher-order QCD and QED corrections and non-oblique contributions. The ratios of form factors can easily be computed with automated tools for one-loop calculations such as `FeynArts/FormCalc` [41–43]. This way, no approximations beyond the one-loop approximation are made for the new-physics contributions. In particular, non-oblique contributions are included correctly this way. The application of our method to other new-physics models is straightforward.

We find that the EWPOs are sensitive to the fourth-generation mixing angles θ_{24} and θ_{34} . The constraints imposed by Γ_{had} and M_W are generally as strong as those obtained from flavour physics. In the region of parameter space that is currently favoured by direct searches for Higgs bosons and fourth-generation quarks the non-oblique corrections to the electroweak observables are of the order of a few permille and should therefore be included in a global fit.

Acknowledgement. The authors would like to thank Alexander Lenz and Ulrich Nierste for fruitful discussions and thorough proof reading.

J.R. is supported by DFG Sonderforschungsbereich SFB/TR 9 “Computergestützte Theoretische Teilchenphysik”. P.G. is supported by DFG SFB/TR9. M.W. is partially supported by project DFG NI 1105/2-1.

A Parametrisation of the CKM Matrix

The CKM matrix of the SM3 depends on three mixing angles θ_{12} , θ_{13} and θ_{23} and one complex phase δ_{13} . Its elements are given by

$$V_{\text{CKM}} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{13}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{13}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{13}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{13}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{13}} & c_{23}c_{13} \end{pmatrix}, \quad (18)$$

where $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$. The SM4 CKM matrix is parametrised by three additional mixing angles θ_{14} , θ_{24} and θ_{34} and two additional phases δ_{14} and δ_{24} . In terms of these parameters, it is then written as

$$V_{\text{CKM4}} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} & V_{ub'} \\ V_{cd} & V_{cs} & V_{cb} & V_{cb'} \\ V_{td} & V_{ts} & V_{tb} & V_{tb'} \\ V_{t'd} & V_{t's} & V_{t'b} & V_{t'b'} \end{pmatrix} \quad (19)$$

with

$$\begin{aligned} V_{ud} &= c_{12}c_{13}c_{14} \quad , \quad V_{us} = c_{13}c_{14}s_{12} \quad , \quad V_{ub} = c_{14}s_{13}e^{-i\delta_{13}} \quad , \\ V_{ub'} &= s_{14}e^{-i\delta_{14}} \quad , \quad V_{cb'} = c_{14}s_{24}e^{-i\delta_{24}} \quad , \quad V_{tb'} = c_{14}c_{24}s_{34} \quad , \quad V_{t'b'} = c_{14}c_{24}c_{34} \quad , \\ V_{cd} &= -c_{23}c_{24}s_{12} + c_{12}(-c_{24}s_{13}s_{23}e^{i\delta_{13}} - c_{13}s_{14}s_{24}e^{i(\delta_{14}-\delta_{24})}) \quad , \\ V_{cs} &= c_{12}c_{23}c_{24} + s_{12}(-c_{24}s_{13}s_{23}e^{i\delta_{13}} - c_{13}s_{14}s_{24}e^{i(\delta_{14}-\delta_{24})}) \quad , \\ V_{cb} &= c_{13}c_{24}s_{23} - s_{13}s_{14}s_{24}e^{i(\delta_{14}-\delta_{13}-\delta_{24})} \quad , \\ V_{td} &= -s_{12}(-c_{34}s_{23} - c_{23}s_{24}s_{34}e^{i\delta_{24}}) \\ &\quad + c_{12}(-c_{13}c_{24}s_{14}s_{34}e^{i\delta_{14}} - s_{13}e^{i\delta_{13}}(c_{23}c_{34} - s_{23}s_{24}s_{34}e^{i\delta_{24}})) \quad , \\ V_{ts} &= c_{12}(-c_{34}s_{23} - c_{23}s_{24}s_{34}e^{i\delta_{24}}) \\ &\quad + s_{12}(-c_{13}c_{24}s_{14}s_{34}e^{i\delta_{14}} - s_{13}e^{i\delta_{13}}(c_{23}c_{34} - s_{23}s_{24}s_{34}e^{i\delta_{24}})) \quad , \\ V_{tb} &= -c_{24}s_{13}s_{14}s_{34}e^{i(\delta_{14}-\delta_{13})} + c_{13}(c_{23}c_{34} - s_{23}s_{24}s_{34}e^{i\delta_{24}}) \quad , \\ V_{t'd} &= -s_{12}(-c_{23}c_{34}s_{24}e^{i\delta_{24}} + s_{23}s_{34}) \\ &\quad + c_{12}(-c_{13}c_{24}c_{34}s_{14}e^{i\delta_{14}} - s_{13}e^{i\delta_{13}}(-c_{34}s_{23}s_{24}e^{i\delta_{24}} - c_{23}s_{34})) \quad , \\ V_{t's} &= c_{12}(-c_{23}c_{34}s_{24}e^{i\delta_{24}} + s_{23}s_{34}) \\ &\quad + s_{12}(-c_{13}c_{24}c_{34}s_{14}e^{i\delta_{14}} - s_{13}e^{i\delta_{13}}(-c_{34}s_{23}s_{24}e^{i\delta_{24}} - c_{23}s_{34})) \quad , \\ V_{t'b} &= -c_{24}c_{34}s_{13}s_{14}e^{i(\delta_{14}-\delta_{13})} + c_{13}(-c_{34}s_{23}s_{24}e^{i\delta_{24}} - c_{23}s_{34}) \quad . \end{aligned} \quad (20)$$

For $\theta_{14} = \theta_{24} = \theta_{34} = \delta_{14} = \delta_{24} = 0$ the SM4 CKM matrix assumes block-diagonal form with the SM3 CKM matrix in the first 3×3 block:

$$V_{\text{CKM4}} = \begin{pmatrix} & & & 0 \\ & V_{\text{CKM}} & & 0 \\ & & & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (21)$$

References

- [1] G. C. Branco and L. Lavoura, *Nuclear Physics B* **278**, 738 (1986), ISSN 0550-3213.
- [2] P. H. Frampton, P. Hung, and M. Sher, *Phys.Rept.* **330**, 263 (2000), [arXiv:hep-ph/9903387 \[hep-ph\]](#).
- [3] F. del Aguila, M. Perez-Victoria, and J. Santiago, *JHEP* **09**, 011 (2000), [arXiv:hep-ph/0007316](#).
- [4] G. Barenboim, F. Botella, and O. Vives, *Nucl.Phys.* **B613**, 285 (2001), [arXiv:hep-ph/0105306 \[hep-ph\]](#).
- [5] B. Holdom, *Phys. Rev.* **D54**, 721 (1996), [arXiv:hep-ph/9602248](#).
- [6] M. Maltoni, V. Novikov, L. Okun, A. N. Rozanov, and M. Vysotsky, *Phys.Lett.* **B476**, 107 (2000), [arXiv:hep-ph/9911535 \[hep-ph\]](#).
- [7] H.-J. He, N. Polonsky, and S.-f. Su, *Phys.Rev.* **D64**, 053004 (2001), [arXiv:hep-ph/0102144 \[hep-ph\]](#).
- [8] V. Novikov, L. Okun, A. N. Rozanov, and M. Vysotsky, *Phys.Lett.* **B529**, 111 (2002), [arXiv:hep-ph/0111028 \[hep-ph\]](#).
- [9] V. Novikov, L. Okun, A. N. Rozanov, and M. Vysotsky, *JETP Lett.* **76**, 127 (2002), [arXiv:hep-ph/0203132 \[hep-ph\]](#).
- [10] G. D. Kribs, T. Plehn, M. Spannowsky, and T. M. Tait, *Phys.Rev.* **D76**, 075016 (2007), [arXiv:0706.3718 \[hep-ph\]](#).
- [11] V. A. Novikov, A. N. Rozanov, and M. I. Vysotsky, *Phys. Atom. Nucl.* **73**, 636 (2010), [arXiv:0904.4570 \[hep-ph\]](#).
- [12] B. Holdom *et al.*, *PMC Phys.* **A3**, 4 (2009), [arXiv:0904.4698 \[hep-ph\]](#).
- [13] J. Alwall *et al.*, *Eur. Phys. J.* **C49**, 791 (2007), [arXiv:hep-ph/0607115](#).
- [14] M. Bobrowski, A. Lenz, J. Riedl, and J. Rohrwild, *Phys.Rev.* **D79**, 113006 (2009), [arXiv:0902.4883 \[hep-ph\]](#).
- [15] M. S. Chanowitz, *Phys. Rev.* **D79**, 113008 (2009), [arXiv:0904.3570 \[hep-ph\]](#).
- [16] W.-S. Hou, Y.-Y. Mao, and C.-H. Shen, *Phys.Rev.* **D82**, 036005 (2010), [arXiv:1003.4361 \[hep-ph\]](#).
- [17] O. Eberhardt, A. Lenz, and J. Rohrwild, *Phys.Rev.* **D82**, 095006 (2010), [arXiv:1005.3505 \[hep-ph\]](#).

- [18] W.-S. Hou, M. Nagashima, and A. Soddu, *Phys.Rev.* **D76**, 016004 (2007), [arXiv:hep-ph/0610385 \[hep-ph\]](#).
- [19] A. Soni, A. K. Alok, A. Giri, R. Mohanta, and S. Nandi, *Phys.Lett.* **B683**, 302 (2010), [arXiv:0807.1971 \[hep-ph\]](#).
- [20] A. Soni, A. K. Alok, A. Giri, R. Mohanta, and S. Nandi, *Phys.Rev.* **D82**, 033009 (2010), [arXiv:1002.0595 \[hep-ph\]](#).
- [21] A. J. Buras, B. Duling, T. Feldmann, T. Heidsieck, C. Promberger, *et al.*, *JHEP* **1009**, 106 (2010), [arXiv:1002.2126 \[hep-ph\]](#).
- [22] A. J. Buras, B. Duling, T. Feldmann, T. Heidsieck, and C. Promberger, *JHEP* **1009**, 104 (2010), [arXiv:1006.5356 \[hep-ph\]](#).
- [23] H. Lacker and A. Menzel, *JHEP* **1007**, 006 (2010), [arXiv:1003.4532 \[hep-ph\]](#).
- [24] J. Erler and P. Langacker, *Phys.Rev.Lett.* **105**, 031801 (2010), [arXiv:1003.3211 \[hep-ph\]](#).
- [25] “Search for $t\bar{t}$ pair production in lepton+jets channel,” CMS Physics Analysis Summary (2011), [CMS-PAS-EXO-11-051](#).
- [26] “Search for a Heavy Bottom-like Quark in 1.14 fb of pp Collisions at $\sqrt{s} = 7$ TeV,” CMS Physics Analysis Summary (2011), [CMS-PAS-EXO-11-036](#).
- [27] X. Ruan and Z. Zhang(2011), [arXiv:1105.1634 \[hep-ph\]](#).
- [28] J. F. Gunion(2011), [arXiv:1105.3965 \[hep-ph\]](#).
- [29] K. Belotsky, D. Fargion, M. Khlopov, R. Konoplich, and K. Shibaev, *Phys.Rev.* **D68**, 054027 (2003), [arXiv:hep-ph/0210153 \[hep-ph\]](#).
- [30] A. Rozanov and M. Vysotsky, *Phys.Lett.* **B700**, 313 (2011), [arXiv:1012.1483 \[hep-ph\]](#).
- [31] W.-Y. Keung and P. Schwaller, *JHEP* **1106**, 054 (2011), [arXiv:1103.3765 \[hep-ph\]](#).
- [32] S. Cetin, T. Cuhadar-Donszelmann, M. Sahin, S. Sultansoy, and G. Unel(2011), [arXiv:1108.4071 \[hep-ph\]](#).
- [33] L. M. Carpenter(2011), [arXiv:1110.4895 \[hep-ph\]](#).
- [34] M. S. Chanowitz, *Phys. Rev. D*, 035018 (2010), [arXiv:1007.0043 \[hep-ph\]](#).
- [35] S. Nandi and A. Soni(2010), [arXiv:1011.6091 \[hep-ph\]](#).
- [36] A. K. Alok, A. Dighe, and D. London(2010), * Temporary entry *, [arXiv:1011.2634 \[hep-ph\]](#).

- [37] M. E. Peskin and T. Takeuchi, *Phys. Rev.* **D46**, 381 (1992).
- [38] I. Maksymyk, C. P. Burgess, and D. London, *Phys. Rev.* **D50**, 529 (1994), [arXiv:hep-ph/9306267](#).
- [39] D. Y. Bardin *et al.*, *Comput. Phys. Commun.* **133**, 229 (2001), <http://www-zeuthen.desy.de/theory/research/zfitter/index.html>, [arXiv:hep-ph/9908433](#).
- [40] A. B. Arbuzov *et al.*, *Comput. Phys. Commun.* **174**, 728 (2006), [arXiv:hep-ph/0507146](#).
- [41] T. Hahn and M. Rauch, *Nucl. Phys. Proc. Suppl.* **157**, 236 (2006), [arXiv:hep-ph/0601248](#).
- [42] T. Hahn, *Comput. Phys. Commun.* **140**, 418 (2001), [arXiv:hep-ph/0012260](#).
- [43] T. Hahn and M. Perez-Victoria, *Comput. Phys. Commun.* **118**, 153 (1999), [arXiv:hep-ph/9807565](#).
- [44] (2010), [arXiv:1012.2367 \[hep-ex\]](#).
- [45] K. Nakamura *et al.* (Particle Data Group), *J.Phys.G* **G37**, 075021 (2010).
- [46] G. Degrandi, P. Gambino, and A. Vicini, *Phys.Lett.* **B383**, 219 (1996), [arXiv:hep-ph/9603374 \[hep-ph\]](#).
- [47] G. Degrandi and P. Gambino, *Nucl.Phys.* **B567**, 3 (2000), [arXiv:hep-ph/9905472 \[hep-ph\]](#).
- [48] A. Freitas, W. Hollik, W. Walter, and G. Weiglein, *Phys.Lett.* **B495**, 338 (2000), 13 pages: error in the numerical evaluation of a specific two-loop vacuum integral corrected Report-no: CERN-TH/2000-194 Journal-ref: Phys.Lett. B495 (2000) 338-346, [arXiv:hep-ph/0007091 \[hep-ph\]](#).
- [49] A. Freitas, W. Hollik, W. Walter, and G. Weiglein, *Nucl.Phys.* **B632**, 189 (2002), [arXiv:hep-ph/0202131 \[hep-ph\]](#).
- [50] M. Awramik and M. Czakon, *Phys.Lett.* **B568**, 48 (2003), 11 pages, 5 figures, Tab. 3 corrected, reference added, [arXiv:hep-ph/0305248 \[hep-ph\]](#).
- [51] M. Awramik, M. Czakon, A. Onishchenko, and O. Veretin, *Phys.Rev.* **D68**, 053004 (2003), [arXiv:hep-ph/0209084 \[hep-ph\]](#).
- [52] K. Chetyrkin, J. H. Kuhn, and M. Steinhauser, *Phys.Rev.Lett.* **75**, 3394 (1995), [arXiv:hep-ph/9504413 \[hep-ph\]](#).
- [53] K. Chetyrkin, J. H. Kuhn, and M. Steinhauser, *Nucl.Phys.* **B482**, 213 (1996), [arXiv:hep-ph/9606230 \[hep-ph\]](#).

- [54] M. Faisst, J. H. Kuhn, T. Seidensticker, and O. Veretin, *Nucl.Phys.* **B665**, 649 (2003), [arXiv:hep-ph/0302275 \[hep-ph\]](#).
- [55] M. Awramik, M. Czakon, A. Freitas, and G. Weiglein, *Phys.Rev.Lett.* **93**, 201805 (2004), [arXiv:hep-ph/0407317 \[hep-ph\]](#).
- [56] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, and A. Vicere, *Phys.Lett.* **B288**, 95 (1992), [arXiv:hep-ph/9205238 \[hep-ph\]](#).
- [57] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, and A. Vicere, *Nucl.Phys.* **B409**, 105 (1993).
- [58] J. Fleischer, O. Tarasov, and F. Jegerlehner, *Phys.Rev.* **D51**, 3820 (1995).
- [59] G. Degrandi, P. Gambino, and A. Sirlin, *Phys.Lett.* **B394**, 188 (1997), [arXiv:hep-ph/9611363 \[hep-ph\]](#).
- [60] A. Djouadi and C. Verzegnassi, *Phys.Lett.* **B195**, 265 (1987).
- [61] B. A. Kniehl, *Nucl.Phys.* **B347**, 86 (1990).
- [62] F. Halzen and B. A. Kniehl, *Nucl.Phys.* **B353**, 567 (1991).
- [63] B. A. Kniehl and A. Sirlin, *Nucl.Phys.* **B371**, 141 (1992).
- [64] B. A. Kniehl and A. Sirlin, *Phys.Rev.* **D47**, 883 (1993).
- [65] A. Djouadi and P. Gambino, *Phys.Rev.* **D49**, 3499 (1994), [arXiv:hep-ph/9309298 \[hep-ph\]](#).
- [66] G. Montagna, O. Nicrosini, F. Piccinini, and G. Passarino, *Comput. Phys. Commun.* **117**, 278 (1999), [arXiv:hep-ph/9804211](#).
- [67] D. Bardin *et al.*, in *Reports of the working group on precision calculations for the Z resonance*, pp. 7–162, contributed to Workshop Group on Precision Calculations for the Z Resonance (2nd meeting held Mar 31, 3rd meeting held Jun 13), Geneva, Switzerland, 14 Jan 1994, [CERN 95-03](#).
- [68] A. Czarnecki and J. H. Kuhn, *Phys. Rev. Lett.* **77**, 3955 (1996), [arXiv:hep-ph/9608366](#).
- [69] R. Harlander, T. Seidensticker, and M. Steinhauser, *Phys. Lett.* **B426**, 125 (1998), [arXiv:hep-ph/9712228](#).
- [70] M. Bohm, H. Spiesberger, and W. Hollik, *Fortsch. Phys.* **34**, 687 (1986).
- [71] M. Böhm, A. Denner, and H. Joos, *Gauge Theories of the Strong and Electroweak Interaction* (B. G. Teubner, 2001).
- [72] A. Sirlin, *Phys.Rev.* **D22**, 971 (1980).