

Hypothesis of photonic portal in the problem of dark matter and classical nature of gravitation

WOJCIECH KRÓLIKOWSKI

Institute of Theoretical Physics, Faculty of Physics, University of Warsaw

Hoza 69, 00-681 Warszawa, Poland

Abstract

Within our model of "photonic portal" to the hypothetical hidden sector of the Universe, an option is considered, where the massive sterile bosons, mediating in this sector and described by the antisymmetric-tensor field $A_{\mu\nu}$ (of dimension one), are gauged by a vector field χ_μ (of dimension zero). When the Einsteinian gravity is switched on into this model, then it is argued that the metric $g_{\mu\nu}(x)$ of spacetime is proportional to $\langle \chi_\mu(x) \chi_\nu(x) \rangle_{\text{vac}}$, being in this way classical in its nature, not the subject of quantization within an overall quantum theory including gravitation. A Lagrangian density (generally dependent on χ_μ and its first and second derivatives) is tentatively proposed in such a form, where χ_μ appears also outside of $A_{\mu\nu}$, spoiling the trivial gauge invariance of our earlier model with respect to the gauging field χ_μ . If the cosmological constant λ is neglected, then the term with χ_μ outside of $A_{\mu\nu}$ vanishes.

PACS numbers: 14.80.-j , 04.50.+h , 95.35.+d

May 2011

Recently, we speculated [1] on the existence of a hidden sector of the Universe, consisting of some sterile particles (Standard-Model neutral), *e.g.* of spin-0 bosons ("sterons"), spin-1/2 fermions ("sterinos") and nongauge mediating bosons ("A bosons") described by an antisymmetric-tensor field $A_{\mu\nu}$ (of dimension one). The A bosons were assumed to be weakly coupled to steron-photon pairs and antisterino-sterino pairs by the interaction Lagrangian densities

$$-\frac{1}{2}\sqrt{f}\varphi F_{\mu\nu}A^{\mu\nu} \quad (1)$$

and

$$-\frac{1}{2}\zeta\sqrt{f}\bar{\psi}\sigma_{\mu\nu}\psi A^{\mu\nu}, \quad (2)$$

respectively. Here, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the Standard-Model electromagnetic field (of dimension two), while \sqrt{f} and $\zeta\sqrt{f}$ denote two dimensionless small coupling constants. We assumed that

$$\varphi = \langle\varphi\rangle_{\text{vac}} + \varphi_{\text{ph}}, \quad (3)$$

where $\langle\varphi\rangle_{\text{vac}} \neq 0$ is a spontaneously nonzero vacuum expectation value of the steron field φ . The kinetic Lagrangian density of A bosons is built up only from $A_{\mu\nu}$ (of dimension one):

$$-\frac{1}{4}[(\partial_\rho A_{\mu\nu})(\partial^\rho A^{\mu\nu}) - M^2 A_{\mu\nu}A^{\mu\nu}] \quad (4)$$

with M standing for a mass scale of A bosons, expected to be large.

The field equations for $F_{\mu\nu}$ and $A_{\mu\nu}$, provided by the interactions (1) and (2), are

$$\partial^\nu [F_{\mu\nu} + \sqrt{f}\varphi A_{\mu\nu}] = -j_\mu, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad (5)$$

and

$$(\square - M^2)A_{\mu\nu} = -\sqrt{f}(\varphi F_{\mu\nu} + \zeta\bar{\psi}\sigma_{\mu\nu}\psi), \quad (6)$$

where j_μ is the Standard-Model electric current. Eqs. (5) give us the form assumed by Maxwell's equations in the presence of our hidden-sector.

In the first Eq. (5) the total electric source-current is

$$j_\mu + \partial^\nu (\sqrt{f} \varphi A_{\mu\nu}), \quad (7)$$

where the second term does not contribute to the total electric charge because of its four-divergence structure:

$$\int d^3x [j_0 + \partial^k (\sqrt{f} \varphi A_{0k})] = \int d^3x j_0 = Q, \quad (8)$$

so it gets a magnetic character.

We called the coupling (1) of photons to the hidden sector of the Universe "photonic portal" (to the hidden sector), in contrast to the popular "Higgs portal" [2]. Such a hidden sector is conjectured to be responsible for the cold dark matter [1].

If there exists a vector field χ_μ (of dimension zero) gauging the sterile mediating field $A_{\mu\nu}$ (of dimension one),

$$A_{\mu\nu} = \partial_\mu \chi_\nu - \partial_\nu \chi_\mu, \quad (9)$$

(cf. Appendix in Ref. [3]) and if, in addition, χ_μ still does not appear outside of $A_{\mu\nu}$ in the total Lagrangian (nor in a subsidiary condition), our model of hidden sector remains essentially unchanged *versus* the previous nongauge case (and is then trivially gauge invariant with respect to χ_μ). It is the matter of experiment to decide, whether the field χ_μ (when it exists) is or is not trivial in the above sense.

When the Einsteinian gravity is switched on in our model, terms including χ_μ may appear also outside of

$$A_{\mu\nu} = \chi_{\nu;\mu} - \chi_{\mu;\nu} \equiv \partial_\mu \chi_\nu - \partial_\nu \chi_\mu \quad (10)$$

in the total Lagrangian and so in the field equation for χ_μ as well. Assume tentatively the following Lagrangian density for the field χ_μ :

$$\begin{aligned} \mathcal{L}^{(\chi)} = & -\frac{1}{4} (A_{\mu\nu;\rho} A^{\mu\nu;\rho} - M^2 A_{\mu\nu} A^{\mu\nu}) - \frac{1}{2} \sqrt{f} (\varphi F_{\mu\nu} + \zeta \bar{\psi} \sigma_{\mu\nu} \psi) A^{\mu\nu} \\ & - \frac{1}{2} \xi M^2 \left[\left(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R^\rho{}_\rho \right) + 8\pi G_N T_{\mu\nu} \right] \chi^\mu \chi^\nu. \end{aligned} \quad (11)$$

Here, $(...)_\rho$ denotes covariant differentiation with respect to x^ρ and $(...)^{\rho\sigma} \equiv g^{\rho\sigma} (...)_\sigma$, $G_N = 1/M_P^2$ ($\hbar = 1 = c$) stands for the Newton gravitational constant and ξ represents a

new dimensionless parameter, while $R_{\mu\nu} \equiv R^\rho_{\mu\nu\rho}$ is the Ricci curvature tensor of spacetime and $T_{\mu\nu}$ – the total energy-momentum tensor of all fields of "matter".

If Einstein gravitational equation

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R^\rho_\rho - \lambda g_{\mu\nu} = -8\pi G_N T_{\mu\nu} \quad (12)$$

or equivalently

$$R_{\mu\nu} + \lambda g_{\mu\nu} = -8\pi G_N \left(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T^\rho_\rho \right) \quad (13)$$

is taken into account, then the Lagrangian density (11) reads

$$\mathcal{L}^{(x)} = -\frac{1}{4} (A_{\mu\nu;\rho} A^{\mu\nu;\rho} - M^2 A_{\mu\nu} A^{\mu\nu}) - \frac{1}{2} \sqrt{f} (\varphi F_{\mu\nu} + \zeta \bar{\psi} \sigma_{\mu\nu} \psi) A^{\mu\nu} - \frac{1}{2} \xi \lambda M^2 \chi_\mu \chi^\mu. \quad (14)$$

Here, λ is the cosmological constant (of dimension two). If $\lambda \neq 0$, the Lagrangian density (14) is not gauge invariant with respect to the gauging field χ_μ , as it is trivially gauge invariant without gravitation. Without gravitation, the Lagrangian density (14) is reduced to the form

$$\mathcal{L}^{(x)} = \mathcal{L}^{(A)} = -\frac{1}{4} [(\partial_\rho A_{\mu\nu})(\partial^\rho A^{\mu\nu}) - M^2 A_{\mu\nu} A^{\mu\nu}] - \sqrt{f} (\varphi F_{\mu\nu} + \zeta \bar{\psi} \sigma_{\mu\nu} \psi) A^{\mu\nu} \quad (15)$$

equal to the sum of Eqs. (1), (2) and (4).

In the Einstein equation (12) or (13), the total energy-momentum tensor of "matter" is given as $(1/2)\sqrt{-g}T_{\mu\nu} = \partial(\sqrt{-g}\mathcal{L})/\partial g^{\mu\nu} - \partial_\rho \partial(\sqrt{-g}\mathcal{L})/\partial \partial_\rho g^{\mu\nu}$, where \mathcal{L} denotes the total Lagrangian density of "matter" ($\mathcal{L} = \mathcal{L}^{(x)} + \dots$) and $g = \det(g_{\mu\nu})$. Then, the variational principle $\delta_g \int d^4V \sqrt{-g}[(R + 2\lambda)/(16\pi G_N) + \mathcal{L}] = 0$ with $R = R^\rho_\rho$ leads to Eq. (12) or (13).

Note that in Eqs. (11) and (12) or (13) [4]

$$R_{\mu\nu} \equiv R^\rho_{\mu\nu\rho} \equiv \Gamma^\rho_{\mu\rho,\nu} - \Gamma^\rho_{\mu\nu,\rho} + \Gamma^\sigma_{\mu\rho} \Gamma^\rho_{\nu\sigma} - \Gamma^\sigma_{\mu\nu} \Gamma^\rho_{\rho\sigma} \quad (16)$$

with $(\dots)_{,\nu} \equiv \partial_\nu(\dots)$, giving the identity

$$\chi^\nu_{;\mu;\nu} - \chi^\nu_{;\nu;\mu} \equiv -R_{\mu\nu} \chi^\nu. \quad (17)$$

Here,

$$\Gamma_{\mu\nu}^{\rho} \equiv \frac{1}{2}g^{\rho\sigma}(g_{\sigma\mu,\nu} + g_{\sigma\nu,\mu} - g_{\mu\nu,\sigma}) \quad (18)$$

and *e.g.*

$$\chi_{\mu;\nu} \equiv \chi_{\mu,\nu} - \Gamma_{\mu\nu}^{\rho}\chi_{\rho} \quad , \quad \chi^{\mu}_{;\nu} \equiv \chi^{\mu}_{,\nu} + \Gamma_{\nu\rho}^{\mu}\chi^{\rho}. \quad (19)$$

In Eq. (12), $g_{\mu\nu}^{;\nu} = 0$ because of $g_{\mu\nu}^{;\rho} = 0$ and $(R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R^{\rho}_{\rho})^{;\nu} = 0$ due to the Bianchi identities, thus $T_{\mu\nu}^{;\nu} = 0$.

The field equation for χ_{μ} following as an Euler-Lagrange equation from the variational principle $\delta_{\chi} \int d^4V \sqrt{-g} \mathcal{L}^{(x)} = 0$ with the tentative Lagrangian density (14) (dependent on χ_{μ} , $A_{\mu\nu}$ and $A_{\mu\nu;\rho}$) is

$$-(A_{\mu\nu;\rho}^{;\rho} + \frac{1}{\sqrt{-g}}g^{\sigma\lambda}\Gamma_{\sigma\lambda}^{\rho}A_{\mu\nu;\rho} + M^2A_{\mu\nu})^{;\nu} + \xi\lambda M^2\chi_{\mu} = -\sqrt{f}(\varphi F_{\mu\nu} + \zeta\bar{\psi}\sigma_{\mu\nu}\psi)^{;\nu}. \quad (20)$$

Here, $g^{\sigma\lambda}\Gamma_{\sigma\lambda}^{\rho} \equiv -\partial_{\sigma}(\sqrt{-g}g^{\sigma\rho}/\sqrt{-g})$ and *e.g.* $A_{\mu\nu}^{;\nu} \equiv \partial^{\nu}(\sqrt{-g}A_{\mu\nu})/\sqrt{-g}$. Without gravitation, Eq. (20) is reduced to the formula

$$(\square - M^2)\partial^{\nu}A_{\mu\nu} = -\sqrt{f}\partial^{\nu}(\varphi F_{\mu\nu} + \zeta\bar{\psi}\sigma_{\mu\nu}\psi) \quad (21)$$

equivalent to Eq. (6) for $A_{\mu\nu}$ when this equation is acted on by ∂^{ν} , where $\partial^{\nu}A_{\mu\nu} = -\partial^{\nu}\partial^{\nu}\chi_{\mu} = \square\chi_{\mu}$ under the condition $\partial^{\nu}\chi_{\nu} = 0$ (*cf.* Appendix in Ref. [3]). Note that the general field equation (20) for χ_{μ} gets the form $(\dots)_{\mu\nu}^{;\nu} = -\xi\lambda M^2\chi_{\mu}$, where $(\dots)_{\mu\nu}^{;\nu} \equiv \partial^{\nu}[\sqrt{-g}(\dots)_{\mu\nu}]/\sqrt{-g}$. If the cosmological constant λ is neglected, it is reduced to the field equation $(\dots)_{\mu\nu}^{;\nu} = 0$ for $A_{\mu\nu}$ or even to $(\dots)_{\mu\nu} = 0$, when $\mathcal{L}^{(x)}$ is not dependent on χ_{μ} *i.e.*, $\mathcal{L}^{(x)} = \mathcal{L}^{(A)}$ (as in the case of Eq. (14) with $\lambda = 0$).

Now, making use of the Lagrangian density (14) or field equation (20), we may quantize all fields of "matter" in our model, keeping the Einsteinian metric $g_{\mu\nu}$ of spacetime still classical (then, $R_{\mu\nu}$ and $T_{\mu\nu}$ in the Einstein equation (12) or (13) are also classical). In this way, the gravity in our model is considered as classical in its nature.

We can stress such a c-number nature of $g_{\mu\nu}(x)$ by showing that – after the above quantization of "matter" – the vacuum expectation value of dimensionless quantum pair $\chi_{\mu}(x)\chi_{\nu}(x)$ is proportional to the (also dimensionless) metric $g_{\mu\nu}(x)$ of spacetime.

In fact, making use of the repers $e_{\mu}^{\rho_0}(x)$ ($\mu = 0, 1, 2, 3$ and $\rho_0 = 0_0, 1_0, 2_0, 3_0$) orthonormalized as

$$e_{\mu}^{\rho_0}(x)e^{\mu\sigma_0}(x) = \eta^{\rho_0\sigma_0} \text{ with } \eta^{\rho_0\sigma_0} = \text{diag}(1, -1, -1, -1) = \eta_{\rho_0\sigma_0}, \quad (22)$$

and having the metric $g_{\mu\nu}(x)$ of spacetime defined through the product

$$g_{\mu\nu}(x) = e_{\mu}^{\rho_0}(x)e_{\nu\rho_0}(x) = e_{\mu}^{\rho_0}(x)e_{\nu}^{\sigma_0}(x)\eta_{\rho_0\sigma_0}, \quad (23)$$

we can represent the field $\chi_{\mu}(x)$ in the framework of the metric in the following form:

$$\chi_{\mu}(x) = e_{\mu}^{\rho_0}(x)\chi_{\rho_0}(x) = e_{\mu\rho_0}(x)\chi^{\rho_0}(x), \quad (24)$$

where the projections $\chi_{\rho_0}(x) = e_{\rho_0}^{\mu}(x)\chi_{\mu}(x)$ of the vector field $(\chi_{\mu}(x))$ onto the repers $(e_{\rho_0}^{\mu}(x))$ satisfy the orthogonality condition

$$\langle \chi_{\rho_0}(x)\chi_{\sigma_0}(x) \rangle_{\text{vac}} = Z\eta_{\rho_0\sigma_0}. \quad (25)$$

A positive Z denotes in Eq. (25) a dimensionless normalization factor requiring a cut-off procedure to treat the distribution $\langle \chi_{\rho_0}(x)\chi_{\sigma_0}(x') \rangle_{\text{vac}}$ at $x - x' \rightarrow 0$. Here, $e_{\mu\rho_0}(x) = \eta_{\rho_0\sigma_0}e_{\mu}^{\sigma_0}(x)$, $\chi^{\rho_0}(x) = \eta^{\rho_0\sigma_0}\chi_{\sigma_0}(x)$, while $e_{\mu\rho_0}(x) = g_{\mu\nu}(x)e_{\rho_0}^{\nu}(x)$. Then, due to the c-number character of the metric, the following relationship holds between the vacuum expectation value of the pair $\chi_{\mu}(x)\chi_{\nu}(x)$ and the metric $g_{\mu\nu}(x)$ of spacetime:

$$\langle \chi_{\mu}(x)\chi_{\nu}(x) \rangle_{\text{vac}} = e_{\mu}^{\rho_0}(x)e_{\nu}^{\sigma_0}(x) \langle \chi_{\rho_0}(x)\chi_{\sigma_0}(x) \rangle_{\text{vac}} = Zg_{\mu\nu}(x). \quad (26)$$

The dimensionless value of Z , related to the unique status of $\chi_{\mu}(x)$ in our model, suggests a distinguished character of the formula (26). For other vector-field structures, *e.g.* the electromagnetic vector field $A_{\mu}(x)$ (of dimension one), we get similar relationships as Eq. (26), but with normalization factors of other dimensions, *e.g.* dimension two. Then, *e.g.* $\langle A_{\mu}(x)A_{\nu}(x) \rangle_{\text{vac}} : \langle \chi_{\mu}(x)\chi_{\nu}(x) \rangle_{\text{vac}}$ is equal to the ratio of both normalization factors of resulting dimension two.

When deriving formally the relationship (26), we considered the metric $g_{\mu\nu}(x)$ as being classical. If the metric of spacetime has to be quantized, then such a relationship

ought to be avoided. Thus, as long as $g_{\mu\nu}(x) = Z^{-1} \langle \chi_\mu(x) \chi_\nu(x) \rangle_{\text{vac}}$, the gravity remains classical in its nature. In such a case, the c-number Einsteinian metric $g_{\mu\nu}(x)$ of spacetime has to collaborate within a quantum theory without a direct quantization of gravity. Then, our model gets a mixed structure containing a classical as well as a quantum part, the latter dynamically not complete (as yet). Of course, the Einstein equation (12) or (13) is here an independent postulate pertaining to the metric of spacetime. It can be seen that the relationship $\langle \chi_\mu(x) \chi_\nu(x) \rangle_{\text{vac}} = Z g_{\mu\nu}(x)$ is a solution to the equation $(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R^\rho_\rho - \lambda g_{\mu\nu} + 8\pi G_N T_{\mu\nu}) \langle \chi^\mu \chi^\nu \rangle_{\text{vac}} = 0$, if and only if the Einstein scalar equation

$$R^\rho_\rho + 4\lambda = 8\pi G_N T^\rho_\rho \quad (27)$$

holds (following from the Einstein equation (12) or (13) when this is satisfied). In fact, evidently $(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R^\rho_\rho - \lambda g_{\mu\nu} + 8\pi G_N T_{\mu\nu}) g^{\mu\nu} = -R^\rho_\rho - 4\lambda + 8\pi G_N T^\rho_\rho$.

In the particular case of $\xi = 0$, Lagrangian densities (11) and (14) are identical and do not contain the field χ_μ outside of $A_{\mu\nu}$ (being then trivially gauge invariant with respect to χ_μ). In this case, λ does not appear in $\mathcal{L}^{(x)} = \mathcal{L}^{(A)}$, though it may be that still $\lambda \neq 0$ in the Einstein equation.

Note that during the formal derivation of relationship (26) no particular dynamics for the vector field $\chi_\mu(x)$ was needed (only its coexistence with classical gravity was required). This makes the relationship (26) an identity for $\chi_\mu(x)$. So, due to the relationship (26), the metric $g_{\mu\nu}(x)$ can be replaced everywhere by the vacuum expectation value $Z^{-1} \langle \chi_\mu(x) \chi_\nu(x) \rangle_{\text{vac}}$. Obviously, the c-number metric $g_{\mu\nu}(x)$ depends functionally on the c-number counterparts of $\chi_\mu(x)$ and other quantum fields of "matter", contained in the classical energy-momentum tensor $T_{\mu\nu}(x)$ standing on the rhs of Einstein equation (12) or (13). Evidently, the content of such a tensor ought to be consistent (in the sense of the considered classical limit) with our experimental knowledge about cold dark matter and its hypothetical hidden-sector origin.

Finally, we can see that in our discussion the bold question is round the corner: could the c-number Einstein equation be closely correlated with the quantum part of our model (when it involves an adequate dynamics for $\chi_\mu(x)$ and other quantum fields), and so be considered as a c-number conclusion from the latter, consistent with general

relativity? In particular, this correlation should be consistent with the relationship $g_{\mu\nu}(x) = Z^{-1} \langle \chi_\mu(x) \chi_\nu(x) \rangle_{\text{vac}}$. In the case of affirmative answer to this question, the quantum part of our model would be dynamically complete. Then, physical roots of the classical-in-nature gravity should be traced within the quantum dynamics of $\chi_\mu(x)$ interacting (very weakly) with all quantum fields.

References

- [1] W. Królikowski, *Acta Phys. Polon.* **B 39**, 1881 (2008); arXiv: 0803.2977 [hep-ph]; *Acta Phys. Polon.* **B 40**, 111 (2009); *Acta Phys. Polon.* **B 40**, 2767 (2009).
- [2] Cf. e.g. J. March-Russell, S.M. West, D. Cumberbath and D. Hooper, *J. High Energy Phys.* **0807**, 058 (2008).
- [3] W. Królikowski, *Acta Phys. Polon.* **B 41**, 1277 (2010).
- [4] Cf. e.g. L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields*, Pergamon Press, New York, 1975; S. Weinberg, *Cosmology*, Oxford University Press, Oxford and New York, 2008; V. Mukhanov, *Physical Foundations of Cosmology*, Cambridge University Press, Cambridge, 2005.