

Generalized polarizability of neutral pions of the process $e^-e^+ \rightarrow \pi^0\pi^0\gamma$ in NJL model

A. I. Ahmadov*

*Joint Institute for Nuclear Research, Dubna, Russia and
Institute of Physics, Azerbaijan National Academy of Sciences, Baku, Azerbaijan*

E. A. Kuraev† and M. K. Volkov‡

Joint Institute for Nuclear Research, Dubna, Russia

(Dated: February 24, 2024)

Differential distributions in the $\pi_0\pi_0\gamma$ system created in the annihilation channel of an electron-positron collision are considered. The energy fractions of the pions (Dalitz-plot) distribution are presented in a general form and in approximation of intermediate vector mesons (excited and ordinary ones). It is pointed out that in relevant experiments the generalized polarizability of the neutral pion can be measured. Numerical illustrations are presented.

I. INTRODUCTION

In a recent paper [1] the process of creation of $\pi^0\omega$ in annihilation of an electron-positron pair was considered in the framework of the Nambu-Jona-Lazinio model. The result obtained turns out to be in very satisfactory agreement with theoretical estimations based on vector meson dominance and experimental measurement of this process with subsequent decay of the ω meson to $\pi_0\gamma$ [2]. In this experiment, the dependence of the total cross section on the total energy of initial electron and positron in the center of mass reference frame was measured.

It is the motivation of this paper to pay attention to the differential distributions in momenta of final particles since they carry information on the properties of the neutral pion such as polarizability.

In this paper, the process of annihilation of $e^-e^+ \rightarrow \gamma^* \rightarrow \pi^0\pi^0\gamma$ is considered. It can be obtained from the process $e^-e^+ \rightarrow \pi^0\omega$, if $\omega \rightarrow \pi^0\gamma$. Hence, we can use the results of [1].

Measuring of the process

$$e^+(p_+) + e^-(p_-) \rightarrow \gamma(q) \rightarrow \pi_0(q_1) + \pi_0(q_2) + \gamma(k) \quad (1)$$

provides a possibility to investigate the crossing process to the Compton scattering on neutral pion in the case when initial photon is of mass shell. Also, the role of excited vector mesons and the generalized polarizabilities of neutral pion can be studied. To article, this process can be useful for research Compton effect.

The dependence of the total cross section on the total energy of the initial electron and positron in the center of mass reference frame first was measured in [2].

This process allows us to describe the process where $e^-e^+ \rightarrow \pi^0\pi^0\gamma$, when $\omega \rightarrow \pi^0\gamma$.

II. INVARIANT STRUCTURE OF THE PROCESS OF $e^-e^+ \rightarrow \pi^0\pi^0\gamma$

In paper [3], the matrix element and the differential cross section was obtained in terms of irreducible tensor structures:

$$M = 8\pi m \sqrt{4\pi\alpha} \frac{1}{s} J^\mu e^\nu [a_1 L_{\mu\nu}^{(1)} + a_2 L_{\mu\nu}^{(2)} + a_3 L_{\mu\nu}^{(3)}], \quad (2)$$

*Electronic address: ahmadov@theor.jinr.ru

†Electronic address: kuraev@theor.jinr.ru

‡Electronic address: volkov@theor.jinr.ru

a_1, a_2 , and a_3 define the properties of polarizability neutral pions, where $J_\mu = \bar{v}(p_+) \gamma_\mu u(p_-)$ is the electromagnetic current,

$$\begin{aligned} L_{\mu\nu}^{(1)} &= (qk)g_{\mu\nu} - k_\mu q_\nu, \\ L_{\mu\nu}^{(2)} &= (kq)Q_\mu Q_\nu - (Qq)(Q_\mu q_\nu + Q_\nu k_\mu) + (Qq)^2 g_{\mu\nu}; \\ L_{\mu\nu}^{(3)} &= (Qq)(q^2 g_{\mu\nu} - q_\mu q_\nu) + Q_\nu(kqq_\mu - q^2 k_\mu), \\ Q &= \frac{1}{2}(q_1 - q_2), \quad q_1^2 = q_2^2 = m^2, \quad k^2 = 0, \quad q^2 = s = 4E^2, \end{aligned} \quad (3)$$

m is the pion mass, E is the energy of electron (positron), and $2E$ is the total energy in the center of mass reference frame. The tensors $L_{\mu\nu}^{(i)}$ obey the gauge conditions

$$L_{\mu\nu}^{(1)} q_\mu = 0; \quad L_{\mu\nu}^{(i)} k_\nu = 0, \quad i = 1, 2, 3. \quad (4)$$

The coefficients a_i are some functions of energy fractions of pions $x_{1,2} = 2q_i q/s$. Moreover, due to Bose statistics they obey the symmetry conditions

$$a_1(x_1, x_2) = a_1(x_2, x_1), \quad a_2(x_1, x_2) = a_2(x_2, x_1), \quad a_3(x_1, x_2) = -a_3(x_2, x_1). \quad (5)$$

The parameter a_1 coincide in static limit $q^2 = 0$ with neutral pion electric polarizability $\alpha_1 = m^3 \alpha_0$, $|\alpha_0| \approx 6.6 \times 10^{-43} \text{ cm}^3$ [4], The parameters a_2, a_3 are some dynamic characteristics of the neutral pion. For the differential cross section it was obtained (in [3] factor 1/16 in the right-hand side of equation was lost):

$$d\sigma^{e\bar{e} \rightarrow \pi_0 \pi_0 \gamma} = \frac{1}{2!} \frac{\alpha}{16\pi^2 m^4} G \frac{d^3 q_1}{E_1} \frac{d^3 q_2}{E_2} \frac{d^3 k}{\omega} \delta^4(p_+ + p_- - q_1 - q_2 - k), \quad (6)$$

with

$$\begin{aligned} G &= c_{11}|\alpha_1|^2 + c_{22}|\alpha_2|^2 + c_{33}|\alpha_3|^2 + 2c_{12}\text{Re}[\alpha_1\alpha_2^*] + 2c_{13}\text{Re}[\alpha_1\alpha_3^*] + 2c_{32}\text{Re}[\alpha_3\alpha_2^*], \\ \alpha_1 &= m^3 a_1; \quad \alpha_2 = m^5 a_2, \quad \alpha_3 = m^5 a_3, \end{aligned} \quad (7)$$

and

$$\begin{aligned} c_{11} &= \frac{1}{E^2} (2\omega^2 - \vec{k}^2); \\ c_{22} &= \frac{1}{E^2 m^4} [8E^2 Q_0^4 + 2\omega Q_0 Q^2 \vec{k} \cdot \vec{Q} - (\omega^2 Q^2 + 4E^2 Q_0^2) \vec{Q}^2 - Q_0^2 Q^2 \vec{k}^2]; \\ c_{33} &= \frac{4}{m^4} [8E^2 Q_0^2 - Q^2 \vec{k}^2 + 4E Q_0 \vec{k} \cdot \vec{Q}]; \\ c_{12} &= \frac{1}{E^2 m^2} [-\omega^2 \vec{Q}^2 + 2Q_0(\omega - E) \vec{k} \cdot \vec{Q} - q_0^2 \vec{k}^2 + 4E\omega Q_0^2]; \\ c_{13} &= \frac{1}{E^2 m^2} [8E^2 \omega Q_0 + 2E\omega \vec{k} \cdot \vec{Q} - 2E Q_0 \vec{k}^2]; \\ c_{23} &= \frac{1}{E^2 M^4} [16E^3 Q_0^3 + 2E(\omega Q^2 + 2EQ_0^2) \vec{k} \cdot \vec{Q} - 4E^2 Q_0 \omega \vec{Q}^2 - 2E Q_0 Q^2 \vec{k}^2]. \end{aligned} \quad (8)$$

Here $\vec{q}_{1,2}, \vec{k} = -(\vec{q}_1 + \vec{q}_2)$ are the two-dimensional vectors in the plane x, y transversal to the beam axis z direction

$$\begin{aligned} Q^2 &= m^2 - \frac{1}{2}s(1-x), \quad Q_0 = \frac{1}{4}(x_1 - x_2)\sqrt{s}; \quad \vec{k}^2 = k_x^2 + k_y^2; \\ \vec{Q}^2 &= \frac{1}{4}[(q_{1x} - q_{2x})^2 + (q_{1y} - q_{2y})^2]; \quad \vec{k} \cdot \vec{Q} = \frac{1}{2}[\vec{q}_2^2 - \vec{q}_1^2]. \end{aligned} \quad (9)$$

Below, we will concentrate our attention on the Dalitz plot distribution over the energy fraction of pions.

The main mechanism of reaction includes two intermediate vector meson states $V = \rho, \omega$:

$$\begin{aligned} \gamma^* &\rightarrow \rho \rightarrow \omega\pi, \quad \omega \rightarrow \gamma\pi^0, \\ \gamma^* &\rightarrow \omega \rightarrow \rho\pi, \quad \rho \rightarrow \gamma\pi^0. \end{aligned} \quad (10)$$

The contribution of other possible channels is negligible [5]

$$\Gamma_{\rho \rightarrow ee} \times B_{\omega \rightarrow \pi\gamma} = 7 \text{KeV} \times 9 \times 10^{-2} >> \Gamma_{\omega \rightarrow ee} \times B_{\rho \rightarrow \pi\gamma} = 0.8 \text{KeV} \times 6 \times 10^{-4}. \quad (11)$$

We do not consider intermediate states with $\sigma(600)$ scalar meson as well as an intermediate state with box-type quark loops. Almost complete cancelation of the relevant contributions, which takes place due to current algebra arguments commuting with the neutral current operators, was shown in [6].

We also remind that the emission of real photon by the initial leptons is strictly forbidden.

III. AMPLITUDE OF THE PROCESS OF $e^- e^+ \rightarrow \gamma^* \rightarrow \rho(\rho') \rightarrow \pi^0 \omega \rightarrow \pi^0 \pi^0 \gamma$

Using [1] we can derive the amplitude

$$M = \frac{\sqrt{(4\pi\alpha)^3 g_{\rho\omega\pi}^2}}{g_\rho^2} \frac{M_\rho^2}{[q^2 - M_\rho^2 + i\Gamma_\rho M_\rho] q^2} J^\mu e(k)^\sigma T_{\mu\sigma} = M_\mu J^\mu, \quad (12)$$

where $g_\rho \approx 6$ is the ρ meson decay constant, M_ρ , and Γ_ρ are the mass and width of the ρ -meson and

$$T_{\mu\sigma} = \frac{V_{q1} V_{k2}}{s(1-x_1) - M_\omega^2 + iM_\omega\Gamma_\omega} (\mu\lambda qq_1)(\lambda\sigma q_2 k) + \frac{V_{q2} V_{k1}}{s(1-x_2) - M_\omega^2 + iM_\omega\Gamma_\omega} (\mu\lambda qq_2)(\lambda\sigma q_1 k), \quad (13)$$

where we use the notation $(abcd) = \epsilon_{\alpha\beta\gamma\sigma} a^\alpha b^\beta c^\gamma d^\sigma$. The quantities V_{qi} , V_{ki} take into account the deviation of the vertex $V_{\mu\nu}^{\rho(q,\mu)\omega(p,\nu)\pi}$ from its point-like approximation $g_{\rho\omega\pi}(\mu\nu qp)$; $g_{\rho\omega\pi} \approx 16(\text{GeV})^{-1}$ will be given in Appendix A.

IV. DALITZ-PLOT DISTRIBUTION

Applying the invariant integration method

$$\int M_\mu (M_\nu)^* d\Gamma = \frac{1}{3} (g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}) \int |M_\lambda|^2 d\Gamma, \quad (14)$$

and using the expression for element of phase volume of the final state

$$d\Gamma = \frac{d^3 q_1}{2E_1} \frac{d^3 q_2}{2E_2} \frac{d^3 k}{2\omega} \frac{1}{(2\pi)^5} \delta^4(p_+ + p_- - q_1 - q_2 - k) = \frac{s\pi^2}{(2\pi)^5} dx_1 dx_2 \theta[(1-x_1)(1-x_2)(x_1+x_2-1) - (m^2/s)(4(1-x_1)(1-x_2) + (x_1-x_2)^2)], \quad (15)$$

we obtain for the Dalitz-plot distribution

$$\begin{aligned} \frac{d\sigma}{\sigma_0 dx_1 dx_2} &= F(x_1, x_2, s); \\ F(x_1, x_2, s) &= \frac{(1-x_1)^2(x-2(1-x_2))}{(1-x_1-M_\rho^2/s)^2} (V_{q1} V_{k2})^2 + \frac{(1-x_2)^2(x-2(1-x_1))}{(1-x_2-M_\rho^2/s)^2} (V_{q2} V_{k1})^2 + \\ &\quad 2 \frac{(1-x_1)(1-x_2)(-1+x^2+x_1x_2)}{(1-x_1-M_\rho^2/s)(1-x_2-M_\rho^2/s)} V_{q1} V_{k2} V_{q2} V_{k1}; \\ \sigma_0 &= \frac{s^2 g_{\rho\omega\pi}^4 \alpha^3}{3(2\pi)^5 g_\rho^4}. \end{aligned} \quad (16)$$

The function $F(x_1, x_2)$ is presented in Fig. 1. Since the set of tensors $L_{\mu\nu}^{(i)}$ is complete, we can present tensor $T_{\mu\nu}$ in the form

$$\begin{aligned} T &= c_1 L^{(1)} + c_2 L^{(2)} + c_3 L^{(3)}, c_i = \frac{\Delta_i}{\Delta}, \\ \Delta &= \det ||L^{(i)} \times L^{(j)}||. \end{aligned} \quad (17)$$




Fig. 1: Function $F(x_1, x_2)$ (see (16)) dependence of x_1, x_2 .

All the necessary convolutions needed to calculate c_i are given in Appendix B. The numerical calculation for $c_i(x_1, x_2)$ is presented in Tables II-IY, with ρ meson exchange in intermediate state. In Tables YI-YIII, we present the contribution of ρ' meson to this quantities.

For completeness we put the Dalitz-plot distribution in terms of the bilinear combinations of c_i :

$$\frac{d\sigma}{dx_1 dx_2} = \frac{1}{2!} \frac{4\alpha}{3s} [|\alpha_1|^2 L_{11} + |\alpha_2|^2 L_{22} + |\alpha_3|^2 L_{33} + 2\text{Re}(\alpha_1 \alpha_2^* L_{12} + \alpha_1 \alpha_3^* L_{13} + \alpha_3 \alpha_2^* L_{32})], \quad (18)$$

with the quantities L_{ij} given in Appendix B.

V. TOTAL CROSS SECTION

For the energy region $300\text{MeV} < \sqrt{s} < 1\text{GeV}$ the main contribution arises from intermediate states with the ρ, ω mesons. The total cross section

$$\sigma_{tot}(s) = \sigma_0(s) \int_0^1 dx_2 \int_{1-x_2}^1 F(x_1, x_2, s) dx_1 \quad (19)$$




Fig. 2: Value $I(s)$ (defined in (16)) as a function of s .

is presented in Fig.4 where the experimental data [2] are presented as well. Neglecting in this region the contributions of general polarizabilities related with a_2, a_3 we can approximate the total cross section in the form

$$\begin{aligned} \sigma_{tot}(s) &\approx \frac{\alpha}{12M^2} (\alpha_0 M^2 \sqrt{s})^2 I(s), \\ I(s) &= \int_0^{1-(4m^2/s)} x^3 \sqrt{1 - \frac{4m^2}{s(1-x)}} dx. \end{aligned} \quad (20)$$

The electrical value of the neutral pion polarizability is estimated as

$$|\alpha_0| \approx 0.5 \times 10^{-43} \text{ cm}^3, \quad (21)$$

which is in reasonable agreement with that obtained in paper of one of us (MKV) $\alpha_0 \approx -0.65 \times 10^{-43} \text{ cm}^3$ [4].

The function $I(s)$ is presented in Fig. 2. The main contribution in the energy range $1 \text{ GeV} < \sqrt{s} < 2 \text{ GeV}$ arises from a transition of the virtual photon to the ordinary ρ and excited ρ' mesons

$$M_\mu \sim \frac{1}{q^2} J^\mu \left[\frac{M_\rho^2}{q^2 - M_\rho^2 + iM_\rho\Gamma_\rho} + C \frac{M_{\rho'}^2}{q^2 - M_{\rho'}^2 + iM_{\rho'}\Gamma_{\rho'}} \right], \quad (22)$$

with the C -relative phase factor.

VI. DISSCUSION AND CONCLUSION

The process $e^-e^+ \rightarrow \pi^0\pi^0\gamma$ was experimentally studied, and the total cross section was measured in [2].

In this paper, we obtained the differential cross section and, in particular, the Dalitz-plot distribution. The matrix element can be expressed in terms of generalized polarizabilities of the neutral pion. Some information on their values can be extracted from the distributions obtained.

In Tables I-IY, the energy range $\sqrt{s} = 800 \text{ MeV}$, where the ρ -meson exchange dominates. In Tables Y-YIII, the energy was $\sqrt{s} = 1.45 \text{ GeV}$, where the ρ' -exchange dominates. In both the cases the interference of the ρ and ρ' exchanges was shown to be negligible.

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	2008.18	84.0845	12.247	5.04132	2.87478	1.92019	0.03241	0.02296	0.0031		
0.90		78.9729	11.3982	4.6616	2.6509	1.776	0.0111	0.0049			
0.80			9.4438	3.7205	2.0541	1.3605					0.00237
0.70				2.538	1.2472	0.7529				0.00324	0.0142
0.60					0.5216	0.1299	0			0.00529	0.0143
0.50						2.033	0.0549	0.2277	0.3276	0.3579	0.3583
0.40							0.3725	0.638	0.8365	0.9036	0.9074
0.30								2.0646	2.4092	2.5266	2.5303
0.20									9.3013	9.3964	9.3494
0.10										95.7964	94.4592
0.05											2709.61

Tab. I: The function of $F(x_1, x_2)$ (see (16)), with ρ meson exchange in intermediate state

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	0	0.0848	0.0932	0.10114	0.29932	0.3297	-0.08686	-0.1687	-0.2145	-0.2417	-0.25034
0.90		0	0.1397	4.8205	0.05259	0.2143	-0.0621	-0.1463	-0.1941	-0.2219	-0.2417
0.80			0	5.6177	-0.03589	0.1204	-0.02098	-0.09602	-0.14471	-0.19409	-0.21447
0.70				0	-0.12207	0.02772	0.02436	-0.03866	-0.09602	-0.14626	-0.16865
0.60					0	-0.10781	0.05765	0.02436	-0.02098	-0.0621	-0.08686
0.50						0	-0.10781	0.02772	0.12039	0.2143	0.3297
0.40							0	-0.12207	-0.03589	0.05258	0.2993
0.30								0	5.6177	4.8205	0.10114
0.20									0	0.13787	0.09317
0.10										0	0.08481
0.05											0

Tab. II: The function of $C_1(x_1, x_2)$ (see (17))

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	-23.5507	-3.2455	-0.05566	1.63001	-3.49016	1.4138	-0.8643	-1.64083	-1.9659	-1.9406	-1.8407
0.90		-5.7549	-5.9489	-246.831	6.8672	5.8997	-1.4455	-2.1446	-2.1798	-1.9809	-1.9406
0.80			-6.9749	-458.986	10.2559	8.3813	-1.7504	-2.3764	-2.26102	-2.1798	-1.9659
0.70				-14.4509	16.8595	10.0958	-1.8649	-2.4164	-2.3764	-2.1464	-1.6408
0.60					-68.7538	11.5783	-1.8016	-1.8649	-1.7504	-1.4455	-0.8643
0.50						0	11.5783	10.0958	8.3813	5.8997	1.4138
0.40							-68.7538	16.8595	10.2559	6.8672	-3.49016
0.30								-14.4509	-458.986	-246.831	1.63001
0.20									-6.9749	-5.9489	-0.5566
0.10										-5.7549	-3.2456
0.05											-23.5507

Tab. III: The function of $C_2(x_1, x_2)$ (see (17))

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	0	-0.55603	-0.54972	-0.54452	-1.46007	-1.49319	0.35126	0.63305	0.71704	0.5373	0
0.90		0	-0.93677	-28.3278	-0.34063	-1.13183	0.24647	0.531003	0.5198	0	-0.5373
0.80			0	-44.4922	0.02245	-1.10212	0.03122	0.2426	0	-0.5198	-0.71704
0.70				0	0.73466	-1.1792	-0.1842	0	-0.2426	-0.531	-0.63305
0.60					0	-1.00014	0	0.1842	-0.0312	-0.2465	-0.3513
0.50						0	1.00014	1.17923	1.10212	1.13183	1.49319
0.40							0	-0.76346	-0.02245	0.34063	1.46007
0.30								0	44.4922	28.3278	0.54452
0.20									0	0.93677	0.54972
0.10										0	0.55603
0.05											0

Tab. IV: The function of $C_3(x_1, x_2)$ (see (17))

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	$8.548 \cdot 10^{-6}$	$7.98 \cdot 10^{-5}$	$8.522 \cdot 10^{-4}$	0.0018	0.00269	0.00322	0.00375	0.00373	0.0035		
0.90		$7.57 \cdot 10^{-5}$	$7.95 \cdot 10^{-4}$	$1.646 \cdot 10^{-3}$	$2.39 \cdot 10^{-3}$	$2.72 \cdot 10^{-3}$	$2.88 \cdot 10^{-3}$	$2.29 \cdot 10^{-3}$	$7.21 \cdot 10^{-4}$		
0.80			0.000734	0.00139	0.00174	0.00137	$8.8 \cdot 10^{-5}$			$5.25 \cdot 10^{-4}$	0.00176
0.70				0.00139	0.0016	0.000099				0.00636	0.0071
0.60					0.002	0.0015	0		0.00097	0.023	0.021
0.50						0.0036	0.0042	0.0079	0.0419	0.0605	0.0493
0.40							0.0149	0.0345	0.1419	0.1418	0.1097
0.30								0.088	0.339	0.2915	0.219
0.20									0.7533	0.5946	0.4374
0.10										1.0366	0.7503
0.05											1.0688

Tab. V: The function of $F(x_1, x_2)$ (see (16)), with ρ' meson exchange in intermediate state

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	0	0.000949	0.0052	0.0127	0.0199	0.046	0.0983	0.212	0.534	0.1897	0.0825
0.90		0	0.01176	1.0559	-0.0188	0.0142	0.0684	0.1816	0.4565	0.1916	0.1897
0.80			0	3.095	-0.119	-0.083	-0.0216	0.1154	0.358	0.456	0.534
0.70				0	-0.1067	-0.0808	-0.032	0.0471	0.1154	0.182	0.212
0.60					0	-0.09114	-0.06059	-0.032	-0.0216	0.0684	0.0983
0.50						0	-0.09114	-0.0808	-0.0831	0.0142	0.0461
0.40							0	-0.1067	-0.1192	-0.0188	0.01995
0.30								0	3.0951	1.0559	0.0127
0.20									0	0.1175	0.0052
0.10										0	0.00095
0.05											0

Tab. VI: The function of $C_1(x_1, x_2)$ (see (17))

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	-0.0174	-0.0394	-0.0488	-0.0496	0.0576	0.0994	0.263	0.6044	1.458	0.4615	0.1846
0.90		-0.164	-0.3466	-16.87	0.417	0.3257	0.4613	0.7912	1.528	0.5207	0.4615
0.80			-1.6412	-78.913	0.9049	0.4274	0.5603	1.0157	1.7026	1.5277	1.4579
0.70				-2.887	1.353	0.4071	0.6085	0.8952	1.0157	0.7912	0.6044
0.60					-10.3302	0.3966	0.5765	0.6085	0.5603	0.4613	0.2633
0.50						0	0.3967	0.4071	0.4274	0.3257	0.099
0.40							-10.3302	1.3531	0.9049	0.4171	0.05758
0.30								-2.887	-78.9132	-16.8713	-0.0496
0.20									-1.6412	-0.3466	-0.04876
0.10										-0.164	-0.0349
0.05											-0.01739

Tab. VII: The function of $C_2(x_1, x_2)$ (see (17))

x_2/x_1	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0.95	0	-0.00188	-0.00924	-0.02039	-0.02943	-0.0627	-0.1242	-0.2505	-0.5945	-0.1737	0
0.90		0	-0.0237	-1.886	0.0314	-0.0179	-0.086	-0.2135	-0.507	0	0.1737
0.80			0	-7.445	0.2545	0.1893	0.1388	0.0713	0	0.5073	0.5945
0.70				0	0.297	0.179	0.08793	0	-0.0712	0.21348	0.2505
0.60					0	0.1885	0	-0.0879	-0.1387	0.086	0.124
0.50						0	-0.1885	-0.179	-0.189	0.0179	0.0627
0.40							0	-0.297	-0.2545	-0.03136	0.0294
0.30								0	7.4449	1.8859	0.0204
0.20									0	0.0237	0.00924
0.10										0	0.00188
0.05											0

Tab. VIII: The function of $C_3(x_1, x_2)$ (see (17))

VII. APPENDIX A

Deviation from the point approximation for the coupling constant $g_{\rho\omega\pi}$ is due to the three-angle loop Feynman integral

$$I_{\mu\nu}(q, q_1) = \frac{4\pi^2 N_c}{(2\pi)^4} \int \frac{d^4 k}{i\pi^2} \frac{Sp}{(k^2 - M^2)((k + q_1)^2 - M^2)((k + q_1 - q)^2 - M^2)},$$

$$Sp = \frac{1}{4} Sp(\hat{k} + M)\gamma_\nu(\hat{k} + \hat{q}_1 + M)\gamma_\mu(\hat{k} + \hat{q}_1 - \hat{q} + M)\gamma_5, \quad (23)$$

with M -constituent quark mass $M \approx 280 MeV$. The standard procedure of joining the denominators performing the loop momentum integration leads to

$$I_{\mu\nu}(q, q_1) = g_{\rho\omega\pi}(\mu\nu q(q - q_1))/g_\rho^2 V_{q,1}, g_\rho \approx 6, \quad (24)$$

with

$$g_{\rho\omega\pi} = \frac{g_\rho^2 N_c}{8\pi^2 M} \approx \frac{16}{GeV}, \quad (25)$$

and

$$V_{q,1} = 2M^2 \int \frac{d^3 x \delta(x_1 + x_2 + x_3 - 1)}{M^2 - q^2 x_2 x_3 - q_1^2 x_1 x_2 - (q - q_1)^2 x_1 x_3}. \quad (26)$$

Note that in the limit of a heavy quark mass we obtain $V_{q1} = 1$. In the approximation of zero pion mass we have for quantities used above

$$\begin{aligned} V_{q1} &= \frac{2M^2}{sx_1} \int_0^1 \frac{dz}{z} \ln \left| \frac{1 - z(1-z)s(1-x_1)/M^2}{1 - z(1-z)s/M^2} \right|; \\ V_{k2} &= -\frac{2M^2}{s(1-x_1)} \int_0^1 \frac{dz}{z} \ln |1 - z(1-z)s(1-x_1)/M^2|. \end{aligned} \quad (27)$$

The remaining quantities V_{q2}, V_{k1} can be obtained from these expressions by the replacement $x_1 \rightarrow x_2$. To provide the QCD principle of quark confinement, we neglect the possible imaginary parts of the three-angle amplitude.

VIII. APPENDIX B

The convolutions of the tensors $L^{(i)} \times L^{(j)}$ are (indices suppressed):

$$\begin{aligned} L_{11} &= \frac{1}{s^2} L^{(1)} \times L^{(1)} = \frac{1}{2} x^2; \\ L_{22} &= \frac{1}{s^4} L^{(2)} \times L^{(2)} = \frac{1}{128} [8\bar{x}^2\bar{x}_1\bar{x}_2 + (x_1 - x_2)^2]; \\ L_{33} &= \frac{1}{s^4} L^{(3)} \times L^{(3)} = \frac{1}{16} [(1+x)(x_1 - x_2)^2 + x^2\bar{x}]; \\ L_{12} &= \frac{1}{s^3} L^{(1)} \times L^{(2)} = \frac{1}{16} [-x^2\bar{x} + (x_1 - x_2)^2]; \\ L_{13} &= \frac{1}{s^3} L^{(1)} \times L^{(3)} = \frac{1}{4} x(x_1 - x_2); \\ L_{23} &= \frac{1}{s^4} L^{(2)} \times L^{(3)} = \frac{1}{32} (x_1 - x_2)^3. \end{aligned} \quad (28)$$

The conversion of the tensors $L^{(i)}$ with the tensor T are

$$\begin{aligned} L^{(1)} \times T &= \frac{s^2}{4} g_{\rho\omega\pi} \left[\frac{x\bar{x} - \bar{x}_2}{1 - (M_\omega^2/s)} V_{q1} V_{k2} + (x_1 \leftrightarrow x_2) \right]; \\ L^{(2)} \times T &= \frac{s^3}{32} g_{\rho\omega\pi} \left[\frac{x\bar{x}\bar{x}_2 - (x - x_2)(x_1 - x_2)^2}{1 - (M_\omega^2/s)} V_{q1} V_{k2} + (x_1 \leftrightarrow x_2) \right]; \\ L^{(3)} \times T &= \frac{s^3}{8} g_{\rho\omega\pi} \left[\frac{(x - x_2)(x_1 - x_2) - \bar{x}_2\bar{x}}{1 - (M_\omega^2/s)} V_{q1} V_{k2} - (x_1 \leftrightarrow x_2) \right]. \end{aligned} \quad (29)$$

Acknowledgments

We are grateful to A.B.Arbuzov for useful discussions. This work was supported by RFBR, grant no. 10-02-01295a. This work was also supported by the Heisenberg–Landau program grant HLP-2010-06, and JINR–Belorus–2010 grant.

- [1] M. Volkov, A. Arbuzov and E. Kuraev, Phys. Rev. C 83 (2011) 048201, arXiv: [hep-ph: 1012.2455]
- [2] R. R. Akhmetishin et al , Phys. Letters B 562(2003),173.
- [3] S. I. Eidelman, E. A. Kuraev, Preprint 85-101, Novosibirsk, INP, 1985.
- [4] M. K. Volkov, , (Particles and Nuclei), v 17, N3, p434 (1986).
- [5] Journal of Physics G, v 33, (2006), Review of particle Physics.
- [6] A. Dorokhov,J. Hufner,M. Volkov and S. Klevansky, Z. Phys. C75(1997), 122;
E. Kuraev and M. Volkov, Yad. Fiz. 62,(1999),133.