

## Testing New TeV-scale Seesaw Mediators at the LHC

---

Ivica Picek<sup>\*,1</sup> and Branimir Radovčić<sup>1,†</sup>

<sup>1</sup>*Department of Physics, University of Zagreb, Croatia*

*E-mail:* picek@phy.hr, bradov@phy.hr

We propose TeV-scale Dirac fermions producing Majorana masses of the known neutrinos via tree-level seesaw, different from standard type I and III seesaw. The employed weak five-plet with nonzero hypercharge leads to new seesaw formula  $m_\nu \sim v^6/M^5$  and to empirical masses  $m_\nu \sim 10^{-1}$  eV for  $M \sim$  TeV new states. For a limited range of the parameter space, where  $M \leq$  a few 100 GeV, the proposed mechanism is testable at the LHC via characteristic decays of Dirac type heavy leptons, produced by a Drell-Yan fusion.

*35th International Conference of High Energy Physics - ICHEP2010,  
July 22-28, 2010  
Paris France*

---

<sup>\*</sup>Speaker.

<sup>†</sup>This work is supported by the Croatian Ministry of Science, Education and Sports under Contract No. 119-0982930-1016.


Our model [1] is based on the standard model group (SMG)  $SU(3)_C \times SU(2)_L \times U(1)_Y$  content extended by vectorlike Dirac 5-plets of leptons,  $\Sigma_{L,R} = (\Sigma^{+++}, \Sigma^{++}, \Sigma^+, \Sigma^0, \Sigma^-)_{L,R} \sim (1, 5, 2)$ , which form a Dirac mass term  $\mathcal{L}_{mass} = -M_\Sigma \bar{\Sigma}_L \Sigma_R + H.c.$ . They, in conjunction with additional scalar four-plets  $\Phi_1 = (\Phi_1^0, \Phi_1^-, \Phi_1^{--}, \Phi_1^{---})$  and  $\Phi_2 = (\Phi_2^+, \Phi_2^0, \Phi_2^-, \Phi_2^{--}) \sim (1, 4, -3)$  and  $(1, 4, -1)$  respectively, build the gauge invariant Yukawa terms

$$\mathcal{L}_Y = Y_1 \bar{l}_L \Sigma_R \Phi_1 + Y_2 \bar{\Sigma}_L (l_L)^c \Phi_2^* + H.c. . \quad (1)$$

Due to the induced  $vevs$   $v_{\Phi_1}$  and  $v_{\Phi_2}$  of new scalar fields, the light neutrino acquires a mass

$$m_\nu \sim \frac{Y_1 Y_2 v_{\Phi_1} v_{\Phi_2}}{M_\Sigma} = \frac{Y_1 Y_2 \lambda_1 \lambda_2 v^6}{M_\Sigma \mu_{\Phi_1}^2 \mu_{\Phi_2}^2} \sim v^6 / M^5, \quad (2)$$

corresponding to seesaw mechanism generated by dimension nine operator shown in the figure.



The new physics scale  $\Lambda_{NP}$  of our model can be estimated from eq. (2) for some reasonable values of Yukawa and quartic coupling strengths  $\lambda_1$  and  $\lambda_2$ , say  $Y_1 \sim Y_2 \sim \lambda_1 \sim \lambda_2 \sim 10^{-2}$ . Assuming degenerate all high scale mass parameters, we obtain  $\Lambda_{NP} \simeq 580 \text{ GeV}$ .

Accordingly, the 5-plet seesaw mediators can be produced by  $W^\pm$ ,  $Z^0$  and  $\gamma$  Drell-Yan processes at the LHC. The associated production of the pairs  $(\Sigma^{+++}, \bar{\Sigma}^{++})$ ,  $(\Sigma^{++}, \bar{\Sigma}^+)$ ,  $(\Sigma^+, \bar{\Sigma}^0)$ ,  $(\Sigma^0, \bar{\Sigma}^-)$  via a charged current would be a crucial test of the five-plet nature of new leptons. Some highest production rates are displayed in the enclosed table for  $\sqrt{s} = 14 \text{ TeV}$  (to be reduced by roughly a factor of 5 for the present LHC energy of  $\sqrt{s} = 7 \text{ TeV}$ ). They are higher than for type III seesaw mediators, because of both additional and enhanced couplings to the gauge bosons.

The produced five-plet states might be recognized via characteristic decays addressed in detail elsewhere [2]. Since these states are characterized by small mass splitting within a multiplet [3], their cascade decays are suppressed. A distinguished triply charged fermion which doesn't mix with SM leptons has a characteristic decay,  $\Sigma^{+++} \rightarrow W^+ W^+ l^+$ . The singly charged and neutral states mix with SM particles, so that the produced pairs lead to

$$\Sigma^+ \bar{\Sigma}^0 \rightarrow l^+ Z l^+ W^- \rightarrow l^+ l^+ + 4 \text{ jets} , \quad \Sigma^0 \bar{\Sigma}^0 \rightarrow l^\pm W^\mp l^\pm W^\mp \rightarrow l^\pm l^\pm + 4 \text{ jets} , \quad (3)$$

the lepton number violating decays with same sign dileptons and the jets as an appealing signature.

## References

- [1] I. Picek and B. Radovčić, Phys. Lett. **B 687** 338 (2010) [0911.1374 [hep-ph]].
- [2] K. Kumerički, I. Picek and B. Radovčić, [in preparation].
- [3] M. Cirelli, N. Fornengo and A. Strumia, Nucl. Phys. **B753** (2006) 178 [hep-ph/0512090].