

Spin and orbital angular momentum of the tensor gauge field

Xiang-Song Chen^{1,2,3,*}, Ben-Chao Zhu¹, and Niall Ó Murchadha^{4†}

¹*Department of Physics, Huazhong University of
Science and Technology, Wuhan 430074, China*

²*Joint Center for Particle, Nuclear Physics and Cosmology, Nanjing 210093, China*

³*Kavli Institute for Theoretical Physics China,
Chinese Academy of Science, Beijing 100190, China*

⁴*Physics Department, University College Cork, Cork, Ireland*

(Dated: April 3, 2024)

Abstract

Following the recent studies of the trickiness in spin and orbital angular momentum of the vector gauge fields, we perform here a parallel analysis for the tensor gauge field, which has certain relation to gravitation. Similarly to the vector case, we find a nice feature that after removing all gauge degrees of freedom the angular momentum of the tensor gauge field vanishes for a stationary system. This angular momentum also shows a one-parameter invariance over the infinitely many ways of complete gauge fixing for the tensor field. The tensor gauge coupling, however, does exhibit a critical difference from the vector gauge coupling that it may induce intrinsic interaction terms into the spatial translation and rotation generators, leaving none of the ten Poincaré generators interaction-free.

PACS numbers: 11.15.-q, 04.20.Cv

Recently, the old problem of spin and orbital angular momentum of the gauge field [1] has revived considerably along two lines. One is the usage of photon orbital angular momentum in laser beams [2], the other is the study of gluon contribution to the nucleon spin [3]. The trickiness in these studies is that the gauge degrees of freedom make it hard to unambiguously construct a canonical quantity like spin, and much controversy arose [4–10]. In the debate of how to properly define a meaningful spin and angular momentum for the gauge field, Chen *et al* found a nice feature that the angular momentum of the vector gauge field can be made vanishing for a stationary system [6]. This feature is physically reasonable and leads to simple pictures of spin structure for atoms and heavy hadrons [6], thus can serve as a guidance or criteria in proper identification of the angular momentum for the gauge field. In this paper, we perform a parallel analysis for the tensor gauge field, and discuss the remarkable similarities and differences in comparison to the vector case.

We consider a symmetric tensor field $h_{\mu\nu}$ with a linear gauge transformation

$$h_{\mu\nu}(x) \rightarrow h'_{\mu\nu}(x) = h_{\mu\nu}(x) + \partial_\mu \xi_\nu(x) + \partial_\nu \xi_\mu(x), \quad (1)$$

where $\xi_\mu(x)$ are four arbitrary gauge parameters. We consider a general model with $h_{\mu\nu}$ coupled to an external conserved source $T^{\mu\nu}(x)$, and require the model be invariant under the gauge transformation in (1). Restricted to quadratic terms in first derivatives, the Lagrangian density of such a model is essentially unique up to irrelevant total divergences [11]:

$$\begin{aligned} \mathcal{L} = \frac{1}{4}(\partial_\mu h^\alpha{}_\alpha \partial^\mu h^\beta{}_\beta - \partial_\mu h_{\alpha\beta} \partial^\mu h^{\alpha\beta} + 2\partial_\mu h^{\mu\alpha} \partial^\nu h_{\nu\alpha} \\ - 2\partial_\mu h^\alpha{}_\alpha \partial_\nu h^{\mu\nu}) + \frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu}. \end{aligned} \quad (2)$$

This can be regarded as the weak-field limit of Einstein's general relativity, with $h_{\mu\nu}$ the metric perturbation and $T^{\mu\nu}$ the energy-momentum tensor of matter. But in this paper we just consider a most general case, and do not assign any specific physical contents to $h_{\mu\nu}$ and $T^{\mu\nu}$.

Given the Lagrangian, we can proceed to construct the angular momentum of the tensor gauge field $h_{\mu\nu}(x)$. We take the canonical expression

$$J_{ij} = \int d^3x \frac{\partial \mathcal{L}}{\partial \dot{h}^{\mu\nu}} [(x_j \partial_i - x_i \partial_j) h^{\mu\nu} - i(\Sigma_{ij})^{\mu\nu}{}_{\alpha\beta} h^{\alpha\beta}]. \quad (3)$$

[Conventions: an over dot denotes time derivative, Greek indices run 0-3, Latin indices run 1-3. Summation is assumed for repeated indices, even when two spatial indices are

both upstairs or downstairs. This would cause no trouble since we take the metric $\eta_{\mu\nu} = \text{diag.}(-1, 1, 1, 1)$.] $\frac{\partial \mathcal{L}}{\partial h^{\mu\nu}} \equiv \Pi_{\mu\nu}$ is the momentum conjugate of $h^{\mu\nu}$, and $(\Sigma_{ij})^{\mu\nu}_{\alpha\beta}$ is the spin matrix governing the Lorentz transformation of $h^{\mu\nu}$. The angular momentum pseudovector is $J_k = \frac{1}{2}\epsilon_{ijk}J_{ij}$, which acts as the rotation generator.

Under an infinitesimal Lorentz transformation,

$$x'^\mu = \Lambda^\mu_\nu x^\nu, \quad \Lambda^\mu_\nu = \delta^\mu_\nu + \omega^\mu_\nu, \quad \omega^{\mu\nu} = -\omega^{\nu\mu}, \quad (4)$$

the tensor field transforms as

$$h'^{\mu\nu} = \Lambda^\mu_\alpha \Lambda^\nu_\beta h^{\alpha\beta} \simeq h^{\mu\nu} + (\delta^\mu_\alpha \omega^\nu_\beta + \delta^\nu_\beta \omega^\mu_\alpha) h^{\alpha\beta}. \quad (5)$$

Casting the field variation into the form

$$\delta h^{\mu\nu} = \frac{i}{2} \omega^{\rho\sigma} (\Sigma_{\rho\sigma})^{\mu\nu}_{\alpha\beta} h^{\alpha\beta}, \quad (6)$$

we can read out the spin matrix to be

$$i(\Sigma_{\rho\sigma})^{\mu\nu}_{\alpha\beta} = \delta^\mu_\alpha (\delta^\nu_\rho \eta_{\sigma\beta} - \delta^\nu_\sigma \eta_{\rho\beta}) + \delta^\nu_\beta (\delta^\mu_\rho \eta_{\alpha\sigma} - \delta^\mu_\sigma \eta_{\alpha\rho}). \quad (7)$$

The momentum conjugates are

$$\Pi_{00} = \frac{1}{2} \partial_i h_{0i}, \quad (8a)$$

$$\Pi_{0i} = \frac{1}{2} (\partial_i h_{00} - \partial_i h + 2\partial_j h_{ij}), \quad (8b)$$

$$\Pi_{ij} = \frac{1}{2} [\dot{h}_{ij} + \delta_{ij} (\partial_k h_{0k} - \dot{h})]. \quad (8c)$$

Here $h \equiv h_{ii}$ is the spatial trace. [We remark that we have identified \dot{h}_{0i} with \dot{h}_{i0} , but not \dot{h}_{ij} with \dot{h}_{ji} , thus we are going to sum over both (ij) and (ji) , but not $(i0)$.]

It is now straightforward to compute the angular momentum tensor J_{ij} . The spin part is found to be

$$\begin{aligned} S_{ij} &\equiv \int d^3x \Pi_{\mu\nu} i(\Sigma_{ij})^{\mu\nu}_{\alpha\beta} h^{\alpha\beta} \\ &= \frac{1}{2} \int d^3x [2(\dot{h}_{jk} h_{ik} - \dot{h}_{ik} h_{jk}) \\ &\quad + h_{0j} (2\partial_k h_{ik} + \partial_i h_{00} - \partial_i h) \\ &\quad - h_{0i} (2\partial_k h_{jk} + \partial_j h_{00} - \partial_j h)]. \end{aligned} \quad (9)$$

The orbital part is found to be

$$\begin{aligned}
L_{ij} &\equiv \int d^3x \Pi_{\mu\nu} (x_j \partial_i - x_i \partial_j) h^{\mu\nu} \\
&= \frac{1}{2} \int d^3x [\dot{h}_{kl} (x_j \partial_i - x_i \partial_j) h_{kl} \\
&\quad - (2\partial_l h_{kl} + \partial_k h_{00} - \partial_k h) (x_j \partial_i - x_i \partial_j) h_{0k} \\
&\quad + \partial_k h_{0k} (x_j \partial_i - x_i \partial_j) h_{00} \\
&\quad + (\partial_k h_{0k} - \dot{h}) (x_j \partial_i - x_i \partial_j) h].
\end{aligned} \tag{10}$$

For comparison, we quote the corresponding expressions for the vector gauge field A^μ , denoted by a superscript A .

$$\mathcal{L}^A = \frac{1}{2} (\partial_\mu A_\nu \partial^\nu A^\mu - \partial_\mu A_\nu \partial^\mu A^\nu) + e A_\mu j^\mu, \tag{11}$$

$$\begin{aligned}
S_{ij}^A &= \int d^3x [\dot{A}_j A_i - \dot{A}_i A_j \\
&\quad + \partial_j A^0 A_i - \partial_i A^0 A_j],
\end{aligned} \tag{12}$$

$$\begin{aligned}
L_{ij}^A &= \int d^3x [\dot{A}_k (x_j \partial_i - x_i \partial_j) A_k \\
&\quad + \partial_k A^0 (x_j \partial_i - x_i \partial_j) A_k].
\end{aligned} \tag{13}$$

We note the following similarities and differences between S_{ij} , L_{ij} , J_{ij} and S_{ij}^A , L_{ij}^A , $J_{ij}^A \equiv S_{ij}^A + L_{ij}^A$:

(i) They are all gauge-dependent. Such a gauge-dependence has long obscured the physical meanings of photon spin and orbital angular momentum [1].

(ii) They all contain terms that involve no time derivative, and thus can survive for a stationary configuration. We will loosely call these terms “static”, though they can certainly be time-dependent as well.

(iii) S_{ij} and L_{ij} appear much more complicated than S_{ij}^A and L_{ij}^A . A major cause is that \dot{A}^0 drops out in \mathcal{L}^A , but \dot{h}_{00} and \dot{h}_{0i} survive in \mathcal{L} (though not quadratically).

(iv) L_{ij} contains a novel trace term with \dot{h} .

In common textbooks on classical electrodynamics, it is popular to discuss angular momentum of a static electromagnetic field. But this notion is really peculiar, since the electromagnetic field is massless, and should possess no momentum when “not moving”. Indeed, it was show in Ref. [6] that the total angular momentum of the vector gauge field can be constructed to vanish identically for a stationary system. When adopting the above

gauge-dependent expressions, this feature occurs *in and only in* the Coulomb gauge. This phenomenon is fairly delicate and needs some elaboration. First, the static terms in S_{ij}^A and L_{ij}^A sum to be

$$\begin{aligned} & \int d^3x [\partial_j A^0 A_i - \partial_i A^0 A_j + \partial_k A^0 (x_j \partial_i - x_i \partial_j) A_k] \\ &= - \int d^3x A^0 (x_j \partial_i - x_i \partial_j) (\partial_k A_k). \end{aligned} \quad (14)$$

Thus, in Coulomb gauge, $\vec{\partial} \cdot \vec{A} = 0$, the static terms in J_{ij}^A vanish and J_{ij}^A simplifies to

$${}^C J_{ij}^A = \int d^3x [\dot{A}_j A_i - \dot{A}_i A_j + \dot{A}_k (x_j \partial_i - x_i \partial_j) A_k]^C. \quad (15)$$

The superscript C denotes imposition of Coulomb gauge. Each term now contains a time-derivative. However, there remains a gap to claim that ${}^C J_{ij}^A$ vanishes for a stationary system: The stationary condition only means that the gauge-invariant physical observables (like the electric current j^μ or electromagnetic field $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$) are time-independent, while the gauge-potential A^μ may contain spurious (nonphysical) time-dependence [12]. This gap is closed by noting that in Coulomb gauge A^μ can be expressed in terms of $F^{\mu\nu}$ [13]:

$${}^C A^\mu = \frac{1}{\vec{\partial}^2} \partial_i F^{i\mu}. \quad (16)$$

Hence, in Coulomb gauge, A^μ is time-independent if $F^{\mu\nu}$ is, and Eq. (15) dictates that ${}^C J_{ij}^A$ vanishes for a stationary system.

Eq. (16) shows a delicate *dual* relation between gauge-fixed and gauge-invariant expressions, as we carefully discussed in [14]: If one solely looks at the right-hand-side of Eq. (16), one can in principle forget all about Coulomb gauge, and define $\frac{1}{\vec{\partial}^2} \partial_i F^{i\mu}$ as a gauge-invariant “physical field” \hat{A}^μ . (Certainly this \hat{A}^μ agrees with A^μ in Coulomb gauge, and its spatial part \hat{A}_i is just the transverse field $A_i^\perp = A_i - \partial_i \frac{1}{\vec{\partial}^2} \vec{\partial} \cdot \vec{A}$.) Analogously, in Eq. (15), if one substitutes ${}^C A_i$ with the explicit expression in Eq. (16) (this is equivalent to replacing ${}^C A_i$ with \hat{A}_i), then in the final expression for J_{ij}^A one can again forget all about Coulomb gauge, and regard the expression as the definition of a gauge-invariant \hat{J}_{ij}^A , which then vanishes identically for a stationary system, in any gauge for A^μ .

As we explained in [14], such a dual relation is only possible if the gauge-fixing is indeed complete. The special role of Coulomb gauge (for a vector field) is exactly that it completely removes the gauge degrees of freedom under a trivial boundary condition. Thus the finding

of Ref. [6] is that the physical degrees of freedom of the vector gauge field contribute no angular momentum for a stationary system. Remarkably, we find that the same feature can be demonstrated for the much more complicated tensor gauge field.

Following the hints from the vector case, we look at the canonical expressions in Eqs. (9) and (10), and examine their properties by applying a complete gauge constraint on $h_{\mu\nu}$. Such a complete tensor gauge condition, however, is not unique [14]. It can take a general form:

$$\partial_i h_{0i} + a \partial_0 h_{ii} = 0, \quad \partial_i h_{ji} + b \partial_j h_{ii} = 0. \quad (17)$$

The parameters a, b can take any value except $b = -1$, which is excluded because $\partial_i h_{ji} - \partial_j h_{ii}$ has a gauge-invariant divergence and thus is unable to fix any gauge. That the constraints in (17) make a complete gauge condition can be seen in two ways [14]. First, (17) permits no more gauge freedom; and second, the gauge-transformation parameter ξ_μ that brings $h_{\mu\nu}$ to the gauge (17) is unique. The special properties of some particular choices of a, b are discussed in [14]. Till the end of our derivation, we will see an interesting one-parameter gauge-invariance for the angular momentum of the tensor gauge field.

Taking the gauge condition in (17), and applying some slight algebra, we find the simplified expressions:

$$S_{ij}^{(ab)} = \frac{1}{2} \int d^3x [2(\dot{h}_{jk} h_{ik} - \dot{h}_{ik} h_{jk}) + (h_{00} - (2b+1)h)(\partial_j h_{0i} - \partial_i h_{0j})]^{(ab)}. \quad (18)$$

$$L_{ij}^{(ab)} = \frac{1}{2} \int d^3x [\dot{h}_{kl}(x_j \partial_i - x_i \partial_j) h_{kl} - (2ab + 2a + 1) \dot{h}(x_j \partial_i - x_i \partial_j) h - (h_{00} - (2b+1)h)(\partial_j h_{0i} - \partial_i h_{0j})]^{(ab)}. \quad (19)$$

The superscript $^{(ab)}$ denotes imposition of the gauge in (17). The static terms cancel exactly between $S_{ij}^{(ab)}$ and $L_{ij}^{(ab)}$, and the total $J_{ij}^{(ab)}$ becomes

$$J_{ij}^{(ab)} = \frac{1}{2} \int d^3x [2(\dot{h}_{jk} h_{ik} - \dot{h}_{ik} h_{jk}) + \dot{h}_{kl}(x_j \partial_i - x_i \partial_j) h_{kl} - (2ab + 2a + 1) \dot{h}(x_j \partial_i - x_i \partial_j) h]^{(ab)}. \quad (20)$$

Similarly to Eq. (15), each term in $J_{ij}^{(ab)}$ contains a time-derivative. But as we remarked above, to conclude that $J_{ij}^{(ab)}$ vanishes for a stationary system, we still need to show that

$h_{ij}^{(ab)}$ cannot induce spurious time-dependence. This property can be inferred from our recent careful examination of tensor gauge conditions [13, 14]:

$$h_{ij}^{(ab)} = f_{ij} - \frac{1+2b}{2(1+b)} \frac{1}{\bar{\partial}^2} (\partial_i \partial_j f_{kk}), \quad (21a)$$

$$h_{0j}^{(ab)} = f_{0j} - \frac{1+a+b}{2(1+b)} \frac{1}{\bar{\partial}^2} (\partial_0 \partial_j f_{kk}), \quad (21b)$$

$$h_{00}^{(ab)} = f_{00} - \frac{1+2a}{2(1+b)} \frac{1}{\bar{\partial}^2} (\partial_0^2 f_{kk}). \quad (21c)$$

Here $f_{\mu\nu} \equiv 2 \frac{1}{\bar{\partial}^2} R_{\mu i i \nu}$, and $R_{\mu\rho\sigma\nu}$ is the linearized Riemann curvature. For completeness and future reference, we have displayed all ten components of $h_{\mu\nu}^{(ab)}$.

Eqs. (21) indicate clearly that $h_{\mu\nu}^{(ab)}$ is time-independent if the gauge-invariant $R_{\rho\sigma\mu\nu}$ is, hence $J_{ij}^{(ab)}$ vanishes for a stationary system. We thus proved the same nice feature as in the vector case that the physical degrees of freedom of the tensor gauge field carry no static angular momentum. Moreover, as in the vector case, one can also define the right-hand-side of Eqs. (21) as a gauge-invariant physical field $\hat{h}_{\mu\nu}$, and forget all about the gauge in (17). With this $\hat{h}_{\mu\nu}$, one can define a gauge-invariant \hat{J}_{ij} by replacing h_{ij} in Eq. (20) with \hat{h}_{ij} , and this \hat{J}_{ij} vanishes identically under the stationary condition, in any gauge for $h_{\mu\nu}$.

The expression in Eq. (20) is not yet the final story, as it has not reached the art of Eq. (15), where ${}^C A_i = A_i^\perp$ represents the two dynamical (propagating) components of the vector field. In Eq. (20) the trace $h^{(ab)}$ is non-dynamical, as revealed by its equation of motion [14]:

$$\bar{\partial}^2 h^{(ab)} = -\frac{\kappa}{1+b} T_{00}. \quad (22)$$

The instantaneous feature of the Laplacian operator $\bar{\partial}^2$ means that $h^{(ab)}$ is completely dictated by the source. Namely, $h^{(ab)}$ is not an independent dynamical quantity that can propagate. An important implication of this fact is that $h_{ij}^{(ab)}$, with a nonzero trace, is not fully dynamical either. Furthermore, by the gauge condition in (17), the spatial divergence of $h_{ij}^{(ab)}$ is non-dynamical as well. To get the purely dynamical component of $h_{ij}^{(ab)}$, we thus need to extract its transverse-traceless (TT) part h_{ij}^{TT} [15]. This h_{ij}^{TT} is completely invariant under gauge transformation in (1). It is the counter part of A_i^\perp for the vector field. Ref. [14] gives how h_{ij}^{TT} relates to $h_{ij}^{(ab)}$:

$$h_{ij}^{(ab)} = h_{ij}^{TT} + \frac{1+b}{2} \delta_{ij} h^{(ab)} - \frac{1+3b}{2} \frac{1}{\bar{\partial}^2} \partial_i \partial_j h^{(ab)}. \quad (23)$$

Inserting this into Eq. (20), and using Eq. (22) for $h^{(ab)}$, we obtain:

$$\begin{aligned}
J_{ij}^{(ab)} = & \frac{1}{2} \int d^3x [2(\dot{h}_{jk}^{TT} h_{ik}^{TT} - \dot{h}_{ik}^{TT} h_{jk}^{TT}) \\
& + \dot{h}_{kl}^{TT} (x_j \partial_i - x_i \partial_j) h_{kl}^{TT} \\
& - \kappa^2 (\frac{1}{2} + 2\frac{a-b}{1+b}) \frac{\dot{T}_{00}}{\partial^2} (x_j \partial_i - x_i \partial_j) \frac{T_{00}}{\partial^2}].
\end{aligned} \tag{24}$$

The last term in Eq. (24) is a bit special and calls for attention. It contains all the dependence of $J_{ij}^{(ab)}$ on the two gauge parameters a, b , but through a single factor $(\frac{1}{2} + 2\frac{a-b}{1+b})$. $J_{ij}^{(ab)}$ thus possesses a one-parameter invariance: $(\frac{1}{2} + 2\frac{a-b}{1+b})$ can take a universal value λ for any $a = b + (\frac{\lambda}{2} - \frac{1}{4})(1+b)$. One interesting example is $a = b$, which gives $\lambda = \frac{1}{2}$. The most attractive choice might be $\lambda = 0$ for any $a = \frac{1}{4}(3b - 1)$. With $\lambda = 0$, Eq. (24) reduces to the same form as for a free field in the absence of source, and mimics exactly Eq. (15), whose form is unaltered by the presence of source.

The gauge with $a = \frac{1}{4}(3b - 1)$, however, is not necessarily consistent with quantum Lorentz invariance. As Weinberg elaborated in [16], by canonical quantization of tensor gauge field with only physical degrees of freedom, Lorentz invariance of S-matrix requires a delicate matching between the Hamiltonian and propagator. This matching can be achieved in some particular gauge. E.g., Weinberg found $a = -\frac{2}{3}$ and $b = -\frac{1}{3}$, which however does not fall into the class of $a = \frac{1}{4}(3b - 1)$.

With $(\frac{1}{2} + 2\frac{a-b}{1+b}) \neq 0$, the last term in Eq. (24) is then intrinsic and novel. The appearance of the coupling constant κ means that this term represents an interaction effect. On the other hand, it is entirely expressed in terms of the source and should apparently be counted as part of the source angular momentum. One should note, however, that such a term is absent for a free source which does not couple to the tensor gauge field. Therefore, the presence of such a term seems to indicate that, unlike the vector gauge coupling in the standard model of particle physics, the tensor gauge coupling induces extra term into the angular momentum of the system. In other words, the tensor gauge coupling modifies the rotation generator of the system.

Exactly analogous situation can be demonstrated for the spatial translation generator (or the momentum) of the tensor gauge field:

$$\begin{aligned}
\vec{P} \equiv & - \int d^3x \Pi_{\mu\nu} \vec{\partial} h^{\mu\nu} = -\frac{1}{2} \int d^3x [\dot{h}_{kl} \vec{\partial} h_{kl} - \dot{h} \vec{\partial} h \\
& + 2\partial_k h_{0k} \vec{\partial} h - 2\partial_l h_{kl} \vec{\partial} h_{0k}].
\end{aligned} \tag{25}$$

Here we also find static terms, which seem to imply that a stationary tensor field can possess momentum. But after applying the constraint (17) to remove all gauge degrees of freedom, \vec{P} simplifies to

$$\begin{aligned}\vec{P}^{(ab)} &= -\frac{1}{2} \int d^3x [\dot{h}_{kl} \vec{\partial} h_{kl} - (2ab + 2a + 1) \dot{h} \vec{\partial} h]^{(ab)} \\ &= -\frac{1}{2} \int d^3x [\dot{h}_{kl}^{TT} \vec{\partial} h_{kl}^{TT} - \kappa^2 (\frac{1}{2} + 2\frac{a-b}{1+b}) \frac{\dot{T}_{00}}{\vec{\partial}^2} \vec{\partial} \frac{T_{00}}{\vec{\partial}^2}].\end{aligned}\quad (26)$$

This is clearly zero for a stationary system, showing that a static, massless tensor gauge field possesses no physical momentum. The second expression in Eq. (26) results from extracting the TT part and using Eq. (22), and we find the same factor $(\frac{1}{2} + 2\frac{a-b}{1+b})$ as in Eq. (24).

In comparison, the momentum expression for the vector gauge field is

$$\vec{P}^A \equiv - \int d^3x \frac{\partial \mathcal{L}^A}{\partial \dot{A}_k} \vec{\partial} A_k = - \int d^3x (\dot{A}_k + \partial_k A^0) \vec{\partial} A_k. \quad (27)$$

In Coulomb gauge, this reduces to

$$^C \vec{P}^A = - \int d^3x [\dot{A}_k \vec{\partial} A_k]^C = - \int d^3x \dot{A}_k^\perp \vec{\partial} A_k^\perp. \quad (28)$$

We see again that unlike the vector gauge coupling, the tensor gauge coupling induces an interaction term in the spatial translation generator. This seems to imply that with the tensor gauge coupling the ten Poincaré generators are all “bad” (in the sense of containing interaction), while for the vector gauge coupling only four generators (for time translation and Lorentz boost) are bad, and six generators (for spatial translation and rotation) remain “good” (interaction-free) [17]. This implication, however, is not decisive, since we have not included dynamical part for the source; and the subject needs further careful investigation.

This work is supported by the China NSF Grants 10875082 and 11035003. X.S.C. is also supported by the NCET Program of the China Education Department.

* Electronic address: cxs@hust.edu.cn

† Electronic address: niall@ucc.ie

- [1] See, e.g., J.M. Jauch and F. Rohrlich, *The Theory of Photons and Electrons* (Springer-Verlag, Berlin 1976); V.B. Berestetskii, E.M. Lifshitz, and L.P. Pitaevskii, *Quantum Electrodynamics* (Pergamon, Oxford 1982), 2nd ed.

- [2] For a review, see, e.g., S.J. van Enk and G. Nienhuis, Phys. Rev. A **76**, 053825 (2007).
- [3] For recent reviews, see e.g., F. Myhrer and A.W. Thomas, J. Phys. G **37**, 023101 (2010); M. Burkardt, C.A. Miller, and W.D. Nowak, Rept. Prog. Phys. **73**, 016201 (2010).
- [4] X.S. Chen, X.F. Lü, W.M. Sun, F. Wang, and T. Goldman, Phys. Rev. Lett. **100**, 232002 (2008).
- [5] X.S. Chen, W.M. Sun, X.F. Lü, F. Wang, and T. Goldman, Phys. Rev. Lett. **103**, 062001 (2009).
- [6] X.S. Chen, W.M. Sun, F. Wang, and T. Goldman, Phys. Rev. D **83**, 071901 (2011); Phys. Lett. B **700**, 21 (2011).
- [7] M. Wakamatsu, Phys. Rev. D **83**, 014012 (2011); *ibid.* **81**, 114010 (2010); Eur. Phys. J. A **44**, 297 (2010); arXiv:1012.0612.
- [8] Y.M. Cho, M.L. Ge, P. Zhang, arXiv:1010.1080.
- [9] E. Leader, arXiv:1101.5956.
- [10] Y. Hatta, arXiv:1101.5989.
- [11] T. Padmanabhan, Int. J. Mod. Phys. D **17**, 367 (2008).
- [12] For example, the Lorentz gauge $\vec{\partial} \cdot \vec{A} + \partial_t A^0 = 0$ does not strictly reduce to Coulomb gauge for a stationary system, because one can assign a time-dependent A^0 in Lorentz gauge while leaving $F^{\mu\nu}$ time-independent.
- [13] X.S. Chen and B.C. Zhu, Phys. Rev. D **83**, 084006 (2011); *ibid.* **83**, 061501 (2011).
- [14] X.S. Chen and B.C. Zhu, arXiv:1101.2809.
- [15] R. Arnowitt, S. Deser, and C.W. Misner, in *Gravitation*, L. Witten ed. (Wiley, New York, 1962), Chapter 7 (posted as arXiv:gr-qc/0405109).
- [16] S. Weinberg, Phys. Rev. **138**, B988 (1965).
- [17] S. Weinberg, *The Quantum Theory of Fields*, Vol. I (Cambridge, New York, 1995).