A NEW PROOF FOR THE EXISTENCE OF AN EQUIVARIANT ENTIRE SOLUTION CONNECTING THE MINIMA OF THE POTENTIAL FOR THE SYSTEM $\Delta u - W_u(u) = 0$

NICHOLAS D. ALIKAKOS

ABSTRACT. Recently, Giorgio Fusco and the author in [1] studied the system $\Delta u - W_u(u) = 0$ for a class of potentials that possess several global minima and are invariant under a general finite reflection group, and established existence of equivariant solutions connecting the minima in certain directions at infinity, together with an estimate. In this paper a new proof is given which, in particular, avoids the introduction of a pointwise constraint in the minimization process.

1. INTRODUCTION

The study of the system

(1)
$$\Delta u - W_u(u) = 0, \text{ for } u : \mathbb{R}^n \to \mathbb{R}^n,$$

where $W : \mathbb{R}^n \to \mathbb{R}$ and $W_u := (\partial W/\partial u_1, \ldots, \partial W/\partial u_n)^{\top}$, under symmetry hypotheses on the potential W was initiated in Bronsard, Gui, and Schatzman [3], where existence for the case n = 2 with the symmetries of the equilateral triangle was settled. About twelve years later this work was followed by Gui and Schatzman [11], where the case n = 3 for the symmetry group of tetrahedron was established. The corresponding solutions are known as the *triple junction* and the *quadruple junction* respectively. This class of solutions is characterized by the fact that they connect the N global minima of the potential W, that is,

(2)
$$\lim_{\lambda \to +\infty} u(\lambda \eta_i) = a_i, \text{ for } i = 1, \dots, N,$$

for certain unit vectors $\eta_i \in \mathbb{S}^{n-1}$, where $\mathbb{S}^{n-1} \subset \mathbb{R}^n$ is the unit sphere. These solutions are related to minimal surface complexes, and particularly to the singular points there (see Taylor [17], Dierkes *et al.* [5, 6]) via the blow-down limit $u_{\varepsilon}(x) := u(x/\varepsilon)$ (see Baldo [2]). Recently in [1] certain general hypotheses on Wwere identified and the problem was settled for general dimension n and for any reflection group G on \mathbb{R}^n .

In this paper we want to give a new derivation of this result, which is based on a positivity property of the gradient flow associated to (1) and comparison arguments involving subharmonic functions, ingredients already existing in [1], but now supplemented with a Kato-type inequality and the De Giorgi oscillation lemma. The present paper is self-contained. Our hope is that this simpler proof will be more adaptable to the general case of a potential W without symmetry requirements. In order to bring out clearly the underlying ideas, we refrain from any generalization which could complicate the technical part.

The author was supported by Kapodistrias grant No. 15/4/5622 at the University of Athens.

1.1. Notation. We denote by B_R the ball of radius R > 0 centered at the origin and by $W_{\rm E}^{1,2}(B_R;\mathbb{R}^n)$ the subspace of *equivariant maps*, that is, u(gx) = gu(x), for all $g \in G$ and $x \in \mathbb{R}^n$. We also denote by $\langle \cdot, \cdot \rangle$ the Euclidean inner product, by $|\cdot|$ the Euclidean norm, and by $d(x, \partial D)$ the distance of x from ∂D . In the case of finite groups G, the notation |G| stand for the number of elements of the group.

We denote the functional associated to (1) by

(3)
$$J(u) = \int_{\mathbb{R}^n} \left\{ \frac{1}{2} |\nabla u|^2 + W(u) \right\} \mathrm{d}x.$$

A Coxeter group is a finite subgroup of the orthogonal group $O(\mathbb{R}^n)$, generated by a set of reflections. A reflection $\gamma \in G$ is associated to the hyperplane

$$\pi_{\gamma} = \{ x \in \mathbb{R}^n \mid \langle x, \eta_{\gamma} \rangle = 0 \}$$

via

$$\gamma x = x - 2 \langle x, \eta_{\gamma} \rangle \eta_{\gamma}, \text{ for } x \in \mathbb{R}^n,$$

where $\eta_{\gamma} \in \mathbb{S}^{n-1}$ is a unit vector. Every finite subgroup of $O(\mathbb{R}^n)$ has a fundamental region¹, that is, a subset $F \subset \mathbb{R}^n$ with the following properties:

- (i) F is open and convex,
- (ii) $F \cap gF = \emptyset$, for $I \neq g \in G$, where I is the identity,
- (iii) $\mathbb{R}^n = \bigcup \{ g\overline{F} \mid g \in G \}.$

We choose the orientation of η_{γ} so that $F \subset \mathcal{P}_{\gamma}^+$, where $\mathcal{P}_{\gamma}^+ = \{x \in \mathbb{R}^n \mid \langle x, \eta_{\gamma} \rangle > 0\}$. Then, we have

(4)
$$F = \cap_{\gamma \in \Gamma} \mathcal{P}_{\gamma}^+,$$

where $\Gamma \subset G$ is the set of all reflections in G. Given $a \in G$, the *stabilizer* of a, denoted by G_a , is the subgroup of G that fixes a.

1.2. The theorem ([1]). We begin with the hypotheses.

Hypothesis 1 (N nondegenerate global minima). The potential W is of class C^2 and satisfies $W(a_i) = 0$, for i = 1, ..., N, and W > 0 on $\mathbb{R}^n \setminus \{a_1, ..., a_N\}$. Furthermore, there holds $v^{\top} \partial^2 W(u) v \ge 2c^2 |v|^2$, for $v \in \mathbb{R}^n$ and $|u - a_i| \le \bar{q}$, for some $c, \bar{q} > 0$, and for i = 1, ..., N.

Hypothesis 2 (Symmetry). The potential W is invariant under a finite reflection group G acting on \mathbb{R}^n (Coxeter group), that is,

(5)
$$W(gu) = W(u), \text{ for all } g \in G \text{ and } u \in \mathbb{R}^n.$$

Moreover, we assume that there exists M > 0 such that $W(su) \ge W(u)$, for $s \ge 1$ and |u| = M.

We seek *equivariant* solutions of system (1), that is, solutions satisfying

(6)
$$u(gx) = gu(x)$$
, for all $g \in G$ and $x \in \mathbb{R}^n$

Hypothesis 3 (Location and number of global minima). Let $F \subset \mathbb{R}^n$ be a fundamental region of G. We assume that \overline{F} (the closure of F) contains a single global minimum of W, say a_1 , and let G_{a_1} be the subgroup of G that leaves a_1 fixed. Then, as it follows by the invariance of W, the number of the minima of W is

(7)
$$N = \frac{|G|}{|G_{a_1}|}$$

¹See [10] or [14].

Hypothesis 4 (Q-monotonicity). We restrict ourselves to potentials W for which there is a continuous function $Q : \mathbb{R}^n \to \mathbb{R}$ that satisfies

(8)
$$Q(u+a_1) = |u| + H(u)$$

where $H: \mathbb{R}^n \to \mathbb{R}$ is a C^2 function such that H(0) = 0 and $H_u(0) = 0$, and

- (9a) Q is convex,
- (9b) $Q(gu) = Q(u), \text{ for } u \in \mathbb{R}^n, g \in G_{a_1},$

(9c)
$$Q(u+a_1) = |u| + H(u), \text{ in a neighborhood of } u = 0,$$

(9d) $Q(u) > 0, \text{ on } \mathbb{R}^n \setminus \{a_1\},$

and, moreover,

(10) $\langle Q_u(u), W_u(u) \rangle \ge 0, \text{ in } D \setminus \{a_1\},\$

where we have set

(11)
$$D := \operatorname{Int} \left(\bigcup_{g \in G_{a_1}} g\overline{F} \right).$$

Theorem 1.1 ([1]). Under Hypotheses 1-4, there exists an equivariant classical solution to system (1) such that

- (i) $|u(x) a_1| \leq K e^{-kd(x,\partial D)}$, for $x \in D$ and for positive constants k, K,
- (ii) $u(\overline{F}) \subset \overline{F}$ and $u(D) \subset D$.

In particular, u connects the $N = |G|/|G_{a_1}|$ global minima of W in the sense that

$$\lim_{\lambda \to +\infty} u(\lambda g\eta) = ga_1, \text{ for all } g \in G,$$

uniformly for η in compact subsets of $D \cap \mathbb{S}^{n-1}$.

2. The extended Kato inequality

We begin by presenting a straightforward extension of the classical Kato inequality. We follow the presentation in [13, p. 85]. Let $\hat{Q} : \mathbb{R}^m \to \mathbb{R}$ be a continuous function satisfying the following assumptions.

- (i) \hat{Q} is convex,
- (ii) $\hat{Q} > 0$ and $\hat{Q}_u \neq 0$, for $u \neq 0$,
- (iii) $\hat{Q} = |u| + H(u)$, for a C^2 function $H : \mathbb{R}^m \to \mathbb{R}$, such that H(0) = 0 and $H_u(0) = 0$.

Lemma 2.1. Let $u \in L^{\infty}(\mathbb{R}^n; \mathbb{R}^m)$ and suppose² that the distributional Laplacian $\Delta u \in L^1_{loc}(\mathbb{R}^n; \mathbb{R}^m)$. Then,

(12)
$$\Delta \hat{Q}(u) \ge \langle \Delta u, \hat{Q}_u(u) \rangle,$$

in the distributional sense, with the definition

$$\hat{Q}_u(u) := \begin{cases} \nabla_u \hat{Q}(u), & \text{for } u \neq 0, \\ 0, & \text{for } u = 0. \end{cases}$$

²The fact that u should be in $u \in L^{\infty}(\mathbb{R}^n; \mathbb{R}^m)$ was pointed out to us by Panagiotis Smyrnelis. If H is assumed globally Lipschitz, then $u \in L^1_{loc}(\mathbb{R}^n; \mathbb{R}^m)$ suffices.

Remarks. The well-known Kato inequality for functions $u \in L^1_{loc}(\mathbb{R}^n; \mathbb{C})$ states that

(13)
$$\Delta |u| \ge \operatorname{Re}[(\operatorname{sgn} u)\Delta u],$$

in the distributional sense. The choice $|u| = \sqrt{u\bar{u}}$ and

$$\operatorname{sgn} u = \begin{cases} 0, & \text{for } u = 0, \\ \bar{u}/|u|, & \text{for } u \neq 0, \end{cases}$$

is a special case, for $\hat{Q}(u) = |u|$ and $\operatorname{Re}[u\overline{v}] = \langle u, v \rangle$. Also, under the hypothesis $u \in W^{1,2}_{\operatorname{loc}}(\mathbb{R}^n;\mathbb{R}^m) \cap L^{\infty}(\mathbb{R}^n;\mathbb{R}^m)$ we note that $\hat{Q}(u(\cdot)) \in W^{1,2}_{\operatorname{loc}}(\mathbb{R}^n;\mathbb{R}^m)$ (see [15, p, 54] or [7, p. 130]). Therefore, (12) holds in $W^{1,2}_{\operatorname{loc}}(\mathbb{R}^n;\mathbb{R}^m)$.

Proof. We utilize the summation convention. We first establish

(14)
$$\Delta \hat{Q}(u) \ge \langle \Delta u, \hat{Q}_u(u) \rangle$$

for $u \in C^{\infty}$, except where $\hat{Q}(u)$ is not differentiable. Set

$$\hat{Q}_{\varepsilon}(u(x)) = \sqrt{\hat{Q}^2(u(x)) + \varepsilon^2}, \text{ for } \varepsilon > 0;$$

then,

(15)
$$\hat{Q}_{\varepsilon} \hat{Q}_{\varepsilon,i} = \hat{Q} \hat{Q}_{,u_k} u_{k,i}, \text{ where } \hat{Q}_{\varepsilon,i} := \frac{\partial}{\partial x_i} \hat{Q}_{\varepsilon}(u(x)),$$

and

(16)
$$(\hat{Q}_{\varepsilon,i})^2 = \left(\frac{\hat{Q}}{\hat{Q}_{\varepsilon}}\right)^2 (\hat{Q}_{,u_k}u_{k,i})^2 \le (\hat{Q}_{,u_k}u_{k,i})^2.$$

Hence,

(17)
$$\hat{Q}_{\varepsilon,i}\,\hat{Q}_{\varepsilon,i} \le (\hat{Q}_{,u_k}u_{k,i})(\hat{Q}_{,u_k}u_{k,i})$$

therefore

(18)
$$|\nabla_x \hat{Q}_{\varepsilon}|^2 \le |(\nabla_x u)^\top \hat{Q}_u|^2$$

Moreover,

(19)
$$(\hat{Q}_{\varepsilon} \, \hat{Q}_{\varepsilon,i})_{,i} = \langle \Delta u, \hat{Q} \, \hat{Q}_u \rangle + \hat{Q} \langle (\partial^2 \hat{Q}) u_{,i}, u_{,i} \rangle + |(\nabla_x u)^\top \hat{Q}_u|^2,$$

where $\partial^2 \hat{Q}$ is the Hessian of \hat{Q} and $u_{i} = (u_{1,i}, \ldots, u_{m,i})$. By convexity it follows that

$$|\nabla_{x}\hat{Q}_{\varepsilon}|^{2} + \hat{Q}_{\varepsilon}\,\Delta\hat{Q}_{\varepsilon} \ge \langle\Delta u, \hat{Q}\,\hat{Q}_{u}\rangle + |(\nabla_{x}u)^{\top}\hat{Q}_{u}|^{2},$$

from which, by (18),

(20)
$$\Delta \hat{Q}_{\varepsilon} \ge \left\langle \Delta u, \frac{\hat{Q} \, \hat{Q}_{u}}{\hat{Q}_{\varepsilon}} \right\rangle$$

At points of smoothness we can take the limit $\varepsilon \to 0$ and obtain (14).

We proceed by mollification. Let $w \in C^{\infty}(\mathbb{R}^n)$, with $w \ge 0$ and $\int w(x) dx = 1$. For $\delta > 0$ we define $w_{\delta}(x) = \delta^{-n} w(\delta^{-1}x)$ and set

$$I_{\delta}u := w_{\delta} * u, \text{ for } \delta > 0.$$

Then, $I_{\delta}u \to u$ and $\Delta(I_{\delta}u) \to \Delta u$ in L^1 , as $\delta \to 0$. Applying (20) to $I_{\delta}u$ we have

(21)
$$\Delta \hat{Q}_{\varepsilon}(I_{\delta}u) \ge \left\langle \Delta(I_{\delta}u), \frac{\frac{\partial}{\partial u}(\frac{1}{2}\hat{Q}^{2})(I_{\delta}u)}{\hat{Q}_{\varepsilon}(I_{\delta}u)} \right\rangle.$$

Taking $\delta \to 0$ and utilizing that \hat{Q}^2 is everywhere differentiable and that the fraction inside the inner product in (21) is bounded $(L^{\infty}$ requirement for $u(\cdot)$), by the dominated convergence theorem we have

(22)
$$\Delta \hat{Q}_{\varepsilon}(u) \ge \left\langle \Delta u, \frac{\frac{\partial}{\partial u}(\frac{1}{2}\hat{Q}^{2})(u)}{\hat{Q}_{\varepsilon}(u)} \right\rangle.$$

Finally, we pass to the limit in \mathcal{D}' as $\varepsilon \to 0$.

3. The gradient flow and positivity ([1])

We define the set of *positive maps* (in the class of equivariant Sobolev maps)

(23)
$$\mathcal{U}^{\operatorname{Pos}} := \left\{ u \in W^{1,2}_{\operatorname{E}}(B_R; \mathbb{R}^n) \mid u(\overline{F_R}) \subset \overline{F} \right\}$$

and the set of strongly positive maps

(24)
$$\mathcal{U}_0^{\text{Pos}} := \left\{ u \in W_{\text{E}}^{1,2}(B_R; \mathbb{R}^n) \mid u(F_R) \subset F \right\},$$

where $F_R = F \cap B_R$. Here R > 0 and clearly the sets \mathcal{U}^{Pos} and $\mathcal{U}_0^{\text{Pos}}$ depend on R. We will utilize the gradient flow

(25)
$$\begin{cases} \frac{\partial u}{\partial t} = \Delta u - W_u(u), & \text{in } B_R \times (0, \infty), \\ \frac{\partial u}{\partial n} = 0, & \text{on } \partial B_R \times (0, \infty), , \\ u(x, 0) = u_0(x), & \text{in } B_R, \end{cases}$$

where $\partial/\partial n$ is the normal derivative. We note that by Hypothesis 2

(26)
$$\langle -W_u(u), u \rangle \leq 0, \text{ for } |u| = M.$$

We will consider initial conditions in (25) satisfying in addition

$$\|u_0\|_{L^{\infty}(B_R;\mathbb{R}^n)} \le M.$$

Since W is C^2 (cf. Hypothesis 1), the results in [12, Ch. 3, §3.3, §3.5] apply and provide a unique solution to (25) in $C(0, \infty; W_{\rm E}^{1,2}(B_R; \mathbb{R}^n))$, which for t > 0, as a function of x, is in $C^{2+\alpha}(\overline{B_R}; \mathbb{R}^n)$), for some $0 < \alpha < 1$. Moreover, the solution satisfies the estimate

(28)
$$\|u(\cdot,t)\|_{L^{\infty}(B_R;\mathbb{R}^n)} \leq M, \text{ for } t \geq 0.$$

This follows from (26), (27), and by well-known invariance results [16, Ch. 14, §B].

Theorem 3.1 ([1]). Let W be a C^2 potential satisfying Hypothesis 2. If $u_0 \in \mathcal{U}^{\text{Pos}}$ and $||u_0||_{L^{\infty}(B_R;\mathbb{R}^n)} \leq M$, then

$$u(\cdot, t; u_0) \in \mathcal{U}^{\operatorname{Pos}}, \text{ for } t \ge 0,$$

and, moreover,

$$u(\cdot,t;u_0) \in \mathcal{U}_0^{\mathrm{Pos}}, \text{ for } t > 0, \text{ provided } u_0(F_R) \cap F \neq \emptyset.$$

 $\mathbf{5}$

Proof. Let $u: B_R \to \mathbb{R}^n$ be an equivariant map. We will prove that u is a positive map if and only if

(29)
$$u(\overline{(\mathcal{P}_{\gamma}^+)_R}) \subset \overline{\mathcal{P}_{\gamma}^+}, \text{ for all } \gamma \in \Gamma,$$

where $(\mathcal{P}_{\gamma}^{+})_{R} = \mathcal{P}_{\gamma}^{+} \cap B_{R}$. Suppose that (29) holds. Then

$$u(\overline{F_R}) = u(\cap_{\gamma \in \Gamma} (\overline{\mathcal{P}_{\gamma}^+})_R) \subset \cap_{\gamma \in \Gamma} u(\overline{(\mathcal{P}_{\gamma}^+)_R}) \subset \cap_{\gamma \in \Gamma} \overline{\mathcal{P}_{\gamma}^+} = \overline{F}.$$

Hence, u is positive. Conversely, suppose that u is a positive equivariant map on B_R . Then, equivalently, u_e defined by

(30)
$$u_{\mathbf{e}}(x) := \begin{cases} u(x), & \text{for } x \in B_R \\ 0, & \text{for } x \in \mathbb{R}^n \setminus B_R \end{cases}$$

is a positive equivariant map on \mathbb{R}^n . For any $g \in G$, we have from equivariance and positivity,

(31)
$$u_{\mathbf{e}}(g(\overline{F})) = g(u_{\mathbf{e}}(\overline{F})) \subset g(\overline{F}).$$

Now pick a $\gamma \in \Gamma$ and take an $x \in \mathcal{P}_{\gamma}^+$ and fix it. There is a $g \in G$, denoted by g_x , such that $x \in g_x(\overline{F})$ and $g_x(F)$ is also a fundamental region. Since for each fundamenal region F' and for each reflection γ we have either $F' \subset \mathcal{P}_{\gamma}^+$ or $F' \subset -\mathcal{P}^+_{\gamma}$, we conclude that

(32)
$$g_x(\overline{F}) \subset \overline{\mathcal{P}_{\gamma}^+}.$$

Thus, by (31), $u_{\rm e}(\overline{\mathcal{P}_{\gamma}^+}) \subset \overline{\mathcal{P}_{\gamma}^+}$, and so (29) follows. Now consider (25) with $u_0 \in \mathcal{U}^{\rm Pos}$. By the regularizing property of the equation the solution is classical for t > 0 and by (26) it exists globally in time and belongs to $C(0, +\infty; W^{1,2}_{\mathrm{E}}(B_R; \mathbb{R}^n)) \cap C^1(0, +\infty; C^{2+\alpha}(\overline{B_R}; \mathbb{R}^n))$, for some $0 < \alpha < 1$ (see [12]). Consider a reflection $\gamma \in \Gamma$ and set

$$\begin{aligned} \zeta(x,t) &= \langle u(x,t,u_0), \eta_\gamma \rangle, \text{ on } B_R \times (0,\infty), \\ \zeta_0(x) &= \langle u_0(x), \eta_\gamma \rangle, \text{ on } B_R. \end{aligned}$$

By taking the inner product of equation (25) with η_{γ} , we obtain

(33)
$$\begin{cases} \frac{\partial \zeta}{\partial t} = \Delta \zeta + c\zeta, & \text{in } B_R \times (0, \infty), \\ \frac{\partial \zeta}{\partial n} = 0, & \text{on } \partial B_R \times (0, \infty), \\ \zeta(\cdot, 0) = \zeta_0, \end{cases}$$

where we have set

$$c(x,t) = \frac{\langle W_u(u(x,t,u_0),\eta_\gamma)}{\zeta(x,t)}.$$

From the equivariance of $u(\cdot, t, u_0)$ and $W_u(\gamma u) = \gamma W_u(u)$ it follows that

(34)
$$\zeta(x,t) = -\zeta(\gamma x,t), \text{ in } B_R \times (0,\infty),$$

 $c(x,t) = c(\gamma x,t), \text{ in } B_R \times (0,\infty).$ (35)

From the symmetry of W we also have that $u \in \pi_{\gamma}$ implies $W_u(u) \in \pi_{\gamma}$. From this we deduce

(36)
$$\langle W_u(u), \eta_\gamma \rangle = \langle u, \eta_\gamma \rangle \left\langle \int_0^1 W_{uu} \left(u + (s-1) \langle u, \eta_\gamma \rangle \eta_\gamma \right) \eta_\gamma \, \mathrm{d}s, \, \eta_\gamma \right\rangle$$

Thus, the coefficient c(x,t) of ζ in (33) is bounded (actually continuous) on $B_R \times (0,\infty)$. Since u_0 is a positive map, we have $\zeta_0 \geq 0$ for $\langle x, \eta_\gamma \rangle \geq 0$. Therefore, for establishing positivity it is sufficient to show that $\zeta(x,t) \geq 0$, for $x \in B_R^+ = \{x \in B_R \mid \langle x, \eta_\gamma \rangle > 0\}$ and $t \geq 0$. We note that by (34) there holds $\zeta(x,t) = 0$ for $x \in \pi_\gamma \times [0,\infty)$, hence if ζ is a classical solution of (33), we have that $\zeta(x,t)$ is nonnegative on $B_R^+ \times [0,\infty)$ by the maximum principle. For general $\zeta_0 \in W^{1,2}(B_R)$ we approximate via mollification as in [7, §4.2, Thm. 2] and note that positivity and symmetry are preserved by the approximation process, rendering $\zeta_0^{\varepsilon} \in C^{\infty}(B_R) \cap L^{\infty}(B_R)$, with $\zeta_0^{\varepsilon} \to \zeta_0$ in $W^{1,2}(B_R)$. By the classical maximum principle, there holds that $\zeta^{\varepsilon}(x,t) \geq 0$ on $B_R^+ \times [0,\infty)$, and by continuous dependence for (33) in $W^{1,2}(B_R)$ [12, Thm. 3.4.1], we have that $\zeta^{\varepsilon}(\cdot,t) \to \zeta(\cdot,t)$ a.e. in B_R along subsequences $\varepsilon_n \to 0$, hence $\zeta(x,t) \geq 0$ a.e. Finally, since $\zeta(x,t) = 0$ for $x \in \pi_\gamma \times (0,\infty)$ and since $\zeta(\cdot,t) \in C^{2+\alpha}(\overline{B_R})$ for t > 0, the Hopf boundary lemma applies on the smooth part of ∂B_R^+ and renders

$$\zeta(x,t) > 0$$
, in $B_R^+ \times (0,\infty)$,

unless $\zeta(x,t) \equiv 0$, hence unless $\zeta_0(x) \equiv 0$. But the hypothesis $u_0(\overline{F_R}) \cap F \neq \emptyset$ excludes this second option.

4. The minimization

Let $A^R := \{ u \in W^{1,2}(B_R, \mathbb{R}^n) \mid u(\overline{F}_R) \subset \overline{F} \}$ and consider the minimization problem

$$\min_{A^R} J_{B_R}, \text{ where } J_{B_R}(u) = \int_{B_R} \left\{ \frac{1}{2} |\nabla u|^2 + W(u) \right\} dx.$$

We will argue first that the minimizer exists. We redefine W(u) for $|u| \ge M+1$, so that the modified W is C^2 , satisfies $W(u) \ge c^2|u|^2$, for $|u| \ge M+1$ and a constant c, and also W(gu) = W(u), for all $g \in G$. We still denote the modified potential by W and the modified functional by J_{B_R} . We note that the convexity of \overline{F} implies that A^R is convex and closed in $W_{\rm E}^{1,2}(B_R;\mathbb{R}^n)$. The modified functional J_{B_R} satisfies all the properties required by the direct method and, as a result, a minimizer $v_R \in A^R$ exists.

Next we will show that as a consequence of Hypothesis 2 we can produce a minimizer $u_R \in A^R$, which in addition satisfies the estimate $|u_R(x)| \leq M$ (cf. (H3) in [3]). Due to this estimate, the values of W outside $\{|u| \leq M\}$ will not matter in the considerations in the rest of the paper and, therefore, the equation that will be solved is (1) with the original unmodified potential W. Set

(37)
$$u_R(x) = P v_R(x),$$

where Pv equals the projection on the sphere $\{v \in \mathbb{R}^n \mid |v| = M\}$, for points outside the sphere (Pv = Mv/|v|), and equals the identity inside the sphere. Since P is a contraction with respect to the Euclidean norm in \mathbb{R}^n , it follows that $u_R \in$ $W^{1,2}(B_R; \mathbb{R}^n)$, with $|\nabla u_R(x)| \leq |\nabla v_R(x)|$. Furthermore,

$$u_R(gx) = Pv_R(gx) = Pgv_R(x) = gPv_R(x) = gu_R(x),$$

hence $u_R \in W_E^{1,2}(B_R; \mathbb{R}^n)$. Clearly $u_R(\overline{F}) \subset \overline{F}$ and $|u_R(x)| \leq M$, for $x \in B_R$. The fact that u_R is also a minimizer is a consequence of Hypothesis 2 and the

following calculation.

$$\begin{split} J_{B_R}(u) &\geq \int_{B_R} \left\{ \frac{1}{2} |\nabla v_R|^2 + W(v_R) \right\} dx, \text{ for } u \in A^R, \\ &\geq \int_{B_R} \left\{ \frac{1}{2} |\nabla u_R|^2 + W(v_R) \right\} dx \\ &= \int_{|v_R(x)| \leq M} \left\{ \frac{1}{2} |\nabla u_R|^2 + W(u_R) \right\} dx \\ &+ \int_{|v_R(x)| > M} \left\{ \frac{1}{2} |\nabla u_R|^2 + W(v_R) \right\} dx \\ &\geq \int_{|v_R(x)| \leq M} \left\{ \frac{1}{2} |\nabla u_R|^2 + W(u_R) \right\} dx \\ &+ \int_{|v_R(x)| > M} \left\{ \frac{1}{2} |\nabla u_R|^2 + W(M \frac{v_R}{|v_R|}) \right\} dx \\ &= \int_{B_R} \left\{ \frac{1}{2} |\nabla u_R|^2 + W(u_R) \right\} dx, \end{split}$$

where the last inequality follows from Hypothesis 2.

We will be constructing the solution by taking the limit

(38)
$$u(x) = \lim_{R \to \infty} u_R(x).$$

For this purpose, we will need to show that the positivity constraint built in A^R does not affect the Euler–Lagrange equation, and we also need certain estimates, uniform in R, which in particular will imply that the solution is nontrivial.

Lemma 4.1. Let u_R be as above. Then, for R > 1, the following hold.

- (i) $J_{B_R}(u_R) \leq CR^{n-1}$, $||u_R||_{L^{\infty}(B_R;\mathbb{R}^n)} \leq M$, and $Q(u_R(x)) \leq \overline{Q}$, where $\overline{Q} := \max_{|u| \le M} Q(u),$

- (ii) $\Delta u_R W_u(u_R) = 0$, in $W_{\text{loc}}^{1,2}(B_R; \mathbb{R}^n)$, (iii) u_R is positive (cf. (23)), (iv) $\Delta Q(u_R(x)) \ge 0$, in $W_{\text{loc}}^{1,2}(D_R)$, where $D_R := D \cap B_R$ (cf. Hypothesis 4).

Proof. For (i), define

$$u_{\text{aff}}(x) := \begin{cases} d(x; \partial D)a_1, & \text{for } x \in D_R \text{ and } d(x; \partial D) \le 1, \\ a_1, & \text{for } x \in D_R \text{ and } d(x; \partial D) \ge 1, \end{cases}$$

and extend it equivariantly on B_R . Clearly, $u_{\text{aff}} \in A^R$. By the nonnegativity of W and a simple calculation,

(39)
$$0 \le J_{B_R}(u_R) \le \min_{A^R} J_{B_R}(u) < J_{B_R}(u_{\text{aff}}) < CR^{n-1},$$

for some constant C independent of R. The rest of (i) is already known.

For (ii), by Theorem 3.1, we have $u(\cdot, t; u_R) \in A^R$, for $t \ge 0$. Since u_R is a global minimizer of J_{B_R} in A^R , and since $u(\cdot, t; u_R) \in C^1(0, \infty; C^{2+\alpha}(\overline{B_R}))$, a classical

solution to (25) for t > 0, we conclude from

(40)
$$\frac{\mathrm{d}}{\mathrm{d}t}J_{B_R}(u(\cdot,t)) = -\int_{B_R} |u_t|^2 \,\mathrm{d}x$$

that $|u_t(x,t)| = 0$, for all $x \in B_R$ and t > 0. Hence, for t > 0, $u(\cdot,t)$ is satisfying

(41)
$$\Delta u(x,t) - W_u(u(x,t)) = 0$$

By taking $t \to 0+$ and utilizing the continuity of the flow in $W^{1,2}(B_R; \mathbb{R}^n)$ at t = 0, $u(\cdot, \cdot; u_R) \in C([0, \infty); W^{1,2}(B_R; \mathbb{R}^n))$, we obtain (ii).

Since (iii) is already known, we go on to (iv) where we obtain from (41), for t > 0,

$$0 = \langle Q_u(u(x,t)), \Delta u(x,t) \rangle - \langle Q_u(u(x,t)), W_u(u(x,t)) \rangle$$

= $\langle \hat{Q}_u(u(x,t) - a_1), \Delta(u(x,t) - a_1) \rangle - \langle Q_u(u(x,t)), W_u(u(x,t)) \rangle$

where $Q(u) = \hat{Q}(u - a_1)$, while using (12) we continue to obtain

(42)

$$0 \leq \Delta \hat{Q}(u(x,t) - a_1) - \langle Q_u(u(x,t)), W_u(u(x,t)) \rangle$$

$$= \Delta Q(u(x,t)) - \langle Q_u(u(x,t)), W_u(u(x,t)) \rangle$$

$$\leq \Delta Q(u(x,t)),$$

by Theorem 3.1, utilizing $u_R \in \mathcal{U}^{\text{Pos}}$, from which it follows that $u(\overline{D_R}, t) \subset D$, and by Hypothesis 4, particularly (10).

Thus, by the second remark following Lemma 2.1, we have

(43)
$$\Delta Q(u(x,t)) \ge 0, \text{ in } W^{1,2}_{\text{loc}}(D_R), \text{ for } t > 0$$

or, equivalently,

(44)
$$\int_{D_R} \nabla Q(u(x,t)) \nabla \phi(x) \, \mathrm{d}x \le 0, \text{ for all } \phi \ge 0, \phi \in W^{1,2}_{\mathrm{loc}}(D_R).$$

We will argue that

(45)
$$\nabla Q(u(\cdot, t)) \to \nabla Q(u_R(\cdot))$$
, weakly in $L^2(B_R)$, as $t \to 0$,

via which the proof of (iv) will be concluded. We know that

(46)
$$\begin{cases} u(\cdot,t;u_R) \to u_R, \text{ in } W^{1,2}(B_R;\mathbb{R}^n), \text{ as } t \to 0, \\ \|u(\cdot,t;u_R)\|_{L^{\infty}(B_R;\mathbb{R}^n)} \le M. \end{cases}$$

Hence,

 $Q(u(\cdot,t;u_R)) \to Q(u_R), \text{ in } L^2(B_R), \text{ as } t \to 0,$

since Q_u can be taken globally bounded. Thus,

 $\nabla Q(u(\cdot, t; u_R)) \to \nabla Q(u_R)$, in $\mathcal{D}'(B_R)$, as $t \to 0$.

However, $\|\nabla Q(u(\cdot, t; u_R))\|_{L^2(B_R)} < C$ by (46). Therefore (45) is established and the proof is complete.

The consideration in Lemma 4.1, particularly (40), together with the fact that u_R is a global minimizer, show that $u(\cdot, t; u_R)$ is an equilibrium of (25) for t > 0, that is, a time-independent solution satisfying in addition the boundary condition $\partial u/\partial \boldsymbol{n} = 0$. We can therefore replace u_R with this equilibrium which satisfies all the properties of Lemma 4.1 and also is in $C^{2+\alpha}(\overline{B_R};\mathbb{R}^n)$.

Corollary 4.2. We may assume that $u_R \in C^{2+\alpha}(\overline{B_R}; \mathbb{R}^n)$ is an equilibrium of (25) that satisfies all the properties of Lemma 4.1. Then,

$$u_R(\overline{F_R}) \cap \overline{F} \neq \emptyset \quad implies \quad u_R \in \mathcal{U}_0^{\mathrm{Pos}}$$

Proof. This follows from Theorem 3.1 and the fact that u_R is a time-independent solution of (25).

5. The comparison function σ ([1])

We prove three lemmas leading to the construction of a map σ that will play a major role in the derivation of the uniform estimates in R in the following section. We let χ_A be the characteristic function of a set A.

Given numbers $l, \lambda > 0$, set $L = l + \lambda$ and let $\varphi = \chi_{\overline{B_l}} \varphi_1 + \chi_{\overline{B_L} \setminus \overline{B_l}} \varphi_2$, where $\varphi_1 : \overline{B_l} \to \mathbb{R}, \varphi_2 : \overline{B_L} \setminus B_l \to \mathbb{R}$ are defined by

(47)
$$\begin{cases} \Delta \varphi_1 = c^2 \varphi_1, & \text{in } B_l, \\ \varphi_1 = \bar{q}, & \text{on } \partial B_l \end{cases}$$

and

(48)
$$\begin{cases} \Delta \varphi_2 = 0, & \text{in } B_L \setminus \overline{B_l} \\ \varphi_2 = \overline{q}, & \text{on } \partial B_l, \\ \varphi_2 = \overline{Q}, & \text{on } \partial B_L, \end{cases}$$

where c, \bar{q} , and M below are the constants defined in Hypotheses 1 and 2 and

(49)
$$\overline{Q} = \max_{|u| \le M} Q(u)$$

(see Hypothesis 2 and (i) of Lemma 4.1). The map φ is radial, that is, $\varphi_j(x) = \phi_j(|x|)$, for j = 1, 2. Classical properties of Bessel functions imply that $\phi_1 : [0, l] \to \mathbb{R}$ is positive and increasing together with the first derivative ϕ'_1 . The function $\phi_2 : [l, L] \to \mathbb{R}$ is increasing with decreasing first derivative ϕ'_2 , by explicit calculation.

Lemma 5.1. The following hold.

(i) The function $\phi'_1(l)$ is strictly increasing for $l \in (0, +\infty)$ and

(50)
$$\lim_{l \to +\infty} \phi_1'(l) = c\bar{q}$$

(ii) There exists a strictly increasing function $h: (0, +\infty) \to (0, +\infty)$ such that

(51)
$$\phi_1(r) \le e^{h(l)(r-l)}\phi_1(l), \text{ for } r \in [0, l],$$

and $\lim_{l\to+\infty} h(l) = c$.

(iii) There is a constant C_0 , independent of l, such that

(52)
$$\phi_1''(r) \le C_0, \text{ for } r \in [0, l].$$

Proof. Statements (i) and (ii) are proved in [8, Lemma 2.4]. For (iii) note that

(53)
$$\phi_1'' = c^2 \phi_1 - \frac{n-1}{r} \phi_1' \le c^2 \psi_1 \le c^2 \bar{q},$$

since ϕ_1 is increasing and bounded by \bar{q} .

A NEW PROOF FOR THE EXISTENCE OF AN EQUIVARIANT ENTIRE SOLUTION 11

An explicit computation yields, for $r \in [l, L]$,

(54)
$$\phi_2'(r) = \begin{cases} \frac{\overline{Q} - \overline{q}}{r \log(L/l)}, & \text{for } n = 2, \\ (n-2)\frac{l^{n-2}(\overline{Q} - \overline{q})}{r^{n-1}(1 - (l/L)^{n-2})}, & \text{for } n > 2. \end{cases}$$

Lemma 5.2. The following hold.

(i) Let the ratio l/L be fixed. Then,

(55)
$$\lim_{l \to +\infty} \phi_2'(l) = 0.$$

(ii) Let the difference L − l = λ be fixed. Then, φ'₂(l) is a decreasing function of l ∈ (0, +∞) and

(56)
$$\lim_{l \to +\infty} \phi_2'(r) = \frac{\overline{Q} - \overline{q}}{\lambda}, \text{ for } r \in [l, l + \lambda].$$

Moreover, there exists a constant C_0 , independent of $l \in [1, +\infty)$, such that

(57)
$$|\phi_2''(r)| \le \frac{C_0}{l}, \text{ for } r \in [l, l+\lambda].$$

Proof. Statement (i) is a straightforward consequence of (54). We prove (ii) for n > 2. The case n = 2 is similar. To show that $\phi'_2(l)$ is decreasing, we prove that the map $f(l) = l(1 - (l/(l + \lambda))^{n-2})$ is increasing. Setting $\xi = l/(l + \lambda)$ we have

$$f'(l) = d(\xi) := 1 - (n-1)\xi^{n-2} + (n-2)\xi^{n-1}$$
, for $\xi \in [0,1)$,

and f'(l) > 0, for $l \in (0, +\infty)$, follows from d(0) = 1, d(1) = 0, and $d'(\xi) < 0$, for $\xi \in (0, 1)$. The limit (56) follows from (54). The last statement of the lemma follows from

$$\phi_2''(r) = -(n-1)\frac{l^{n-1}}{r^n}\phi_2'(l).$$

Let φ be as before and let $\delta > 0$ be a small number. Denote by $\vartheta : B_{l+\delta} \setminus \overline{B_{l-\delta}} \to \mathbb{R}$ the solution of the problem

(58)
$$\begin{cases} \Delta \vartheta = 0, & \text{in } B_{l+\delta} \setminus \overline{B_{l-\delta}}, \\ \vartheta = \varphi, & \text{on } \partial (B_{l+\delta} \setminus \overline{B_{l-\delta}}). \end{cases}$$

We have $\vartheta(x) = \theta(|x|)$, where $\theta : [l - \delta, l + \delta] \to \mathbb{R}$ satisfies

(59)
$$\theta'(r) = \begin{cases} \frac{\phi_2(l+\delta) - \phi_1(l-\delta)}{r\log\frac{l-\delta}{l-\delta}}, & \text{for } n = 2, \\ (n-2)\frac{(l-\delta)^{n-2}(\phi_2(l+\delta) - \phi_1(l-\delta))}{r^{n-1}(1 - (\frac{l-\delta}{l+\delta})^{n-2})}, & \text{for } n > 2. \end{cases}$$

Lemma 5.3. There exist positive constants l_0 , λ , δ , $\bar{q}' < \bar{q}$, δ' , μ , such that $l \ge l_0$, $L = l + \lambda$ implies

(i) $\phi'_1(l) > \phi'_2(l) + \mu$, (ii) $\vartheta < \varphi$, in $B_{l+\delta} \setminus \overline{B_{l-\delta}}$,

NICHOLAS D. ALIKAKOS

(iii) The map $\sigma : \overline{B_L} \to \mathbb{R}$ defined by $\sigma = \chi_{B_{l-\delta} \cup (\overline{B_L} \setminus \overline{B_{l+\delta}})} \varphi + \chi_{\overline{B_{l+\delta}} \setminus B_{l-\delta}} \vartheta$ satisfies

(60)
$$\sigma \leq \bar{q}' < \bar{q}, \ in \ \overline{B_{l+\delta'}}.$$

Proof. Letting the ratio $\rho = l/L$ be fixed, then (50) and (55) imply that there is an l_0 such that (i) holds for $l = l_0$ and some $\mu > 0$. Fixing $\lambda = l_0((l/\rho) - 1)$, then (i) holds for all $l \ge l_0$. This follows from Lemmas 5.1 and (ii) of Lemma 5.2, which imply that $\phi'_1(l)$ is increasing and $\phi'_2(l)$ is decreasing for fixed λ . From (59), the relation

$$\phi_2(l+\delta) - \phi_1(l-\delta) = (\phi'_2(l) + \phi'_1(l))\delta + o(\delta),$$

which holds uniformly in l since $\phi_1(l) = \phi_2(l) = \bar{q}$, and

$$\log \frac{l+\delta}{l-\delta} = 2\frac{\delta}{l} + o(\delta), \qquad \left(\frac{l-\delta}{l+\delta}\right)^{n-2} = 1 - 2(n-2)\frac{\delta}{l} + o(\delta),$$

it follows that

(61)
$$\left| \theta'(r) - \frac{1}{2} (\phi'_2(l) + \phi'_1(l)) \right| \le C\delta, \text{ for } r \in [l - \delta, l + \delta],$$

(62)
$$|\theta''| \le \frac{C}{l}, \text{ for } r \in [l - \delta, l + \delta]$$

for some constant C > 0, independent of $l \in [l_0, +\infty)$. From (i) and (61), and the bounds on $\phi_1'', \phi_2'', \theta''$, it follows that there is a small $\delta > 0$, independent of $l \in [l_0, +\infty)$, such that

$$\begin{cases} \theta'(r) < \phi'_1(r), & \text{for } r \in [l-\delta, l], \\ \theta'(r) > \phi'_2(r), & \text{for } r \in [l, l+\delta]. \end{cases}$$

This and $\theta(l-\delta) = \phi_1(l-\delta)$, $\theta(l+\delta) = \phi_2(l+\delta)$, prove (ii). The existence of the number $\bar{q}' < \bar{q}$ and $0 < \delta' < \delta$, independent of $l \in [l_0, +\infty)$, follows by the same arguments and from the existence of the limits (50) and (56).

6. Uniform estimates in R

In this section we will make use of special notation. We denote by $B_R(x_R)$ the ball of radius R > 0 centered at x_R . As before, B_R denotes the ball of radius R > 0 centered at the origin and $D_{4R} = D \cap B_{4R}$, with x_R a point in D_{4R} such that $B_R(x_R) \subset D_{4R}$. The function u_{4R} is the minimizer for the functional $J_{B_{4R}}$ in Corollary 4.2.

Set

(63)
$$v_R(x) := \frac{Q(u_{4R}(x)) - \bar{q}/2}{\overline{Q} - \bar{q}/2}, \text{ for } x \in B_R(x_R),$$

where \bar{q} as in Hypothesis 1, \overline{Q} as in Lemma 4.1, Hypothesis 2, with $\overline{Q} > \bar{q}/2$. We will also rescale the dependent variable via $y = (x - x_R)/R$ and define

(64)
$$\hat{v}_R(y) := v_R(Ry + x_R) = v_R(x), \text{ for } y \in \hat{B}_1$$

where $\hat{B}_1 := \{y \in \mathbb{R}^n \mid |y| < 1\}, B_1 := \{x \in \mathbb{R}^n \mid |x| < 1\}, \text{ and } B_R^+(x_R) := \{x \in B_R(x_R) \mid v_R(x) \ge 0\}, B_R^-(x_R) := \{x \in B_R(x_R) \mid v_R(x) \le 0\}, \text{ and analogously,} \hat{B}_1^+ := \{y \in \hat{B}_1 \mid \hat{v}_R(y) \ge 0\}, B_1^- := \{y \in \hat{B}_1 \mid \hat{v}_R(y) \le 0\}.$ Notice that \hat{B}_1^+, \hat{B}_1^- depend on R.

12

By definition

(65)
$$Q(u_{4R}(x)) \ge \frac{\bar{q}}{2}, \text{ on } B_R^+(x_R)$$

By positivity ((iii) of Lemma 4.1) and equivariance, there holds $u_{4R}(B_R(x_R)) \subset u_{4R}(D_{4R}) \subset \overline{D}$. Hence,

(66)
$$W(u_{4R}(x)) \ge \varepsilon_0(\bar{q}) > 0, \text{ on } B_R^+(x_R),$$

since a_1 is the unique zero of W in \overline{D} (Hypotheses 3, 4).

Lemma 6.1. The following estimate holds for the Lebesgue measure of \hat{B}_1^- .

(67)
$$|\hat{B}_1^-| \ge |\hat{B}_1| - \frac{C}{\varepsilon_0(\bar{q})R},$$

where C is a constant depending only on the constant C in (i) of Lemma 4.1 and the dimension n.

Proof. We have

$$CR^{n-1} \ge \int_{B_{4R}} W(u_{4R}(x)) \,\mathrm{d}x \quad (by (i) \text{ of Lemma 4.1})$$
$$\ge \int_{B_R^+(x_R)} W(u_{4R}(x)) \,\mathrm{d}x \quad (by \ W \ge 0)$$
$$\ge \varepsilon_0(\bar{q}) \ |B_R^+(x_R)| \quad (by \ (66)).$$

Therefore,

$$\frac{C}{R} \ge \varepsilon_0(\bar{q}) \frac{|B_R^+(x_R)|}{R^n} = \varepsilon_0(\bar{q}) |\hat{B}_1^+|,$$

hence

$$|\hat{B}_1^-| = |\hat{B}_1| - |\hat{B}_1^+| \ge |\hat{B}_1| - \frac{C}{\varepsilon_0(\bar{q})R}.$$

Remark. The lemma above is a direct consequence of the basic integral estimate in (i) Lemma 4.1 and the assumption that in D the potential has a unique zero. Estimate (67) states that the minimizer $u_{4R}(x)$ on a set of large measure in $B_R(x_R)$ is close to a_1 , the zero of W, for $R \to \infty$.

The point in the next lemma is that the subharmonicity of $Q(u_{4R}(x))$ in D (by (iv) of Lemma 4.1) via a classical result of De Giorgi (see Appendix) allows us to obtain a pointwise estimate in the ball $B_{R/2}(x_R)$ of half the radius.

Lemma 6.2. Fix a $\delta \in (0,1)$. Then, for R large enough such that

(68)
$$1 - \frac{1}{\varepsilon_0(\bar{q})} \frac{C}{R} \frac{1}{c_0} \ge 1 - \delta,$$

we have the estimate $% \left({{{\left({{{\left({{{\left({{{\left({{{\left({{{c}}}} \right)}} \right.}$

(69)
$$\sup_{B_{R/2}(x_R)} Q(u_{4R}(x)) \le \frac{\bar{q}}{2} + \mu(1-\delta) \left(\overline{Q} - \frac{\bar{q}}{2}\right),$$

where $\mu(\cdot)$ is defined in the Appendix, with $\mu(1-\delta) < 1$.

Here C is the constant in Lemma 6.1 and c_0 is the volume of the unit ball in \mathbb{R}^n .

Proof. Note that $\Delta_y \hat{v}_R \ge 0$ in \hat{B}_1 and $\hat{v}_R \le 1$, in \hat{B}_1 , by (iv) and (i) of Lemma 4.1 respectively, and moreover

$$\frac{|\hat{B}_{1}^{-}|}{|\hat{B}_{1}|} \ge 1 - \frac{1}{\varepsilon_{0}(\bar{q})} \frac{C}{R} \frac{1}{c_{0}} \ge 1 - \delta$$

by (67). Hence, by the lemma in the Appendix,

$$\sup_{\hat{B}_{1/2}} \hat{v}_R(y) \le \mu(1-\delta) < 1,$$

which is equivalent to (69).

Next we will iterate. The number δ is fixed in Lemma 6.2 and we select k as the minimal integer with the property

(70)
$$\frac{\bar{q}}{2} + (\mu(1-\delta))^k \left(\overline{Q} - \frac{\bar{q}}{2}\right) < \bar{q}.$$

Clearly k depends only on δ . Finally we choose $R_0 = R_0(\delta)$ such that

(71)
$$1 - \frac{1}{\varepsilon_0(\bar{q})} \frac{C}{R} \frac{1}{|\hat{B}_{1/2^k}|} \ge 1 - \delta, \text{ for } R \ge R_0,$$

with C as in Lemma 6.2.

From now on, R, in the definition of A^R and in the definition of the minimizer u_R , is assumed to satisfy (71), and free otherwise. For such an R we define

$$\begin{cases} \overline{Q}_0 := \overline{Q}, \\ \overline{Q}_i := \frac{\overline{q}}{2} + \mu(a^*) \left(\overline{Q}_{i-1} - \frac{\overline{q}}{2} \right), \\ v_i(x) := \frac{Q(u_{4R}(x)) - \overline{q}/2}{\overline{Q}_{i-1} - \overline{q}/2}, \text{ for } x \in B_{R/2^i}(x_R), \\ \hat{v}_i(y) := v_i(Ry), \text{ for } y \in \hat{B}_{1/2^i}, \end{cases}$$

for i = 1, 2, ..., k and $a^* = 1 - \delta$.

We notice that (71) implies all the corresponding inequalities for i = 1, 2, ..., kand, particular, (68).

Lemma 6.3. For an integer $k = k(\delta)$, as in (70), and for $R \ge R_0(\delta)$, as in (71), the following estimate holds.

(72)
$$\sup_{B_{R/2^k}(x_R)} Q(u_{4R}(x)) \le \frac{\bar{q}}{2} + (\mu(a^*))^k \left(\overline{Q} - \frac{\bar{q}}{2}\right) < \bar{q}.$$

Proof. We make the simple observation that

(73)
$$\sup_{B_{R/2^{i}}(x_{R})} Q(u_{4R}(x)) \leq \frac{\bar{q}}{2} + (\mu(a^{*}))^{i} \left(\overline{Q} - \frac{\bar{q}}{2}\right)$$

holds for i = 1, 2, ..., k.

We note that for i = 1 this is just (69). Let us establish (73) for i = 2. We may assume that $k \ge 3$ since otherwise we have the estimate we need, hence

 $\overline{Q}_1 > \overline{q}/2$. We have $Q(u_{4R}(x)) \leq \overline{Q}_1$, on $B_{R/2}(x_R)$, by (69), and $Q(u_{4R}(x)) \geq \overline{q}/2$, on $B_{R/2}^+(x_R)$, by definition. Hence,

$$CR^{n-1} \ge \int_{B_{R}^{+}(x_{R})} W(u_{4R}(x)) \, \mathrm{d}x$$

$$\ge \int_{B_{R/2}^{+}(x_{R})} W(u_{4R}(x)) \, \mathrm{d}x \quad (\text{cf. Proof of Lemma 6.1})$$

$$\ge \varepsilon_{0}(\bar{q}) |B_{R/2}^{+}(x_{R})| \quad (\text{by (66)}).$$

Therefore,

$$\frac{C}{R} \ge \varepsilon_0(\bar{q}) \frac{|B_{R/2}^+(x_R)|}{R^n} = \varepsilon_0(\bar{q}) |\hat{B}_{1/2}^+|,$$

hence,

$$|\hat{B}_{1/2}^{-}| = |\hat{B}_{1/2}| - |\hat{B}_{1/2}^{+}| \ge |\hat{B}_{1/2}| - \frac{C}{\varepsilon_0(\bar{q})R}.$$

It follows that

(74)
$$\frac{|\bar{B}_{1/2}|}{|\hat{B}_{1/2}|} \ge 1 - \frac{C}{\varepsilon_0(\bar{q})R|\hat{B}_{1/2}|} \ge 1 - \delta,$$

by (71).

On the other hand, $\Delta_y \hat{v}_2(y) \ge 0$ in $\hat{B}_{1/2}$ and $\hat{v}_2 \le 1$, in $\hat{B}_{1/2}$, hence, by the lemma in the Appendix,

$$\sup_{\hat{B}_{1/2^2}} \hat{v}_2(y) \le \mu(a^*)$$

which equivalently gives

$$\sup_{B_{R/2^2}(x_R)} Q(u_{4R}(x)) \le \frac{\bar{q}}{2} + \mu(a^*) \left(\overline{Q}_1 - \frac{\bar{q}}{2}\right)$$

or

$$\sup_{B_{R/2^2}(x_R)} Q(u_{4R}(x)) \le \frac{\bar{q}}{2} + \mu(a^*)^2 \left(\overline{Q} - \frac{\bar{q}}{2}\right).$$

By repeating this process for $i = 3, \ldots, k$, we obtain (72).

So far we have established that

(75)
$$\sup_{B_{R^*}(x_R)} Q(u_{4R}(x)) \le \bar{q},$$

where $R^* = R/2^k$, for $R \ge R_0$, and an integer k independent of R. Utilizing the comparison function σ in Section 5 it is possible to show that the ball $B_{R^*}(x_R)$ in the supremum in (75) can be replaced by a large set D_R^* which includes all of D_{4R} with the exception of a strip along the boundary ∂D of width d_0 independent of R, for $R \ge R_0$, that is,

(76)
$$D_R^* \supset \{x \in D_{4R} \mid d(x, \partial D_{4R}) \ge d_0\},$$

for some $d_0 > 0$, which depends on l_0 in Lemma 5.3.

Lemma 6.4. The following estimate holds.

(77)
$$\sup_{D_R^*} Q(u_{4R}) \le \bar{q},$$

where D_R^* has the properties stated above.

Proof. First we note that by Hypotheses 1, 4,

$$\langle Q_u(u), W_u(u) \rangle \ge c^2 Q(u), \text{ for } |u - a_1| \le \bar{q},$$

which implies, via Lemma 2.1, (ii) of Lemma 4.1, and (75), the estimate

(78)
$$\Delta Q(u_{4R}) \ge \langle \Delta u_{4R}, Q_u(u_{4R}) \rangle = \langle W_u(u_{4R}), Q_u(u_{4R}) \rangle \ge c^2 Q(u_{4R}),$$

in $W^{1,2}(B_{-n}(x_{-n}))$

in $W_{loc}^{1,2}(B_{R^*}(x_R))$. Next we refer to Section 5. Consider a ball $B_l(\xi)$, tangent to $\partial B_{R^*}(x_R)$ and with its center ξ inside $B_{R^*}(x_R)$, and also consider the concentric ball $B_L(\xi)$. Notice that $B_l(\xi)$ is the translation of B_l and $B_L(\xi)$ the translation B_L . Similarly consider

the translations of $\varphi_1, \varphi_2, \vartheta$, which we still denote by the same symbols.

We now observe by (47), (75), and (78), that

$$\begin{cases} \Delta \varphi_1 = c^2 \varphi_1, & \text{in } B_l(\xi), \\ \varphi_1 = \bar{q}, & \text{on } \partial B_l(\xi), \end{cases} \quad \begin{cases} \Delta Q(u_{4R}) \ge c^2 Q(u_{4R}), & \text{in } B_l(\xi), \\ Q(u_{4R}) \le \bar{q}, & \text{on } \partial B_l(\xi), \end{cases}$$

hence, by the maximum principle for $W^{1,2}$ solutions (see [9]), we have

(79)
$$Q(u_{4R}) \le \varphi_1, \text{ in } B_l(\xi)$$

Also, by (48), (i) and (iv) of Lemma 4.1, and (75),

$$\begin{cases} \Delta \varphi_2 = 0, & \text{in } B_L(\xi) \setminus \overline{B_l(\xi)}, \\ \varphi_2 = \overline{q}, & \text{on } \partial B_l(\xi), \\ \varphi_2 = \overline{Q}, & \text{on } \partial B_L(\xi), \end{cases} \qquad \begin{cases} \Delta Q(u_{4R}) \ge 0, & \text{in } B_L(\xi), \\ Q(u_{4R}) \le \varphi_2, & \text{on } \partial (B_L(\xi) \setminus \overline{B_l(\xi)}), \end{cases}$$

hence,

(80)
$$Q(u_{4R}) \leq \varphi_2, \text{ in } B_L(\xi) \setminus \overline{B_l(\xi)}.$$

We deduce therefore by Lemma 5.3 that

(81)
$$\begin{cases} Q(u_{4R}) \leq \varphi, & \text{in } B_L(\xi), \\ Q(u_{4R}) \leq \vartheta, & \text{in } B_{l+\delta}(\xi) \setminus \overline{B_{l-\delta}(\xi)}, \\ Q(u_{4R}) \leq \sigma \leq \overline{q}' < \overline{q}, & \text{in } \overline{B_{l+\delta'}(\xi)} \end{cases}$$

Thus, we see from (iii) of (81) that the estimate (75) holds on a set larger than $B_{R^*}(x_R)$. Clearly, by repeating this process we obtain (77).

We are now able to finish the proof of the theorem.

Proof of Theorem 1.1. First note that if
$$q: D_R^* \to \mathbb{R}$$
 is the solution to

(82)
$$\Delta q = c^2 q, \text{ in } D_R^*, q = \overline{q}', \text{ on } D_R^*,$$

then,

(83)
$$q(x) \le K e^{-kd(x,\partial D_R^*)}, \text{ for } x \in D_R^*,$$

for positive constants K, k, independent of R.

Indeed, by the maximum principle, there holds $q \leq \bar{q}'$. It follows that if φ is the solution of equation (82) on the ball with center x and radius $d(x, \partial D_R^*)$ and with boundary condition $\varphi = \bar{q}'$, we have $q \leq \varphi$. This and the estimate (51) in (ii) of Lemma 5.1 imply (83). Moreover, we note that if $d_0 > 0$ is as in (76), then

(84)
$$q(x) \le K e^{-kd(x,\partial D_{4R})}, \text{ in } B_{d_0}(x) \subset D_{4R},$$

since $d(x, \partial D_{4R}) \leq d(x, \partial D_R^*) + d_0$. This last inequality follows from (76) with a new constant K.

Now utilizing (75) and $\Delta Q(u_{4R}) \geq c^2 Q(u_{4R})$, in $W_{\text{loc}}^{1,2}(D_R^*)$, by (78), we obtain by comparing with (82) that $Q(u_{4R}(x)) \leq q(x)$, for $x \in D_R^*$, and, therefore, by (84) and (9c),

(85)
$$|u_{4R}(x) - a_1| \le K e^{-kd(x, \partial D_{4R})}$$

The uniform bound in (i) of Lemma 4.1 and elliptic regularity, via a diagonal argument, allow us to pass to the limit along a subsequence in R and capture a function

$$u(x) = \lim_{R_n \to \infty} u_{R_n}(x).$$

The uniform bound (85) implies that the limit function satisfies the exponential estimate in the theorem and is also a solution to

$$\Delta u - W_u(u) = 0, \text{ in } \mathbb{R}^n,$$

by (ii) of Lemma 4.1 (and the comment after its proof). Clearly, also $u \in \mathcal{U}^{\text{Pos}}$.

Finally we argue the strong positivity for u(x) with respect to D. Take now an open connected set $U \subset D$ which contains some points far enough from ∂D so that by the exponential estimate there holds $u(U) \cap D \neq \emptyset$. Define $\Gamma' := \{\gamma \in \Gamma \mid \pi_{\gamma} \cap D \neq \emptyset\}$; then, $D = \bigcap_{\gamma \in \Gamma \setminus \Gamma'} \mathcal{P}_{\gamma}^+$ (see Lemmas 2.1 and 5.1 in [1]). As in the second part of the proof of Theorem 3.1, particularly (33), set $z(x) = \langle u(x), \eta_{\gamma} \rangle$, for $x \in U$ and $\gamma \in \Gamma \setminus \Gamma'$. Then,

$$\begin{cases} \Delta z + cz = 0, & \text{in } U, \\ z \ge 0, & \text{in } U. \quad (\text{by positivity}) \end{cases}$$

By a well-known variant of the strong maximum principle, there holds z > 0 in U, unless $z \equiv 0$. Triviality is excluded by $u(U) \cap D \neq \emptyset$ above. From this, strong positivity follows.

The proof is complete.

Appendix

We state a special case of the *De Giorgi oscillation lemma* which was originally established for general elliptic operators $\mathcal{L}u := \operatorname{div}(A(x)\nabla u)$, with bounded, measurable coefficients (see [4, p. 195]).

Lemma (De Giorgi). Consider the ball $\hat{B}_1 = \{y \in \mathbb{R}^n \mid |y| \le 1\}$ and a function $\hat{v} = \hat{v}(y)$ which satisfies

- (i) $\Delta_y \hat{v} \ge 0$, in $W_{\text{loc}}^{1,2}(\hat{B}_1)$,
- (ii) $\hat{v} \leq 1$, in \hat{B}_1 ,

(iii) $|\hat{B}_1^-|/|\hat{B}_1| \ge a^* > 0$, for $\hat{B}_1^- = \{y \in \hat{B}_1 \mid \hat{v}(y) \le 0\}$,

where $|\Omega|$ is the Lebesgue measure of the set Ω . Then,

$$\sup_{\hat{B}_{1/2}} \hat{v} \le \mu(a^*) < 1$$

Notice that the statement of the lemma is invariant under the scaling $y \to \lambda y$, for $\lambda > 0$, hence \hat{B}_1 can be replaced by \hat{B}_{λ} and $\hat{B}_{1/2}$ by $\hat{B}_{\lambda/2}$, without affecting $\mu(a^*)$.

NICHOLAS D. ALIKAKOS

ACKNOWLEDGMENTS

Thanks are due to Giorgio Fusco for discussions on the contents of the paper, to Panagiotis Smyrnelis for pointing out two errors in a former version of the manuscript, and to Apostolos Damialis for his suggestions for improving the presentation.

References

- [1] N. D. Alikakos and G. Fusco. Entire solutions to equivariant elliptic systems with variational structure. To appear in *Arch. Rat. Mech. Anal.*
- [2] S. Baldo. Minimal interface criterion for phase transitions in mixtures of Cahn-Hilliard fluids. Ann. Inst. Henri Poincaré, Anal. Non Linéaire 7 No. 2 (1990), pp. 67–90.
- [3] L. Bronsard, C. Gui, and M. Schatzman. A three-layered minimizer in R² for a variational problem with a symmetric three-well potential. *Comm. Pure. Appl. Math.* **49** No. 7 (1996), pp. 677–715.
- [4] L. Caffarelli and S. Salsa. A geometric approach to free boundary problems. Graduate Studies in Mathematics 68, American Mathematical Society, Providence, RI, 2005.
- [5] U. Dierkes, S. Hildenbrandt, A. Küster, and O. Wohlrab. *Minimal surfaces I. Boundary value problems*. Grundlehren der mathematischen Wissenschaften **295**, Springer-Verlag, Berlin, 1992.
- [6] U. Dierkes, S. Hildenbrandt, A. Küster, and O. Wohlrab. *Minimal surfaces II. Boundary regularity*. Grundlehren der mathematischen Wissenschaften **296**, Springer-Verlag, Berlin, 1992.
- [7] L. C. Evans and R. F. Gariepy. Measure theory and fine properties of functions. CRC Press, Boca Raton, FL, 1992.
- [8] G. Fusco, F. Leonetti, and C. Pignotti. A uniform estimate for positive solutions of semilinear elliptic equations. Trans. Amer. Math. Soc. 363 No. 8 (2011), pp. 4285–4307.
- D. Gilbarg and N. S. Trudinger. Elliptic partial differential equations of second order. Grundlehren der mathematischen Wissenschaften 224, Springer-Verlag, Berlin, revised second edition, 1998.
- [10] L. C. Grove and C. T. Benson. *Finite reflection groups*. Graduate Texts in Mathematics 99, Springer-Verlag, Berlin, second edition, 1985.
- [11] C. Gui and M. Schatzman. Symmetric quadruple phase transitions. Ind. Univ. Math. J. 57 No. 2 (2008), pp. 781–836.
- [12] D. Henry. Geometric theory of semilinear parabolic equations. Lecture Notes in Mathematics 840, Springer-Verlag, Berlin-Heidelberg-New York, 1981.
- [13] P. D. Hislop and I. M. Sigal. Introduction to spectral theory with applications to Schrödinger operators. Applied Mathematical Sciences 113, Springer-Verlag, New York, 1996.
- [14] J. E. Humphreys Reflection groups and Coxeter groups. Cambridge Studies in Advanced Mathematics 29, Cambridge University Press, Cambridge, 1992.
- [15] D. Kinderlehrer and G. Stampacchia. An introduction to variational inequalities and their applications. Pure and Applied Mathematics 88, Academic Press, New York, 1980.
- [16] J. Smoller. Shock waves and reaction-diffusion equations. Grundlehren der Mathematischen Wissenschaften 258, Springer-Verlag, Berlin, second edition, 1994.
- [17] J. E. Taylor. The structure of singularities in soap-bubble-like and soap-film-like minimal surfaces. Ann. Math. 103 (1976), pp. 489–539.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ATHENS, PANEPISTEMIOPOLIS, 15784 ATHENS, GREECE AND INSTITUTE FOR APPLIED AND COMPUTATIONAL MATHEMATICS, FOUNDATION OF RE-SEARCH AND TECHNOLOGY – HELLAS, 71110 HERAKLION, CRETE, GREECE

E-mail address: nalikako@math.uoa.gr