

Calculating the renormalisation group equations of a SUSY model with **Susyno**

Renato M. Fonseca

Centro de Física Teórica de Partículas, Instituto Superior Técnico,
Universidade Técnica de Lisboa, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal

E-mail address: renato.fonseca@ist.utl.pt

Abstract

Susyno is a Mathematica package dedicated to the computation of the 2-loop renormalisation group equations of a supersymmetric model based on any gauge group (the only exception being multiple $U(1)$ groups) and for any field content.

1 Introduction

Supersymmetry (SUSY) is an elegant and well known extension of the Standard Model (SM) which aims at solving or softening some of the SM theoretical and experimental shortcomings. Supersymmetric models based on Grand Unified Theories (GUTs) offer the appealing feature of being described by a unique gauge group and coupling constant; furthermore, the potential new superfields, as well as their dynamics, can provide answers to fundamental problems. For example $SO(10)$ based models offer a natural explanation for neutrino masses and mixings via a type-I seesaw. This occurs since the spinor representation of the group contains a SM singlet superfield (with the appropriate quantum numbers). Even larger and/or more complex groups are often considered, for example E_6 in string inspired models.

The analysis of the theoretical and phenomenological implications of SUSY GUT models requires a careful study of the evolution of the fundamental parameters from the high-energy scale down to the electroweak scale, at which observables are computed and constraints applied. As such, knowledge of the renormalisation group equations (RGEs) is necessary. Although the RGEs of several models (e.g., MSSM, NMSSM) are already known [1, 3], for other SUSY extensions of the SM complicated general equations must be used [1, 2].

Here we describe **Susyno**, a Mathematica-based package that addresses this issue. The program takes as input the gauge group, the representations (i.e., the chiral superfield content), the number of flavours/copies¹ of each representation, and any abelian discrete symmetries (e.g., R-parity). **Susyno** then constructs the most general superpotential and soft-SUSY-breaking Lagrangian consistent with the field content and symmetries imposed. Once these elements have been derived, **Susyno** then computes the 2-loop β -functions of all the parameters of the model, which is its main output.

¹Throughout this document, the word *flavours* is meant to convey the sense of *copies* or *repetitions* of a given representation/field.

There is another Mathematica package, **SARAH** [4], which works in a similar way for models based on $SU(n)$ gauge factor groups. However, **Susyno** is prepared to accept as working input any gauge group that does not contain more than one $U(1)$ factor².

This document is organised as follows. Section 2 explains how to install the program and run a first simple example (the MSSM). Sections 3 and 4 explain how to prepare the input and how to read and interpret the output, also using as example the MSSM case. Finally, Section 5 summarises the tests conducted to validate the code.

2 Installation and quick start

Susyno works on Windows, Linux and Mac OS provided that Mathematica 7 (or a latter version) is installed. The program is obtainable from

<http://web.ist.utl.pt/renato.fonseca/susyno.html>

The files *LieGroups.m*, *SusyRGEs.m* and *Susyno.m* are the core of the program. These and other auxiliary files can be found inside the folder *Susyno*, which must be extracted from the downloaded *Susyno-1.1.tar* file to a location that is visible to Mathematica. Typing `$Path` in Mathematica will show a complete list of acceptable locations. A good choice is to place the whole folder (not just its contents) in

(Mathematica base directory)/AddOns/Applications

(note that in a Windows system the slashes “/” must be replaced by backslashes “\”). The package can be loaded by typing

```
<< Susyno`
```

in Mathematica’s front end. A text message is returned, informing that a built-in help system provides a detailed description of the program and its functions (see A). A tutorial is also included.

The **Susyno** lines below allow a simple and easy first run: the example consists of a possible way of writing the MSSM input.

```
normalisation = Sqrt[3/5];
Q = {U1->1/6 normalisation, SU2->{1}, SU3->{1, 0}, NFlavours->3, DiscreteSym->-1};
u = {U1->-2/3 normalisation, SU2->{0}, SU3->{0, 1}, NFlavours->3, DiscreteSym->-1};
d = {U1->1/3 normalisation, SU2->{0}, SU3->{0, 1}, NFlavours->3, DiscreteSym->-1};
L = {U1->-1/2 normalisation, SU2->{1}, SU3->{0, 0}, NFlavours->3, DiscreteSym->-1};
e = {U1->normalisation, SU2->{0}, SU3->{0, 0}, NFlavours->3, DiscreteSym->-1};
Hu = {U1->1/2 normalisation, SU2->{1}, SU3->{0, 0}, NFlavours->1, DiscreteSym->1};
Hd = {U1->-1/2 normalisation, SU2->{1}, SU3->{0, 0}, NFlavours->1, DiscreteSym->1};

model = {Q, u, d, L, e, Hu, Hd};
BetaFunctions2L[model];
```

²Multiple $U(1)$ groups lead to “ $U(1)$ mixing” [5] which requires special care [6].

Evaluation of this simple code generates the 2-loop β -functions of the model (MSSM in this case). Notice that no external input or output files are used - everything happens on Mathematica's front end.

3 Defining a model

A SUSY model contains two building blocks: a superpotential and a soft-SUSY-breaking Lagrangian. For a simple gauge group, a general superpotential can be written as

$$W = \frac{1}{6}Y^{ijk}\Phi_i\Phi_j\Phi_k + \frac{1}{2}\mu^{ij}\Phi_i\Phi_j + L^i\Phi_i, \quad (1)$$

with Φ denoting a chiral superfield, and where Y , μ and L are dimensionless, mass, and mass² parameters, respectively. A generic soft-SUSY-breaking Lagrangian reads

$$- \mathcal{L}_{soft} = \left(\frac{1}{6}h^{ijk}\phi_i\phi_j\phi_k + \frac{1}{2}b^{ij}\phi_i\phi_j + s^i\phi_i + \text{h.c.} \right) + (m^2)_j^i\phi_i\phi_j^* + \left(\frac{1}{2}M\lambda\lambda + \text{h.c.} \right) \quad (2)$$

where λ is a gaugino field and ϕ denotes scalar fields. M (m^2) correspond to gaugino (scalar) soft breaking masses, while h , b and s denote trilinear, bilinear and linear soft breaking parameters. `Susyno` requires as input the list of chiral superfields (just *fields* from now on) of a model:

```
model={field1, field2,...};
```

For each field, the user only needs to provide its defining elements. The syntax is

```
field={g1->g1_rep, g2->g2_rep, ..., NFlavours->nf, DiscreteSym->q};
```

where the values to be filled are the following:

- **g1, g2, ...** (gauge factor groups);
- **g1_rep, g2_rep, ...** (representations of the field under each of the gauge factor groups);
- **nf** (number of flavours);
- **q** (charge under some discrete abelian symmetry).

To obtain the β -functions, the user must call `BetaFunctions1L` (or `BetaFunctions2L` for 2-loop results):

```
BetaFunctions1L[model]
BetaFunctions2L[model]
```

or

```
BetaFunctions1L[model,Verbose->False];
BetaFunctions2L[model,Verbose->False];
```

if we do not wish to have the results printed on Mathematica's front end.

With this information the program then builds internally the superpotential and the soft-SUSY-breaking Lagrangian using an algorithm to automatically name the parameters of the model (see Section 4). Assuming that the model contains parameters named $X1, X2, \dots$, the program's output is

$$\begin{aligned} \{\{X1, \beta_{X1}^{(1)}\}, \{X2, \beta_{X2}^{(1)}\}, \dots\} & \quad (\text{BetaFunctions1L}) \\ \{\{X1, \beta_{X1}^{(1)}, \beta_{X1}^{(2)}\}, \{X2, \beta_{X2}^{(1)}, \beta_{X2}^{(2)}\}, \dots\} & \quad (\text{BetaFunctions2L}) \end{aligned}$$

that is, a set of sublists each containing a parameter Xi , and the one-loop (one- and two-loop) β -functions for its RG evolution: $\beta_{Xi}^{(1)}$ ($\beta_{Xi}^{(1)}, \beta_{Xi}^{(2)}$).

In the following, we will look at each of the input elements mentioned above in greater detail.

3.1 Gauge factor groups

The program needs to know the model's gauge group³, which it extracts from each `field` object mentioned above. Therefore the gauge factor groups should be presented in the same order in all fields. In addition, $U(1)$'s must always come first. For the MSSM (for completeness, the superpotential and soft-SUSY-breaking Lagrangian of the MSSM are presented in B) we could write each field as

```
field = {U1->U1_charge, SU2->SU2_rep, SU3->SU3_rep, ...};
```

We may change the order of $SU(2)$ and $SU(3)$ but the $U(1)$ factor must always come first. Any group can be used: all the possible gauge factor groups, as well as their corresponding `Susyno` input are collected in Table 1.

Gauge factor group	Susyno input
$U(1)$	U1
$SU(n)$	SU2, SU3, SU4, SU5, ...
$SO(n)$	S03, S05, S06, S07, ...
$Sp(2n)$	SP2, SP4, SP6, ...
G_2	G2
F_4	F4
E_6, E_7, E_8	E6, E7, E8

Table 1: Gauge factor groups

Keep in mind that despite its generality, `Susyno` only assigns to variables a finite number of groups (for instance variable `S0100` is not set). Should this ever become a problem, there is an easy method of circumventing it, discussed in C.

3.2 Representations

As mentioned before, `Susyno` is designed to accept an arbitrary field content so any representation of any gauge group is allowed. For $U(1)$ factor groups, all we need are the hypercharges of each field, which are just numbers⁴. On the other hand, the representations of simple gauge groups

³We emphasise here that we are actually dealing with algebras, not groups. Nevertheless, we will adopt the common practice in high-energy physics of using the word *group* for both these concepts.

⁴Notice that, while it is possible to have more than one hypercharge (i.e. more than one $U(1)$ factor group), the formalism of `Susyno` is based on [1, 2], which do not include the effects of a possible $U(1)$ mixing.

must be specified by their Dynkin coefficients (see C for details). In Table 2 we list some of the representations of SU(2), SU(3), SU(5) and SO(10) (Dynkin coefficients and corresponding dimensions).

Group	Representation	
	<i>Dynkin coefficients</i>	<i>Dimension (Name)</i>
SU(2)	{0}	1 (Trivial/Singlet)
	{1}	2 (Fundamental/Doublet)
	{2}	3 (Adjoint/Triplet)
SU(3)	{0,0}	1 (Trivial/Singlet)
	{1,0}	3 (Fundamental)
	{0,1}	$\bar{\mathbf{3}}$ (Anti-fundamental)
SU(5)	{1,1}	8 (Adjoint)
	{0,0,0,0}	1 (Trivial/Singlet)
	{1,0,0,0}	5 (Fundamental)
	{0,0,0,1}	$\bar{\mathbf{5}}$ (Anti-fundamental)
	{0,1,0,0}	10
	{2,0,0,0}	15
	{0,0,0,2}	$\bar{\mathbf{15}}$
SO(10)	{1,0,0,1}	24 (Adjoint)
	{0,0,0,0,0}	1 (Trivial/Singlet)
	{1,0,0,0,0}	10 (Fundamental)
	{0,0,0,0,1}	16 (Spinor)
	{0,0,0,1,0}	$\bar{\mathbf{16}}$ (Spinor's conjugate)
	{0,1,0,0,0}	45 (Adjoint)
	{2,0,0,0,0}	54
	{0,0,1,0,0}	120
	{0,0,0,0,2}	126
	{0,0,0,2,0}	$\bar{\mathbf{126}}$

Table 2: List of some of the representations of SU(2), SU(3), SU(5) and SO(10)

To understand how the MSSM was specified in the example of Section 2, we just need the following information from Table 2:

- the Dynkin coefficients of the trivial and fundamental representations of SU(2): {0} and {1};
- the Dynkin coefficients of the trivial, fundamental (**3**) and anti-fundamental ($\bar{\mathbf{3}}$) representations of SU(3): {0,0}, {1,0}, {0,1}.

Hence

```

Q = {U1->1/6 normalisation, SU2->{1}, SU3->{1, 0}, ...};
u = {U1->-2/3 normalisation, SU2->{0}, SU3->{0, 1}, ...};
d = {U1->1/3 normalisation, SU2->{0}, SU3->{0, 1}, ...};
L = {U1->-1/2 normalisation, SU2->{1}, SU3->{0, 0}, ...};
e = {U1->normalisation, SU2->{0}, SU3->{0, 0}, ...};
Hu = {U1->1/2 normalisation, SU2->{1}, SU3->{0, 0}, ...};
Hd = {U1->-1/2 normalisation, SU2->{1}, SU3->{0, 0}, ...};

```

Notice that **Susyno** accepts any normalisation of the hypercharges. In the sample code of Section 2, we used the usual $\sqrt{\frac{3}{5}}$ factor (from an embedding of the MSSM in an SU(5) based model).

3.3 Number of flavours and abelian discrete symmetries

Many models contain repetitions of some of the representations of the gauge group. Instead of considering them distinct fields, we usually view these as different flavours of a single field and **Susyno** needs to know how many flavours there are in order to compute the β -functions. The user should use the syntax `NFlavours->...` for this purpose. For example, in the MSSM we have

```
Q = {..., NFlavours->3, ...};
u = {..., NFlavours->3, ...};
d = {..., NFlavours->3, ...};
L = {..., NFlavours->3, ...};
e = {..., NFlavours->3, ...};
Hu = {..., NFlavours->1, ...};
Hd = {..., NFlavours->1, ...};
```

Let us consider another example: if we were to modify the MSSM to include m copies of H_u and H_d , we would write

```
Hu = {..., NFlavours->m, ...};
Hd = {..., NFlavours->m, ...};
```

Finally, there are models in which we must forbid some couplings and in order to do that some abelian discrete symmetry is introduced. These symmetries are also a defining element of these models and as such each field's charge must be passed on to the program using the expression `DiscreteSym->....`. The MSSM is one such case; there is a Z_2 symmetry known as R-parity so we should write

```
Q = {..., DiscreteSym->-1};
u = {..., DiscreteSym->-1};
d = {..., DiscreteSym->-1};
L = {..., DiscreteSym->-1};
e = {..., DiscreteSym->-1};
Hu = {..., DiscreteSym->1};
Hd = {..., DiscreteSym->1};
```

For the most general (R-parity violating) MSSM the first 5 lines must be replaced by

```
Q = {..., DiscreteSym->1};
u = {..., DiscreteSym->1};
d = {..., DiscreteSym->1};
L = {..., DiscreteSym->1};
e = {..., DiscreteSym->1};
```

Note that the expressions `NFlavours->...` and `DiscreteSym->...` can be placed anywhere along the definition of each field. They can even be omitted: if no `NFlavours` is written, the program will use the symbolic value `nf[i]` where i is the order by which a field is given; if there is no `DiscreteSym` then `Susyno` will consider that the field transforms trivially (charge equal to 1).

4 The output of Susyno

Once all the input is provided, `Susyno` automatically builds the Lagrangian of a model, asking neither for names of fields nor names of parameters. On the one hand, inputting a model becomes very easy, since it is not even necessary to know the exact number of its parameters. On the other hand, this renders the output hard to read (and hence not particularly user-friendly), since the names of the parameters are chosen by the program.

Still, provided we know the conventions used by the program, the notation used by `Susyno` can be manually changed by the user (after the code is executed). We notice that in the present version of `Susyno` there are no custom built-in functions to export the results. Users who wish to do so must do it manually or with the help of Mathematica's built-in functions `CForm` and `FortranForm`.

4.1 Naming of parameters

`Susyno` assigns names to the parameters of a model in such a way that the user can identify which fields they are multiplying:

```
y[field1, field2, field3, InvIndex, <flav1>, <flav2>, <flav3>]
mu[field1, field2, InvIndex, <flav1>, <flav2>]
l[field1, InvIndex, <flav1>]
h[field1, field2, field3, InvIndex, <flav1>, <flav2>, <flav3>]
b[field1, field2, InvIndex, <flav1>, <flav2>]
s[field1, InvIndex, <flav1>]
m2[field1, field2, InvIndex, <flav1>, <flav2>]
```

A few comments concerning the above (output) tensors are in order:

- `y`, `mu`, `l`, `h`, `b`, `s` and `m2` can be easily identified with the different types of couplings and dimensionful parameters of the superpotential and the soft-SUSY-breaking Lagrangian;
- `field1`, `field2`, `field3` are the indices of the fields entering a given coupling, or soft breaking term. For example, with `model={Q, u, d, L, e, Hu, Hd}` we have $Q=1$, $u=2$, $d=3$, $L=4$, $e=5$, $Hu=6$, $Hd=7$. The up-quark Yukawa couplings would then be `y[1,2,6,...]`;
- There is the possibility that the product of 3 representations, $R_1 \otimes R_2 \otimes R_3$, contains more than one invariant. Therefore we need `InvIndex=1,2,...` to distinguish them. This is rare though, so in most cases (e.g., the MSSM) `InvIndex=1` for all parameters. Notice that in linear and bilinear terms, R_1 and $R_1 \otimes R_2$, this problem does not arise since there is at most one invariant. However, for consistency `Susyno` still uses `InvIndex` in this case, although setting its value to 1;
- `<flav1>`, `<flav2>`, `<flav3>` are the flavour indices of `field1`, `field2`, `field3`. If any of these fields has only one flavour, the corresponding index is omitted. Consider again the example of the up-quark Yukawa couplings: we would have `y[1,2,6,1,i,j,k]` where $i =$

flavour of Q , j = flavour of u , k = flavour of H_u . Yet H_u only has one flavour so the correct parameter name is `y[1,2,6,1,i,j]`.

Additionally, there are also the coupling constants and the gaugino masses:

`g[1], g[2], ...`
`M[1], M[2], ...`

A list containing the matching of MSSM's standard notation parameters and **Susyno** assignments can be found in `B`.

4.2 The full Lagrangian as used by the program

Knowing the full Lagrangian can be important. As an example, consider a possible normalisation issue: how can we be sure that **Susyno** is correctly taking $\mu H_u \cdot H_d$ as `mu[6,7,1] H_u · H_d`, and not `-mu[6,7,1] H_u · H_d`, `2mu[6,7,1] H_u · H_d` or any other multiple? The way to solve such problems related to the normalisation of the parameters is to check the Lagrangian. This can be done using the **ShowLagrangian** function:

ShowLagrangian[model]

This function returns the list of tensors $\{Y, \mu, L, h, b, s, m^2\}$. Note that **Susyno** never expands the flavour indices of these tensors so the first, second and third indices are assumed to carry symbolic flavours i (or none), j (or none), k (or none)⁵. The argument (`model`) is the same as in the **BetaFunctions1L** and **BetaFunctions2L** functions. There is also a **Verbose** option (set to **True** by default) that prints the different parts of W and \mathcal{L}_{soft} :

- $\frac{1}{6} Y^{ijk} \Phi_i \Phi_j \Phi_k$ (Y part), $\frac{1}{2} \mu^{ij} \Phi_i \Phi_j$ (μ part), $L^i \Phi_i$ (L part);
- $\frac{1}{6} h^{ijk} \phi_i \phi_j \phi_k$ (H part), $\frac{1}{2} b^{ij} \phi_i \phi_j$ (B part), $s^i \phi_i$ (S part) and $(m^2)_j^i \phi_i \phi_j^*$ (M2 part).

ShowLagrangian is completely independent of **BetaFunctions1L** and **BetaFunctions2L**, so it can be called at any time to generate the superpotential and soft-SUSY-breaking Lagrangian of a given model.

Finally, notice that in order to build \mathcal{L} it is necessary to assume a particular basis for the representations/fields of the model. Quantities such as the β -functions clearly do not depend on this choice. However, it is important to know that **Susyno** can use unconventional basis for the representations. The following is a good example. According to the program, two fields, $A = \{a[1], a[2], a[3]\}$ and $B = \{b[1], b[2], b[3]\}$, transforming as **3** and $\bar{\mathbf{3}}$ of $SU(3)$ will form the invariant $a[1]b[3] - a[2]b[2] + a[3]b[1]$, not $a[1]b[1] + a[2]b[2] + a[3]b[3]$. Both are valid expressions as long as they are used in a consistent way - it is just a matter of basis choice. Thus, when analysing the output of **ShowLagrangian** we must distinguish between irrelevant basis choices and relevant normalisation conventions for the parameters. The β -functions are only sensitive to the latter ones.

⁵These flavour indices do not show up if the particular entry of the tensors we are considering is unflavoured. For example the entry of the Y tensor related to the fields Q , d , H_d will depend on i (flavour of the first field, Q) and j (flavour of the second field, d) but not on k since the third field (H_d) is unflavoured. Notice also that exchanging for example H_d with Q corresponds to having a different entry of the Y tensor: this is because the flavour of Q (now the third field) is assumed to be symbolically k instead of i .

5 Tests/validation of Susyno

The output of `Susyno` was confronted with the analysis of some models available in the literature. In particular, the RGEs generated by `Susyno` were compared with the results for the MSSM [1], the R-parity violating MSSM [7], the general NMSSM [3] and some SU(5)-based models [8].

In general the program's RGEs are consistent with the results collected in the above publications, but in some cases differences were found. These have been collected in the program's webpage.

Acknowledgments

The author would like to thank Jorge Romão and Ana M. Teixeira for the many suggestions and encouragement given. This work was supported by the *Fundação para a Ciência e a Tecnologia* under the grant SFRH/BD/47795/2008.

A List of available functions

As in most programs, the code `Susyno` is spread over many internal functions. Due to their nature, some of these functions may be useful on their own, and they were thus built in a user-friendly way, and are documented.

Below is the list of functions that can be called directly by the user in Mathematica's front-end, followed by a brief description. The built-in help system describes in detail how to use them.

- `Adjoint` - Computes the Dynkin coefficients of the adjoint representation of a group;
- `BetaFunctions1L` - Computes the 1-loop β -functions of a SUSY model;
- `BetaFunctions2L` - Computes the 2-loop β -functions of a SUSY model;
- `CanonicalForm` - Simplifies an expression written in Einstein's notation;
- `Casimir` - Computes the quadratic Casimir of a representation;
- `CartanMatrix` - Computes the Cartan matrix of any simple group;
- `DimR` - Computes the dimension of a representation;
- `Invariants` - Computes (in some basis) the invariant combination(s) of a product of one, two or three representations;
- `ListContract` - Efficiently calculates traces of multi-index sparse tensors;
- `PositiveRoots` - Computes the positive roots of a group;
- `ReduceRepProduct` - Decomposes a direct product representation in its irreducible parts [9];
- `RepMatrices` - Computes (in some basis) the explicit representation matrices;
- `ShowLagrangian` - Generates the Lagrangian (W and \mathcal{L}_{soft}) of a SUSY model;
- `Weights` - Computes the weights of a representation, including degeneracy.

Parameter	Susyno's notation
g_1, g_2, g_3	$g[1], g[2], g[3]$
M_1, M_2, M_3	$M[1], M[2], M[3]$
$(Y_u)_{ij}$	$y[1,2,6,1,j,i]$
$(Y_d)_{ij}$	$y[1,3,7,1,j,i]$
$(Y_e)_{ij}$	$y[4,5,7,1,j,i]$
μ	$mu[6,7,1]$
$(h_u)_{ij}$	$h[1,2,6,1,j,i]$
$(h_d)_{ij}$	$h[1,3,7,1,j,i]$
$(h_e)_{ij}$	$h[4,5,7,1,j,i]$
B	$b[6,7,1]$
$(m_Q^2)_{ij}$	$m2[1,1,1,j,i]$
$(m_u^2)_{ij}$	$m2[2,2,1,i,j]$
$(m_d^2)_{ij}$	$m2[3,3,1,i,j]$
$(m_L^2)_{ij}$	$m2[4,4,1,j,i]$
$(m_e^2)_{ij}$	$m2[5,5,1,i,j]$
$m_{H_u}^2$	$m2[6,6,1]$
$m_{H_d}^2$	$m2[7,7,1]$

Table 1: Parameters of the MSSM assuming the field ordering $\{Q,u,d,L,e,H_u,H_d\}$ and the group ordering $\{U1,SU2,SU3\}$.

B Parameters of the MSSM

For completeness, we include here the superpotential and soft-SUSY-breaking Lagrangian of the MSSM, following the conventions of [1].

$$\begin{aligned}
W &= \hat{u}Y_u\hat{Q} \cdot \hat{H}_u + \hat{d}Y_d\hat{Q} \cdot \hat{H}_d + \hat{e}Y_e\hat{L} \cdot \hat{H}_d + \mu\hat{H}_u \cdot \hat{H}_d \\
-\mathcal{L}_{soft} &= \left(\tilde{u}h_u\tilde{Q} \cdot H_u + \tilde{d}h_d\tilde{Q} \cdot H_d + \tilde{e}h_e\tilde{L} \cdot H_d + BH_u \cdot H_d + \text{h.c.} \right) \\
&\quad + \tilde{Q}^\dagger m_Q^2 \tilde{Q} + \tilde{L}^\dagger m_L^2 \tilde{L} + \tilde{u}m_u^2 \tilde{u}^\dagger + \tilde{d}m_d^2 \tilde{d}^\dagger + \tilde{e}m_e^2 \tilde{e}^\dagger + m_{H_u}^2 H_u^\dagger H_u + m_{H_d}^2 H_d^\dagger H_d \\
&\quad + \left(\frac{1}{2} M_a \lambda_a \lambda_a + \text{h.c.} \right)
\end{aligned} \tag{3}$$

In the above, and as usual, hats denote superfields and tildes the scalar component of each chiral superfield. The contraction of two SU(2) doublets (denoted by \cdot) follows the convention $H_u \cdot H_d = H_u^+ H_d^- - H_u^0 H_d^0$. In addition to the parameters entering W and \mathcal{L}_{soft} , one still has the gauge couplings g_a . Table 1 summarises the matching of the physical notation of Eqs. (3,4) with the one used by Susyno.

C More on groups and representations

A simple complex Lie algebra is specified by a Cartan matrix [10] which is a square matrix with diagonal entries all equal to 2 and off-diagonal entries equal to 0, -1, -2 or -3. For example, the

Cartan matrix of $SO(10)$ is

$$\begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 2 \end{pmatrix} \quad (5)$$

The Cartan matrix is the ingredient required by **Susyno** whenever a group is being asked as input (the exception is $U(1)$, which is taken to be just **U1** or **{}**). Although the user can manually provide the Cartan matrices, the program already has a built-in function, **CartanMatrix**, that can compute them for any simple group. The syntax is

```
CartanMatrix["group name",familyIndex]
```

where "group name" can be "SO", "SU", "SP", "G", "F" or "E" and familyIndex should be an integer. For example

```
CartanMatrix["SO",10]
```

will return the above 5×5 Cartan matrix. This function will work for virtually any simple group. However, this is still a heavy syntax, so the program assigns to some variables (**SU2**, **SU3**, **SU5**, **S010**, etc.) the corresponding Cartan matrices (e.g., **S010** is the same as **CartanMatrix**["SO",10]). For the case of $SO(n)$, $SU(n)$ with $n > 32$ and $Sp(2m)$ with $m > 16$ no variables were set, and the **CartanMatrix** function should be used.

As mentioned in the text, **Susyno** requires that each field be specified as a list of hypercharges and representations under the simple gauge factor groups. We will focus now on this last element: representations of simple groups.

A representation of a (simple) group is often labeled by its dimension (e.g., **3** of $SU(3)$, **54** of $SO(10)$, ...) but this can be ambiguous. To avoid the problem, we should use Dynkin coefficients which are a list of n non-negative integers. This number n is known as the group rank and is equal to the size of the (square) Cartan matrix of the group. For instance, since the Cartan matrix of $SO(10)$ is 5×5 , all representations of the group are of the form

```
{i1,i2,i3,i4,i5}
```

(i1,..., i5 are non-negative integers).

There are functions in **Susyno** that compute properties of the representations (e.g., **DimR** calculates the dimension of a representation, **ReduceRepProduct** reduces products of representations) and they are documented in the built-in help files (see also A). These should be enough to allow recognising a representation by its Dynkin coefficients; however, should the user wish to consult lists of representations, these are available in the literature [11]. Some of the most frequently used representations were shown in table 2.

D Renormalisation group equations of a general model

Susyno uses the generic two loop RGEs of [1, 2]. For completeness these results are reproduced here (for a model based on a simple gauge group G).

In general a parameter X evolves according to the equation

$$\frac{d}{dt}X = \frac{1}{16\pi^2}\beta_X^{(1)} + \frac{1}{(16\pi^2)^2}\beta_X^{(2)} \quad (6)$$

where $t = \log Q$ (Q being the renormalisation scale parameter). The β -functions for the several parameters (gauge coupling constants, superpotential parameters and soft-SUSY-breaking parameters) are as follows.

Gauge coupling constants

$$\beta_g^{(1)} = g^3 [S(R) - 3C(G)] \quad (7)$$

$$\beta_g^{(2)} = g^5 \left\{ -6[C(G)]^2 + 2C(G)S(R) + 4S(R)C(R) \right\} - g^3 \frac{C(k)}{d(G)} Y^{ijk} Y_{ijk} \quad (8)$$

Superpotential parameters

$$[\beta_Y^{(1)}]^{ijk} = Y^{ijp} \gamma_p^{(1)k} + (k \leftrightarrow i) + (k \leftrightarrow j) \quad (9)$$

$$[\beta_Y^{(2)}]^{ijk} = Y^{ijp} \gamma_p^{(2)k} + (k \leftrightarrow i) + (k \leftrightarrow j) \quad (10)$$

$$[\beta_\mu^{(1)}]^{ij} = \mu^{ip} \gamma_p^{(1)j} + (j \leftrightarrow i) \quad (11)$$

$$[\beta_\mu^{(2)}]^{ij} = \mu^{ip} \gamma_p^{(2)j} + (j \leftrightarrow i) \quad (12)$$

$$[\beta_L^{(1)}]^i = L^p \gamma_p^{(1)i} \quad (13)$$

$$[\beta_L^{(2)}]^i = L^p \gamma_p^{(2)i} \quad (14)$$

where

$$\gamma_i^{(1)j} = \frac{1}{2} Y_{ipq} Y^{jpq} - 2g^2 \delta_i^j C(i) \quad (15)$$

$$\begin{aligned} \gamma_i^{(2)j} = & -\frac{1}{2} Y_{imn} Y^{npq} Y_{pqr} Y^{mrj} + g^2 Y_{ipq} Y^{jpq} [2C(p) - C(i)] \\ & + 2\delta_i^j g^4 [C(i)S(R) + 2C(i)^2 - 3C(G)C(i)] \end{aligned} \quad (16)$$

Soft-SUSY-breaking Lagrangian parameters

$$\beta_M^{(1)} = g^2 [2S(R) - 6C(G)] M \quad (17)$$

$$\begin{aligned} \beta_M^{(2)} = & g^4 \left[-24C(G)^2 + 8C(G)S(R) + 16S(R)C(R) \right] M + 2g^2 \frac{C(k)}{d(G)} \left(h^{ijk} - M Y^{ijk} \right) Y_{ijk} \\ & \quad (18) \end{aligned}$$

$$\left[\beta_h^{(1)}\right]^{ijk} = \frac{1}{2}h^{ijl}Y_{lmn}Y^{mnk} + Y^{ijl}Y_{lmn}h^{mnk} - 2\left(h^{ijk} - 2MY^{ijk}\right)g^2C(k) + (k \leftrightarrow i) + (k \leftrightarrow j) \quad (19)$$

$$\begin{aligned} \left[\beta_h^{(2)}\right]^{ijk} = & -\frac{1}{2}h^{ijl}Y_{lmn}Y^{npq}Y_{pqr}Y^{mrk} - Y^{ijl}Y_{lmn}Y^{npq}Y_{pqr}h^{mrk} - Y^{ijl}Y_{lmn}h^{npq}Y_{pqr}Y^{mrk} \\ & + \left(h^{ijl}Y_{lpq}Y^{pqk} + 2Y^{ijl}Y_{lpq}h^{pqk} - 2MY^{ijl}Y_{lpq}Y^{pqk}\right)g^2[2C(p) - C(k)] \\ & + \left(2h^{ijk} - 8MY^{ijk}\right)g^4[C(k)S(R) + 2C(k)^2 - 3C(G)C(k)] + (k \leftrightarrow i) + (k \leftrightarrow j) \end{aligned} \quad (20)$$

$$\left[\beta_b^{(1)}\right]^{ij} = \frac{1}{2}b^{il}Y_{lmn}Y^{mnj} + \frac{1}{2}Y^{ijl}Y_{lmn}b^{mn} + \mu^{il}Y_{lmn}h^{mnj} - 2(b^{ij} - 2M\mu^{ij})g^2C(i) + (i \leftrightarrow j) \quad (21)$$

$$\begin{aligned} \left[\beta_b^{(2)}\right]^{ij} = & -\frac{1}{2}b^{il}Y_{lmn}Y^{pqn}Y_{pqr}Y^{mrj} - \frac{1}{2}Y^{ijl}Y_{lmn}b^{mr}Y_{pqr}Y^{pqn} - \frac{1}{2}Y^{ijl}Y_{lmn}\mu^{mr}Y_{pqr}h^{pqn} \\ & - \mu^{il}Y_{lmn}h^{npq}Y_{pqr}Y^{mrj} - \mu^{il}Y_{lmn}Y^{npq}Y_{pqr}h^{mrj} + 2Y^{ijl}Y_{lpq}(b^{pq} - \mu^{pq}M)g^2C(p) \\ & + \left(b^{il}Y_{lpq}Y^{pqj} + 2\mu^{il}Y_{lpq}h^{pqj} - 2\mu^{il}Y_{lpq}Y^{pqj}M\right)g^2[2C(p) - C(i)] \\ & + (2b^{ij} - 8\mu^{ij}M)g^4[C(i)S(R) + 2C(i)^2 - 3C(G)C(i)] + (i \leftrightarrow j) \end{aligned} \quad (22)$$

$$\left[\beta_s^{(1)}\right]^i = \frac{1}{2}Y^{ipq}Y_{pqr}s^r + h^{ipq}Y_{pqr}L^r + \mu^{ir}Y_{rpq}b^{pq} + 2Y^{imn}(m^2)_m\mu_{nl} + h^{ipq}b_{pq} \quad (23)$$

$$\begin{aligned} \left[\beta_s^{(2)}\right]^i = & 2g^2C(q)Y^{ipq}Y_{pqr}s^r - \frac{1}{2}Y^{irn}Y_{npq}Y^{pqm}Y_{mrl}s^l - 4g^2C(q)(Y^{ipq}M - h^{ipq})Y_{pql}L^l \\ & - (Y^{imn}Y_{npq}h^{pqr}Y_{rml} + h^{imn}Y_{npq}Y^{pqr}Y_{rml})L^l - 4g^2C(q)\mu^{il}Y_{lpq}(\mu^{pq}M - b^{pq}) \\ & - \mu^{il}(Y_{lmn}h^{npq}Y_{pqr}\mu^{rm} + Y_{lmn}Y^{npq}Y_{pqr}b^{rm}) + 4g^2C(q)[2Y^{ipq}\mu_{pq}|M|^2 - Y^{ipq}b_{pq}M \\ & - h^{ipq}\mu_{pq}M^* + h^{ipq}b_{pq} + Y^{ipq}(m^2)_p\mu_{rq} + Y^{ipr}(m^2)_r\mu_{pq}] - Y^{imn}Y_{npq}h^{pql}b_{lm} \\ & - h^{imn}Y_{npq}Y^{pql}b_{lm} - Y^{imn}h_{npq}h^{pql}\mu_{lm} - h^{imn}h_{npq}Y^{pql}\mu_{lm} - Y^{imn}(m^2)_m\mu_{lr}Y^{rpq}Y_{pqn} \\ & - Y^{imn}Y_{npq}Y^{pql}(m^2)_l\mu_{rm} - Y^{imn}(m^2)_nY_{lpq}Y^{pqr}\mu_{rm} - 2Y^{imn}Y_{npq}(m^2)_rY^{rpl}\mu_{lm} \end{aligned} \quad (24)$$

$$\left[\beta_{m^2}^{(1)}\right]_i^j = \frac{1}{2}Y_{ipq}Y^{pqn}(m^2)_n^j + \frac{1}{2}Y^{jnpq}Y_{pqn}(m^2)_i^n + 2Y_{ipq}Y^{jpr}(m^2)_r^q + h_{ipq}h^{jpr} - 8\delta_i^j|M|^2g^2C(i) + 2g^2\mathbf{t}_i^{Aj}\text{Tr}(\mathbf{t}^A m^2) \quad (25)$$

$$\begin{aligned} \left[\beta_{m^2}^{(2)}\right]_i^j = & -\frac{1}{2}(m^2)_i^lY_{lmn}Y^{mrj}Y_{pqr}Y^{pqn} - \frac{1}{2}(m^2)_l^jY^{lmn}Y_{mri}Y^{pqr}Y_{pqn} - Y_{ilm}Y^{jnm}(m^2)_r^lY_{npq}Y^{rpq} \\ & - Y_{ilm}Y^{jnm}(m^2)_n^rY_{rpq}Y^{lpq} - Y_{ilm}Y^{jnr}(m^2)_n^lY_{pqr}Y^{pqm} - 2Y_{ilm}Y^{jln}Y_{npq}Y^{mpr}(m^2)_r^q \\ & - Y_{ilm}Y^{jln}h_{npq}h^{mpq} - h_{ilm}h^{jln}Y_{npq}Y^{mpq} - h_{ilm}Y^{jln}Y_{npq}h^{mpq} - Y_{ilm}h^{jln}h_{npq}Y^{mpq} \\ & + \left[(m^2)_i^lY_{lpq}Y^{jpr} + Y_{ipq}Y^{lpq}(m^2)_l^j + 4Y_{ipq}Y^{jpl}(m^2)_l^q + 2h_{ipq}h^{jpr} - 2h_{ipq}Y^{jpr}M \right. \\ & \left. - 2Y_{ipq}h^{jpr}M^* + 4Y_{ipq}Y^{jpr}|M|^2\right]g^2[C(p) + C(q) - C(i)] - 2g^2\mathbf{t}_i^{Aj}(\mathbf{t}^A m^2)_r^lY_{lpq}Y^{rpq} \\ & + 8g^4\mathbf{t}_i^{Aj}\text{Tr}[\mathbf{t}^A C(r)m^2] + \delta_i^jg^4|M|^2[24C(i)S(R) + 48C(i)^2 - 72C(G)C(i)] \\ & + 8\delta_i^jg^4C(i)\left\{\text{Tr}[S(r)m^2] - C(G)|M|^2\right\} \end{aligned} \quad (26)$$

A few comments and clarifications are still in order concerning the variables appearing in the different β -functions:

- $Y_{ijk} = (Y^{ijk})^*$, $h_{ijk} = (h^{ijk})^*$, $\mu_{ij} = (\mu^{ij})^*$ and $b_{ij} = (b^{ij})^*$;
- $d(G)$ = Dimension of the adjoint representation of group G ;
- $C(i)$ = Quadratic Casimir invariant of the representation of the chiral superfield with index i ;
- $C(G)$ = Quadratic Casimir invariant of the adjoint representation of group G ;
- $S(R)$ = Dynkin index summed over all chiral multiplets. However $S(R)C(R)$ should be interpreted as the sum of Dynkin indices weighted by the quadratic Casimir invariant;
- t^A = Representation matrices under the gauge group G . Terms with t^A are only relevant for U(1) groups;
- In $\beta_{m^2}^{(1)}$ and $\beta_{m^2}^{(2)}$, the traces should be understood as traces over all chiral superfields.

If the gauge group is not simple but rather a direct product of simple groups and U(1) factors, the equations shown here must be adapted. The modifications are fairly straightforward, and can be found in [1].

References

- [1] S. P. Martin and M. T. Vaughn, Phys. Rev. D **50** (1994) 2282 [Erratum-ibid. D **78** (2008) 039903] [arXiv:hep-ph/9311340].
- [2] Y. Yamada, Phys. Rev. D **50** (1994) 3537 [arXiv:hep-ph/9401241].
- [3] U. Ellwanger, C. Hugonie and A. M. Teixeira, Phys. Rept. **496** (2010) 1 [arXiv:0910.1785 [hep-ph]].
- [4] F. Staub, Comput. Phys. Commun. **182** (2008) 808 [arXiv:hep-ph/0806.0538].
- [5] F. del Aguila, G.D. Coughlan and M. Quirós, Nucl. Phys. B **307** (1988) 633;
M. x. Luo and Y. Xiao, Phys. Lett. B **555** (2003) 279 [arXiv:hep-ph/0212152];
I. Jack and D. R. T. Jones, Phys. Rev. D **63** (2001) 075010 [arXiv:hep-ph/0010301].
- [6] R. M. Fonseca, M. Malinsky, W. Porod and F. Staub, [arXiv:1107.2670 [hep-ph]].
- [7] B.C. Allanach, A. Dedes and H.K. Dreiner, Phys. Rev. D **60** (1999) 056002 [arXiv:hep-ph/9902251].
- [8] F. Borzumati and T. Yamashita, Prog. Theor. Phys. **124** (2009) 761 [arXiv:hep-ph/0903.2793].
- [9] D.M. Snow, ACM Trans. Math. Softw. **16** (1990) 94; ACM Trans. Math. Softw. **19** (1993) 95.
- [10] R.N. Cahn, *Semi-Simple Lie Algebras and Their Representations* (Dover Publications, 2006).
- [11] R. Slansky, Phys. Rept. **79** (1981) 1.