A new method for the determination of the growth rate from galaxy redshift surveys

Adi Nusser¹, Enzo Branchini², and Marc Davis³

¹Physics Department and the Asher Space Science Institute-Technion, Haifa 32000, Israel

²Department of Physics, Università Roma Tre, Via della Vasca Navale 84, 00146, Rome, Italy

INFN Sezione di Roma 3, Via della Vasca Navale 84, 00146, Rome, Italy

INAF, Osservatorio Astronomico di Brera, Milano, Italy

³ Departments of Astronomy & Physics, University of California, Berkeley, CA. 94720

ABSTRACT

Given a redshift survey of galaxies with measurements of apparent magnitudes, we present a novel method for measuring the growth rate $f(\Omega)$ of cosmological linear perturbations. We use the galaxy distribution within the survey to solve for the peculiar velocity field which depends in linear perturbation theory on $\beta = f(\Omega)/b$, where b is the bias factor of the galaxy distribution. The recovered line-of-sight peculiar velocities are subtracted from the redshifts to derive the distances, which thus allows an estimate of the absolute magnitude of each galaxy. A constraint on β is then found by minimizing the spread of the estimated magnitudes from their distribution function. We apply the method to the all sky K = 11.25 Two-MASS Redhsift Survey (2MRS) and derive $\beta = 0.35 \pm 0.1$ at $z \sim 0$, remarkably consistent with our previous estimate from the velocity-velocity comparison. The method could easily be applied to subvolumes extracted from the SDSS survey to derive the growth rate at $z \sim 0.1$. Further, it should also be applicable to ongoing and future spectroscopic redshift surveys to trace the evolution of $f(\Omega)$ to $z \sim 1$. Constraints obtained from this method are entirely independent from those obtained from the two-dimensional distortion of $\xi(s)$ and provide an important check on $f(\Omega)$, as alternative gravity models predict observable differences.

Subject headings: Cosmology: dark matter - cosmology: large scale structure of the universe

1. Introduction

Large scale density perturbations in the Universe are gravitationally unstable and grow via linear theory. The growing mode of large scale density perturbations, D(a), is characterized by the more observationally relevant growth rate

$$f(\Omega) = \frac{d \ln D}{d \ln a} , \qquad (1)$$

where a is the scale factor of the Universe and Ω is the matter density parameter. It is found that the growth index $\gamma = d \ln f / d \ln \Omega$ is very well approximated by $\gamma = 0.55 + 0.05[1 + w(z = 1)]$ (Linder 2005) for a cosmological background dominated by dark energy

with an equation of state, $P = w\rho c^2$. The growth rate is not only sensitive to the background cosmology, but also to the theory of gravitation invoked as the driver for structure formation. Geometric \mathbb{R}^n (e.g. Gannouji et al. 2009), and Dvali-Gabadadze-Porrati (DGP) (Dvali et al. 2000; Wei 2008) gravity models give substantially different behaviors of f. Models with dark sector long range forces, in addition to gravity, even introduce a scale dependence into f (e.g. Keselman et al. 2010).

Here we present a new method for constraining $f(\Omega)$ from redshift surveys of galaxies with measured apparent magnitudes. Redshifts of galaxies systematically differ from the actual distances by the line-of-sight components of their peculiar velocities. Hence, the directly *measurable* intrinsic luminosities or absolute magnitudes of galaxies inferred from the observed flux using redshifts rather than distances will show

¹E-mail: adi@physics.technion.ac.il

²E-mail: branchin@fis.uniroma3.it

³E-mail: mdavis@berkeley.edu

larger spread than the true values.

Gravitational instability theory allows a prediction of the peculiar velocity field from observed galaxy distribution given $f(\Omega)$ and the biasing relation between galaxies and mass. The method uses this predicted velocity field to get distances for deriving true absolute magnitudes. Constraints on f can then be derived by minimizing the scatter of the estimated absolute magnitudes from a reference distribution. Since the galaxy distribution could be biased relative to the mass density field, the constraints on f are degenerate with the assumed biasing. Adopting linear biasing, $\delta_g = b\delta_m$, between the galaxy number density contrast, δ_q , and the mass density contrast, δ_m , yields constraints on $\beta = f/b$ that are independent from those obtained from the apparent anisotropy in the observed galaxy clustering (Kaiser 1988).

In §2 we describe the method in detail, presenting general expressions and deriving the relevant approximations. In §3 we offer general analytic assessments of the method. In §4 we apply the method to the 2MRS of galaxies limited to magnitude K = 11.25 (Huchra et al. 2005). We conclude in §5 with a general discussion of the results and of the prospects for the application of the method to future data.

2. The method

This section is largely based on Nusser et al. (2011). We are given a flux limited survey of galaxies with observed apparent magnitudes $m < m_l$, angular positions, and redshifts cz (in km s⁻¹). Let r (also in km s⁻¹) be the luminosity distance to a galaxy in the sample. For simplicity of notation and description we assume here that the distance and spatial extent of the survey are small so that r is well approximate by the physical distance. Therefore, cz = r + v where $v = \hat{r} \cdot \mathbf{u}$ is the line-of-sight component of three dimensional peculiar velocity \mathbf{u} of the galaxy. The results can readily be extended to the general case once we specify the underlying cosmological model.

The method relies on a prediction of v from the observed distribution of galaxies in the survey. Since r = cz - v this prediction allows an estimate of the true absolute magnitude,

$$M = m - 15 - 5\log r = M_0 - 5\log(1 - v/cz) \quad (2)$$

where the measurable absolute magnitude $M_0 = m - 15 - 5 \log cz$ is determined from observations. Because the peculiar velocity of a galaxy is uncorrelated with its true absolute magnitude, constraints on the underlying velocity field can be derived by demanding that the distribution of the magnitudes, M, is consistent with a reference distribution function (i.e. lumi-

nosity function). The equations of gravitational instability theory relate the underlying mass density contrast, δ_m , to the peculiar velocity field **u**. Further, we will use the linearized equations in which the relation solely depends on $f(\Omega)$. For linear biasing, $\delta_q = b\delta_m$, between mass and galaxies the appearance of f is replaced by the single parameter $\beta = f/b$. Therefore, the method presented here will focus on constraints on β only. More sophisticated models for the velocity field involving additional parameters will not be discussed here. In principle, the constraint can be obtained without resorting to the luminosity function simply by minimizing the variance of $M_0 - 5\log(1 - v(\beta)/cz)$ with respect to β . However, much tighter constraints are obtained from the full distribution. We define the luminosity function, $\Phi(M)$, expressed in terms of the absolute magnitudes, as the number density of galaxies per unit magnitude. The probability, $P(M_0|cz, v)$, of observing a galaxy having an observed magnitude M_0 in a flux limited sample, depends on its redshift, cz, and its radial peculiar velocity, v, and is well approximated as (Nusser et al. 2011)

$$P(M_0|cz,v) = \frac{\Phi(M)}{\int_{-\infty}^{M_l} \Phi(M) dM} , \qquad (3)$$

where $M_l = M_{0l} - 5 \log(1 - v/cz)$ and $M_{0l} = m_l - 15 - 5 \log cz$. The expression is valid as long as the relative errors on the measured redshift are small ($\sigma_{cz}/cz \ll 1$) The probability distribution of the whole sample of galaxies is the product of the single probabilities

$$P_s = \prod_i P(M_{0i} | cz_i, v_i) , \qquad (4)$$

where *i* runs over all galaxies of the sample. Given a form for $\Phi(M)$, the parameter β is then constrained by maximizing P_s in which the dependence on β is via the $v(\beta)$ as inferred from the spatial distribution of galaxies. In principle one could use the "nonparamatric" fit methods (Efstathiou et al. 1988; Davis & Huchra 1982) to approximate $\Phi(M)$. However, here we will only apply the method to the 2MRS sample which is reasonably approximated by a Schechter luminosity function (Westover 2007). Therefore, we assume $\Phi(M)$ is well approximated by a Schechter form (Schechter 1980)

$$\Phi(M) = 0.4 \ln(10) \Phi^* 10^{0.4(\alpha+1)(M_*-M)} \\ \times \exp\left(-10^{0.4(M_*-M)}\right) .$$
 (5)

The normalization Φ_* does not concern us here. The shape parameters M_* and α generally depend on the galaxies' type, redshift and band of observation. In terms of the luminosity $(M = -2.5 \log L + const)$, this function acquires the simpler form

$$\Phi(M(L)) = 0.4\ln(10)\Phi^* \left(\frac{L}{L_*}\right)^{1+\alpha} \exp\left(-\frac{L}{L_*}\right) .$$
(6)

Inserting all this expression into (3) gives

$$P(M_0|cz;v) = \frac{0.4\ln(10)\left(\frac{L}{L_*}\right)^{1+\alpha} e^{-L/L_*}}{\Gamma(1+\alpha, L_l/L_*)}$$
(7)

where $L/L_* = (1 - v/cz)^2 10^{0.4(M_* - M_0)}$, $L_l/L_* = (1 - v/cz)^2 10^{0.4(M_* - M_0)}$.

To summarize, given the observed absolute magnitudes M_0 for a sample of objects, a Schechter model for the luminosity function and linear theory prediction for the underlying velocity field $v(\beta)$, we determine β , α and M_* by minimizing Eq. 4.

3. General assessments of the method

There are three sources of error which affect the derivation of β in the method presented here. a) "shotnoise" error resulting from the finite number galaxies, b) cosmic variance due to variations in the large scale structure in volumes of the universe comparable in size to the volume probed by the redshift survey at hand and c) inaccuracies in the peculiar velocity reconstruction. In this section we offer a general assessment of the applicability of the method to data at higher redshifts. To compensate for the degrading of the signal $(\sim v/cz)$ with redshift, surveys with large number of galaxies need to be invoked. We will only consider here shot-noise error. Peculiar velocity reconstruction errors and cosmic variance (of surveys with similar volumes) scale with redshift in a similar manner to the signal and hence their corresponding relative error will not depend on redshift.

To the list of errors above we do not add biases introduced by adopting a specific form for the luminosity function since, as we will argue in §3.2, these biases are expected to be insignificant. Further, the issue becomes completely irrelevant for future surveys where the number of galaxies is large enough to allow the use of "nonparametric fit" techniques for modeling the luminosity function, relaxing the need for specific parametric forms.

3.1. Sensitivity to "shot-noise" and redshift

Assume a cubic¹ region at high redshift cz containing a flux limited sample of N galaxies with redshifts

 $cz_i \approx cz$. From the galaxy distribution in this region, derive the peculiar velocity field as a function of β . Consider a Schechter form for the luminosity function and assume cz is large enough that the limiting luminosity L_l of a galaxy that could be observed is significantly larger than L_* . In the limit $L_l \gg L_*$, the Γ function in the expression for $P(M_0|cz;v)$ in (7) is well approximated as $\Gamma(1 + \alpha, L_l/L_*) = (L_l/L_*)^{\alpha} \exp(-L_l/L_*)$ so that,

$$P(M_0|cz;v) = 0.4\ln(10)\frac{L}{L_*}\left(\frac{L}{L_l}\right)^{\alpha} e^{-(L-L_l)/L_*}.$$
 (8)

This approximation will allow an analytic expression for β by minimizing $-\ln P_s = -\sum \ln P(M_{0i}|cz_i, v(\beta))$. For simplicity, we further approximate $v_i(\beta) =$ $F(\beta)v_{1i}$ where v_1 is the line of sight velocity reconstructed with $\beta = 1$. For linear velocity reconstruction from the galaxy distribution in real space $F(\beta) = \beta$. But for reconstruction from redshift space data a significantly better scaling is $F = 2.5\beta/(1 + 1.5\beta)$ (Davis et al. 2011). This is not an exact result, but it suffices here since we are only interested in a general assessment of the expected error on β . Since β appears only via $F(\beta)$, we will perform the minimization with respect to F and write the result in terms of β at the end of the calculation. Using $L = (1 - Fv_{1i}/cz_i)^2 L_0$ and $L_l = (1 - Fv_{1i}/cz_i)^2 L_{0l}$ we get

$$\frac{\partial \ln P_s}{\partial F} = \tag{9}$$

$$\frac{\partial}{\partial F} \sum \left[\ln \left(1 - F \frac{v_{1i}}{cz_i} \right)^2 - \frac{L_{0i} - L_{0li}}{L_*} \left(1 - F \frac{v_{1i}}{cz_i} \right)^2 \right]$$

$$= 2 \sum \left[\frac{\Delta L_i}{L_*} \frac{v_{1i}}{cz_i} - F \left(\frac{v_{1i}}{cz_i} \right)^2 \left(1 + \frac{\Delta L_i}{L_*} \right) \right],$$

where $\Delta L_i = L_{0i} - L_{0li}$ and in the last step we have neglected $O(v_1/cz)^2$ terms and assumed that the Hubble flow-like $\sum v_{1i}/cz_i$ is negligible compared to the other terms. The 1σ shot-noise error on F is, therefore,

$$\delta F = \left(\frac{-2}{\partial^2 \ln P / \partial F^2}\right)^{1/2}$$
(10)
=
$$\left[\sum \left(\frac{v_{1i}}{cz_i}\right)^2 \left(1 + \frac{\Delta L_i}{L_*}\right)\right]^{-1/2}.$$

This expression can be easily estimated when the luminosity, L_0 , is computed from the actual distances, i.e. $cz_i = r_i$. This means that L_0 is the true intrinsic luminosities and, therefore, ΔL and v_1 are uncorrelated. Further, in the limit $L_l \gg L_*$, the average $\langle \Delta L/L_* \rangle$ is unity. From $F = 2.5\beta/(1 + 1.5\beta)$ we get, $\delta\beta \approx 0.4(1+1.5\beta)\delta F$. With all this, equation (10)

¹Any survey geometry with boundary conditions allowing robust velocity reconstruction will suffice for our purposes.

gives

$$\delta\beta = 0.4(1+1.5\beta)(2N)^{-1/2} \frac{c\tilde{z}}{\sigma_{v_1}}$$
(11)
= 0.044(1+1.5\beta) $\left(\frac{10^5}{N}\right)^{1/2} \frac{\tilde{z}}{0.1} \frac{600}{\sigma_{v_1}}$,

where $\tilde{z} = \langle 1/z^2 \rangle^{-1/2}$, and σ_{v1} is the rms value of v_1 in km s⁻¹. The analysis of Davis et al. (2011) gives, on large scales, $\sigma_{v1} \sim 600 \text{ km s}^{-1}$ for $\beta = 1$, at z = 0.

The scaling $\delta\beta \propto N^{-1/2}\tilde{z}$ in Eq. 11 should be approximately valid also when the condition $L_l \gg L_*$ is not strictly satisfied. For 2MRS this scaling gives roughly the same error on β using distant galaxies with $cz > 4000 \text{ km s}^{-1}$ as using galaxies with the lower redshifts .

3.2. Sensitivity to assumed form of the luminosity function

It is instructive to assess the sensitivity of β on the assumed form of the luminosity function. We give here a simple example in which the assumed luminosity function differs greatly from the true form, yet the estimate for β is unbiased. Consider an ideal volume-limited sample of galaxies with a true luminosity function of a Schechter form with $\alpha = 0$, i.e. an exponential distribution. Let is try to recover β assuming a gaussian form for the luminosity function, $P(M_0|cz, v) \propto \sigma_L^{-1}(1 - Fv_1)^2 e^{-(L-L_m)^2/2\sigma_L^2}$, where, like in the previous section, $v(\beta) = F(\beta)v_1$. Minimizing the quantity, $-\sum \ln P(M_{0i}|cz_i, Fv_{1i})$, with respect to L_m , σ_L and F, yields, respectively,

$$\begin{array}{rcl}
0 &=& L_m - \langle L \rangle , & (12) \\
0 &=& \sigma_L^2 - \langle (L - L_m)^2 \rangle , \\
0 &=& \sum_i \left[\frac{\frac{v_{1i}}{cz_i}}{1 - F \frac{v_{1i}}{cz_i}} - \frac{L_0(L_i - L_m)}{\sigma_L^2} \frac{v_{1i}}{cz_i} (1 - F \frac{v_{1i}}{cz_i})^3 \right].
\end{array}$$

For simplicity we further assume that L_0 is computed with $cz_i = r_i$ so that the true solution $\beta = 0$ (the generalization to cases where the true β is different from zero is trivial). Hence, L_0 are equal to the true luminosities and, therefore, follow an exponential distribution for which $\langle L_0 \rangle = L_*$ and $\langle L_0^2 \rangle = 2L_*^2$. Further, there is no correlation between L_0 and $\frac{v_{1i}}{cz_i}$, meaning that the average of products of powers of $(1 - Fv_{1i}/cz_i)$ and L_0 is the product of the averages. In the limit $N \to \infty$, straightforward algebraic manipulation then yields $F(\beta) = 0$, $L_m = L_*$, and $\sigma_L^2 = L_*^2$. Therefore, in this example, where the assumed and true luminosity functions differ grossly, the best fit β is unbiased. This is not surprising since the underlying principle of the method is a reduction in the spread of L. Therefore, the assumed form of the luminosity function should only affect the weighting given to galaxies in a certain luminosity range rather, than the best fit β . Assuming a wrong luminosity function increases the random error on β but does not introduce any systematic biases.

4. Application to 2MRS

In this section we apply the method outlined above to the all sky 2MRS consisting of 23,200 galaxies down to the magnitude K = 11.25. Details about the catalog, including the precise completeness, sky coverage and selection effects can be found in (Huchra et al. 2005). The preparation of the catalog for the purpose of the application of the method is is done similarly to Davis et al. (2011). The peculiar velocity field is derived from the galaxy distribution in the 2MRS for an array of β values using the linear theory methodology of Nusser & Davis (1994). The derived velocity field is robust within $cz < 10^4 \text{ km s}^{-1}$, above that redshift discreteness effects become important. Hence we limit the analysis to the 18,000 galaxies with $cz < 10^4 \text{ km s}^{-1}$. In the derivation of the velocity fields, the galaxy distribution is smoothed with a gaussian window of constant width of 400 km s^{-1} . To further remove strong nonlinearities the derived threedimensional velocity fields are smoothed with an Gaussian window of constant width, R_s . Linear theory recovers the flow pattern reasonably well even at $\delta \lesssim 1$, but not beyond (Nusser et al. 1991; Branchini et al. 2002a). Therefore, although the peculiar velocity field is predicted from the distribution of all galaxies, in the maximization of P_s to assess the robustness of the method we remove galaxies in regions with density contrast higher than δ_{cut} as listed in Table 1 for both values of R_s . Using the expression (7) we minimize $-\ln P_s = -\sum_i \ln P(M_{0i}|cz_i; v(\beta))$ (the summation is over all galaxies) with respect to β and the Schechter parameters, α and M_* .

4.1. Error estimation based on mock 2MRS catalogs

The overall expected errors in β , including possible biases, are based on mock catalogs designed to match the general properties of the 2MRS. For this purpose we use 135 2MRS mock catalogs very similar to those compiled by Davis et al. (2011). These catalogs are extracted from a parent mock catalog of the Two Micron All Sky Survey (2MASS) (Skrutskie et al. 2006). The parent catalog is generated from the Millennium simulation (Springel et al. 2005) using semi-analytic galaxy formation models (De Lucia & Blaizot 2007). All 135 2MRS catalogs satisfy the following conditions. a) The central "observer" in each mock is selected to reside in a galaxy with a quiet velocity field within 500 km s⁻¹, similar to the observed universe, b) The motion of the central galaxy is 500 to 700 km s⁻¹, and c) The density in the environment of the local group, averaged over a sphere of 400 km/s radius , is less than twice the normal. These conditions select central observers that are similar to the conditions of our own Local Group.

The luminosity function in each mock is approximated by a Schechter form. Using the galaxy distribution in each mock, the corresponding peculiar velocity field is generated for an array of β values. The mean and standard deviation of those best values from the 135 catalogs is $\beta = 0.49 \pm 0.13$. The cosmological density parameters of the Millennium simulation are $\Omega\,=\,0.25$ and $\Lambda\,=\,0.75$ for matter and cosmological constant. This yields $f = \Omega^{0.55} = 0.47$ (Linder 2005). A calculation of the rms galaxy density fluctuations of the mocks yields $b = 1 \pm 0.1$ for the mocks. The 1σ rms scatter of the best fit β values is 0.13. This scatter reflects "shot-noise" errors due to the finite number of galaxies, cosmic variance due to the limited volume covered by the 2MRS, and inaccuracies in the reconstruction of the peculiar velocity field by means of linear theory. Shot-noise is subdominant when the method is applied to all 2MRS galaxies within 10^4 km s^{-1} . Since P_s contains no information about cosmic variance and reconstruction errors, the width of $-2\ln P_s$ versus β , reflects shotnoise errors only. From the width of $-2\ln P_s$ we get 1σ shot-noise error of $\delta\beta_{sn} \approx 0.055$ in the application to 2MRS within $cz < 10^4 \text{ km s}^{-1}$. Cosmic variance error could be estimated by running the method with the actual velocity field of galaxies in the mock. This amounts to $\delta\beta_{cv} \approx 0.09$. Adding errors in quadratures, we infer a velocity reconstruction error of $\delta \beta_{vrec} \approx 0.1$, comparable to $\delta\beta_{cv}$.

We will apply the method to various cuts of the 2MRS. All corresponding errors listed in Table 1 are based on similar cuts taken from the mocks. Shotnoise errors, $\delta\beta_{sn}$, will be treated as (nearly) independent of β (see §3.1), while cosmic variance, $\delta\beta_{cv}$ and velocity reconstruction errors, $\delta\beta_{vrec}$, are assumed to be proportional to β , because of the β dependence in the reconstructed field.

4.2. β from the real 2MRS

The results are presented in Table 1 and the top panel of Fig. 1. The plotted quantity, $\Delta \chi^2$, is $-2\ln P_s$ evaluated as a function of β minus its value at the minimum. In the figure, the parameters (α , M_*) are fixed at their best fit values. The overall errors listed in Table 1 are based on the analysis of the mocks in §4.1. The 1σ shot-noise, δ_{sn} , given by the width of the curves at $\Delta\chi^2 = 1$ are significantly smaller than the corresponding total errors quoted in the Table 1, i.e. $\delta\beta_{sn}$ is not the main source of error for the 2MRS catalog. Table 1 lists results only for $R_s =$ $6h^{-1}$ Mpc, however, an application of the method with $R_s = 10h^{-1}$ Mpc and $12h^{-1}$ Mpc yields consistent results within the 1σ errors in the table. The derived β for both choices of δ_{cut} are consistent with Davis et al. (2011) and the values of α and M_* agree well with Westover (2007).

It is useful to examine how the best fit β changes when in the maximization of P_s we include only distance galaxies with $cz > 4000 \text{ km s}^{-1}$. The results are given in the Table 1 and in the bottom panel of Fig. 1 showing $\Delta \chi^2$ versus β for this case. Shot-noise as indicated by width of curves increases with respect to the full sample (see top panel of the same figure). The constraints from the distant cut are still reasonably tight and, within the total 1σ errors, are fully consistent with those obtained from the whole sample.

We also applied the method to galaxies in the Northern and Southern Galactic hemispheres, separately. The results, listed in the Table 1, show a non-negligible difference between the derived β in the two hemispheres. The corresponding error on β is based on the application of the method to "Northern" and "Southern" hemispheres in the mocks. We attribute the difference between the North and the South to cosmic variance and confirm that it is consistent with the mocks. For each mock catalog we compute β_{north} and β_{south} and find an rms value $< (\beta_{north} - \beta_{south})^2 >^{1/2} = 0.2$.

5. Discussion and Conclusions

We have presented a new method to determine β from galaxy redshift surveys. The method is entirely independent of distance indicators such as the Tully-Fisher relation and of analyses of anisotropic correlation functions, $\xi(r_p, \pi)$ (e.g. Kaiser 1987). As a preliminary application of the method, we have resorted to the K=11.25 flux limited 2MRS all sky survey. In the maximization procedure of $P(M_0|cz, v(\beta))$, galaxies in dense regions should be excluded since the flow pattern in these regions is not well recovered by linear theory. For our adopted density contrast cut $\delta_{cut} = 1$, we get a best fit $\beta = 0.35 \pm 0.1$ for velocities smoothed with a gaussian window of $6h^{-1}$ Mpc in width. However, the results obtained with $\delta_{cut} = 2$ and $\delta_{cut} = \infty$ are consistent with this best fit value and are reported in the Table 1. These constraints on β agree very well with those of Davis et al. (2011) who compared the peculiar velocities of the Spiral Field I Band (SFI++) catalog of spiral galaxies (Masters et al. 2006; Springob et al.

Table 1: Derived parameters, $\beta = f/b$, α and M_* for various cuts of the 2MRS. Galaxies in regions with density contrast larger than δ_{cut} and redshifts less than cz_{cut} are excluded from the maximization of P_s . The quoted errors on β are based on mock catalogs and include shot-noise, cosmic variance and inaccuracies in the peculiar velocity reconstruction. The results given here correspond to $R_s = 6h^{-1}$ Mpc.

sky	cz_{cut}	δ_{cut}	fraction	(α, M_*)	$\beta \pm 1\sigma$ error
	${\rm kms^{-1}}$				
all	0	∞	100%	(-0.92, -23.14)	0.30 ± 0.10
all	0	2	79%	(-0.90, -23.19)	0.31 ± 0.10
all	0	1	58%	(-0.89, -23.14)	0.35 ± 0.10
all	4000	2	61%	(-0.84, -23.15)	0.26 ± 0.15
all	4000	1	44%	(-0.81, -23.09)	0.29 ± 0.15
north	0	2	41%	(-0.88, -23.16)	0.23 ± 0.10
north	0	1	30%	(-0.85, -23.09)	0.26 ± 0.11
south	0	2	38%	(-0.93, -23.20)	0.40 ± 0.17
south	0	1	29%	(-0.93, -23.17)	0.41 ± 0.17

2007) with the velocity field predicted from the 2MRS.

There are three sources of uncertainties which contribute to the error budget on the estimated β :

- (1) "shot-noise" due to the finite number of galaxies
- (2) cosmic variance which reflects the variation of the large scale structure in random volumes comparable in size to the volume covered by the data set under consideration.
- (3) inaccuracies in the linear methodology for reconstructing the peculiar velocity from the galaxy distribution.

For the 2MRS sample of \sim 18,000 galaxies within $cz = 10^4 \text{ km s}^{-1}$, shot-noise is subdominant. Increasing the number of galaxies in the sample without probing larger volumes will not tighten the constraint significantly. Cosmic variance can only be reduced by pushing toward deeper and larger surveys. The main galaxy sample of the Sloan Digital Sky Survey (SDSS) (Strauss et al. 2002) already offers this opportunity. Let us consider the $\sim 7500 \text{ deg}^2$ patch around the Northern Galactic cap in the SDSS-DR7 release (Abazajian et al. 2009). A shell with $\Delta z = 0.1$ centered at $z \sim 0.135$, close to the peak of the galaxy dN/dz has a comoving volume ~ 20 times larger than that of the 2MRS and it could be divided into ~ 12 independent cubes of $200h^{-1}$ Mpc, each containing as many galaxy as 2MRS. Applying our method to each of them would dramatically decrease cosmic variance and errors in the reconstructed velocities. According to Eq. 11 the now dominant shot-noise error would be $\delta\beta \sim 0.06$, twice as small as in the 2MRS case, as shown in Table 2.

Shot-noise errors increase linearly with redshift and they will be the limiting factor in limiting the precision of constraining β from the method presented in this paper. Next-generation, large redshift surveys like Boss, BigBoss and EUCLID will allow an application of our method to measure β out to $z \sim 1$. We have applied Eq. 11 to estimate the expected shot noise error on β for all these surveys. The results are listed in Table 2. The ongoing SDSS-III (BOSS) survey will target highly luminous galaxies with a nearly constant number density $n \sim 3 \cdot 10^{-4} h^3 \text{ Mpc}^{-3}$ (Eisenstein et al. 2011) over the redshift range [0.2-0.6]. Because of the relatively low number density of objects our method will measure β with uncertainties twice as large as for the 2MRS case. The BigBOSS survey will expand Boss, both in sky coverage and depth. The expected number of ELGs (Schlegel et al. 2011) will be large enough to decrease the errors on β significantly. In fact, the reduction of the relative errors will be even larger since, in this redshift rage, β is an increasing function of z. Finally, the EUCLID survey (Laureijs 2009) will reduce errors even further in the redshift range z = [0.7 - 1.0].

How do the β estimates obtained with our method compare with those obtained from the analysis of the anisotropy pattern of the two-point correlation function $\xi(r_p, \pi)$? At low (z < 0.2) redshift the $\xi(r_p, \pi)$ method has been applied to SDSS (Tegmark et al. 2006) and 2dF (Hawkins et al. 2003) allowing to estimate β with an error $\delta\beta = 0.1 - 0.15$. The application of our method to 2MRS already gives β with a similar precision and the upcoming application to SDSS-II data (Bernardi et al. 2003) will allow us to estimate β with a precision sufficient to test the va-



Fig. 1.— Top: Curves of $\Delta \chi^2$ as a function of β for $R_s = 6h^{-1}$ Mpc and $10h^{-1}$ Mpc, and $\delta_{cut} = 2$ and 1, as indicated in the figure. The width of the curves reflects uncertainties due to the finite number of galaxies ("shot-noise") and do not include cosmic variance and inaccuracies in the peculiar velocity reconstruction. Bottom: The same as the top panel, but excluding nearby galaxies with $cz < 4000 \text{ km s}^{-1}$ in the minimization of P_s . The widths of curves here are substantially larger than in the top panel due to the smaller number of galaxies and the larger redshifts.

lidity of popular alternative gravity model, like the 5-dimensional brane-world of Dvali et al. (2000); Wei (2008) as well as the possibility of a coupling between the dark energy and the dark matter sectors (di Porto & Amendola 2008).

The first measurement of the growth rate at larger redshifts has been performed by Guzzo et al. (2008). From the observed $\xi(r_p, \pi)$ of VVDS galaxies they obtained $\beta(z = 0.77) = 0.70 \pm 0.26$. More recently, the Wiggle-z experiment (Blake et al. 2011) has measured the normalized growth rate $f(\Omega)\sigma_8$ in 4 redshift bins to z = 0.9 with an error $\delta f(\Omega)\sigma_8 \sim 0.1$. This is already comparable with the expected performance of our method on future datasets. Indeed, our method does not compare favorably with the analysis of galaxy clustering in future surveys. In the case of EUCLID, the goal is to estimate the growth rate from $\xi(r_p,\pi)$ with a precision of 0.01 at z = 1, if the rms mass fluctuation σ_8 can be determined accurately (Song & Percival 2009). In this case our alternative method to measure β will constitute an effective way to keep systematic errors below $\delta\beta = 0.1$. It is important to note that if galaxy bias will only be constrained at the 10% level then both methods, the analysis of $\xi(r_p, \pi)$ and the one proposed here, will constrain the growth rate $f(\Omega)$ with similar precision. Further, on-going and planned redshift surveys will deliver velocity dispersion for all the elliptical galaxies will be obtained. Using the Faber-Jackson relation (Faber & Jackson 1976), between luminosity and velocity dispersion of elliptical galaxies, in conjunction with our method will produce even tighter constraints. Additional constraints on f could also be obtained from the expected large scale supernova survey (e.g. Bhattacharya et al. 2011).

Quasi-linear (i.e. the mildly non-linear regime) dynamical reconstruction methods offer a substantial improvement in the accuracy of the predicted peculiar velocities. In particular, the Fast Action Method (Nusser & Branchini 2000; Branchini et al. 2002b) adaptation of Peebles' least action principle (Shava et al. 1995) is fast and easy to implement. This method is significantly better than linear theory for the reconstruction of the peculiar velocity field on small scales and also in dense regions. The linear theory relation between mass and velocity solely depends on the growth rate $f(\Omega) = \Omega^{\gamma}$. In the quasi-linear regime, there is an additional explicit dependence on Ω , raising the possibility of separate constraints on Ω and the growth index γ . However, this does not seem a promising route to break the $\Omega - \gamma$ degeneracy since the explicit Ω dependence is very weak (~ $\Omega^{0.2}$) (Nusser & Colberg 1998).

Another improvement could be achieved by use of

Table 2: Expected shot-noise errors on β estimated from Eq. 11 for present and future galaxy redshift surveys. To minimize evolutionary effects all errors are estimated in bins $\Delta z = 1$. The reference $\beta(z)$ function has been obtained by considering the growth rate of a Λ CDM model of Linder (2005) and a linear bias $b(z) = \sqrt{(1+z)}$ as in Di Porto et al. (2011). The references to the surveys' parameters are given in the text.

Survey	sky	$\langle z \rangle$	$N_{\rm gals.}$	$\delta\beta$
	deg^2			
SDSS-II	7500	0.135	$3 \cdot 10^5$	0.07
Boss	10000	0.25	$9\cdot 10^4$	0.18
Boss	10000	0.35	$3\cdot 10^5$	0.17
BigBoss	14000	0.45	$5\cdot 10^5$	0.16
BigBoss	14000	0.55	$1.3\cdot 10^6$	0.14
BigBoss	14000	0.65	$2\cdot 10^6$	0.13
EUCLID	15000	0.70	$3.5\cdot 10^6$	0.11
EUCLID	15000	0.80	$6.5\cdot 10^6$	0.09
EUCLID	15000	0.90	$7\cdot 10^6$	0.09
EUCLID	15000	1.00	$7.5\cdot 10^6$	0.10
EUCLID	15000	1.10	$7\cdot 10^6$	0.11

nonparametric fit techniques for modeling the galaxy luminosity distribution. Although the Schechter form is a good approximation for the 2MRS, it is likely less successful for larger and deeper data sets. A variety of such nonparametric fit methods exist (e.g. Efstathiou et al. 1988; Davis & Huchra 1982) could easily be incorporated in the method presented here.

Angular coherent photometric mis-calibrations are likely to contaminate the data at some level. However, as long as the density field inferred from the galaxy distribution is not significantly affected, the method should yield an unbiased β . This is because the underlying velocity field should be uncorrelated with observational mis-calibrations. If systematic biases introduce serious spurious modes in the density field then it is doubtful if the data could be useful for any analysis of large scale clustering.

6. Acknowledgments

We thank Gerard Lemson for help in generating the mock 2MRS catalogs. This work was supported by THE ISRAEL SCIENCE FOUNDATION (grant No.203/09), the German-Israeli Foundation for Research and Development, the Asher Space Research Institute and by the WINNIPEG RESEARCH FUND. MD acknowledges the support provided by the NSF grant AST-0807630. EB acknowledges the support provided by MIUR PRIN 2008 'Dark energy and cosmology with large galaxy surveys' and by Agenzia Spaziale Italiana (ASI-Uni Bologna-Astronomy Dept. 'Euclid-NIS' I/039/10/0) EB also thanks the Physics Department and the Asher Space Science Institute-Technion for the kind hospitality.

REFERENCES

- Abazajian, K. N., et al. 2009, ApJ. S, 182, 543
- Bernardi, M., et al. 2003, AJ, 125, 1817
- Bhattacharya, S., Kosowsky, A., Newman, J. A., & Zentner, A. R. 2011, Physical Review D., 83, 043004
- Blake, C., et al. 2011, MNRAS, 834
- Branchini, E., Eldar, A., & Nusser, A. 2002a, MNRAS, 335, 53
- —. 2002b, MNRAS, 335, 53
- Davis, M., & Huchra, J. 1982, ApJ, 254, 437
- Davis, M., Nusser, A., Masters, K. L., Springob, C., Huchra, J. P., & Lemson, G. 2011, MNRAS, 413, 2906
- De Lucia, G., & Blaizot, J. 2007, MNRAS, 375, 2
- di Porto, C., & Amendola, L. 2008, Physical Review D., 77, 083508
- Di Porto, C., Amendola, L., & Branchini, E. 2011, ArXiv e-prints
- Dvali, G., Gabadadze, G., & Porrati, M. 2000, Physics Letters B, 485, 208
- Efstathiou, G., Ellis, R. S., & Peterson, B. A. 1988, MNRAS, 232, 431
- Eisenstein, D. J., et al. 2011, ArXiv e-prints
- Faber, S. M., & Jackson, R. E. 1976, ApJ, 204, 668

- Gannouji, R., Moraes, B., & Polarski, D. 2009, JCAP, 2, 34
- Guzzo, L., et al. 2008, Nature, 451, 541
- Hawkins, E., et al. 2003, MNRAS, 346, 78
- Huchra, J., et al. 2005, in Astronomical Society of the Pacific Conference Series, Vol. 329, Nearby Large-Scale Structures and the Zone of Avoidance, ed. A. P. Fairall & P. A. Woudt, 135–+
- Kaiser, N. 1987, MNRAS, 227, 1
- -. 1988, MNRAS, 231, 149
- Keselman, J. A., Nusser, A., & Peebles, P. J. E. 2010, Physical Review D., 81, 063521
- Laureijs, R. 2009, ArXiv e-prints
- Linder, E. V. 2005, Physical Review D., 72, 043529
- Masters, K. L., Springob, C. M., Haynes, M. P., & Giovanelli, R. 2006, ApJ, 653, 861
- Nusser, A., & Branchini, E. 2000, MNRAS, 313, 587
- Nusser, A., Branchini, E., & Davis, M. 2011, ApJ, 735, 77
- Nusser, A., & Colberg, J. M. 1998, MNRAS, 294, 457
- Nusser, A., & Davis, M. 1994, ApJL, 421, L1
- Nusser, A., Dekel, A., Bertschinger, E., & Blumenthal, G. R. 1991, ApJ, 379, 6
- Schechter, P. L. 1980, Astronomical Journal, 85, 801
- Schlegel, D., et al. 2011, ArXiv e-prints
- Shaya, E. J., Peebles, P. J. E., & Tully, R. B. 1995, ApJ, 454, 15
- Skrutskie, M. F., et al. 2006, AJ, 131, 1163
- Song, Y.-S., & Percival, W. J. 2009, JCAP, 10, 4
- Springel, V., et al. 2005, Nature, 435, 629
- Springob, C. M., Masters, K. L., Haynes, M. P., Giovanelli, R., & Marinoni, C. 2007, ApJ. S, 172, 599
- Strauss, M. A., et al. 2002, AJ, 124, 1810
- Tegmark, M., et al. 2006, Physical Review D., 74, 123507
- Wei, H. 2008, Physics Letters B, 664, 1
- Westover, M. 2007, PhD thesis, Harvard University

This 2-column preprint was prepared with the AAS ${\rm IAT}_{\rm E}{\rm X}$ macros v5.2.