Heavy-flavor dynamics in nucleus–nucleus collisions: from RHIC to LHC

M. Monteno¹, W.M. Alberico^{2,1}, A. Beraudo^{3,4}, A. De Pace¹, A. Molinari^{2,1}, M. Nardi¹ and F. Prino¹

¹Istituto Nazionale di Fisica Nucleare, Sezione di Torino,
via P. Giuria 1, I-10125 Torino, Italy
²Dipartimento di Fisica Teorica dell'Università di Torino,
via P. Giuria 1, I-10125 Torino, Italy
³Centro Studi e Ricerche *Enrico Fermi*, Piazza del Viminale 1, Roma, Italy
⁴Physics Department, Theory Unit, CERN, CH-1211 Genève 23, Switzerland

E-mail: monteno@to.infn.it

Abstract. The stochastic dynamics of c and b quarks in the fireball created in nucleus–nucleus collisions at RHIC and LHC is studied employing a relativistic Langevin equation, based on a picture of multiple uncorrelated random collisions with the medium. Heavy-quark transport coefficients are evaluated within a pQCD approach, with a proper HTL resummation of medium effects for soft scatterings. The Langevin equation is embedded in a multi-step setup developed to study heavy-flavor observables in pp and AA collisions, starting from a NLO pQCD calculation of initial heavy-quark yields, complemented in the nuclear case by shadowing corrections, k_T -broadening and nuclear geometry effects. Then, only for AA collisions, the Langevin equation is solved numerically in a background medium described by relativistic hydrodynamics. Finally, the propagated heavy quarks are made hadronize and decay into electrons. Results for the nuclear modification factor R_{AA} of heavy-flavor hadrons and electrons from their semi-leptonic decays are provided, both for RHIC and LHC beam energies.

Heavy-flavor electron spectra, measured in Au–Au collisions at $\sqrt{s_{NN}}=200~{\rm GeV}$ by PHENIX [1, 2] and STAR [3] experiments at RHIC, have displayed a large suppression with respect to pp collisions, comparable in amount to the one observed for charged hadrons. Models considering medium-induced gluon radiation [4, 5] as the dominant energy loss mechanism for heavy quarks propagating in QGP come up against difficulties in reproducing such results.

These findings gave a great boost to calculations taking in consideration the role of collisional energy loss [6, 7]. In some models the heavy-quark propagation in QGP is described through a Langevin stochastic equation [8, 9, 10], assuming that heavy-quark spectrum modifications arise from the cumulated effect of many uncorrelated random collisions with the medium.

In our approach we use a relativistic Langevin equation [11], describing the time evolution of the heavy-quark momentum:

$$\frac{\Delta p^i}{\Delta t} = -\eta_D(p)p^i + \xi^i(t) \tag{1}$$

that involves a *deterministic* friction term and a *stochastic* noise term $\xi^{i}(t)$, completely determined by its two-point temporal correlator:

$$\langle \xi^{i}(t)\xi^{j}(t')\rangle = b^{ij}(\boldsymbol{p})\delta(t-t'), \text{ with } b^{ij}(\boldsymbol{p}) \equiv \kappa_{L}(p)\hat{p}^{i}\hat{p}^{j} + \kappa_{T}(p)(\delta^{ij} - \hat{p}^{i}\hat{p}^{j})$$
 (2)

The latter involves the transport coefficients $\kappa_T(p) \equiv \frac{1}{2} \frac{\langle \Delta p_T^2 \rangle}{\Delta t}$ and $\kappa_L(p) \equiv \frac{\langle \Delta p_L^2 \rangle}{\Delta t}$, which are evaluated according to the procedure presented in [12]. We introduce an intermediate cutoff $|t|^* \sim m_D^2$ $(t \equiv (P'-P)^2)$ to separate hard and soft scatterings. The contribution of hard collisions $(|t| > |t|^*)$ is evaluated through a pQCD calculation of the processes $Q(P)q_{i/\bar{i}} \to Q(P')q_{i/\bar{i}}$ and $Q(P)g \to Q(P')g$. On the other hand for soft collisions $(|t| < |t|^*)$ a resummation of medium effects is provided by the Hard Thermal Loop approximation, with $\alpha_s(\mu)$ evaluated at a scale $\mu \propto T$. The final result [12] is given by $\kappa_{T/L}(p) = \kappa_{T/L}^{\text{hard}}(p) + \kappa_{T/L}^{\text{soft}}(p)$. According to the scale μ at which $\alpha_s(\mu)$ is evaluated to calculate $\kappa_{T/L}^{\text{hard}}$, we devised two different sets of calculations, referred to in the following as HTL1 (for $\mu \propto T$) and HTL2 (for $\mu = |t|$).

The Langevin simulation tool is embedded in a full setup to calculate heavy-flavor observables in pp and AA collisions, divided into the following independent steps [12]:

- (i) A sample of c and b quarks is generated using POWHEG [13], a code which implements pQCD at NLO accuracy, with CTEQ6M PDFs as input. For AA collisions, EPS09 nuclear corrections to PDFs are employed [14]; then, heavy quarks are distributed in the transverse plane according to the nuclear overlap function $T_{AB}(x,y) \equiv T_A(x+b/2,y)T_B(x-b/2,y)$ corresponding to the selected impact parameter b; a p_T -broadening correction to heavy-quark momenta is also included.
- (ii) At a given proper-time τ_0 an iterative procedure is started, only for AA collisions, to follow the stochastic evolution of the heavy quarks in the plasma until hadronization: the Langevin transport coefficients are evaluated at each step according to the local 4-velocity and temperature T(x) of the expanding background medium, as provided by hydrodynamic codes: both ideal and viscous fluid scenarios were tested [15, 16, 17].
- (iii) Heavy quarks are made fragment into hadron species, sampled according to c and b branching fractions taken from Refs. [18, 19], while their momenta are sampled from a Peterson fragmentation function [20], with $\epsilon = 0.04$ and 0.005 for c and b respectively. Finally, each heavy-quark hadron is forced to decay into electrons with PYTHIA [21], using updated tables of branching ratios based on Ref. [19].

The effects of the Langevin evolution of c and b quarks in AA collisions resulting from our calculations are studied through the nuclear modification factor $R_{AA}(p_T) \equiv (dN/dp_T)^{AA}/\langle N \rangle_{\text{coll}} (dN/dp_T)^{pp}$ and the elliptic flow coefficient $v_2(p_T) \equiv \langle \cos(2\phi) \rangle_{p_T}$ of




Figure 1. (left) $R_{AA}(p_T)$ of heavy-flavor electrons in minimum-bias Au–Au collisions at RHIC (0 – 92% of the inelastic cross section); (right) R_{AA} obtained by integrating the electron yields over the indicated momentum ranges, as a function of N_{part} . In both panels our predictions (with HTL1 calculation) are compared to PHENIX data [2].

final-state heavy-quark hadrons or decay-electrons. Here we will display only results obtained using viscous hydrodynamics for some representative values of the input parameters (τ_0 and the QCD scale μ), among those fully explored in [12].

In Fig. 1 our findings for the R_{AA} of heavy-flavor electrons in Au–Au collisions at RHIC ($\sqrt{s}_{NN}=200~{\rm GeV}$), obtained with the calculation HTL1 by assuming viscous hydrodynamics, $\tau_0=1~{\rm fm/c}$ and $\mu=3\pi T/2$, are compared to PHENIX [2] data. In the left panel we observe a general agreement of $R_{AA}(p_T)$ with the PHENIX results on a minimum-bias data sample (0-92% of the inelastic cross section) for $p_T \gtrsim 3~{\rm GeV/c}$. In the right panel, displaying the R_{AA} obtained by integrating the electron yields above a given p_T and plotted versus N_{part} , the centrality dependence of the PHENIX data is shown to be nicely reproduced, except for the most-peripheral centrality bin. We do not show here the results obtained for the heavy-flavor electron $v_2(p_T)$ [12], that appear to underestimate the PHENIX minimum-bias data. However, a more detailed treatment of hadronization, including also the coalescence mechanism, should enhance both v_2 and R_{AA} at $p_T \lesssim 3~{\rm GeV/c}$.

In Fig. 2 we show our predictions for the $R_{AA}(p_T)$ of heavy-flavor electrons (left panel) and of D, B mesons (right panel) in Pb–Pb collisions at LHC ($\sqrt{s}_{NN}=2.76$ TeV), calculated by selecting the 0-10% most-central events, under the hypothesis of viscous hydrodynamics and for two different choices (HTL1 or HTL2) of the μ scale in the calculation of $\kappa_{T/L}^{hard}$. General features of the R_{AA} of heavy-flavor electrons appear similar to those observed at RHIC at the same centrality [12], with a stronger suppression of both charm and bottom contributions. However, results obtained with the calculation HTL2 display a weaker quenching for $p_T > 3$ GeV/c. As regards D and B suppression, slighter differences between HTL1 and HTL2 are observed, and at higher p_T .

The main goal of our study was to deliver a weak-coupling calculation to be used




Figure 2. The R_{AA} of heavy-flavor electrons (left) and D, B mesons (right) in Pb–Pb collisions at LHC ($\sqrt{s}_{NN}=2.76$ TeV) for the 0-10% most-central events obtained with the HTL1 (solid line) and HTL2 (dashed line) calculations.

as a benchmark for advanced studies or less conventional scenarios. The capability to accommodate the electron spectra observed at RHIC for $p_T \gtrsim 3 \text{ GeV/c}$ strengthens the hypothesis that heavy-quark collisional energy loss must be taken into proper account. Moreover, we will soon be able to test our predictions against first data with Pb-Pb collisions delivered by LHC experiments [22, 23, 24].

References

- [1] Adare A et al. (PHENIX Collaboration) 2007 Phys. Rev. Lett. 98 172301
- [2] Adare A et al. (PHENIX Collaboration) 2010 Preprint arXiv:nucl-ex/1005.1627
- [3] Abelev B I et al. (STAR Collaboration) 2011 Phys. Rev. Lett. 106 159902
- [4] Djordjevic M and Gyulassy M 2004 Nucl. Phys. A 733 265
- Armesto N, Salgado C A and Wiedemann U A 2004 Phys. Rev. D 69 114003
- [6] van Hees H, Greco V and Rapp R 2006 Phys. Rev. C 73 034913
- [7] Uphoff J et al. 2011 Preprint arXiv:hep-ph/1104.2437
- [8] Moore G D and Teaney D 2005 Phys. Rev. C **71** 064904
- [9] Gossiaux P B, Bierkandt R and Aichelin J 2009 Phys. Rev. C 79 044906
- [10] Akamatsu Y, Hatsuda T and Hirano T 2009 Phys. Rev. C 79 054907
- [11] Beraudo A et al. 2009 Nucl. Phys. A 831 59
- [12] Alberico W M et al. 2011 Eur. Phys. J. C **71** 1666
- [13] Frixione S, Nason P and Ridolfi G 2007 J. High Energy Phys. JHEP 0709 (2007) 126
- [14] Eskola K J, Paukkunen H and Salgado C A 2009 J. High Energy Phys. JHEP 0904 (2009) 065
- [15] Kolb P F, Sollfrank J and Heinz U 2000 Phys. Rev. C 62 054909
- [16] Romatschke P and Romatschke U 2007 Phys. Rev. Lett. 99 172301
- [17] Luzum M and Romatschke P 2008 Phys. Rev. C 78 034915
- [18] Chekanov S et al. (ZEUS Collaboration) 2005 Eur. Phys. J. C 44 351
- [19] Nakamura N et al. (Particle Data Group) 2010 J. Phys. G 37 075021
- [20] Peterson C et al. 1983 Phys. Rev. D 27 105
- [21] Sjostrand T, Mrenna S and Skands P Z 2006 J. High Energy Phys. JHEP 0605 (2006) 026
- [22] Masciocchi S (for the ALICE Collaboration) 2011 These proceedings
- [23] Rossi A (for the ALICE Collaboration) 2011 These proceedings
- [24] Dahms T (for the CMS Collaboration) 2011 These proceedings