

Some properties of the moment estimator of shape parameter for the gamma distribution

Piotr Nowak

Mathematical Institute, University of Wrocław

Pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland

E-mail: nowak@math.uni.wroc.pl

Abstract

Exact distribution of the moment estimator of shape parameter for the gamma distribution for small samples is derived. Order preserving properties of this estimator are presented.

Keywords: moment estimator, stochastic ordering, student's ratio, dispersive ordering.

2010 MSC: 60E15, 62E15.

1 Introduction and preliminaries

It is well known that the gamma distribution has wide application, for example in meteorology to describe the distribution of rainfall.

The moment estimators $\hat{\alpha}$ and $\hat{\lambda}$ of the parameters α and λ of the gamma distribution with density

$$f(x; \alpha, \lambda) = \frac{1}{\Gamma(\alpha)\lambda^\alpha} x^{\alpha-1} \exp(-x/\lambda), \quad x > 0, \alpha > 0, \lambda > 0$$

for a random sample $X_1, \dots, X_n$ are

$$(1.1) \quad \hat{\alpha} = \frac{\bar{X}^2}{S^2} \quad \text{and} \quad \hat{\lambda} = \frac{S^2}{\bar{X}},$$

where $\bar{X}$ and S^2 are sample mean and sample biased variance respectively.

In this paper we are mainly interested in order preserving property of the estimator $\hat{\alpha}$. In general, we say that the estimator $\hat{\theta}$ based on the sample $X_1, \dots, X_n$ from population with density $f(\cdot; \theta)$ is increasing in θ with respect to the order $\prec$ if $\hat{\theta}_1 \prec \hat{\theta}_2$, whenever $\theta_1 < \theta_2$, where $\hat{\theta}_1$ ($\hat{\theta}_2$) is the estimator based on sample from density $f(\cdot; \theta_1)$ ($f(\cdot; \theta_2)$). We shall consider when $\prec$ is one of the following orders: $\leq_{\text{st}}$, $\leq_{\text{disp}}$, $\leq^*$. For details and definitions of these stochastic orders we refer the reader to Shaked and Shanthikumar [5].

Nowak [3] proved the following theorem.

Theorem 1. *The estimators $\hat{\alpha}$ and $\hat{\lambda}$ are stochastically increasing respectively in α and λ .*

Deriving exact distributions of moment estimators $\hat{\alpha}$ and $\hat{\lambda}$ is very tedious for $n > 2$. Now we find the distribution of $\hat{\alpha}$ for $n = 2$. First we prove the more general fact.

Lemma 1. *Suppose that $X_1 \sim G(\lambda, \alpha_1)$ and $X_2 \sim G(\lambda, \alpha_2)$, X_1 and X_2 are independent random variables. Then $Z = \frac{S^2}{\bar{X}^2}$ has density of the form*

$$h(z; \alpha_1, \alpha_2) = 1/B(\alpha_1, \alpha_2) 2^{-(\alpha_1 + \alpha_2)} z^{-1/2} \times \\ \times [(1 - \sqrt{z})^{\alpha_1 - 1} (1 + \sqrt{z})^{\alpha_2 - 1} + (1 + \sqrt{z})^{\alpha_1 - 1} (1 - \sqrt{z})^{\alpha_2 - 1}].$$

Corollary 1. *Let $X_1, X_2 \sim G(\lambda, \alpha)$, $Y \sim Ex(1)$, X_1 , X_2 and Z are independent random variables. Then*

$$(1.2) \quad \frac{X_1^2 + X_2^2}{(X_1 + X_2)^2} =_{st} \frac{(X_1 + Y)^2 + X_2^2}{(X_1 + Y + X_2)^2}.$$

Problem 1. *Equation (1.2) yields some interesting property of the exponential distribution. There arises a question: Does this equation characterize the exponential distribution?*

Corollary 2. *For $n = 2$ the statistics $1/\hat{\alpha}$ has the beta distribution $B(1/2, \alpha)$. Hence $\hat{\alpha}$ has monotone likelihood ratio.*

Now we examine dispersive ordering of the $\hat{\alpha}$ estimator. First we notice that $1/\hat{\alpha}$ is not dispersive monotone in α . We conclude it from Theorem 3.B.14 of Shaked and Shanthikumar [5] since the support of the estimator $1/\hat{\alpha}$ is the finite interval $(0, n - 1)$.

Theorem 2. *The estimator $1/\hat{\alpha}$ is not dispersively monotone in α .*

We now consider dispersive ordering of the estimator $\hat{\alpha}$. From previous theorem it does not follow that $\hat{\alpha}$ is not dispersive monotone in α , though the function $1/x$ is decreasing and convex for $x > 0$. We should notice that Theorem 3.B.10(b) of Shaked and Shanthikumar [5] is not valid here. For example, in the following theorem we prove that for $n = 2$ the estimator $\hat{\alpha}$ is dispersively increasing in α .

Theorem 3. *For $n = 2$ the estimator $\hat{\alpha}$ is dispersively increasing in α .*

Proof. Let f^α denotes density of the estimator $\hat{\alpha}$. We know that $1/\hat{\alpha}$ has the beta distribution $\mathcal{B}(1/2, \alpha)$, so

$$f^\alpha(x) = \frac{1}{B(1/2, \alpha)} x^{-3/2} (1 - 1/x)^{\alpha-1}, \quad x > 1.$$

In order to prove that $\hat{\alpha}_1 \leq_{\text{disp}} \hat{\alpha}_2$ whenever $\alpha_1 < \alpha_2$ it suffices to prove that

$$S^-(f_c^{\alpha_1}(x+1) - f^{\alpha_2}(x+1)) \leq 2 \quad \text{for every } c > 0$$

with the sign sequence being $-, +, -$ in the case of equality and $\hat{\alpha}_1 \leq_{\text{st}} \hat{\alpha}_2$ (see Theorem 2.6 and Remark 2.1 of Shaked [4]).

Fix $c > 0$. When $x > c$ then

$$\begin{aligned} f_c^{\alpha_1}(x+1) - f^{\alpha_2}(x+1) &= \frac{1}{B(1/2, \alpha_2)} (x+1)^{-3/2} (1 - 1/(x+1))^{\alpha_2-1} \times \\ &\times [A(x+1)^{3/2} (x-c+1)^{-3/2} (1 - 1/(x-c+1))^{\alpha_1-1} / (1 - 1/(x+1))^{\alpha_2-1} - 1], \end{aligned}$$

where $A = B(1/2, \alpha_2)/B(1/2, \alpha_1)$. Thus $S^-(f_c^{\alpha_1} - f^{\alpha_2}) = S^-(h(x))$, where

$$\begin{aligned} h(x) &= \log A + 3/2 \log(x+1) - 3/2 \log(x-c+1) + \\ &+ (\alpha_1 - 1) \log(1 - 1/(x-c+1)) - (\alpha_2 - 1) \log(1 - 1/(x+1)). \end{aligned}$$

Now we show that h has exactly one maximum when $\alpha_1 > 1$ and the sign sequence is $-, +, -$. In the other case h is decreasing. After calculating we have that $h'(x) = 0$ if and only if $-w(x)/(2x(x+1)(x-c)(x-c+1)) = 0$, where

$$w(x) = (2\alpha_2 - 2\alpha_1 + 3c)x^2 + (2\alpha_2 - 2\alpha_1 + 4c - 4\alpha_2 c - 3c^2)x + 2c - 2\alpha_2 c - 2c^2 + 2\alpha_2 c^2.$$

So we must show that w has only one root in the interval (c, ∞) if $\alpha_1 > 1$ and no roots when $\alpha_1 \in (0, 1]$. Define $\bar{w}(x) = w(x+c)$. Therefore

$$\bar{w}(x) = (2(\alpha_2 - \alpha_1) + 3c)x^2 + (2(\alpha_2 - \alpha_1) + 4c(1 - \alpha_1) + 3c^2)x + 2(1 - \alpha_1)(c + c^2).$$

It is easy to see that for $\alpha_1 > 1$, $\bar{w}$ has two different roots x_1 and x_2 that $x_1 x_2 < 0$ and for $\alpha_1 \in (0, 1]$ $\bar{w}$ has no roots in $(0, \infty)$. At the end we must show that the sign sequence is $-, +, -$ when $S^-(h) = 2$. For $\alpha_1 > 1$

$$\lim_{x \rightarrow c^+} h(x) = -\infty, \quad \lim_{x \rightarrow \infty} h(x) = \log A < 0.$$

Combining these above facts with Theorem 1 we end the proof.

Remark 1. The above theorem does not hold in general for $n \geq 3$, but direct calculating is impossible due to occurrence of hyper elliptic integrals. For example, numerical calculations show, that for $n = 3$, $\hat{\alpha}_1 \not\leq_{\text{disp}} \hat{\alpha}_2$ if $\alpha_1 = 1/5$ and $\alpha_2 = 1/4$. Then $G^{-1}(x) - F^{-1}(x)$ has the local maximum at $x \approx 0.72$ and the local minimum at $x \approx 0.85$.

Deriving exact distribution when $n = 3$ is more tedious. We should add, that the estimator $\hat{\alpha}$ is closely related to the student statistic $\sqrt{n}\bar{X}/S$. The properties of this statistic was very intensively studied in the literature, especially under non-normal conditions. First, we find joint distribution of the vector $(\bar{X}, S)$. One can proof the following lemma, see for example Craig [2].

Lemma 2. *If f is a density on $(0, \infty)$, then the joint distribution $(\bar{X}, S)$ for $n = 3$ is given by formula*

$$F(\bar{x}, s) = \begin{cases} 18s \int_{\bar{x}-s\sqrt{2}}^{\bar{x}+s\sqrt{2}} \frac{1}{R} f(x_1) f\left(\frac{3\bar{x}-x_1+R}{2}\right) f\left(\frac{3\bar{x}-x_1-R}{2}\right) dx_1, & 0 \leq s \leq \bar{x}\sqrt{2}/2, \\ 18s \left[\int_0^{\frac{3\bar{x}-\sqrt{6s^2-3\bar{x}^2}}{2}} + \int_{\frac{3\bar{x}+\sqrt{6s^2-3\bar{x}^2}}{2}}^{\bar{x}+s\sqrt{2}} \right] \frac{1}{R} f(x_1) f\left(\frac{3\bar{x}-x_1+R}{2}\right) f\left(\frac{3\bar{x}-x_1-R}{2}\right) dx_1, & \bar{x}\sqrt{2}/2 \leq s \leq \bar{x}\sqrt{2}, \end{cases}$$

where $R = \sqrt{6s^2 - 3(x_1 - \bar{x})^2}$.

If we have the joint distribution $(\bar{X}, S)$ it is easy to derive the distribution of the statistics $T = \bar{X}/S = \sqrt{\hat{\alpha}}$ from the formula

$$f_T(t) = \int u F(ut, u) du.$$

Proposition 1. *For $n = 3$ the cumulative distribution function of T for the exponential distribution with density $f(x) = e^{-x}$, $x > 0$ is given by*

$$F(t) = \begin{cases} 1 - \frac{2\pi}{3\sqrt{3}t^2}, & t \geq \sqrt{2}, \\ -\frac{\sqrt{2-t^2}}{\sqrt{3}t} + \frac{\pi}{3\sqrt{3}t^2} - \frac{2 \arcsin\left(\frac{t}{\sqrt{2}}\right)}{\sqrt{3}t^2} + 1, & \frac{\sqrt{2}}{2} \leq t \leq \sqrt{2}. \end{cases}$$

On the other hand it is not easy to derive in the general the distribution of T . For example, if $\alpha \in (0, 1)$ above integrals can be calculated only by numerical methods. For $n = 3$, after a bit algebra we can also derive the distribution of T for the gamma distribution with shape parameter $\alpha = 2$.

Proposition 2. *For $n = 3$ the cumulative distribution function of T for the gamma distribution with density $g(x) = xe^{-x}$, $x > 0$ is given by*

$$G(t) = \begin{cases} 1 - \frac{10\pi(4t^2-3)}{27\sqrt{3}t^4}, & t \geq \sqrt{2}, \\ \frac{\sqrt{2-t^2}(-33t^4-13t^2+8)+5\pi t(4t^2-3)-30t(4t^2-3) \arcsin\left(\frac{t}{\sqrt{2}}\right)}{27\sqrt{3}t^5} + 1, & \frac{\sqrt{2}}{2} \leq t \leq \sqrt{2}. \end{cases}$$

From previous propositions it follows, that the statistic T is not monotone with respect to the star order $\leq_*$, that is the function $G^{-1}F(x)/x$ is not monotone in $x > 0$. To see it, let us calculate:

$$\begin{aligned} 1) \left. \frac{G^{-1}F(x)}{x} \right|_{x=\frac{11}{10}} &= \frac{10}{33} \sqrt{\frac{11 \left(220\pi + \sqrt{10\pi(-891\sqrt{79}-560\pi+16200 \arccos(\frac{11}{10\sqrt{2}}))} \right)}{33\sqrt{79}+200\pi-600 \arccos(\frac{11}{10\sqrt{2}})}} \approx 1.32686 \\ 2) \left. \frac{G^{-1}F(x)}{x} \right|_{x=\frac{\sqrt{6}}{2}} &= \sqrt{\frac{40\pi+2\sqrt{10\pi(22\pi-27\sqrt{3})}}{27\sqrt{3}+18\pi}} \approx 1.31502 \\ 3) \left. \frac{G^{-1}F(x)}{x} \right|_{x=\sqrt{2}} &= \frac{\sqrt{40+2\sqrt{130}}}{6} \approx 1.32081 \end{aligned}$$

Since the star order is preserved under increasing function, we have the following corollary.

Corollary 3. *For $n = 3$ the estimator $\hat{\alpha}$ is not increasing in α with respect to the star order.*

The star order is closely related to the dispersive order, since

$$X \leq^* Y \quad \text{iff} \quad \log X \leq_{\text{disp}} \log Y.$$

Using the same technique as in Theorem 3 we can proof the following proposition.

Proposition 3. *For $n = 2$ the estimator $\hat{\alpha}$ is increasing in α with respect to the star order.*

Applying the property that for nonnegative random variables such $X \leq_{\text{st}} Y$ and $X \leq^* Y$ implies the ordering $X \leq_{\text{disp}} Y$ (see, for example, Ahmed et al. [1]) we can also deduce from Theorem 1 and Proposition 3 that $\hat{\alpha}$ is dispersively monotone for $n = 2$.

References

- [1] A. N Ahmed, A. Alzaid, J. Bartoszewicz, S. C. Kocher, *Dispersive and super-additive ordering*, Advances in Applied Probability 18, (1986), 1019–1022.
- [2] Allen T. Craig, *The simultaneous distribution of mean and standard deviation in small samples*, The Annals of Mathematical statistics, Vol 3, No. 2, 1932, p.126–140.
- [3] P. Nowak, *Stochastic comparisons of moment estimators of gamma distribution parameters*, to appear in Applicationes Mathematicae.
- [4] M. Shaked, *Dispersive ordering of distributions*, J. Appl. Prob. 19 (1982), 310–320.
- [5] M. Shaked, J. G. Shanthikumar, *Stochastic Orders*, Springer Verlag, New York, 2007.