

Gluon distribution in proton at soft and hard pp collisions

G.I.Lykasov^{1,4}, V.A.Bednyakov¹, A.A.Grinyuk¹, M.Poghosyan^{2,3}
and A.G.Dolbilov¹

¹ *JINR, Dubna, Moscow region, 141980, Russia,*

² *Torino University, Torino, Italy*

³ *CERN, Geneva, Switzerland*

⁴ *E-mail: lykasov@jinr.ru*

Abstract

We analyze the inclusive spectra of hadrons produced in pp collisions at high energies in the mid-rapidity region within the soft QCD and perturbative QCD assuming the possible creation of the soft gluons at low intrinsic transverse momenta k_t . From the best description of the LHC data we found the parametrization of the unintegrated gluon distribution which at low k_t is different from the one obtained within the perturbative QCD.

1. Introduction

As is well known, hard processes involving incoming protons, such as deep-inelastic lepton-proton scattering (DIS), are described using the scale-dependent PDFs. A distribution like this is usually calculated as a function of the Bjorken variable x and the square of the four-momentum transfer $q^2 = -Q^2$, integrated over the parton transverse momentum k_t . However, for semi-inclusive processes, such as inclusive jet production in DIS, electroweak boson production [1], etc., the parton distributions unintegrated over the transverse momentum k_t are more appropriate. The theoretical analysis of the unintegrated quark and gluon PDFs can be found, for example, in [2]-[5]. According to [3], the gluon distribution function $g(k_t)$ at fixed Q^2 has the very interesting behaviour at small $x \leq 0.01$, it increases very fast starting from almost zero values of k_t . In contrast, the quark distribution $q(k_t)$ is almost constant in the whole region of k_t up to $k_t \sim 100$ GeV/ c and much smaller than $g(x)$. This parametrization of the PDFs was obtained in [3] within the leading order (LO) and next-to-leading order of QCD (NLO) at large Q^2 from the known (DGLAP-evolved [6]) parton densities determined from the global data analysis. At small values of Q^2 the nonperturbative effects should be included to get the PDFs. The nonperturbative effects can arise from the complex structure of the QCD vacuum. For example, within the instanton approach [4] the very fast increase of the unintegrated gluon distribution function at $0 \leq k_t \leq 0.5$ GeV/ c and $Q^2 = 1$ (GeV/ c)² is also shown. These results stimulated us to assume, that the unintegrated gluon distribution in the proton can be included by analyzing also the soft hadron production in pp collisions. In this paper we analyze the inclusive spectra of the hadrons produced in pp collisions at LHC energies in the mid-rapidity

region including the possible creation of soft gluons in the proton. We estimate the unintegrated gluon distribution function (UGDF) at low intrinsic transverse momenta $k_t \leq 1.5 - 1.6$ GeV/c and its parameters extract from the best description of the LHC data at low transverse momenta p_t of the produced hadrons. We also show that our UGDF is similar to the UGDF obtained in [2, 5] at large values of k_t .

2. Inclusive spectra of hadrons in pp collisions

2.1 Quark-gluon string model (QGSM)

Let us analyze the hadron production in pp collisions within the QGSM [7] or the dual parton model (DPM) [8] including the transverse motion of quarks and diquarks in colliding protons [9, 10]. As is known, the cylinder-type graphs presented in Fig.1 make the main contribution to this process. The left diagram of Fig.1, the so-called one-cylinder graph, corresponds to the cut one-pomeron graph and the right diagram of Fig.1, the so-called multi-cylinder graph, corresponds to the cut n -pomeron graphs. The general form for the invariant inclusive hadron spectrum




Figure 1: The one-cylinder graph (left) and the multi-cylinder graph (right)

within the QGSM is the following [7]:

$$\rho(x, p_t) \equiv E \frac{d\sigma}{d^3\mathbf{p}} = \sum_{n=1}^{\infty} \sigma_n(s) \phi_n(x, p_t) , \quad (1)$$

where $E, \mathbf{p}$ are the energy and three-momentum of the produced hadron h in the l.s. of colliding protons respectively; x, p_t are the Feynman variable and the transverse momentum of h ; σ_n is the cross section for production of the n -pomeron chain (or $2n$ quark-antiquark strings) decaying into hadrons, calculated within the “eikonal approximation” [11] the function $\phi_n(x, p_t)$ has the following form [9]:

$$\phi_n(x, p_t) = \int_{x^+}^1 dx_1 \int_{x_-}^1 dx_2 \psi_n(x, p_t; x_1, x_2) , \quad (2)$$

where

$$\psi_n(x, p_t; x_1, x_2) =$$

$$\begin{aligned}
& F_{qq}^{(n)}(x_+, p_t; x_1) F_{q_v}^{(n)}(x_-, p_t; x_2) / F_{q_v}^{(n)}(0, p_t) + \\
& + F_{q_v}^{(n)}(x_+, p_t; x_1) F_{qq}^{(n)}(x_-, p_t; x_2) / F_{qq}^{(n)}(0, p_t) + \\
& 2(n-1) F_{q_s}^{(n)}(x_+, p_t; x_1) F_{\bar{q}_s}^{(n)}(x_-, p_t; x_2) / F_{q_s}^{(n)}(0, p_t) .
\end{aligned} \tag{3}$$

and $x_{\pm} = 0.5(\sqrt{x^2 + x_t^2} \pm x)$, $x_t = 2\sqrt{(m_h^2 + p_t^2)}/s$,

$$\begin{aligned}
& F_{\tau}^{(n)}(x_{\pm}, p_t; x_{1,2}) = \\
& \int d^2 k_t \tilde{f}_{\tau}^{(n)}(x_{\pm}, k_t) \tilde{G}_{\tau \rightarrow h} \left(\frac{x_{\pm}}{x_{1,2}}, \tilde{k}_t; p_t \right) ,
\end{aligned} \tag{4}$$

$$\begin{aligned}
& F_{\tau}^{(n)}(0, p_t) = \\
& \int_0^1 dx' d^2 k_t \tilde{f}_{\tau}^{(n)}(x', k_t) \tilde{G}_{\tau \rightarrow h}(0, p_t) = \tilde{G}_{\tau \rightarrow h}(0, p_t) .
\end{aligned} \tag{5}$$

Here τ means the flavour of the valence (or sea) quark or diquark, $\tilde{f}_{\tau}^{(n)}(x', k_t)$ is the quark distribution function depending on the longitudinal momentum fraction x' and the transverse momentum k_t in the n -pomeron chain; $\tilde{G}_{\tau \rightarrow h} = z \tilde{D}_{\tau \rightarrow h}(z, \tilde{k}_t; p_t)$, $\tilde{D}_{\tau \rightarrow h}(z, \tilde{k}_t; p_t)$ is the fragmentation function of a quark (antiquark) or diquark of flavour τ into a hadron h .

2..2 Hadron production in the mid-rapidity region

. According to the Abramovski-Gribov-Kancheli cutting rules (AGK) [12], at mid-rapidity only Mueller-Kancheli type diagrams contribute to the inclusive spectrum of hadrons. It was shown in [9, 13] that at $x = 0$ the function $F_{\tau}^{(n)}(x_{\pm} \simeq 0, p_t, x_{1,2})$ does not depend on n . Therefore, the function $\phi_n(0, p_t)$ becomes proportional to n . In accordance with [13], the inclusive spectrum corresponding to the contribution of quarks at $x = 0$ has the following simple form:

$$\begin{aligned}
\rho_q(0, p_t) &= \tilde{\phi}_q(0, p_t) \sum_{n=1}^{\infty} n \sigma_n(s), \\
& g(s/s_0)^{\Delta} \tilde{\phi}_q(0, p_t),
\end{aligned} \tag{6}$$

where $\tilde{\phi}_q(0, p_t)$ is the so-called vertex function in the Mueller-Kancheli type diagram which is determined by the density distribution of the particles produced in the central region; $\Delta = 0.12$, $g=21$ mb and σ_{nd} is the nondiffractive cross section due to the exchange of any number cut-pomerons.

According to Eqs.(2-3), the sea quarks contribute to ϕ_n and the inclusive spectrum at $n \geq 2$. Assuming possible creation of soft gluons in the proton, which are split into $q\bar{q}$ pairs and should vanish at the zero intrinsic transverse momentum

($k_t \sim 0$), one can obtain the following energy dependence for the gluon contribution $\rho_g(0, p_t)$ to the inclusive spectrum [13]:

$$\rho_g(x=0, p_t) = \sum_{n=2}^{\infty} \sigma_n(s) \phi_n^g(0, p_t) \equiv (g(s/s_0)^\Delta - \sigma_{nd}) \tilde{\phi}_g(0, p_t) , \quad (7)$$

Finally, we can present the inclusive spectrum at $x \simeq 0$ in the following form:

$$\rho(p_t) = \rho_q(x=0, p_t) + \rho_g(x=0, p_t) . \quad (8)$$

The functions $\tilde{\phi}_q(0, p_t)$ and $\tilde{\phi}_g(0, p_t)$ are fixed from the fit of the experimental data on the inclusive spectra of the charged hadrons produced in the central pp collisions at different initial energies. So, we found the following parametrization for $\tilde{\phi}_q(0, p_t)$ and $\tilde{\phi}_g(0, p_t)$ [13]:

$$\begin{aligned} \tilde{\phi}_q(0, p_t) &= A_q \exp(-b_q p_t) \\ \tilde{\phi}_g(0, p_t) &= A_g \sqrt{p_t} \exp(-b_g p_t), \end{aligned} \quad (9)$$

charged particles at $y = 0$: where $A_q = 4.78 \pm 0.16$ (GeV/c) $^{-2}$, $b_q = 7.24 \pm 0.11$ (GeV/c) $^{-1}$ and $A_g = 1.42 \pm 0.05$ (GeV/c) $^{-2}$; $b_g = 3.46 \pm 0.02$ (GeV/c) $^{-1}$.

2..3 Gluon distribution in proton within GBW model

The conventional QGSM or DPM model does not include the distribution of gluons in the proton. However, as is well known, at large transverse momenta p_t of hadrons the gluons in the proton play very important role in description of the experimental data. Therefore, one can assume that the contribution of the gluon distribution in the proton to the inclusive spectrum of the produced hadrons slowly appears when p_t increases and it will be sizable at high values of p_t . This assumption is also confirmed by the increase of the UGDF in the proton at $x \sim 0$ as a function of the internal transverse momentum k_t when k_t grows [2]-[5].

According to Refs.[2, 5], the UGDF, as a function of k_t at some value of Q_0 and low x is presented in the following form:

$$xg(x, k_t, Q_0) = C_0 R_0^2(x) k_t^2 \exp\left(-R_0^2(x) k_t^2\right) , \quad (10)$$

where $C_0 = 3\sigma_0 / (4\pi^2\alpha_s(Q_0))$, $R_0(x) = \text{GeV}^{-1}(x/x_0)^{\lambda/2}$, x_0, λ and σ_0 are defined in [2, 5]; $\alpha_s(Q_0)$ is the QCD coupling constant. To get the UGDF at low k_t we assume the possible creation of the soft gluons in the proton which appears at nonzero k_t . We calculated the gluon contribution $\tilde{\phi}_g(0, p_t)$ entering into Eq.(7) as the cut graph of the one-pomeron exchange in the gluon-gluon interaction (Fig.2, right) using the splitting of the gluons into the $q\bar{q}$ pair. Then the calculation was made in a way similar to the calculation of the sea quark contribution to the inclusive spectrum




Figure 2: The one-pomeron exchange graph between two gluons in the elastic pp scattering (left) and the corresponding cut graph (right).

within the QGSM, see Eq.(3) at $n = 2$. Calculating the diagram of Fig.2 (right) we assumed the following form for the $xg(x, k_t, Q_0)$:

$$xg(x, k_t, Q_0) = C_0 C_1 (1 - x)^{b_g} \times \left(R_0^2(x) k_t^2 + C_2 (R_0(x) k_t)^a \right) \exp(-R_0(x) k_t) , \quad (11)$$

where C_2, a, b_g and λ are the parameters which were found from the best description of data. As it will be shown in the next section, the parametrization given by Eq.(11) results in the satisfactory description of the inclusive spectra of hadrons produced in pp collisions at the mid-rapidity region and low transverse momenta $p_t \leq 1.5$ GeV/c. At higher values of p_t the gluon contribution $\tilde{\phi}_g(0, p_t)$ given by Eq.(9) is decreasing very fast and, in fact, does not contribute to the hadron spectrum at large p_t . Therefore, one can find the parametrization for the UGDF that is very close to the GBW parametrization [2, 5] at large k_t and coincides with the UGDF given by Eq.(11) at low k_t . It is the following:

$$xg(x, k_t, Q_0) = C_0 C_3 (1 - x)^{b_g} \times \left(R_0^2(x) k_t^2 + C_2 (R_0(x) k_t)^a \right) \exp \left(-R_0(x) k_t - d(R_0(x) k_t)^3 \right) , \quad (12)$$

The coefficients C_1, C_3 are found from the normalization

$$xg(x, Q_0^2) = \int_0^{Q_0^2} dk_t^2 xg(x, k_t, Q^2) \quad (13)$$

3. Results and discussion

The results of our calculations of the charged hadron inclusive spectrum $(1/N_{ev})d^3N/dydp_t^2$, where N_{ev} is the total number of the inelastic events, in the mid-rapidity at $\sqrt{s}=7$ TeV are presented in Fig.(3). The solid curve corresponds to the quark contribution $\rho_q(x=0, p_t)$ given by Eq.(6), the long-dash line is the gluon

contribution $\rho_g(x=0, p_t)$ (7) to the inclusive spectrum; the dotted curve corresponds to the sum of both contributions, see Eqs.(8). One can see that the conventional quark contribution $\rho_q(y=0, p_t)$ is able to describe the data up to $p_t \leq 1$ GeV/c, whereas the inclusion of the gluon contribution allows us to extend the range of good description up to 2 GeV/c. At larger values of p_t the contribution of hard processes is not negligible. Therefore, we calculated the inclusive spectrum at $p_t \geq 2$ GeV/c within the leading order perturbative QCD (LO PQCD) [15]-[22], see the details in [13]. The short-dash curve in Fig.3 (top) corresponds to our LO PQCD. From the best fit of the inclusive p_t -spectra of the charged hadrons produced in pp collisions at $x \simeq 0$ presented in Fig.3 (bottom) by the short dash line we found the following parameter values entering into the form for $xg(x, k_t, Q_0)$ given by Eq.(11):

$$a = 0.7; C_2 \simeq 2.3; \lambda = 0.22; b_g = 12; d = 0.2; C_3 = 0.3295$$

and

$$C_1 = \frac{2\pi}{\Gamma(4)} (1 + C_2 \Gamma(a+2)/\Gamma(4))^{-1} .$$

In Fig.4 the UGDF $xg(x, k_t, Q_0)/C_0$ as a function of k_t at $x = x_0$ and $Q_0 = 1$.GeV/c is presented. The solid line (GBW UGDF) is the GBW parametrization [2, 5] given by Eq.(10) divided by C_0 ; the dashed curve (UGDF1) corresponds to the UGDF given by Eq.(11) and the dotted line (UGDF2) corresponds to the UGDF given by Eq.(12) which is very close to the GBW UGDF at $k_t \geq 1.5$ GeV/c. It is seen from Fig.4 that the parametrization UGDF2 is different from the GBW UGDF mainly at $k_t \leq 1.4$ GeV/c.

4. Conclusion

We assume that the contribution of the gluon distribution in the proton to the inclusive spectrum of the produced hadrons slowly appears when p_t increases and it will be sizable at high values of p_t . This assumption is also confirmed by the increase of the unintegrated gluon distribution in the proton at $x \sim 0$ as a function of the internal transverse momentum k_t when k_t grows [3, 4, 2, 5].

Therefore, to illustrate this hypothesis we fit the experimental data on the inclusive spectra of charged particles produced in the central pp collisions at energies larger than the ISR starting by the sum of the quark contribution ρ_q given by Eq.(6) and the gluon contribution ρ_g (see Eq.(7)). The parameters of this fit do not depend on the initial energy in that energy interval. From the best fit of the LHC data on the inclusive spectra of the charged hadrons produced in the mid-rapidity pp collisions at low p_t we found a new parametrization of the unintegrated gluon distribution in the proton at small values of the intrinsic momentum k_t , which is similar to the UGDF obtained in [2, 5] at large k_t . The inclusion of the hard pp collision within LO QCD allows us to describe rather satisfactorily the data at higher values of p_t .




Figure 3: The inclusive spectrum of charged hadron as a function of p_t (GeV/c) in the central rapidity region ($y = 0$) at $\sqrt{s} = 7$ TeV and wide region of p_t (top) and the same spectrum at $p_t \leq 1.6$ GeV/c (bottom) compared with the CMS [14] which are very close to the ATLAS data..




Figure 4: The unintegrated gluon distribution $xg(x, k_t, Q_0)/C_0$ as a function of k_t at $x = x_0$ and $Q_0 = 1. \text{GeV/c}$. The solid line is the GBW UGDF [2, 5], the dashed curve (UGDF1) corresponds to Eq.(11) and the dotted line (UGDF2) corresponds to Eq.(12) which is very close to the GBW UGDF at $k_t \geq 1.5 \text{ GeV/c}$.

Acknowledgements

The authors are very grateful to A.Bakulev, V.Cavazini, A.Dorokhov, A.V.Efremov, F.Francavilla, C. Gwenlan, H.Jung, V.Kim, B.Kniehl, N.Kochelev, E.A.Kuraev, A.V.Lipatov, L.N.Lipatov, T.Lomtadze, M.Mangano, C.Merino, S.V.Mikhailov, E.Nurse, F.Palla, E.Pilkington, C.Royon, M.G.Ryskin, E.Sarkisyan-Grinbaum, O.V.Teryaev, Yu.Shabelski and N.P.Zotov for very useful discussions and comments. This work was supported in part by the Russian Foundation for Basic Research, project No:11-02-01538-a.

References

- [1] G. Watt, A.D. Martin and M.G. Ryskin, Eur.Phys.J., C31 (2003) 73 [arXiv:hep-ph/0306169]; *ibid* Phys.Rev. D70 (2004) 014012 [Erratum-*ibid*. D70 (2004) 079902] [arXiv:hep-ph/0309096].
- [2] K.Golec-Biernat, M.Wusthof, Phys.Rev D60 (1999) 114023.

- [3] A. D. Martin, M.G. Ryskin and G. Watt, Eur.Phys.J., C66 (2010) 163 [arXiv:hep-ph/0909.5529].
- [4] N.I. Kochelev, Phys.Lett., B426 (1998) 149;
- [5] H. Jung, Proc. of the DIS'2004, Strbaske' Pleso, Slovakia, arXiv:0411287 [hep-ph].
- [6] V.N. Gribov and L.N. Lipatov, Sov.J.Nucl.Phys. 15 (1972) 438; G. Altarelli and G. Parisi, Nucl.Phys. B 126 (1997) 298; Yu.L. Dokshitzer, Sov.Phys. JETP, 46 (1977) 641.
- [7] A.B.Kaidalov, Z.Phys., C12 (1982) 63; Sarveys High Energy Phys., 13 (1999) 265. A.B.Kaidalov, O.I.Piskunova, Z.Phys., C30 (1986) 145; Yad.Fiz., 43 (1986) 1545.
- [8] A.Capella, U.Sukhatme, C.J.Tan, J,Tran Thanh Van, Phys.Rev. D**36**, 109 (1987); *ibid* Adv.Ser.Direct.High Energy Phys.2:428-480,1988.
- [9] G.I.Lykasov, M.N.Sergeenko, Z.Phys.C56 (1992) 697; *ibid* Z.Phys., C52 (1991) 635; *ibid* Z.Phys.C70 (1996) 455.
- [10] V.A. Bednyakov, G.I. Lykasov and V.V. Lyubushkin, Europhys.Lett. 92 (2010) 31001; arXiv:hep-ph/1005.0559.
- [11] K.A.Ter-Martirosyan, Sov.J.Nucl.Phys. 44 (1986) 817.
- [12] V.Abramovsky, V.N.Gribov, O.Koncheli, Sov.J.Nucl.Phys., 18 (1973) 308 .
- [13] V.A. Bednyakov, A.A. Grinyuk, G.I. Lykasov, M. Poghosyan, arXiv:11040532 [hep-ph].
- [14] CMS Colalaboration, Vadrn Kachatryan, et al., Phys.Rev.Lett., 105 (2010) 022002; arXiv:1005.3299 [hep-ex]; CMS-QCD, CERN-PH-EP-2010-003, Feb.2010.
- [15] A.V.Efremov, Sov.J.Nucl.Phys., 19 (1974) 176.
- [16] P.Nasson, S.Dawson & R.K.Ellis, Nucl.Phys., B303 (1988) 607; B327 49 (1989); B335 (1989) 260(E).
- [17] R.D.Field,R.P.Feyman, Phys.Rev., D15 (1977) 2590.
- [18] R.P.Feyman, R.D.Field, and G.C.Fox, Nucl.Phys., B128 (1977) 1.
- [19] R.P.Feyman, R.D.Field, and G.C.Fox, Phys.Rev., D18 (1977) 3320.
- [20] M.L.Mangano, Physics-Uspekhi, 53 (2010) 109.

- [21] A.Sherstner, R.Torne, Eur.Phys.J C55 (2008) 553.
- [22] S.Albino, B.A.Kniehl, G.Kramer (AKK08), Nucl.Phys. B803 (2008) 42.