

# Hiding the Higgs Boson from Prying Eyes

Ernest Ma

*Department of Physics and Astronomy, University of California,  
Riverside, California 92521, USA*

## Abstract

There are two simple ways that the Higgs boson  $H$  of the Standard Model (SM) may be more difficult to observe than expected at the Large Hadron Collider (LHC) or the Tevatron. One is well-known, i.e.  $H$  decays invisibly, into dark-matter scalar particles for example. The other is that  $H$  mixes with a heavy singlet scalar  $S$  which couples to new colored fermions and scalars. Of the two mass eigenstates, the light one could (accidentally) have a suppressed effective coupling to two gluons, and the heavy one could be kinematically beyond the reach of the LHC.

The one Higgs boson [1]  $H$  of the Standard Model (SM) of particle interactions is expected to be produced by gluon-gluon fusion at the Large Hadron Collider (LHC) and be observed through its decay into  $ZZ$ ,  $WW$ , and other channels. Recent reported data [2, 3, 4] have excluded the following mass ranges at 95% confidence level:

$$\text{ATLAS : } \quad 146 - 232, \quad 256 - 282, \quad 296 - 466 \text{ GeV}, \quad (1)$$

$$\text{CMS : } \quad 145 - 216, \quad 226 - 288, \quad 310 - 340 \text{ GeV}, \quad (2)$$

$$\text{TEVATRON : } \quad 156 - 177 \text{ GeV}. \quad (3)$$

Combined with the LEP II bound [5] of  $m_H > 114.4 \text{ GeV}$ , this leaves only a small window for its observation. Whereas more data could eventually find  $H$  through its rare decay mode to two photons, it is perhaps a good time now to consider how  $H$  may be hidden from view because of either its decay or its production.

A first possibility is that  $H$  decays significantly into invisible channels, thereby diminishing its branching fractions into observable final states. This is a very old idea [6] and has many different model realizations. One recent example is the model of a dark (inert) scalar doublet [7], where the Standard Model is extended to include a second scalar doublet, which is odd under an exactly conserved  $Z_2$  symmetry [8]. If the neutral member  $\eta^0 = (\eta_R + i\eta_I)/\sqrt{2}$  of this doublet is split so that  $m_R < m_I$  by at least the order of 100 keV, then  $\eta_R$  is a good dark-matter candidate [9, 10]. For the latest discussion on this model, see Ref. [11]. If  $2m_R < m_H$ , then the invisible decay of  $H$  into these dark-matter scalars will suppress its branching fractions to other particles, as already discussed in detail a few years ago [12]. The effect is especially significant below the  $WW$  threshold and could suppress the  $\gamma\gamma$  branching fraction by as much as a factor of three.

A second possibility is a new proposal. The idea is very simple. Suppose there is a scalar singlet  $S$  which couples to new colored fermions and scalars. In that case, both  $H$  and  $S$  will couple to two gluons through loops. Let the  $Hgg$  amplitude be  $A_H$  and the  $Sgg$  amplitude be

$A_S$ , then  $A_H$  is dominated by the  $t$ -quark loop, and  $A_S$  comes from the new colored fermions and scalars. Take for example  $A_S = 3A_H$ . Now if  $H$  mixes with  $S$ , the linear combination  $H' = (3H - S)/\sqrt{10}$  would not couple to two gluons, and would not be produced at the LHC by gluon-gluon fusion. If  $H'$  also happens to be a mass eigenstate, then it could hide from being seen at the LHC even if its mass is 170 GeV (above the  $WW$  threshold). The orthogonal combination  $S' = (3S + H)/\sqrt{10}$  has an enhanced coupling to two gluons, but it is presumably heavy because it is mostly a singlet, and could be kinematically beyond the reach of the LHC.

Consider the scalar potential of the SM doublet  $\Phi = (\phi^+, \phi^0)$  and a real singlet  $S$ :

$$V = \mu_1^2 \Phi^\dagger \Phi + \frac{1}{2} \lambda_1 (\Phi^\dagger \Phi)^2 + \frac{1}{2} \mu_2^2 S^2 + \frac{1}{3} \mu_3 S^3 + \frac{1}{4} \lambda_2 S^4 + \mu_4 S \Phi^\dagger \Phi + \frac{1}{2} \lambda_3 S^2 \Phi^\dagger \Phi. \quad (4)$$

Let  $\langle \phi^0 \rangle = v$  and  $\langle S \rangle = u$ , then the minimum of  $V$  is determined by

$$0 = v(2\mu_1^2 + 2\lambda_1 v^2 + \lambda_3 u^2 + 2\mu_4 u), \quad (5)$$

$$0 = u(\mu_2^2 + \lambda_2 u^2 + \lambda_3 v^2 + \mu_3 u) + \mu_4 v^2. \quad (6)$$

The  $2 \times 2$  mass-squared matrix spanning the physical scalars  $H$  and  $S$  is given by

$$\mathcal{M}^2 = \begin{pmatrix} 2\lambda_1 v^2 & \sqrt{2}(\lambda_3 u + \mu_4)v \\ \sqrt{2}(\lambda_3 u + \mu_4)v & 2\lambda_2 u^2 + \mu_3 u - \mu_4 v^2/u \end{pmatrix}. \quad (7)$$

Let the mass eigenstates of the above be  $H' = H \cos \theta - S \sin \theta$  and  $S' = S \cos \theta + H \sin \theta$ , with eigenvalues  $m_1^2$  and  $m_2^2$ , then Eq. (7) may be rewritten as

$$\mathcal{M}^2 = \begin{pmatrix} m_1^2 \cos^2 \theta + m_2^2 \sin^2 \theta & (m_2^2 - m_1^2) \sin \theta \cos \theta \\ (m_2^2 - m_1^2) \sin \theta \cos \theta & m_1^2 \sin^2 \theta + m_2^2 \cos^2 \theta \end{pmatrix}. \quad (8)$$

As an example, let  $\sin \theta = 1/\sqrt{10}$ ,  $\cos \theta = 3/\sqrt{10}$ ,  $u = 2\sqrt{2}v = 492.4$  GeV, where  $v = 174.1$  GeV, we then obtain  $m_1 = 170$  GeV and  $m_2 = 500$  GeV for the choice  $\lambda_1 = 0.84$ ,  $\lambda_2 = 0.47$ ,  $\lambda_3 = 0.55$ , and  $\mu_3 = \mu_4 = 0$ . This demonstrates the numerical viability of this proposal.

It has been assumed that  $S$  couples to new colored fermions and scalars. This is of course model-dependent, but a necessary condition is to have  $A_S = A(S \rightarrow gg)$  a few times larger

than  $A_H = A(H \rightarrow gg)$ . Now  $A_H$  is dominated by the  $t$  quark which is a fundamental triplet under  $SU(3)_C$  and is proportional to  $(\sqrt{2}v)^{-1}$ . Suppose  $A_S$  comes from a colored fermion octet  $Q$  with the coupling  $S\bar{Q}Q$ , then it is proportional to  $u^{-1}$  but its color factor of 3 is 6 times that of the  $t$  quark. Hence for the above choice of  $u = 2\sqrt{2}v$ ,  $A_S \simeq 3A_H$  is realized. The allowed mass term  $\bar{Q}Q$  would change the details of the above, but may be forbidden by a  $Z_2$  symmetry under which  $S$  and  $Q_L$  are odd, but  $Q_R$  is even.

More realistically,  $H'$  is unlikely to decouple from  $gg$  entirely. In that case, the suppression (or enhancement if  $\sin \theta < 0$ ) factor in  $H'$  production at the LHC is  $(\cos \theta - (A_S/A_H) \sin \theta)^2$ . On the other hand, depending on the choice of new colored fermions and scalars, there is also a contribution from  $A(S \rightarrow \gamma\gamma)$  to  $H'$  decay. This means that the branching fraction of  $H'$  to  $\gamma\gamma$  would also not be the same as in the SM. If a particle is discovered at the LHC in the  $\gamma\gamma$  channel below 145 GeV, but with a branching fraction different from what is expected from the SM, especially if it is *greater*, it may be due to this effect. The presence of the octet  $Q$  may also be relevant in gauge-coupling unification [13] without supersymmetry.

In conclusion, the existence of the Higgs boson may be hidden from view at present because of a variety of scenarios, some of which have been discussed recently [14, 15, 16, 17]. In this paper, two simple ways are considered: the presence of light dark-matter scalars which affects the decay or an accidental cancellation between  $A(H \rightarrow gg)$  and  $A(S \rightarrow gg)$  in  $H - S$  mixing which affects the production. In the latter case, an increase from the present  $E_{cm} = 7$  TeV to 14 TeV at the LHC in the future would produce  $S'$  easily, and the decay  $S' \rightarrow H'H'$  would be a spectacular signature for discovering  $H'$ .

I thank Bohdan Grzadkowski and Maria Krawczyk for a stimulating “Scalars 2011” Conference in Warsaw (August 2011) which led to this work. My research is supported in part by the U. S. Department of Energy under Grant No. DE-AC02-06CH11357.

## References

- [1] For a review, see for example A. Djouadi, Phys. Rep. **457**, 1 (2008).
- [2] A. Nisati, Talk at the XXV International Symposium on Lepton Photon Interactions at High Energies (Mumbai, India), August 2011.
- [3] V. Sharma, Talk at the XXV International Symposium on Lepton Photon Interactions at High Energies (Mumbai, India), August 2011.
- [4] M. Verzocchi, Talk at the XXV International Symposium on Lepton Photon Interactions at High Energies (Mumbai, India), August 2011.
- [5] Particle Data Group: K. Nakamura *et al.*, J. Phys. G: Nucl. Part. Phys. **37**, 075021 (2010).
- [6] R. E. Shrock and M. Suzuki, Phys. Lett. **B110**, 250 (1982).
- [7] E. Ma, Phys. Rev. **D73**, 077301 (2006).
- [8] N. G. Deshpande and E. Ma, Phys. Rev. **D18**, 2574 (1978).
- [9] R. Barbieri, L. J. Hall, and V. S. Rychkov, Phys. Rev. **D74**, 015007 (2006).
- [10] L. Lopez Honorez, E. Nezri, J. F. Oliver, and M. H. G. Tytgat, JCAP **0702**, 028 (2007).
- [11] L. Lopez Honorez and C. E. Yaguna, JCAP **1101**, 002 (2011).
- [12] Q.-H. Cao, E. Ma, and G. Rajasekaran, Phys. Rev. **D76**, 095011 (2007).
- [13] E. Ma, Phys. Lett. **B625**, 76 (2005).
- [14] F. Bonnet, M. B. Gavela, T. Ota, and W. Winter, arXiv:1105.5140 [hep-ph].
- [15] K. Ghosh, B. Mukhopadhyaya, and U. Sarkar, Phys. Rev. **D84**, 015017 (2011) [arXiv:1105.5837 [hep-ph]].
- [16] X.-G. He and G. Valencia, arXiv:1108.0222 [hep-ph].
- [17] X.-G. He and J. Tandean, arXiv:1109.1277 [hep-ph].