Sigma term and strangeness content of octet baryons

S. Dürr[a,b], Z. Fodor[a,b,c], T. Hemmert[d], C. Hoelbling[a], J. Frison[e], S.D. Katz[a,c], S. Krieg[a,b], T. Kurth[a], L. Lellouch[e], T. Lippert[a,b], A. Portelli[e], A. Ramos[e], A. Schäfer[d], K.K. Szabó[a]

[a]Bergische Universität Wuppertal, Gaussstr. 20, D-42119 Wuppertal, Germany
[b]Jülich Supercomputing Centre, Forschungszentrum Jülich, D-52425 Jülich, Germany
[c]Institute for Theoretical Physics, Eötvös University, H-1117 Budapest, Hungary
[d]Universität Regensburg, Universitätsstr. 31, D-93053 Regensburg, Germany
[e]Centre de Physique Théorique[1], Case 907, CNRS Luminy, F-13288 Marseille, France

arXiv:1109.4265v3 [hep-lat] 27 Feb 2016

Abstract

By using lattice QCD computations we determine the sigma terms and strangeness content of all octet baryons by means of an application of the Hellmann-Feynman theorem. In addition to polynomial and rational expressions for the quark-mass dependence of octet members, we use $SU(3)$ covariant baryon chiral perturbation theory to perform the extrapolation to the physical up and down quark masses. Our $N_f = 2 + 1$ lattice ensembles include pion masses down to about 190 MeV in large volumes ($M_\pi L \gtrsim 4$), and three values of the lattice spacing. Our main results are the nucleon sigma term $\sigma_{\pi N} = 39(4)(^{+18}_{-7})$ and the strangeness content $y_N = 0.20(7)(^{+13}_{-17})$. Under the assumption of validity of covariant baryon χPT in our range of masses one finds $y_N = 0.276(77)(^{+90}_{-62})$.

1 Introduction

The nucleon sigma term, and the nucleon strangeness content are phenomenologically important quantities that are not directly accessible to experiment. They are related to $\pi - N$ and $K - N$ scattering lengths, to the quark-mass ratio m_s/m_{ud} (where $m_{ud} = (m_u + m_d)/2$), to the hadron spectrum and even to counting rates in Higgs boson searches. They also play a key role in the direct detection of dark matter (DM): sigma terms relate the effective DM - nucleon coupling to the fundamental Lagrangian parameters that couple the DM particle to the quarks. A precise first principle determination of these quantities is thus very interesting. The sigma terms are defined as

$$\sigma_{\pi N} = \langle N(p)|m_{ud}(\bar{u}u + \bar{d}d)(0)|N(p)\rangle \tag{1a}$$
$$\sigma_{\bar{s}sN} = \langle N(p)|m_s\bar{s}s(0)|N(p)\rangle \tag{1b}$$

The usual way of computing $\sigma_{\pi N}$ is by using $\pi - N$ scattering data. $\sigma_{\pi N}$ cannot be directly obtained in this way, but available data can be extrapolated to the Cheng-Dashen point, although the machinery involved in this determination is complicated. χPT can then be used to reliably extrapolate from the Cheng-Dashen point to the chiral limit [1, 2].

Such an analysis was carried out in the 1980's [1, 2], resulting in a value of $\sigma_{\pi N} = 45 \pm 8$ MeV. A later study claims to obtain a higher value [3], $\sigma_{\pi N} = 64 \pm 7$ MeV which is almost 2 standard deviations away. Also it has been pointed that the uncertainties associated with these determinations are already affecting the interpretation of (direct) dark matter search experiments [4], although lattice simulations are already helping to change this situation [5].

Sigma-terms of other octet members are also of phenomenological interest. Recently they have been used in the context of the hadron resonance gas model for estimating quark-mass

[1]CPT is research unit UMR 6207 of the CNRS and of Aix-Marseille univ. and Univ. Sud Toulon-Var.

effects in QCD thermodynamics calculations [6]. Moreover, as we will see, an analysis of all octet members can be further used to constrain the strangeness content of the nucleon.

In this paper we compute the sigma terms of the octet baryons $\sigma_{\pi X}, \sigma_{\bar{s}sX}$ for $X = N, \Lambda, \Sigma, \Xi$. The Feynman-Hellman theorem applied to QFT relates the sigma terms to the dependence of baryon masses with respect to quark masses through:

$$\sigma_{\pi X} = m_{ud}\frac{\partial M_X}{\partial m_{ud}} \tag{2a}$$

$$\sigma_{\bar{s}sX} = m_s\frac{\partial M_X}{\partial m_s} \tag{2b}$$

This opens the possibility of computing the octet sigma terms via the variation of octet masses with respect to quark masses in a lattice simulation, which is the approach that we use here. A preliminary account of this work has been given in [7]. There are other interesting quantities directly related to the previously mentioned sigma terms that are also useful for phenomenology

$$y_X = \frac{2\langle X(p)|\bar{s}s(0)|X(p)\rangle}{\langle X(p)|(\bar{u}u+\bar{d}d)(0)|X(p)\rangle} \tag{3a}$$

$$f_{udX} = \frac{m_{ud}\langle X(p)|(\bar{u}u+\bar{d}d)(0)|X(p)\rangle}{M_X} \tag{3b}$$

$$f_{\bar{s}sX} = \frac{m_s\langle X(p)|\bar{s}s(0)|X(p)\rangle}{M_X}. \tag{3c}$$

The first is called the strangeness content, and the other two are sometimes referred to as "*dimensionless sigma terms*". The latter two are directly related to the conversion of the fundamental DM-quark coupling to the effective DM-nucleon coupling.

2 Simulation details and ensembles

The gauge and fermionic actions, as well as the algorithms used are described in [8, 9]. Here it suffices to mention that we simulate QCD with two degenerate light quarks and a heavier strange quark, and that we use tree level improved Wilson fermions.

To set the lattice spacing corresponding to each value of the coupling β and fix the quark masses to their physical values, we use M_π, M_K, and M_Ω. We extrapolate the values of aM_π, aM_K, aM_Ω to the point where the ratios aM_π/aM_Ω and aM_K/aM_Ω agree with the experimental values[2].

As shown in Table 1 at $\beta = 3.3$ and $\beta = 3.7$ the strange quark mass is held fixed, whereas for $\beta = 3.57$ we simulate at three different values of m_s to have some lever arm to perform the small extrapolation to $m_s^{\rm phys}$ (see also Fig. 1). Our data sets cover a wide range of pion masses from $M_\pi \sim 190$ MeV up to $M_\pi \sim 680$ MeV, although in this analysis we only use ensembles with $M_\pi < 550$ MeV.

On every ensemble we measure the octet masses M_N, M_Λ, M_Σ and M_Ξ with valence quark masses equal to sea quark masses (only the unitary theory is considered). It is worth mentioning that these ensembles have previously been used to accurately predict the light hadron spectrum [9], including the masses of the octet baryons.

[2]Experimental inputs are corrected for isospin breaking and electromagnetic effects according to [9].

β	am_{ud}	am_s	$L^3\times T$	Trajectories	aM_π	aM_K
3.3	-0.0960	-0.057	$16^3\times 32$	10000	0.4115(6)	0.4749(6)
	-0.1100	-0.057	$16^3\times 32$	1450	0.322(1)	0.422(1)
	-0.1200	-0.057	$16^3\times 64$	4500	0.2448(9)	0.3826(6)
	-0.1233	-0.057	$24^3\times 64$	2000	0.2105(8)	0.3668(6)
	-0.1233	-0.057	$32^3\times 64$	1300	0.211(1)	0.3663(8)
	-0.1265	-0.057	$24^3\times 64$	2100	0.169(1)	0.3500(7)
3.57	-0.0318	0,-0.010	$24^3\times 64$	1650,1650	0.2214(7),0.2178(5)	0.2883(7),0.2657(5)
	-0.0380	0,-0.010	$24^3\times 64$	1350,1550	0.1837(7),0.1778(7)	0.2720(6),0.2469(6)
	-0.0440	0,-0.007	$32^3\times 64$	1000,1000	0.1348(7),0.1320(7)	0.2531(6),0.2362(7)
	-0.0483	0,-0.007	$48^3\times 64$	500,1000	0.0865(8),0.0811(5)	0.2401(8),0.2210(5)
3.7	-0.007	0.0	$32^3\times 96$	1100	0.2130(4)	0.2275(4)
	-0.013	0.0	$32^3\times 96$	1450	0.1830(4)	0.2123(3)
	-0.020	0.0	$32^3\times 96$	2050	0.1399(3)	0.1920(3)
	-0.022	0.0	$32^3\times 96$	1350	0.1273(5)	0.1882(4)
	-0.025	0.0	$40^3\times 96$	1450	0.1021(4)	0.1788(4)

Table 1: Parameters of our simulations. The errors quoted here are purely statistical. These results correspond to one of the 18 two-point function, time fit intervals that we use in our estimate of systematic uncertainties. In this particular analysis, the scales at $\beta = 3.3, 3.57, 3.7$ are $a^{-1} = 1616(20)$ MeV, $2425(27)$ MeV, $3142(37)$ MeV, respectively.

3 Chiral extrapolation

As already stated, we simulate QCD for values of the light quark masses larger than the physical values, whereas the strange quark is close to its physical value with some lever arm to perform a small extrapolation. Thus our data requires an extrapolation in the light quark masses and an interpolation in m_s.

For this, we need to describe the quark-mass dependence of octet baryons. We will always use the tree level $SU(3)$ chiral relation to express the quark-mass dependence through the meson mass dependence via

$$\begin{aligned} m_{ud} &\propto M_\pi^2 \\ m_s &\propto M_{\bar{s}s}^2 = 2M_K^2 - M_\pi^2 \,. \end{aligned} \tag{4}$$

To study this mass dependence and the associated model uncertainty, we consider two very different approaches [9, 10]. First we consider a regular expansion in quark masses around a non singular point where none of the quark masses vanish. For this purpose we define the expansion variables

$$\begin{aligned} \Delta_\pi &= \frac{M_\pi^2 - (M_\pi^{\rm cen})^2}{M_\Omega^2} \\ \Delta_{\bar{s}s} &= \frac{M_{\bar{s}s}^2 - \left(M_{\bar{s}s}^{\rm phys}\right)^2}{M_\Omega^2}, \end{aligned} \tag{5}$$

where $(M_\pi^{\rm cen})^2 = \frac{1}{2}\left[\left(M_\pi^{\rm phys}\right)^2 + \left(M_\pi^{\rm cut}\right)^2\right]$, and $M_\pi^{\rm cut}$ denotes the mass of the heaviest pion included in the fit.




Figure 1: Overview of our simulation points in terms of M_π and $\sqrt{2M_K^2 - M_\pi^2}$. The former gives a measure of the isospin averaged up and down quark mass while the latter determines the strange quark mass. The symbols refer to the three lattice spacings, and the physical point is marked with a cross. Error bars are statistical only.

Note that changing the expansion point in polynomial-like formulae is simply a reshuffling of the coefficients. When interested in checking the $SU(3)$ symmetric line $m_{ud} = m_s$ it is far more convenient to define our expansion variables as

$$\begin{aligned} \Delta_\pi &= \left(\frac{M_\pi}{M_\Omega}\right)^2 \\ \Delta_{\bar{s}s} &= \left(\frac{M_{\bar{s}s}}{M_\Omega}\right)^2, \end{aligned} \tag{6}$$

where the $SU(3)$ symmetric line is characterized by $\Delta_\pi = \Delta_{\bar{s}s}$. Because the extrapolation in m_s is small, a linear term in $\Delta_{\bar{s}s}$ suffices

Second we consider a singular expansion around the $SU(3)$ chiral point $m_{ud} = m_s = 0$, which allows for a much more constrained expansion. Low-energy processes in QCD have been extensively studied in the framework of chiral perturbation theory. The seminal work [11] and its success in explaining meson observables, have pushed the study of baryons within the same framework. However baryon χPT is far more involved. The main difficulty is that baryon masses M_B are not small in comparison with the scale of chiral symmetry breaking ($\Lambda_\chi \sim 1$ GeV), and Weinberg's power counting theorem [12] fails: higher-order loop corrections contribute with powers of M_B/Λ_χ and are no longer small. This results in a very slow (practical) convergence of the series [13]. This convergence is partially improved by treating the baryons as heavy degrees of freedom, in what is known as the heavy baryon chiral perturbation theory [14]. Although it is reasonable for dealing with the mass dependence of baryons with light quarks, experience seems to show that observables that depend on the strange quark are not well described in this framework (see for example [15–17]). A number of possibilities have been proposed to improve the convergence and include the strange quark in the analysis. The various flavors of cutoff BχPT (see [18] and references therein), or covariant BχPT [19] are some examples. In this work we explore this last possibility. The interested reader can consult the review [20] and references therein.

3.1 Regular expansions

We have several possibilities if we decide to expand around a regular point. We can treat any octet member $X = N, \Lambda, \Sigma, \Xi$ as independent and use a Taylor expansion. Using the variables defined in Eq. (5) our results are well described by the ansatz

$$M_X = M_0^X + \alpha_1^X \Delta_\pi + \alpha_2^X \Delta_\pi^2 + \beta^X \Delta_{\bar{s}s}. \tag{7}$$

One can also use a Padé like functional form

$$M_X = \frac{M_0^X}{1 - \alpha_1^X \Delta_\pi - \alpha_2^X \Delta_\pi^2 - \beta^X \Delta_{\bar{s}s}}. \tag{8}$$

In these expansions, $M_0^X, \alpha_{1,2}^X$ and β^X are the 16 fitting parameters.

In principle all octet masses should be degenerate along the $SU(3)$ symmetric line $m_{ud} = m_s$. We can choose to impose this constraint in our functional form. In fact our data is well fitted by the following $SU(3)$ symmetric regular expansion

$$M_X = M_0 + \alpha_1^X \Delta_\pi + \alpha_2^X \Delta_\pi^2 + \left(C_1 - \alpha_1^X\right) \Delta_{\bar{s}s} + \left(C_2 - \alpha_2^X\right) \Delta_\pi \Delta_{\bar{s}s}, \tag{9}$$

or the corresponding Padé like ansatz

$$M_X = \frac{M_0}{1 - \alpha_1^X \Delta_\pi - \alpha_2^X \Delta_\pi^2 - \left(C_1 - \alpha_1^X\right) \Delta_{\bar{s}s} - \left(C_2 - \alpha_2^X\right) \Delta_\pi \Delta_{\bar{s}s}}. \tag{10}$$

Now the fitting parameters have been reduced to 11: $M_0, \alpha_{1,2}^X$ and the coefficients C_1 and C_2 that give the dependence of octet masses along the $SU(3)$ symmetric line $m_{ud} = m_s$.

Note that Taylor-like and Padé-like ansatze differ in higher-order terms in Δ_π and $\Delta_{\bar{s}s}$. Thus the difference in physical results obtained by using these two functional forms measures higher-order contributions to the mass expansion.

3.2 Covariant BχPT

Details of the quark-mass dependence of the octet members in $SU(3)$ baryon χPT can be found in Appendix A. Here we run the index $X = N, \Lambda, \Sigma, \Xi$ over octet members, and index $\alpha = \pi, K, \eta$ over the mesons.

Octet masses to NLO involve the nonlinear function

$$h(x) = -\frac{x^3}{4\pi^2} \left\{ \sqrt{1 - \left(\frac{x}{2}\right)^2} \arccos \frac{x}{2} + \frac{x}{2} \log x \right\}. \tag{11}$$

The masses of the octet baryons are given by

$$M_X = M_0 - 4c_{\pi X} M_\pi^2 - 4c_{\bar{s}sX} M_{\bar{s}s}^2 + \sum_{\alpha=\pi,K,\eta} \frac{g_{\alpha X}}{F_\alpha^2} M_0^3 h\left(\frac{M_\alpha}{M_0}\right) + 4d_\pi M_\pi^4 + 4d_{\bar{s}s} M_{\bar{s}s}^4, \tag{12}$$

where $c_{\pi X}, c_{\bar{s}sX}$ are functions of the three low-energy constants (LEC), b_0, b_d, b_f, and the $g_{\alpha X}$ can be written in terms of the axial coupling g_A and the ratio of couplings ξ (as shown in Tabs. 2 and 3). Finally $d_\pi, d_{\bar{s}s}$ parametrize higher-order corrections.

α	$g_{\alpha N}$	$g_{\alpha\Lambda}$	$g_{\alpha\Sigma}$	$g_{\alpha\Xi}$
π	$\frac{3}{4}g_A^2$	$\frac{1}{(1+\xi)^2}g_A^2$	$\frac{(1+6\xi^2)}{3(1+\xi)^2}g_A^2$	$\frac{3(1-\xi)^2}{4(1+\xi)^2}g_A^2$
K	$\frac{(5-6\xi+9\xi^2)}{6(1+\xi)^2}g_A^2$	$\frac{(1+9\xi^2)}{3(1+\xi)^2}g_A^2$	$\frac{(1+\xi^2)}{(1+\xi)^2}g_A^2$	$\frac{(5+6\xi+9\xi^2)}{6(1+\xi)^2}g_A^2$
η	$\frac{(1-3\xi)^2}{12(1+\xi)^2}g_A^2$	$\frac{1}{3(1+\xi)^2}g_A^2$	$\frac{1}{3(1+\xi)^2}g_A^2$	$\frac{(1+3\xi)^2}{12(1+\xi)^2}g_A^2$

Table 2: Meson-loop couplings as a function of the πN coupling g_A and the quantity ξ. In the chiral limit $g_A = D + F$ and $\xi = F/D$, as described in Appendix A.

X	N	Λ	Σ	Ξ
c_X^{π}	$-2(2b_0 + b_D + b_F)$	$-4(b_0 + b_D/3)$	$-4(b_0 + b_D)$	$-2(2b_0 + b_D - b_F)$
$c_X^{\bar{s}s}$	$-2(b_0 + b_D - b_F)$	$-2(b_0 + 4b_D/3)$	$-2b_0$	$-2(b_0 + b_D + b_F)$

Table 3: Leading order octet quark-mass dependence, $c_{\pi X}$ and $c_{\bar{s}sX}$, as a function of the LEC b_0, b_D, b_F (see Appendix A for more details).

To the order at which we are working only two pseudo-Goldstone boson masses are linearly independent, being related through the Gell-Mann-Okubo relation $3M_\eta^2 = 4M_K^2 - M_\pi^2$.

When fitting lattice data to the chiral formula, it is important not to break the $SU(3)$ symmetry built into this expression. In that sense it is important that all octet masses become degenerate when $M_{\bar{s}s} = M_\pi$. This rules out the possibility of fixing the meson decay constants $F_{\pi,K,\eta}$ to their physical values. Here we choose to fix all of them to the common value $F_\pi^{\rm phys} = 92.2$ MeV. This is correct to the order at which we are working. Note that we have repeated the fits including the NLO terms for the decay constants and obtained results compatible within statistical errors. We have also confirmed that the quality of fits decreases if a non $SU(3)$ symmetric ansatz is imposed by fixing each meson decay constant to its physical value.

The axial coupling g_A is well known from phenomenology. The most precise value at the physical point is $g_A = 1.2695(29)$ [21], and is expected to be close to the value in the chiral limit $D + F$. The ratio of couplings ξ is not well determined experimentally, but there are two preferred phenomenological scenarios: $\xi = 2/3$ and $\xi \sim 0.5$ [22].


It makes sense to try our χPT fits both fixing $(g_A, \xi) = (1.2695, 2/3)$ or allowing them to be free. Regarding the higher-order contributions given by $d_\pi, d_{\bar{s}s}$, we can also choose to either include them in the fit, or not. In terms of how well our data are described (i.e. fit quality) the four possibilities are equally reasonable options. In Fig. 2, we show one possible χPT fit, where we have fitted g_A, ξ and we have not included the higher-order terms in the fit.

4 Cutoff, finite volume effects and excited-state contributions


We combine our chiral extrapolation with the continuum extrapolation. It has been demonstrated in different contexts [8, 9, 23, 24] that a smeared clover action is very close to be nonperturbatively $\mathcal{O}(a)$-improved with cutoff effects being $\mathcal{O}(a^2)$. Nevertheless we cannot rule out the possibility that our results have linear discretization effects.

Guided by our experience [9] in determining light hadron masses we parametrize cutoff effects with the substitution

$$M_X \longrightarrow M_X(1 + c_X a^p) \tag{13}$$



(a) Fit in the M_π^2, M_X^2 plane. Data have been corrected to the physical value of $M_{\bar{s}s}$ for a better visualisation



(b) Fit in the $M_{\bar{s}s}^2, M_X^2$ plane. Data have been corrected to the physical value of M_π for a better visualisation

Figure 2: Example of a χPT fit corresponding to a particular choice of fitting interval for the correlators. In this particular case we choose to fit g_A and ξ, and not to include higher-order terms. We have only used data with $M_\pi < 410$ MeV. The correlated $\chi^2 = 38.5$ for 34 degrees of freedom, yields a fit quality of ~ 0.27.

with $p = 1, 2$. The cutoff effects are small enough in our data that they make it difficult to distinguish a and a^2. In fact their coefficients are, for the most part, compatible with zero within statistical errors. Thus we will also include fits without any cutoff corrections altogether.

Stable hadron masses have leading finite-volume effects $\propto e^{-M_\pi L}$ [25]. In all of our ensembles the bound $M_\pi L \gtrsim 4$ is maintained, implying negligible finite-volume effects. The detailed analysis of [9], with additional ensembles in smaller volumes, shows that finite-volume corrections are below the statistical accuracy of our data. Thus we do not add finite-volume corrections to our analysis here.

To estimate the possible contamination by excited states in the extraction of the masses from the correlators we use 18 fitting time intervals: $t_{\rm min}/a = 5$ or 6 for $\beta = 3.3$, $t_{\rm min}/a = 7, 8$ or 9 for $\beta = 3.57$ and $t_{\rm min}/a = 10, 11$ or 12 for $\beta = 3.7$ [9].


5 Determination of the uncertainties

Our complete analysis includes a total of 8 formulae to extrapolate to the physical mass point: Four of them are regular expansions either imposing or not the $SU(3)$ flavor constraint, and either in the Taylor-like form or Padé-like form. The other four functional forms are derived from $SU(3)$ covariant baryon χPT, where we can choose to parametrize or not higher-order contributions, and to fit g_A and ξ or fix g_A to its physical value $g_A = 1.2695(29)$ [21] and ξ to a reasonable phenomenological value[3] $\xi = 2/3$ [22].


To have more control over the uncertainties associated with higher-order terms in the mass extrapolation, we impose two different pion mass cuts: $M_\pi < 410$ MeV and $M_\pi < 550$ MeV.

As explained in Sec. 4 we consider three possibilities to parametrize cutoff effects which are compatible with our data: either assume that they are absent, or parametrize them as $\mathcal{O}(a)$ or $\mathcal{O}(a^2)$. Finite-volume corrections are not considered because they are below our statistical

[3] We have checked that using $\xi = 0.5$ (the other common phenomenological scenario) leads to very similar results.



(a) Final distribution of values for $\sigma_{\pi N}$ and the contributions of different pion mass cuts.



(b) Final distribution of values for $\sigma_{\pi N}$ and the contributions of different functional forms.

Figure 3: Distribution of values for $\sigma_{\pi N}$ in background (yellow), and how different pion mass cuts (left) and functional forms (right) shape it.

accuracy. Finally we repeat the full analysis with 18 different time-fitting intervals for the correlators to estimate excited state contributions.

This strategy leads to $8 \times 2 \times 3 \times 18 = 864$ different procedures to estimate the physical value of each of the sigma terms of the octet members. The data are fitted (taking correlations into account) using all of the previously explained procedures. Each of the 864 results is weighted with the fit quality [4] (p-value or Q-value) to produce a distribution of values. The median (typical result of our analysis) is taken as our final value. The $16^{\underline{\text{th}}}/84^{\underline{\text{th}}}$ percentiles (i.e. the values which, in a Gaussian distribution, correspond to $\pm 1\sigma$ deviations from the median) yield the systematic uncertainty of our computation (see Fig. 3). This systematic uncertainty measures how different parametrizations for mass extrapolation, cutoff effects and excited-state contributions affect the final result. The statistical uncertainty is determined by bootstraping the whole procedure 2000 times, and computing the variance of the medians.

It is important to note here three points:

- Most of the previously mentioned fits are "good." Our average fit quality is 0.34, and this does not vary significantly over our different procedures. For example fits with $M_\pi < 410$ MeV have an average fit quality of 0.38 whereas for $M_\pi < 550$ MeV it is 0.29 (see Fig. 3(a)). For covariant BχPT, Taylor and Padé functional forms the p-values are respectively $0.30, 0.47$ and 0.29 (see Fig. 3(b)). The fit quality is almost insensitive to how we choose to parametrise the cutoff effects, or what time intervals we use to fit correlators. Although there is some variation, we cannot rule out any of our fits based on how well our data are described.

- Adding more variations does not increase the systematic uncertainty. For example adding a pion mass cut of $M_\pi < 680$ MeV (i.e. including all our ensembles in the fit), the covariant BχPT functional form gives fit qualities of 2×10^{-6}. Even if they are included in our analysis, these fits do not contribute to our distribution of values or to our final results and estimates of systematic uncertainties.

- On the other hand, *eliminating* some of the analysis typically results in a compatible final value but with a smaller systematic uncertainty. For example had we chosen to

[4] The confidence of fit is defined as $Q = \int_{\chi^2}^{\infty} dz\ P(z,d)$ where $P(z,d) = z^{d/2-1}e^{-z/2}/[2^{d/2}\Gamma(d/2)]$ is the probability distribution function to obtain $\chi^2 = z$ in a fit with d degrees of freedom.

	$\sigma_{\pi X}$ [MeV]	$\sigma_{\bar{s}sX}$ [MeV]	y_X	f_{udX}	$f_{\bar{s}sX}$
N	$39(4)(^{+18}_{-7})$	$67(27)(^{+55}_{-47})$	$0.20(7)(^{+13}_{-17})$	$0.042(5)(^{+21}_{-8})$	$0.072(29)(^{+60}_{-50})$
Λ	$29(3)(^{+11}_{-5})$	$180(26)(^{+48}_{-77})$	$0.51(15)(^{+48}_{-27})$	$0.027(3)(^{+10}_{-5})$	$0.165(25)(^{+47}_{-63})$
Σ	$23(3)(^{+19}_{-3})$	$245(29)(^{+50}_{-72})$	$0.82(21)(^{+87}_{-39})$	$0.019(3)(^{+17}_{-3})$	$0.210(25)(^{+46}_{-62})$
Ξ	$15(2)(^{+8}_{-3})$	$312(32)(^{+72}_{-77})$	$1.7(5)(^{+1.9}_{-0.7})$	$0.012(2)(^{+6}_{-2})$	$0.240(26)(^{+58}_{-61})$

Table 4: Final results for all octet members. The first two columns give the baryon octet sigma terms (Eq. (1)), the third one the strangeness content of the octet member (Eq. (3a)), and the last two the dimensionless sigma terms (Eqs. (3b), (3c)). The first error is statistical, the second one systematic.

perform our analysis only with $M_\pi < 550$ MeV the systematic uncertainty of $\sigma_{\pi N}$ would have dropped by almost 50%. (see Fig. 3(a)).

6 Results

Following the method described in the previous section, we arrive at the results quoted in Table 4.

For the nucleon sigma term our result agrees with the "canonical" determination coming from $\pi - N$ scattering data [2] of $\sigma_{\pi N} = 45 \pm 8$ MeV, but with a larger uncertainty, and disfavors the larger value $\sigma_{\pi N} = 64 \pm 7$ MeV from [3]. It also agrees with recent estimates that use lattice data [15, 26–31, 39] though these assume a particular functional form for the light quark-mass dependence while we consider a whole range of forms. Another study considers a variety of chiral forms, but with pion masses only down to 300 MeV [32], yielding values in the range 42 to 84 MeV, with a statistical accuracy of about 15 MeV. Our value is less consistent with the determinations of [33, 34] that are close to the higher value (of [3]) for the sigma term.

We believe that the agreement within error of our results obtained from extrapolations using very different functional forms enhances the credibility of our final results.

In Table 5 we give the contribution of each source of systematic uncertainty to the final error. Although we show the results for $\sigma_{\pi N}$, the conclusion that the chiral extrapolation dominates the systematic uncertainty of our computation is generally true.

Source of systematic error	error on $\sigma_{\pi N}$ [MeV]
Chiral Extrapolation:	
- Pion mass range	9.0
- Functional form	5.5
Continuum extrapolation	1.9

Table 5: Contribution to the total uncertainty of the different sources of systematic error.

To reduce this source of systematic uncertainty we would need to add ensembles at lower pion masses. From the value of the uncertainties of $\sigma_{\pi N}$ at $M_\pi = 200$ MeV, we can roughly guess the gain that would be obtained by having equally precise data at the physical point. Such data would imply a reduction of around 50% in the systematic error. On the other hand the situation for both $\sigma_{\bar{s}sN}$ and y_N is less clear. Our data set has a much smaller range of strange quark masses than light quark masses. If we add the fact that the contribution of the strange quark to the nucleon mass seems to be small "per se", it is natural that the final

results for both $\sigma_{\bar{s}sN}$ and y_N to show large statistical and systematic uncertainties. Only by adding simulation points for a wider range of strange quark masses one can increase the chances of getting a precise determination of these quantities without making an additional assumption which relates m_s to m_{ud} dependence as in $SU(3)$ baryon χPT.

Regarding the other octet members, the values of the sigma terms agrees with other determinations based on lattice data and a low-energy effective field theory approach [30, 31].

6.1 χPT consistency check

Our objective in this paper is to determine the sigma-terms and strangeness contents of the octet baryons at the physical point. To this end, we generated ensembles close to the physical value of the strange quark mass and as close as possible to the physical light quark mass, with a large enough lever arm in the light sector to perform a credible extrapolation to the physical point.

Such a set of ensembles, however, is not ideally suited to determine the $SU(3)$ χPT LECs, which are defined in the chiral limit. That would require simulations with a strange quark mass approaching the chiral limit. This means that the LECs extracted from our fits have uncertainties which we cannot properly control. Nevertheless, it is worth noting that the values obtained are roughly consistent with general expectations from phenomenology [21] and other lattice studies [30, 31] (see Table 6). This further enhances the credibility of the results obtained at the physical point.

M_0 [GeV]	b_0 [GeV^{-1}]	b_D [GeV^{-1}]	b_F [GeV^{-1}]	g_A	ξ
0.75(15)	−0.71(24)	0.103(60)	−0.359(72)	0.92(13)	0.402(90)

Table 6: Values of the LEC's obtained from our χPT fits. Total errors (statistics and systematics included) are computed in the same way as for the sigma terms. For the reasons noted in the text, we do not consider these errors to be reliable.

7 Conclusions

We computed the sigma terms of light octet baryons by studying their mass dependence on quark masses through the Feynman-Hellman theorem. We also obtained the strangeness content and dimensionless sigma terms for all octet members. We estimated the systematic uncertainties associated with the choice of parametrization for this mass dependence by considering two very different expansion schemes. We considered regular expansions of octet masses, and a singular three-flavor, low-energy effective field theory approach. Within each of the approaches we estimated the impact of higher order contributions. Moreover we varied the range of fitted quark masses, cutoff parametrizations and excited-state contributions.

The most interesting quantities are the ones associated with the nucleon. Our final value for $\sigma_{\pi N} = 39(4)(^{+18}_{-7})$ favors the "low scenario" with $\sigma_{\pi N} = 45 \pm 8$ MeV of [1, 2], but the size of our uncertainties does not allow us to exclude the higher value of [3]. Regarding the strangeness content we obtain a value with large statistical and systematic uncertainties $y_N = 0.20(7)(^{+13}_{-17})$. The dimensionless sigma terms $f_{udN} = 0.042(5)(^{+21}_{-8})$ and $f_{\bar{s}sN} = 0.072(29)(^{+60}_{-50})$ are the quantities of interest for direct DM searches.

This work shows the difficulties associated with obtaining precise values for these form factors in a model independent way. Even with pion masses as low as 190 MeV the mass extrapolation

dominates the systematic uncertainties of our final results. This explicitly shows the risks of using lattice data and only one approach to perform the mass extrapolation. A proper estimation of systematic uncertainties should include the uncertainty associated with the model that is used, and, at least in our case, this is one of the main sources of systematic uncertainty.

For $\sigma_{\pi N}$ the only way to increase the precision of our computation without assuming a model would be to generate data at lower quark masses. Replacing the extrapolation by an interpolation drastically reduces the uncertainty in the final result, as we have explicitly seen by considering the values of the sigma terms at $M_\pi = 200$ MeV.

For the case of the strangeness content of the nucleon and $\sigma_{\bar{s}sN}$, the situation is even more delicate. First, $N_f = 2 + 1$ lattice ensembles usually bracket the physical value of m_s with a small lever arm. This is more than sufficient to determine physical observables, but it is not the ideal setup to estimate how physical quantities change with m_s. Second, the contribution of the strange quark to the nucleon mass is small "per se". In principle the first problem can be solved by measuring the nucleon mass for a larger range of strange quark masses. The second problem is solved if we restrict our analysis to three flavor BχPT. This low-energy effective field theory approach relates the m_{ud} dependence of octet members to its m_s dependence, and thus constrains y_N. The value of the nucleon strangeness content obtained from the covariant BχPT analysis alone has a small relative error $y_N = 0.276(77)(^{+90}_{-62})$, illustrating the success of the idea. However, since we do not have a large enough range of strange quark masses to establish the validity of $SU(3)$ BχPT, this result must be considered model dependent. Note also that a direct calculation of the corresponding disconnected contribution suggests a very small y_N value, barely compatible with the BχPT result [38].

Acknowledgments

Computations were performed using HPC resources from FZ Jülich and from GENCI-[IDRIS/CCRT] (Grant No. 52275) and clusters at Wuppertal and CPT. This work is supported in part by EU Grants No. I3HP, FP7/2007-2013/ERC No. 208740, No. MRTN-CT-2006-035482 (FLAVIAnet), DFG Grant No. FO 502/2, SFB-TR 55, by CNRS Grant GDR No. 2921 and PICS No. 4707.

The authors also thank Martin Jung for pointing out an error in the normalization of the dimensionless sigma terms.

Appendix

A Sketch of three-flavor covariant baryon χPT

In this appendix, we give a brief outline of the derivation of our chiral fit form (12). Working to $\mathcal{O}(p^3)$, we use the standard octet baryon Lagrangians [35, 36]

$$\mathcal{L}^{(1)}_{MB} = \mathrm{Tr}\Big(\mathrm{i}\overline{B}\gamma^\mu D_\mu B - M_0\overline{B}B + \frac{D}{2}\overline{B}\gamma^\mu\gamma_5\{u_\mu, B\} + \frac{F}{2}\overline{B}\gamma^\mu\gamma_5[u_\mu, B]\Big) \tag{14}$$

$$\mathcal{L}^{(2)}_{MB} = \mathrm{Tr}\Big(b_D\overline{B}\{\chi_+, B\} + b_F\overline{B}[\chi_+, B] + \dots\Big) + b_0\mathrm{Tr}\Big(\overline{B}B\Big)\mathrm{Tr}\Big(\chi_+\Big) + \dots \tag{15}$$

$g_{\alpha B}$	N	Λ	Σ	Ξ
π	$\frac{3}{4}(D+F)^2$	D^2	$\frac{1}{3}(D^2+6F^2)$	$\frac{3}{4}(D-F)^2$
K	$\frac{1}{6}(5D^2-6DF+9F^2)$	$\frac{1}{3}(D^2+9F^2)$	(D^2+F^2)	$\frac{1}{6}(5D^2+6DF+9F^2)$
η	$\frac{1}{12}(D-3F)^2$	$\frac{1}{3}D^2$	$\frac{1}{3}D^2$	$\frac{1}{12}(D+3F)^2$

Table 7: Summary of meson-loop couplings $g_{\alpha B}$ in terms of the low-energy constants D, F.

with the chiral tensors

$$
\begin{aligned}
D_\mu B &= \partial_\mu B + [\Gamma_\mu, B] && (16)\\
\Gamma_\mu &= \frac{1}{2}(u^\dagger(\partial_\mu + \ldots)u + u(\partial_\mu + \ldots)u^\dagger) && (17)\\
u_\mu &= \mathrm{i}(u^\dagger(\partial_\mu + \ldots)u - u(\partial_\mu + \ldots)u^\dagger) && (18)\\
\chi_\pm &= u^\dagger\chi u^\dagger \pm u\chi^\dagger u && (19)\\
\chi &= 2B_0(M + \ldots), && (20)
\end{aligned}
$$

where M denotes the (diagonal) quark-mass matrix. To the order we are working, there is no contribution from the $\mathcal{O}(p^3)$ Lagrangian, $\mathcal{L}^{(3)}_{MB}$, to the quark-mass dependence of the octet baryon masses. The coupling constants D, F of Eq. (14) are related to the actual couplings of the nucleon in the chiral limit through $D + F = \lim_{u,d,s\to 0} g_A$ and $F/D = \lim_{u,d,s\to 0}\xi$, whereas M_0 corresponds to the chiral limit value of the mass of the baryon octet. The field $u(x) = \sqrt{U(x)}$ describes a nonlinear matrix representation of the (quasi-) Goldstone boson fields and B is the matrix-valued interpolating field for a spin 1/2 octet baryon. The next-to-leading order low-energy constants b_D, b_F, b_0 govern the leading quark-mass contributions to the mass of a spin 1/2 octet baryon, with $\mathcal{M}$ denoting the quark-mass matrix for three light flavors u, d and s. Furthermore, we note that the parameter B_0 is a measure of the size of the chiral condensate $\langle 0|\bar{q}q|0\rangle$ (in the chiral limit) [37]

$$
B_0 \equiv -\frac{\langle 0|\bar{q}q|0\rangle_{m_q\to 0}}{F_0^2} \tag{21}
$$

where the low-energy constant F_0 is identified with the value of the octet Goldstone boson decay constant in the chiral limit. Finally, for the calculation at hand one needs to know the chiral meson Lagrangian up to $\mathcal{O}(p^2)$ [37]

$$
\mathcal{L}^{(2)}_M = \frac{F_0^2}{4}\mathrm{Tr}\{(\partial_\mu + \ldots)U^\dagger(\partial^\mu + \ldots)U + \chi^\dagger U + \chi U^\dagger\}. \tag{22}
$$

Generalizing the SU(2) calculation of [27] along the lines of [31], we obtain the leading one-loop result for the mass of a nucleon to $\mathcal{O}(p^3)$

$$
\begin{aligned}
M_N = {} & M_0 - 4(b_0 + (b_D + b_F)/2)\bar{M}_\pi^2 - 2(b_0 + b_D - b_F)\bar{M}_{\bar{s}s}^2 \\
& - \sum_{\alpha=\pi,K,\eta} \frac{g_{\alpha N}\,\bar{M}_\alpha^3}{4\pi^2 F_\alpha^2}\Big\{\sqrt{1-\frac{\bar{M}_\alpha^2}{4M_0^2}}\arccos\frac{\bar{M}_\alpha}{2M_0} + \frac{\bar{M}_\alpha}{4M_0}\log\frac{\bar{M}_\alpha^2}{M_0^2}\Big\} + \mathcal{O}(p^4).
\end{aligned} \tag{23}
$$

To the chiral order we are working here, the meson-loop couplings $g_{\pi N}, g_{KN}, g_{\eta N}$ can be expressed in terms of the two SU(3) parameters F and D, as given in Table 7. For generality, we have expressed the Goldstone boson decay constant F_0 entering the loop contributions via three different symbols F_α to account for the individual contribution of the pion, kaon and eta-cloud of the nucleon.

For a fit to lattice QCD data, it is convenient to slightly rewrite (23) without changing the expression at the order at which we are working:

1. We express the GMOR-masses for $\bar{M}_K, \bar{M}_\eta$ in (23) as a function of the GMOR-mass of the pion and of the mass-parameter $M_{\bar{s}s}$ introduced in (4):

$$\bar{M}_K^2 \rightarrow \frac{1}{2}(\bar{M}_\pi^2 + M_{\bar{s}s}^2) \tag{24}$$

$$\bar{M}_\eta^2 \rightarrow \frac{1}{3}(\bar{M}_\pi^2 + 2M_{\bar{s}s}^2) \tag{25}$$

   which amounts to a change of variables.

2. We identify the GMOR-mass $\bar{M}_\pi$ of (4) with the corresponding lowest-lying 0^--boson mass M_π in each simulation. Possible deviations in the quark-mass dependence from the linear GMOR-behavior predicted in (4) can only affect $\mathcal{O}(p^4)$ corrections (23).

3. The Goldstone boson decay constants F_α in (23) are identified with the decay constant $F_\pi^{\rm phys}$. Other assignments compatible with SU(3) symmetry might be chosen as discussed in Sec. 3.2.

4. With the same reasoning as in the previous item, at chiral order $\mathcal{O}(p^3)$ we can identify the chiral limit couplings $\tilde{g}_{\pi N}, \tilde{g}_{KN}, \tilde{g}_{\eta N}$ with the physical couplings $g_{\pi N}, g_{KN}, g_{\eta N}$, resulting in Table 2. The unknown ratio ξ of couplings is either treated as an external input to our fits or is left as a free fit parameter.

With a very similar derivation for the case of the octet hyperons, we finally arrive at our fit function (12).

References

[1] R. Koch, *A New Determination of the pi N Sigma Term Using Hyperbolic Dispersion Relations in the (nu**2, t) Plane*, *Z.Phys.* **C15** (1982) 161–168.

[2] J. Gasser, H. Leutwyler, and M. Sainio, *Sigma term update*, *Phys.Lett.* **B253** (1991) 252–259.

[3] M. Pavan, I. Strakovsky, R. Workman, and R. Arndt, *The Pion nucleon Sigma term is definitely large: Results from a G.W.U. analysis of pi nucleon scattering data*, *PiN Newslett.* **16** (2002) 110–115, [hep-ph/0111066].

[4] J. R. Ellis, K. A. Olive, and C. Savage, *Hadronic Uncertainties in the Elastic Scattering of Supersymmetric Dark Matter*, *Phys. Rev.* **D77** (2008) 065026, [arXiv:0801.3656].

[5] J. Giedt, A. W. Thomas, and R. D. Young, *Dark matter, the CMSSM and lattice QCD*, *Phys.Rev.Lett.* **103** (2009) 201802, [arXiv:0907.4177].

[6] **Wuppertal-Budapest Collaboration** Collaboration, S. Borsanyi *et. al.*, *Is there still any T_c mystery in lattice QCD? Results with physical masses in the continuum limit III*, *JHEP* **1009** (2010) 073, [arXiv:1005.3508].

[7] S. Durr, Z. Fodor, J. Frison, T. Hemmert, C. Hoelbling, *et. al.*, *Sigma term and strangeness content of the nucleon*, *PoS* **LATTICE2010** (2010) 102, [arXiv:1012.1208].

[8] S. Durr *et. al.*, *Scaling study of dynamical smeared-link clover fermions*, *Phys. Rev.* **D79** (2009) 014501, [arXiv:0802.2706].

[9] S. Durr *et. al.*, *Ab Initio Determination of Light Hadron Masses*, *Science* **322** (2008) 1224–1227, [arXiv:0906.3599].

[10] L. Lellouch, *Kaon physics: a lattice perspective*, arXiv:0902.4545.

[11] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory to One Loop*, *Annals Phys.* **158** (1984) 142.

[12] S. Weinberg, *Phenomenological Lagrangians*, *Physica* **A96** (1979) 327. Festschrift honoring Julian Schwinger on his 60th birthday.

[13] J. Gasser, M. Sainio, and A. Svarc, *Nucleons with Chiral Loops*, *Nucl.Phys.* **B307** (1988) 779.

[14] E. E. Jenkins and A. V. Manohar, *Baryon chiral perturbation theory using a heavy fermion Lagrangian*, *Phys.Lett.* **B255** (1991) 558–562.

[15] B. Borasoy and U.-G. Meissner, *Chiral expansion of baryon masses and sigma terms*, *Annals Phys.* **254** (1997) 192–232, [hep-ph/9607432].

[16] J. F. Donoghue, B. R. Holstein, and B. Borasoy, *SU(3) baryon chiral perturbation theory and long distance regularization*, *Phys.Rev.* **D59** (1999) 036002, [hep-ph/9804281].

[17] B. Borasoy, *Sigma terms in heavy baryon chiral perturbation theory revisited*, *Eur.Phys.J.* **C8** (1999) 121–130, [hep-ph/9807453].

[18] R. D. Young, D. B. Leinweber, and A. W. Thomas, *Convergence of chiral effective field theory*, *Prog.Part.Nucl.Phys.* **50** (2003) 399–417, [hep-lat/0212031].

[19] M. Dorati, T. A. Gail, and T. R. Hemmert, *Chiral perturbation theory and the first moments of the generalized parton distributions in a nucleon*, *Nucl.Phys.* **A798** (2008) 96–131, [nucl-th/0703073].

[20] V. Bernard, *Chiral Perturbation Theory and Baryon Properties*, *Prog.Part.Nucl.Phys.* **60** (2008) 82–160, [arXiv:0706.0312].

[21] **Particle Data Group** Collaboration, K. Nakamura *et. al.*, *Review of particle physics*, *J.Phys.G* **G37** (2010) 075021.

[22] T. Cheng and L.-F. Li, *Chiral quark model of nucleon spin flavor structure with SU(3) and axial U(1) breakings*, *Phys.Rev.* **D57** (1998) 344–349, [hep-ph/9701248].

[23] S. Capitani, S. Durr, and C. Hoelbling, *Rationale for UV-filtered clover fermions*, *JHEP* **0611** (2006) 028, [hep-lat/0607006].

[24] R. Hoffmann, A. Hasenfratz, and S. Schaefer, *Non-perturbative improvement of nHYP smeared Wilson fermions*, *PoS* **LAT2007** (2007) 104, [arXiv:0710.0471].

[25] M. Luscher, *Volume Dependence of the Energy Spectrum in Massive Quantum Field Theories. 1. Stable Particle States*, *Commun.Math.Phys.* **104** (1986) 177.

[26] D. B. Leinweber, A. W. Thomas, and R. D. Young, *Physical nucleon properties from lattice QCD*, *Phys.Rev.Lett.* **92** (2004) 242002, [hep-lat/0302020].

[27] M. Procura, T. R. Hemmert, and W. Weise, *Nucleon mass, sigma term and lattice QCD*, *Phys.Rev.* **D69** (2004) 034505, [hep-lat/0309020].

[28] M. Procura, B. Musch, T. Wollenweber, T. Hemmert, and W. Weise, *Nucleon mass: From lattice QCD to the chiral limit*, *Phys.Rev.* **D73** (2006) 114510, [hep-lat/0603001].

[29] H. Ohki, H. Fukaya, S. Hashimoto, T. Kaneko, H. Matsufuru, *et. al.*, *Nucleon sigma term and strange quark content from lattice QCD with exact chiral symmetry*, *Phys.Rev.* **D78** (2008) 054502, [arXiv:0806.4744].

[30] R. Young and A. Thomas, *Octet baryon masses and sigma terms from an SU(3) chiral extrapolation*, *Phys.Rev.* **D81** (2010) 014503, [arXiv:0901.3310].

[31] J. Martin Camalich, L. Geng, and M. Vicente Vacas, *The lowest-lying baryon masses in covariant SU(3)-flavor chiral perturbation theory*, *Phys.Rev.* **D82** (2010) 074504, [arXiv:1003.1929].

[32] A. Walker-Loud, H.-W. Lin, D. Richards, R. Edwards, M. Engelhardt, *et. al.*, *Light hadron spectroscopy using domain wall valence quarks on an Asqtad sea*, *Phys.Rev.* **D79** (2009) 054502, [arXiv:0806.4549].

[33] **European Twisted Mass Collaboration** Collaboration, C. Alexandrou *et. al.*, *Light baryon masses with dynamical twisted mass fermions*, *Phys.Rev.* **D78** (2008) 014509, [arXiv:0803.3190].

[34] **PACS-CS Collaboration** Collaboration, K.-I. Ishikawa *et. al.*, *SU(2) and SU(3) chiral perturbation theory analyses on baryon masses in 2+1 flavor lattice QCD*, *Phys.Rev.* **D80** (2009) 054502, [arXiv:0905.0962].

[35] J. A. Oller, M. Verbeni, and J. Prades, *Meson-baryon effective chiral lagrangians to O(q**3)*, *JHEP* **0609** (2006) 079, [hep-ph/0608204].

[36] M. Frink and U.-G. Meissner, *On the chiral effective meson-baryon Lagrangian at third order*, *Eur.Phys.J.* **A29** (2006) 255–260, [hep-ph/0609256].

[37] J. Gasser and H. Leutwyler, *Low-Energy Expansion of Meson Form-Factors*, *Nucl. Phys.* **B250** (1985) 517–538.

[38] A. Schäfer, talk at Lattice 2011.

[39] R. Horsley, talk at Lattice 2011.